

# PSEUDO-INDEX OF FANO MANIFOLDS AND SMOOTH BLOW-UPS

LAURENT BONAVERO

**Abstract.** Suppose  $\pi : X \rightarrow Y$  is a smooth blow-up along a submanifold  $Z$  of  $Y$  between complex Fano manifolds  $X$  and  $Y$  of pseudo-indices  $i_X$  and  $i_Y$  respectively (recall that  $i_X$  is defined by  $i_X := \min\{-K_X \cdot C \mid C \text{ is a rational curve of } X\}$ ). We prove that  $i_X \leq i_Y$  if  $2 \dim(Z) < \dim(Y) + i_Y - 1$  and show that this result is optimal by classifying the “boundary” cases. As expected, these results are obtained by studying rational curves on  $X$  and  $Y$ .

## 1. STATEMENT OF THE RESULTS

**1.1. Introduction.** When studying surjective morphisms  $f : X \rightarrow Y$  between smooth Fano manifolds  $X$  and  $Y$  of the same dimension, one generally observes that the anti-canonical bundle  $-K_Y$  of  $Y$  is “more positive” than the anti-canonical bundle  $-K_X$  of  $X$ , one of the most important results in this direction being the famous theorem of Lazarsfeld [La83] stating that if  $\mathbb{P}^n \rightarrow Y$  is a surjective morphism from  $\mathbb{P}^n$  to an  $n$ -dimensional manifold  $Y$ , then  $Y \simeq \mathbb{P}^n$ .

For a Fano manifold  $X$  (i.e., a complex manifold with ample anti-canonical line bundle  $-K_X$ ), one defines two integers called the index  $r_X$  and the pseudo-index  $i_X$  of  $X$  by

$$r_X := \max\{m \in \mathbb{N} \mid -K_X = mL \text{ with } L \in \text{Pic}(X)\}$$

and

$$i_X := \min\{-K_X \cdot C \mid C \text{ is a rational curve of } X\}.$$

Of course,  $i_X$  is a multiple of  $r_X$  and many results are known for these numbers. Among others, Fano manifolds of dimension  $n$  with large index (namely bigger than  $n - 2$ ) are classified (see [IP99] for a complete survey on Fano manifolds), the situation being much more complicated for the pseudo-index: one knows that  $i_X \leq n + 1$  by Mori theory, equality holding if and only if  $X \simeq \mathbb{P}^n$  [CMS00].

**1.2. The main result.** Let us start with an easy remark: let  $Y$  be a complex manifold of dimension  $n$ , let  $Z$  be a connected submanifold of  $Y$ , let  $X := B_Z(Y)$  be the blow-up of  $Y$  with center  $Z$  and let  $E$  be the exceptional divisor of  $\pi : X \rightarrow Y$ . We classically have

$$H^2(X, \mathbb{Z}) \simeq H^2(Y, \mathbb{Z}) \oplus \mathbb{Z} \cdot E \text{ and } K_X = \pi^* K_Y + (n - \dim(Z) - 1)E.$$

Therefore, if both  $Y$  and  $X$  are Fano,  $r_X$  is equal to the greatest common divisor of  $r_Y$  and  $n - \dim(Z) - 1$ , which implies in particular that  $r_X \leq r_Y$  and confirms the philosophy described above.

In this Note, we study the behaviour of the pseudo-index with respect to smooth blow-ups. Quite surprisingly, this behaviour depends on the dimension of the center of the blow-up. Our precise results are the following.

---

*Date:* September 2003.

**Key-words :** Fano manifolds, smooth blow-up, rational curves. **A.M.S. classification :** 14J45, 14E30, 14E05.

**Theorem 1.** *Let  $Y$  be a complex manifold of dimension  $n$ , let  $Z$  be a connected submanifold of  $Y$  and let  $X := B_Z(Y)$  be the blow-up of  $Y$  with center  $Z$ . Suppose both  $Y$  and  $X$  are Fano.*

- (i) *If  $2 \dim(Z) < n + i_Y - 1$ , then  $i_X \leq i_Y$ ,*
- (ii) *if  $2 \dim(Z) = n + i_Y - 1$  and  $i_Y \geq 2$ , then  $i_X \leq i_Y$ ,*
- (iii) *if  $\dim(Z) < n/2$ , then  $i_X \leq i_Y$ .*

Of course, (iii) is an obvious consequence of (i) since  $i_Y \geq 1$ . This result says that the pseudo-index has the “expected behaviour” when the center of the blow-up has small dimension. Remark that the case where  $\dim(Z) = 0$  could be proved by looking at the classification given in [BCW02] and the case where  $\dim(Z) = 1$  is Proposition 3.7 of [BCDD03].

Let us now give an example where  $i_X$  is bigger than  $i_Y$ . In the following proposition (as in the whole paper), we do not follow Grothendieck’s convention:  $\mathbb{P}(V)$  denotes the projective space of *lines* of the vector space  $V$ .

**Proposition 1.** *Let  $n := 2m$  be an even integer, let  $\mathcal{E}$  be the following rank  $m + 1$  vector bundle over  $\mathbb{P}^m$ :*

$$\mathcal{E} = \mathcal{O}_{\mathbb{P}^m}^{\oplus m} \oplus \mathcal{O}_{\mathbb{P}^m}(1)$$

*and let  $Y_n$  be the  $n$ -dimensional manifold  $Y_n = \mathbb{P}(\mathcal{E})$ . The trivial rank  $m$ -subbundle of  $\mathcal{E}$  defines a submanifold  $Z_m$  isomorphic to  $\mathbb{P}^{n/2}$  with normal bundle  $N_{Z_m/Y_n}$  isomorphic to  $\mathcal{O}_{\mathbb{P}^m}(-1)^{\oplus m}$ . Finally, let  $\pi_n : X_n = B_{Z_m}(Y_n) \rightarrow Y_n$  be the blow-up of  $Y_n$  along  $Z_m$ . Then  $Y_n$  and  $X_n$  are Fano manifolds of dimension  $n$  if  $n \geq 4$ . Moreover  $i_{Y_n} = 1$ ,  $i_{X_4} = 1$  and  $i_{X_n} = 2$  if  $n \geq 6$ .*

Therefore, the inequalities of Theorem 1 are optimal: for any  $n = 2m \geq 6$ ,  $\pi_n : X_n = B_{Z_m}(Y_n) \rightarrow Y_n$  is a blow-up with smooth connected center between Fano manifolds with  $\dim(Z_m) = \dim(X_n)/2$  and  $i_{X_n} > i_{Y_n}$ .

*Proof of Proposition 1.* Since  $X_n$  and  $Y_n$  are naturally toric manifolds, it is enough to compute the anti-canonical degree of invariant (rational) curves. If  $d$  is a line contained in  $Z_m$ , then  $-K_{Y_n} \cdot d = 1$ , which gives  $i_{Y_n} = 1$ . The Fano manifold  $X_n$  is isomorphic to the  $\mathbb{P}^1$ -bundle  $\mathbb{P}_{\mathbb{P}^{m-1} \times \mathbb{P}^m}(\mathcal{O}_{\mathbb{P}^{m-1} \times \mathbb{P}^m} \oplus \mathcal{O}_{\mathbb{P}^{m-1} \times \mathbb{P}^m}(1, 1))$  over  $\mathbb{P}^{m-1} \times \mathbb{P}^m$  hence  $i_{X_n} \leq 2$  (the  $\mathbb{P}^1$ -fibers having anti-canonical degree equal to 2). Let  $E \simeq \mathbb{P}^{m-1} \times \mathbb{P}^m$  be the exceptional divisor of  $\pi_n$ : the lines contained in a  $\mathbb{P}^{m-1} \times \{*\} \subset E$  have anti-canonical degree equal to  $m - 1$ , hence  $i_{X_4} = 1$  and  $i_{X_n} = 2$  if  $n \geq 6$ . ■

Remark: for  $n \geq 8$ , the previous computations show that the rational curves in  $X_n$  of minimal anti-canonical degree are not mapped by  $\pi_n$  to curves of minimal anti-canonical degree in  $Y_n$ .

Let us now discuss in more details the optimality of Theorem 1 by classifying the “boundary cases”.

**Theorem 2.** *Let  $\pi : X \rightarrow Y$  be a blow-up with smooth connected center  $Z$  between Fano manifolds  $X$  and  $Y$  of dimension  $n$ . If  $2 \dim(Z) = n + i_Y - 1$  then  $i_X \leq i_Y$  unless  $n \geq 6$  is even,  $X = X_n$ ,  $Y = Y_n$  and  $\pi = \pi_n$ .*

**1.3. Some consequences.** The results above have the following consequences when the pseudo-index of  $Y$  is large or in low dimensions.

**Corollary 1.** *Let  $\pi : X \rightarrow Y$  be a blow-up with smooth connected center between Fano manifolds  $X$  and  $Y$  of dimension  $n$ .*

- (i) If  $i_Y > n/3 - 1$ , then  $i_X \leq i_Y$ .
- (ii) If  $i_Y = n/3 - 1$ , then  $i_X \leq i_Y$  unless  $n = 6$ ,  $X = X_6$ ,  $Y = Y_6$  and  $\pi = \pi_6$ .

*Proof of Corollary 1.*

*Proof of (i).* Suppose by contradiction that  $i_X > i_Y$ . Then by Theorem 1(i),  $2 \dim(Z) \geq n + i_Y - 1$ . But the lines contained in the non-trivial fibers of the blow-up are rational curves of anti-canonical degree  $n - 1 - \dim(Z)$ , therefore  $n - 1 - \dim(Z) \geq i_X > i_Y$ , hence

$$n - 1 - i_Y \geq \dim(Z) + 1 \geq n/2 + i_Y/2 + 1/2$$

and  $i_Y \leq n/3 - 1$ , a contradiction.

*Proof of (ii).* Suppose that  $i_Y = n/3 - 1$  and that  $i_X > i_Y$ . The previous computations implies that every inequality occuring in the proof of (i) is an equality. In particular, one has  $2 \dim(Z) = n + i_Y - 1$ . Therefore, Theorem 2 implies that  $n$  is even and  $Y = Y_n$ . In particular,  $i_Y = 1 = n/3 - 1$ , hence  $n = 6$ , which ends the proof. ■

Remark that according to the generalised Mukai conjecture, as stated and studied in [BCDD03], Fano manifolds  $Y$  of dimension  $n \geq 6$  with  $i_Y > n/3 - 1$  should have Picard number  $\rho_Y$  satisfying  $\rho_Y < \frac{3n}{n-6}$ . Corollary 1 has the immediate following corollary.

**Corollary 2.** *Let  $\pi : X \rightarrow Y$  be a blow-up with smooth connected center between Fano manifolds  $X$  and  $Y$  of dimension  $n$ .*

- (i) If  $n \leq 5$ , then  $i_X \leq i_Y$ .
- (ii) If  $n = 6$ , then  $i_X \leq i_Y$  unless  $X = X_6$ ,  $Y = Y_6$  and  $\pi = \pi_6$ .

## 2. PROOFS

**2.1. Proof of Theorem 1.** It is enough to prove assertion (i), since (iii) is an obvious consequence of (i) and (ii) is an immediate consequence of Theorem 2 (note that the Fano manifolds  $Y_n$  have pseudo-index 1).

Let  $\pi : X = B_Z(Y) \rightarrow Y$  be a blow-up with smooth center  $Z$  between Fano manifolds  $X$  and  $Y$ . We will denote by  $E = \pi^{-1}(Z)$  the exceptional divisor of  $\pi$ . The basic idea is very simple: we take a rational curve  $C$  in  $Y$  such that  $-K_Y \cdot C = i_Y$  and we want to show that there is a rational curve  $\tilde{C}$  in  $X$ , mapping surjectively to  $C$  by  $\pi$ , such that  $-K_X \cdot \tilde{C} \leq i_Y$ .

Suppose first that there is a rational curve  $C$  in  $Y$  such that  $-K_Y \cdot C = i_Y$  and such that  $C$  is not contained in  $Z$ . The strict transform  $\tilde{C}$  of  $C$  is a rational curve satisfying  $E \cdot \tilde{C} \geq 0$  and the formula  $-K_X = \pi^*(-K_Y) - (n - 1 - \dim(Z))E$  immediately implies that  $-K_X \cdot \tilde{C} \leq i_Y$ .

Now take a rational curve  $C$  in  $Y$  such that  $-K_Y \cdot C = i_Y$  and assume  $C \subset Z$ . Let us decompose  $N_{Z/Y|C}$  (we allow here a slight abuse of notations, since  $C$  might be a singular rational curve, we should rather write  $\nu^*(N_{Z/Y|C})$  where  $\nu : \mathbb{P}^1 \rightarrow C$  is the normalisation of  $C$ ):

$$N_{Z/Y|C} = \bigoplus_{i=1}^r \mathcal{O}_{\mathbb{P}^1}(a_i)$$

where  $r = n - \dim(Z)$ . The sub-line bundles  $\mathcal{O}_{\mathbb{P}^1}(a_i)$  of  $N_{Z/Y|C}$  define rational curves  $\tilde{C}_i$  of  $E$  satisfying  $-K_X \cdot \tilde{C}_i = -K_Y \cdot C - (r - 1)a_i$ . Therefore, we are done if there exists  $i$  such that  $a_i \geq 0$ . Suppose the contrary, namely that  $a_i \leq -1$  for all  $i$ . Then

$$-K_Z \cdot C = -K_Y \cdot C - \deg(N_{Z/Y|C}) \geq i_Y + \text{rk}(N_{Z/Y|C}) = i_Y + n - \dim(Z).$$

Under the assumption  $2 \dim(Z) < n + i_Y - 1$ , we get  $-K_Z \cdot C > \dim(Z) + 1$  hence, by Mori's bend-and-break lemma (see for example [Deb01], p. 58), the curve  $C$  is numerically equivalent in  $Z$ , hence in  $Y$ , to a connected nonintegral effective rational 1-cycle (passing through 2 arbitrary fixed points of  $C$ ). Each reduced irreducible component of this 1-cycle has  $-K_Y$  anti-canonical degree strictly less than  $i_Y$ , contradiction! ■

Remarks: in the above proof, we only used that  $-K_Y \cdot C > 0$  for any rational curve of  $Y$ . Moreover, the previous proof also shows the following. Let  $Y$  be a complex manifold, let  $Z$  be a connected submanifold of  $Y$  and let  $X := B_Z(Y)$  be the blow-up of  $Y$  with center  $Z$ . Suppose both  $Y$  and  $X$  are Fano and  $i_X > i_Y$ . Then any rational curve  $C$  satisfying  $-K_Y \cdot C = i_Y$  is contained in  $Z$  and for any such curve  $C$ , the vector bundle  $N_{Z/Y|C}^*$  is ample.

**2.2. Proof of Theorem 2.** This proof assumes that the reader has some familiarity with Mori theory, see for example [Deb01] for a nice introduction.

By the remark above, if  $i_X > i_Y$ , any rational curve  $C$  in  $Y$  such that  $-K_Y \cdot C = i_Y$  is contained in  $Z$ . For such a curve  $C$ , which has minimal degree with respect to an ample line bundle, its deformations in  $Z$  containing a given point cover a subvariety of dimension  $\geq -K_Z \cdot C - 1$  (recall that this is an easy consequence of Riemann-Roch formula and the bend-and-break lemma, see for example [Deb01], §6.5). Since the computations in the proof of Theorem 1(i) show that  $-K_Z \cdot C = \dim(Z) + 1$ , the deformations of  $C$  in  $Z$  containing a given point cover  $Z$ . Therefore, the Picard number of  $Z$  is one (see [Ko96], IV 3.13.3). One deduces that any rational curve  $C'$  of  $Z$  is numerically proportional (in  $N_1(Z)$ ) to  $C$ , and since  $C$  has minimal anti-canonical degree in  $Y$ ,  $C'$  satisfies  $-K_Z \cdot C' \geq \dim(Z) + 1$ , hence  $Z \simeq \mathbb{P}^{\dim(Z)}$  by [CMS00]. Finally, the computations above also show that for any line  $d$  in  $Z$ ,  $N_{Z/Y|d} \simeq \mathcal{O}_d(-1)^{\oplus n - \dim(Z)}$ , which implies that  $N_{Z/Y} \simeq \mathcal{O}_{\mathbb{P}^{\dim(Z)}}(-1)^{\oplus n - \dim(Z)}$  by Theorem (3.2.1) in [OSS81].

Let  $E = \mathbb{P}^{\dim(Z)} \times \mathbb{P}^{n - \dim(Z) - 1}$  be the exceptional divisor of  $\pi$ , and let  $\omega$  be a Mori extremal rational curve in  $X$  such that  $E \cdot \omega > 0$  (such a curve exists by the classical following argument: take any curve with strictly positive intersection with  $E$  and decompose it in the Mori cone  $\text{NE}(X)$  as an effective combination of extremal curves, at least one of these curves has strictly positive intersection with  $E$ ). The corresponding Mori contraction  $\varphi_\omega$  satisfies Wiśniewski's inequality [Wi91]:

$$\dim(\text{Exc}(\varphi_\omega)) + \dim(f) \geq n - 1 + i_X$$

where  $\text{Exc}(\varphi_\omega)$  is the locus of contracted curves, and  $f$  is any non-trivial fiber of  $\varphi_\omega$ . Since every contracted curve is proportional to  $\omega$  in  $N_1(X)$  and since  $\mathcal{O}(E)|_E \simeq \mathcal{O}_{\mathbb{P}^{\dim(Z)} \times \mathbb{P}^{n - \dim(Z) - 1}}(-1, -1)$ , none of these curves are contained in  $E$ , therefore any non-trivial fiber  $f$  of  $\varphi_\omega$  satisfies  $\dim(f) = 1$ . Moreover,  $i_X \geq 2$ , therefore  $\text{Exc}(\varphi_\omega) = X$  by Wiśniewski's inequality above and  $\varphi_\omega$  is a fibration which, by Ando's classification [An85], is a smooth  $\mathbb{P}^1$ -bundle over an  $(n-1)$ -dimensional Fano manifold  $X'$ . Moreover,  $i_X = 2$ , hence  $i_Y = 1$ , therefore  $\dim(Z) = n/2$ .

Let us show now that  $E \cdot f = 1$  for any fiber of  $\varphi_\omega$ . Indeed, if  $\mathbb{P}^1 \rightarrow X'$  is a rational curve of  $X'$ , the surface  $S = \mathbb{P}^1 \times_{X'} X$  is a ruled surface, i.e., a Hirzebruch surface, and the exceptional curve of  $S$  is nothing else than  $\mathbb{P}^1 \times_{X'} E$ , which is a section of  $S \rightarrow \mathbb{P}^1$ . Hence  $E \cdot f = 1$  for any fiber of  $\varphi_\omega$ .

One immediately deduces that  $\varphi_\omega : E \rightarrow X'$  is an isomorphism, hence  $X = \mathbb{P}(\mathcal{E})$  for some rank 2 bundle  $\mathcal{E}$  over  $X' \simeq E \simeq \mathbb{P}^{n/2} \times \mathbb{P}^{n/2-1}$ , and  $E$  defines a sub-line bundle

of  $\mathcal{E}$ . Therefore  $\mathcal{E}$  splits and since  $N_{E/X} \simeq \mathcal{O}_{\mathbb{P}^{n/2} \times \mathbb{P}^{n/2-1}}(-1, -1)$ , one deduces that

$$X \simeq \mathbb{P}(\mathcal{O}_{\mathbb{P}^{n/2} \times \mathbb{P}^{n/2-1}} \oplus \mathcal{O}_{\mathbb{P}^{n/2} \times \mathbb{P}^{n/2-1}}(1, 1)),$$

which ends the proof. ■

### 3. SOME COMMENTS AND SOME MORE EXAMPLES

**3.1. On the normal bundle of the center.** The following proposition sheds some light on the example explained in Proposition 1.

**Proposition 2.** *Let  $\pi : X \rightarrow Y$  be a blow-up with smooth connected center  $Z$  between Fano manifolds  $X$  and  $Y$  of dimension  $n$ . Suppose moreover that the conormal bundle  $N_{Z/Y}^*$  is ample. Then,*

- (i) either  $i_X \leq i_Y$ ,
- (ii) or  $i_X = 2$ ,  $i_Y = 1$ ,  $Z$  is a Fano manifold,  $X$  is a  $\mathbb{P}^1$ -bundle over the  $(n-1)$ -dimensional Fano manifold  $\mathbb{P}(N_{Z/Y})$  and  $Y$  is a  $\mathbb{P}^{n-\dim(Z)}$ -bundle over  $Z$ .

(Sketch of) proof. Suppose  $i_X > i_Y$  and let denote by  $E$  the exceptional divisor of  $\pi$ . Since  $N_{Z/Y}^*$  is ample,  $-E|_E$  is also ample and by Grauert's criterion,  $E$  is contractible to a point. Moreover, if  $\omega$  is a Mori extremal rational curve in  $X$  such that  $E \cdot \omega > 0$ , using the same arguments as in the proof of Theorem 2, the corresponding Mori contraction  $\varphi_\omega$  is a  $\mathbb{P}^1$ -bundle over the Fano manifold  $E \simeq \mathbb{P}(N_{Z/Y})$ . Finally,  $Z$  is Fano by [SW90] and one has  $\rho_Y + 1 = \rho_X = \rho_E + 1 = \rho_Z + 2$ , hence  $\rho_Y = \rho_Z + 1$ . This implies that there is at least one Mori extremal curve of  $Y$  which is not contained in  $Z$ . Since  $\pi$  is surjective, the Mori cone  $\text{NE}(Y)$  is generated by the images of Mori extremal curve of  $X$ , which are contained in  $E$ , except for the fibers  $f$  of the  $\mathbb{P}^1$ -bundle structure  $X \rightarrow E$ . This implies that  $\pi(f)$  is extremal in  $Y$  and the corresponding extremal contraction  $\psi : Y \rightarrow W$  is a fibration. But then, the fibers of  $\psi$  have dimension less or equal to  $n - \dim(Z)$ , hence equal to  $n - \dim(Z)$  since  $-K_Y \cdot \pi(f) = n - \dim(Z) + 1$ . Therefore the generic fiber of  $\psi$  is  $\mathbb{P}^{n-\dim(Z)}$ , and finally every fiber of  $\psi$  is  $\mathbb{P}^{n-\dim(Z)}$  and meets  $Z$  transversally at exactly one point (all this is verified since on  $X$ , the extremal contraction associated to  $f$  is a  $\mathbb{P}^1$ -bundle). Finally,  $W \simeq Z$ . ■

**3.2. Some examples.** The previous Proposition implies that if  $\pi : X \rightarrow Y$  is a blow-up with smooth connected center  $Z$  between Fano manifolds  $X$  and  $Y$  with  $i_X > i_Y \geq 2$ , then the conormal bundle  $N_{Z/Y}^*$  is not ample, although its restriction to any rational curve of minimal  $-K_Y$ -degree (recall that such a curve has to be contained in  $Z$ ) is ample as we saw in the proof of Theorem 1 ! It is therefore the good place to give a list of examples (communicated to me by Cinzia Casagrande) of blow-ups  $\pi : X \rightarrow Y$  with smooth connected center  $Z$  between Fano manifolds  $X$  and  $Y$  with  $i_X > i_Y \geq 2$ .

**Examples.** Let  $a$ ,  $d$ ,  $r$  and  $s$  be positive integers, let  $\mathcal{E}$  be the following rank  $r + s$  vector bundle over  $\mathbb{P}^a$ :  $\mathcal{E} = \mathcal{O}_{\mathbb{P}^a}^{\oplus r} \oplus \mathcal{O}_{\mathbb{P}^a}(d)^{\oplus s}$  and let  $\mathcal{F} = \mathcal{O}_{\mathbb{P}^a}(d)^{\oplus s}$  be the rank  $s$  vector subbundle of  $\mathcal{E}$  defined by the  $\mathcal{O}_{\mathbb{P}^a}(d)$ 's factors. Define  $Y := \mathbb{P}(\mathcal{E})$  and  $Z := \mathbb{P}(\mathcal{F}) \simeq \mathbb{P}^a \times \mathbb{P}^{s-1}$ . This submanifold  $Z$  of  $Y$  has codimension  $r$  and normal bundle  $N_{Z/Y}$  equal to  $\mathcal{O}_{\mathbb{P}^a \times \mathbb{P}^{s-1}}(-d, 1)^{\oplus r}$ . Finally, let  $X := B_Z(Y)$  be the blow-up of  $Y$  with center  $Z$ . Easy computations show that  $Y$  is Fano if and only if  $a \geq rd$  and  $X$  is Fano if and only if  $a \geq d$ , and that under these assumptions, one has

$$i_Y = \min(r + s, 1 + a - rd) \text{ and } i_X = \min(r - 1, s + 1, 1 + a - d),$$

which leads to many examples satisfying  $i_X > i_Y \geq 2$ . The example of lowest dimension (namely 10) for such  $X$  and  $Y$  is given when  $(a, d, r, s) = (5, 1, 4, 2)$ . Let us also say that many of these examples lead to  $X$  and  $Y$  satisfying  $r_X < i_X$  and  $r_Y < i_Y$ . In the case  $s = 1$ , the Fano manifolds  $Y$ 's have been considered by Debarre (see [Deb01], §5.11) to construct Fano manifolds of high degree  $(-K_Y)^{\dim(Y)}$ .

**3.3. Minimal degree of free rational curves.** When studying Fano manifolds, one often uses *free* rational curves, which means rational curves  $f : \mathbb{P}^1 \rightarrow X$  such that

$$f^*T_X \simeq \mathcal{O}_{\mathbb{P}^1}(a_1) \oplus \cdots \oplus \mathcal{O}_{\mathbb{P}^1}(a_{\dim X})$$

with all the  $a_i$ 's greater or equal to 0 (see [Deb01], Chapter 4 for details). One may then introduce another invariant:

$$f_X := \min\{-K_X \cdot C \mid C \text{ is a free rational curve of } X\},$$

which is of great importance in Hwang and Mok's recent works. It can be interpreted as the minimal anti-canonical degree of rational curves whose deformations cover an open dense subset of  $X$ . It is an easy exercise to show that if  $f : X \rightarrow Y$  is a surjective morphism between Fano manifolds  $X$  and  $Y$ , then  $f_X \leq f_Y$ . Of course, in any of the examples above where  $i_X > i_Y$ , the rational curves in  $Y$  of minimal anti-canonical degree are not free curves and their deformations do not cover any dense open subset of  $Y$ .

**3.4. A final remark on a related question.** In the above results, the assumption that both  $X$  and  $Y$  are Fano is essential: when  $\pi : X \rightarrow Y$  is a blow-up with smooth connected center  $Z$  between complex manifolds  $X$  and  $Y$ , understanding on which conditions  $X$  Fano (resp.  $Y$  Fano) implies  $Y$  Fano (resp.  $X$  Fano) is a completely different question, whose study has been initiated by Wiśniewski in [Wi91]. In particular, no condition on the dimension of the center is neither necessary nor sufficient (except of course when  $Z$  is a point, see [BCW02] for a complete classification) to get one of the implications above: the examples in §3.2, in the particular case where  $rd > a \geq d$ , give examples of smooth blow-up  $\pi : X \rightarrow Y$  between complex manifolds  $X$  and  $Y$  with  $X$  being Fano and  $Y$  not.

*Thanks to Cinzia Casagrande and Olivier Debarre for their comments on a preliminary version of this Note.*

#### REFERENCES

- [An85] T. Ando. On extremal rays of the higher dimensional varieties. *Invent. Math.* 81, 347–357 (1985).
- [BCW02] L. Bonavero, F. Campana et J.A. Wiśniewski. Variétés complexes dont l'éclatée en un point est de Fano. *C. R. Math. Acad. Sci. Paris* 334, no. 6, 463–468 (2002).
- [BCDD03] L. Bonavero, C. Casagrande, O. Debarre and S. Druel. Sur une conjecture de Mukai. *Comment. Math. Helv.* 78, 601–626 (2003).
- [CMS00] K. Cho, Y. Miyaoka, N. Shepherd-Barron. Characterizations of projective space and applications to complex symplectic manifolds. *Higher dimensional birational geometry (Kyoto, 1997)*, *Adv. Stud. Pure Math.*, 35, Math. Soc. Japan, Tokyo, 1–88 (2002).
- [Deb01] O. Debarre. Higher-Dimensional Algebraic Geometry. *Universitext*, Springer Verlag, (2001).
- [IP99] V.A. Iskovskikh, Yu.G. Prokhorov. Fano varieties. Algebraic geometry, V. *Encyclopaedia Math. Sci.* 47, Springer-Verlag, Berlin, 1999.
- [Ko96] J. Kollár. Rational curves on algebraic varieties. *Ergebnisse der Mathematik und ihre Grenzgebiete. 3 Folge 032*, Springer-Verlag, 1996.

- [La83] R. Lazarsfeld. Some applications of the theory of positive vector bundles. *Complete intersections, Lect. 1st Sess. C.I.M.E., Acireale/Italy 1983, Lect. Notes Math. 1092, 29–61 (1984)*.
- [OSS81] C. Okonek, M. Schneider, H. Spindler. Vector bundles on complex projective spaces. *Progress in Mathematics, 3. Birkhäuser, Boston, Mass., 1980. vii+389 pp.*
- [SW90] M. Szurek, J. Wiśniewski. Fano bundles over  $\mathbb{P}^3$  and  $Q^3$ . *Pac. J. Math. 141, No.1, 197-208 (1990)*.
- [Wi91] J. Wiśniewski. On contractions of extremal rays of Fano manifolds. *J. Reine Angew. Math. 417, 141–157 (1991)*.

---

*Laurent Bonavero. Institut Fourier, UMR 5582, Université de Grenoble 1, BP 74.  
38402 Saint Martin d'Hères. FRANCE  
e-mail : bonavero@ujf-grenoble.fr*