

Learning rational stochastic languages

François Denis, Yann Esposito, Amaury Habrard

Laboratoire d'Informatique Fondamentale de Marseille (L.I.F.) UMR CNRS 6166
{fdenis,esposito,habrard}@cmi.univ-mrs.fr

Abstract Given a finite set of words w_1, \dots, w_n independently drawn according to a fixed unknown distribution law P called a *stochastic language*, an usual goal in Grammatical Inference is to infer an estimate of P in some class of probabilistic models, such as *Probabilistic Automata* (PA). Here, we study the class $\mathcal{S}_{\mathbb{R}}^{rat}(\Sigma)$ of *rational stochastic languages*, which consists in stochastic languages that can be generated by *Multiplicity Automata* (MA) and which strictly includes the class of stochastic languages generated by PA. Rational stochastic languages have minimal normal representation which may be very concise, and whose parameters can be efficiently estimated from stochastic samples. We design an efficient inference algorithm DEES which aims at building a minimal normal representation of the target. Despite the fact that no recursively enumerable class of MA computes exactly $\mathcal{S}_{\mathbb{Q}}^{rat}(\Sigma)$, we show that DEES strongly identifies $\mathcal{S}_{\mathbb{Q}}^{rat}(\Sigma)$ in the limit. We study the intermediary MA output by DEES and show that they compute rational series which converge absolutely to one and which can be used to provide stochastic languages which closely estimate the target.

1 Introduction

In probabilistic grammatical inference, it is supposed that data arise in the form of a finite set of words w_1, \dots, w_n , built on a predefined alphabet Σ , and independently drawn according to a fixed unknown distribution law on Σ^* called a *stochastic language*. Then, an usual goal is to try to infer an estimate of this distribution law in some class of probabilistic models, such as *Probabilistic Automata* (PA), which have the same expressivity as Hidden Markov Models (HMM). PA are identifiable in the limit [6]. However, to our knowledge, there exists no efficient inference algorithm able to deal with the whole class of stochastic languages that can be generated from PA. Most of the previous works use restricted subclasses of PA such as Probabilistic Deterministic Automata (PDA) [5,12]. In the other hand, Probabilistic Automata are particular cases of *Multiplicity Automata*, and stochastic languages which can be generated by multiplicity automata are special cases of *rational languages* that we call *rational stochastic languages*. MA have been used in grammatical inference in a variant of the exact learning model of Angluin [3,1,2] but not in probabilistic grammatical inference. Let us design by $\mathcal{S}_K^{rat}(\Sigma)$, the class of rational stochastic languages over the semiring K . When $K = \mathbb{Q}^+$ or $K = \mathbb{R}^+$, $\mathcal{S}_K^{rat}(\Sigma)$ is exactly the class of stochastic languages generated by PA with parameters in K . But, when $K = \mathbb{Q}$ or $K = \mathbb{R}$, we obtain strictly greater classes which provide several advantages and at least one drawback: elements of $\mathcal{S}_{K^+}^{rat}(\Sigma)$ may have significantly smaller representation in $\mathcal{S}_K^{rat}(\Sigma)$ which is clearly an advantage from a learning perspective; elements of $\mathcal{S}_K^{rat}(\Sigma)$ have a minimal normal representation while such normal representations do not exist for PA; parameters of these minimal representations are directly

related to probabilities of some natural events of the form $u\Sigma^*$, which can be efficiently estimated from stochastic samples; lastly, when K is a field, rational series over K form a vector space and efficient linear algebra techniques can be used to deal with rational stochastic languages. However, the class $\mathcal{S}_{\mathbb{Q}}^{rat}(\Sigma)$ presents a serious drawback: there exists no recursively enumerable subset of MA which exactly generates it [6]. Moreover, this class of representations is unstable: arbitrarily close to an MA which generates a stochastic language, we may find MA whose associated rational series r takes negative values and is not absolutely convergent: the global weight $\sum_{w \in \Sigma^*} r(w)$ may be unbounded or not (absolutely) defined. However, we show that $\mathcal{S}_{\mathbb{Q}}^{rat}(\Sigma)$ is strongly identifiable in the limit: we design an algorithm DEES such that, for any target $P \in \mathcal{S}_{\mathbb{Q}}^{rat}(\Sigma)$ and given access to an infinite sample S drawn according to P , will converge in a finite but unbounded number of steps to a minimal normal representation of P . Moreover, DEES is efficient: it runs within polynomial time in the size of the input and it computes a minimal number of parameters with classical statistical rates of convergence. However, before converging to the target, DEES output MA which are close to the target but which do not compute stochastic languages. The question is: what kind of guarantees do we have on these intermediary hypotheses and how can we use them for a probabilistic inference purpose? We show that, since the algorithm aims at building a minimal normal representation of the target, the intermediary hypotheses r output by DEES have a nice property: they absolutely converge to 1, i.e. $\bar{r} = \sum_{w \in \Sigma^*} |r(w)| < \infty$ and $\sum_{k \geq 0} r(\Sigma^k) = 1$. As a consequence, $r(X)$ is defined without ambiguity for any $X \subseteq \Sigma^*$, and it can be shown that $N_r = \sum_{r(u) < 0} |r(u)|$ tends to 0 as the learning proceeds. Given any such series r , we can efficiently compute a stochastic language p_r , which is not rational, but has the property that $e^{N_r/\bar{r}} \leq p_r(u)/r(u) \leq 1$ for any word u such that $r(u) > 0$. Our conclusion is that, despite the fact that no recursively enumerable class of MA represents the class of rational stochastic languages, MA can be used efficiently to infer such stochastic languages.

Classical notions on stochastic languages, rational series, and multiplicity automata are recalled in Section 2. We study an example which shows that the representation of rational stochastic languages by MA with real parameters may be very concise. We introduce our inference algorithm DEES in Section 3 and we show that $\mathcal{S}_{\mathbb{Q}}^{rat}(\Sigma)$ is strongly identifiable in the limit. We study the properties of the MA output by DEES in Section 4 and we show that they define absolutely convergent rational series which can be used to compute stochastic languages which are estimates of the target.

2 Preliminaries

Formal power series and stochastic languages. Let Σ^* be the set of words on the finite alphabet Σ . The empty word is denoted by ε and the length of a word u is denoted by $|u|$. For any integer k , let $\Sigma^k = \{u \in \Sigma^* : |u| = k\}$ and $\Sigma^{\leq k} = \{u \in \Sigma^* : |u| \leq k\}$. We denote by $<$ the length-lexicographic order on Σ^* . A subset P of Σ^* is *prefixial* if for any $u, v \in \Sigma^*$, $uv \in P \Rightarrow u \in P$. For any $S \subseteq \Sigma^*$, let $pref(S) = \{u \in \Sigma^* : \exists v \in \Sigma^*, uv \in S\}$ and $fact(S) = \{v \in \Sigma^* : \exists u, w \in \Sigma^*, uvw \in S\}$.

Let Σ be a finite alphabet and K a semiring. A *formal power series* is a mapping r of Σ^* into K . In this paper, we always suppose that $K \in \{\mathbb{R}, \mathbb{Q}, \mathbb{R}^+, \mathbb{Q}^+\}$. The set of all formal power series is denoted by $K\langle\langle \Sigma \rangle\rangle$. Let us denote by $supp(r)$ the *support* of r , i.e. the set $\{w \in \Sigma^* : r(w) \neq 0\}$.

A *stochastic language* is a formal series p which takes its values in \mathbb{R}^+ and such that $\sum_{w \in \Sigma^*} p(w) = 1$. For any language $L \subseteq \Sigma^*$, let us denote $\sum_{w \in L} p(w)$ by $p(L)$. The set of all stochastic languages over Σ is denoted by $\mathcal{S}(\Sigma)$. For any stochastic language p and any word u such that $p(u\Sigma^*) \neq 0$, we define the stochastic language $u^{-1}p$ by $u^{-1}p(w) = \frac{p(uw)}{p(u\Sigma^*)} \cdot u^{-1}p$ is called the *residual language* of p wrt u . Let us denote by $res(p)$ the set $\{u \in \Sigma^* : p(u\Sigma^*) \neq 0\}$ and by $Res(p)$ the set $\{u^{-1}p : u \in res(p)\}$. We call *sample* any finite sequence of words. Let S be a sample. We denote by P_S the empirical distribution on Σ^* associated with S . A *complete presentation* of P is an infinite sequence S of words independently drawn according to P . We denote by S_n the sequence composed of the n first words of S . We shall make a frequent use of the Borel-Cantelli Lemma which states that if $(A_k)_{k \in \mathbb{N}}$ is a sequence of events such that $\sum_{k \in \mathbb{N}} Pr(A_k) < \infty$, then the probability that a finite number of A_k occurs is 1.

Automata. Let K be a semiring. A K -multiplicity automaton (MA) is a 5-tuple $\langle \Sigma, Q, \varphi, \iota, \tau \rangle$ where Q is a finite set of states, $\varphi : Q \times \Sigma \times Q \rightarrow K$ is the transition function, $\iota : Q \rightarrow K$ is the initialization function and $\tau : Q \rightarrow K$ is the termination function. Let $Q_I = \{q \in Q \mid \iota(q) \neq 0\}$ be the set of *initial states* and $Q_T = \{q \in Q \mid \tau(q) \neq 0\}$ be the set of *terminal states*. The *support* of an MA $A = \langle \Sigma, Q, \varphi, \iota, \tau \rangle$ is the NFA $supp(A) = \langle \Sigma, Q, Q_I, Q_T, \delta \rangle$ where $\delta(q, x) = \{q' \in Q \mid \varphi(q, x, q') \neq 0\}$. We extend the transition function φ to $Q \times \Sigma^* \times Q$ by $\varphi(q, wx, r) = \sum_{s \in Q} \varphi(q, w, s) \varphi(s, x, r)$ and $\varphi(q, \varepsilon, r) = 1$ if $q = r$ and 0 otherwise, for any $q, r \in Q, x \in \Sigma$ and $w \in \Sigma^*$. For any finite subset $L \subseteq \Sigma^*$ and any $R \subseteq Q$, define $\varphi(q, L, R) = \sum_{w \in L, r \in R} \varphi(q, w, r)$.

For any MA A , let r_A be the series defined by $r_A(w) = \sum_{q, r \in Q} \iota(q) \varphi(q, w, r) \tau(r)$. For any $q \in Q$, we define the series $r_{A,q}$ by $r_{A,q}(w) = \sum_{r \in Q} \varphi(q, w, r) \tau(r)$. A state $q \in Q$ is *accessible* (resp. *co-accessible*) if there exists $q_0 \in Q_I$ (resp. $q_t \in Q_T$) and $u \in \Sigma^*$ such that $\varphi(q_0, u, q) \neq 0$ (resp. $\varphi(q, u, q_t) \neq 0$). An MA is *trimmed* if all its states are accessible and co-accessible. From now, we only consider trimmed MA.

A *Probabilistic Automaton (PA)* is a trimmed MA $\langle \Sigma, Q, \varphi, \iota, \tau \rangle$ s.t. ι, φ and τ take their values in $[0, 1]$, such that $\sum_{q \in Q} \iota(q) = 1$ and for any state $q, \tau(q) + \varphi(q, \Sigma, Q) = 1$. Probabilistic automata generate stochastic languages. A *Probabilistic Deterministic Automaton (PDA)* is a PA whose support is deterministic.

For any class C of multiplicity automata over K , let us denote by $\mathcal{S}_K^C(\Sigma)$ the class of all stochastic languages which are recognized by an element of C .

Rational series and rational stochastic languages. Rational series have several characterization ([11,4,10]). Here, we shall say that a formal power series over Σ is K -rational iff there exists a K -multiplicity automaton A such that $r = r_A$, where $K \in \{\mathbb{R}, \mathbb{R}^+, \mathbb{Q}, \mathbb{Q}^+\}$. Let us denote by $K^{rat} \langle \langle \Sigma \rangle \rangle$ the set of K -rational series over Σ and by $\mathcal{S}_K^{rat}(\Sigma) = K^{rat} \langle \langle \Sigma \rangle \rangle \cap \mathcal{S}(\Sigma)$, the set of *rational stochastic languages* over K . Rational stochastic languages have been studied in [7] from a language theoretical point of view. Inclusion relations between classes of rational stochastic languages are summarized on Fig 1. It is worth noting that $\mathcal{S}_{\mathbb{R}}^{PDA}(\Sigma) \subsetneq \mathcal{S}_{\mathbb{R}}^{PA}(\Sigma) \subsetneq \mathcal{S}_{\mathbb{R}}^{rat}(\Sigma)$.

Let P be a rational stochastic language. The MA $A = \langle \Sigma, Q, \varphi, \iota, \tau \rangle$ is a *reduced representation* of P if (i) $P = P_A$, (ii) $\forall q \in Q, P_{A,q} \in \mathcal{S}(\Sigma)$ and (iii) the set $\{P_{A,q} : q \in Q\}$ is linearly independent. It can be shown that $Res(P)$ spans a finite dimensional vector subspace $[Res(P)]$ of $\mathbb{R} \langle \langle \Sigma \rangle \rangle$. Let Q_P be the smallest subset of $res(P)$ s.t. $\{u^{-1}P : u \in Q_P\}$ spans $[Res(P)]$. It is a finite prefixial subset of Σ^* . Let $A = \langle \Sigma, Q_P, \varphi, \iota, \tau \rangle$ be the MA defined by:




Figure1. Inclusion relations between classes of rational stochastic languages.

- $\iota(\varepsilon) = 1, \iota(u) = 0$ otherwise; $\tau(u) = u^{-1}P(\varepsilon)$,
- $\varphi(u, x, ux) = u^{-1}P(x\Sigma^*)$ if $u, ux \in Q_P$ and $x \in \Sigma$,
- $\varphi(u, x, v) = \alpha_v u^{-1}P(x\Sigma^*)$ if $x \in \Sigma, ux \in (Q_P \Sigma \setminus Q_P) \cap res(P)$ and $(ux)^{-1}P = \sum_{v \in Q_P} \alpha_v v^{-1}P$.

It can be shown that A is a reduced representation of P ; A is called the *prefixial reduced representation* of P . Note that the parameters of A correspond to natural components of the residual of P and can be estimated by using samples of P .

We give below an example of a rational stochastic language which cannot be generated by a PA. Moreover, for any integer N there exists a rational stochastic language which can be generated by a multiplicity automaton with 3 states and such that the smallest PA which generates it has N states. That is, considering rational stochastic language makes it possible to deal with stochastic languages which cannot be generated by PA; it also permits to significantly decrease the size of their representation.

Proposition 1. For any $\alpha \in \mathbb{R}$, let A_α be the MA described on Fig. 2. Let $S_\alpha = \{(\lambda_0, \lambda_1, \lambda_2) \in \mathbb{R}^3 : r_{A_\alpha} \in \mathcal{S}(\Sigma)\}$. If $\alpha/(2\pi) = p/q \in \mathbb{Q}$ where p and q are relatively prime, S_α is the convex hull of a polygon with q vertices which are the residual languages of any one of them. If $\alpha/(2\pi) \notin \mathbb{Q}$, S_α is the convex hull of an ellipse, any point of which, is a stochastic language which cannot be computed by a PA.

Proof (sketch).

Let r_{q_0}, r_{q_1} and r_{q_2} be the series associated with the states of A_α . We have

$$r_{q_0}(a^n) = \frac{\cos n\alpha - \sin n\alpha}{2^n}, r_{q_1}(a^n) = \frac{\cos n\alpha + \sin n\alpha}{2^n} \text{ and } r_{q_2}(a^n) = \frac{1}{2^n}.$$

The sums $\sum_{n \in \mathbb{N}} r_{q_0}(a^n)$, $\sum_{n \in \mathbb{N}} r_{q_1}(a^n)$ and $\sum_{n \in \mathbb{N}} r_{q_2}(a^n)$ converge since $|r_{q_i}(a^n)| = O(2^{-n})$ for $i = 0, 1, 2$. Let us denote $\sigma_i = \sum_{n \in \mathbb{N}} r_{q_i}(a^n)$ for $i = 0, 1, 2$. Check that

$$\sigma_0 = \frac{4 - 2 \cos \alpha - 2 \sin \alpha}{5 - 4 \cos \alpha}, \sigma_1 = \frac{4 - 2 \cos \alpha + 2 \sin \alpha}{5 - 4 \cos \alpha} \text{ and } \sigma_2 = 2.$$

Consider the 3-dimensional vector subspace \mathcal{V} of $\mathbb{R}\langle\langle \Sigma \rangle\rangle$ generated by r_{q_0}, r_{q_1} and r_{q_2} and let $r = \lambda_0 r_{q_0} + \lambda_1 r_{q_1} + \lambda_2 r_{q_2}$ be a generic element of \mathcal{V} . We have $\sum_{n \in \mathbb{N}} r(a^n) = \lambda_0 \sigma_0 + \lambda_1 \sigma_1 + \lambda_2 \sigma_2$. The equation $\lambda_0 \sigma_0 + \lambda_1 \sigma_1 + \lambda_2 \sigma_2 = 1$ defines a plane \mathcal{H} in \mathcal{V} .

Consider the constraints $r(a^n) \geq 0$ for any $n \geq 0$. The elements r of \mathcal{H} which satisfies all the constraints $r(a^n) \geq 0$ are exactly the stochastic languages in \mathcal{H} .

If $\alpha/(2\pi) = k/h \in \mathbb{Q}$ where k and h are relatively prime, the set of constraints $\{r(a^n) \geq 0\}$ is finite: it delimites a convex regular polygon P in the plane \mathcal{H} . Let p be a vertex of P . It can be shown that its residual languages are exactly the h vertices of P and any PA generating p must have at least h states.

If $\alpha/(2\pi) \notin \mathbb{Q}$, the constraints delimitate an ellipse E . Let p be an element of E . It can be shown, by using techniques developed in [7], that its residual languages are dense in E and that no PA can generate p . \square

Matrices. We consider the Euclidan norm on \mathbb{R}^n : $\|(x_1, \dots, x_n)\| = (x_1^2 + \dots + x_n^2)^{1/2}$. For any $R \geq 0$, let us denote by $B(0, R)$ the set $\{x \in \mathbb{R}^n : \|x\| \leq R\}$. The induced norm on the set of $n \times n$ square matrices M over \mathbb{R} is defined by: $\|M\| = \sup\{\|Mx\| : x \in \mathbb{R}^n \text{ with } \|x\| = 1\}$. Some properties of the induced norm: $\|Mx\| \leq \|M\| \cdot \|x\|$ for all $M \in \mathbb{R}^{n \times n}$, $x \in \mathbb{R}^n$; $\|MN\| \leq \|M\| \cdot \|N\|$ for all $M, N \in \mathbb{R}^{n \times n}$; $\lim_{k \rightarrow \infty} \|M^k\|^{1/k} = \rho(M)$ where $\rho(M)$ is the spectral radius of M , i.e. the maximum magnitude of the eigen values of M (Gelfand's Formula).




Figure2. When $\lambda_0 = \lambda_2 = 1$ and $\lambda_1 = 0$, the MA $A_{\pi/6}$ defines a stochastic language P whose prefixed reduced representation is the MA B (with approximate values on transitions). In fact, P can be computed by a PDA and the smallest PA computing it is C .

3 Identifying $S_{\mathbb{Q}}^{rat}(\Sigma)$ in the limit.

Let S be a non empty finite sample of Σ^* , let Q be prefixial subset of $pref(S)$, let $v \in pref(S) \setminus Q$, and let $\epsilon > 0$. We denote by $I(Q, v, S, \epsilon)$ the following set of inequations over the set of variables $\{x_u | u \in Q\}$:

$$I(Q, v, S, \epsilon) = \{|v^{-1}P_S(w\Sigma^*) - \sum_{u \in Q} x_u u^{-1}P_S(w\Sigma^*)| \leq \epsilon | w \in fact(S)\} \cup \{\sum_{u \in Q} x_u = 1\}.$$

Let DEES be the following algorithm:

Input: a sample S

Output: a prefixial reduced MA $A = \langle \Sigma, Q, \varphi, \iota, \tau \rangle$

$Q \leftarrow \{\epsilon\}$, $\iota(\epsilon) = 1$, $\tau(\epsilon) = P_S(\epsilon)$, $F \leftarrow \Sigma \cap pref(S)$

while $F \neq \emptyset$ do {

$v = ux = \text{Min}F$ where $u \in \Sigma^*$ and $x \in \Sigma, F \leftarrow F \setminus \{v\}$
 if $I(Q, v, S, |S|^{-1/3})$ has no solution then{
 $Q \leftarrow Q \cup \{v\}, \iota(v) = 0, \tau(v) = P_S(v)/P_S(v\Sigma^*),$
 $\varphi(u, x, v) = P_S(v\Sigma^*)/P_S(u\Sigma^*), F \leftarrow F \cup \{vx \in \text{res}(P_S) | x \in \Sigma\}$
 else{
 let $(\alpha_w)_{w \in Q}$ be a solution of $I(Q, v, S, |S|^{-1/3})$
 $\varphi(u, x, w) = \alpha_w P_S(v\Sigma^*)$ for any $w \in Q$ }}

Lemma 1. *Let P be a stochastic language and let $u_0, u_1, \dots, u_n \in \text{Res}(P)$ be such that $\{u_0^{-1}P, u_1^{-1}P, \dots, u_n^{-1}P\}$ is linearly independent. Then, with probability one, for any complete presentation S of P , there exist a positive number ϵ and an integer M such that $I(\{u_1, \dots, u_n\}, u_0, S_m, \epsilon)$ has no solution for every $m \geq M$.*

Proof. Let S be a complete presentation of P . Suppose that for every $\epsilon > 0$ and every integer M , there exists $m \geq M$ such that $I(\{u_1, \dots, u_n\}, u_0, S_m, \epsilon)$ has a solution. Then, for any integer k , there exists $m_k \geq k$ such that $I(\{u_1, \dots, u_n\}, u_0, S_{m_k}, 1/k)$ has a solution $(\alpha_{1,k}, \dots, \alpha_{n,k})$. Let $\rho_k = \text{Max}\{1, |\alpha_{1,k}|, \dots, |\alpha_{n,k}|\}$, $\gamma_{0,k} = 1/\rho_k$ and $\gamma_{i,k} = -\alpha_{i,k}/\rho_k$ for $1 \leq i \leq n$. For every k , $\text{Max}\{|\gamma_{i,k}| : 0 \leq i \leq n\} = 1$. Check that

$$\forall k \geq 0, \left| \sum_{i=0}^n \gamma_{i,k} u_i^{-1} P_{S_{m_k}}(w\Sigma^*) \right| \leq \frac{1}{\rho_k k} \leq \frac{1}{k}.$$

There exists a subsequence $(\alpha_{1,\phi(k)}, \dots, \alpha_{n,\phi(k)})$ of $(\alpha_{1,k}, \dots, \alpha_{n,k})$ such that $(\gamma_{0,\phi(k)}, \dots, \gamma_{n,\phi(k)})$ converges to $(\gamma_0, \dots, \gamma_n)$. We show below that we should have $\sum_{i=0}^n \gamma_i u_i^{-1} P(w\Sigma^*) = 0$ for every word w , which is contradictory with the independence assumption since $\text{Max}\{\gamma_i : 0 \leq i \leq n\} = 1$.

Let $w \in \text{fact}(\text{supp}(P))$. With probability 1, there exists an integer k_0 such that $w \in \text{fact}(S_{m_k})$ for any $k \geq k_0$. For such a k , we can write

$$\gamma_i u_i^{-1} P = (\gamma_i u_i^{-1} P - \gamma_i u_i^{-1} P_{S_{m_k}}) + (\gamma_i - \gamma_{i,\phi(k)}) u_i^{-1} P_{S_{m_k}} + \gamma_{i,\phi(k)} u_i^{-1} P_{S_{m_k}}$$

and therefore

$$\left| \sum_{i=0}^n \gamma_i u_i^{-1} P(w\Sigma^*) \right| \leq \sum_{i=0}^n |u_i^{-1}(P - P_{S_{m_k}})(w\Sigma^*)| + \sum_{i=0}^n |\gamma_i - \gamma_{i,\phi(k)}| + \frac{1}{k}$$

which converges to 0 when k tends to infinity. \square

Let P be a stochastic language over Σ , let $\mathcal{A} = (A_i)_{i \in I}$ be a family of subsets of Σ^* , let S be a finite sample drawn according to P , and let P_S be the empirical distribution associated with S . It can be shown [13,9] that for any confidence parameter δ , with a probability greater than $1 - \delta$, for any $i \in I$,

$$|P_S(A_i) - P(A_i)| \leq c \sqrt{\frac{\text{VC}(\mathcal{A}) - \log \frac{\delta}{4}}{\text{Card}(S)}} \quad (1)$$

where $\text{VC}(\mathcal{A})$ is the dimension of Vapnik-Chervonenkis of \mathcal{A} and c is a constant.

When $\mathcal{A} = (\{w\Sigma^*\})_{w \in \Sigma^*}$, $\text{VC}(\mathcal{A}) \leq 2$. Indeed, let $r, s, t \in \Sigma^*$ and let $Y = \{r, s, t\}$. Let u_{rs} (resp. u_{rt}, u_{st}) be the longest prefix shared by r and s (resp. r and t , s

and t). One of these 3 words is a prefix of the two other ones. Suppose that u_{rs} is a prefix of u_{rt} and u_{st} . Then, there exists no word w such that $w\Sigma^* \cap Y = \{r, s\}$. Therefore, no subset containing more than two elements can be shattered by \mathcal{A} .

$$\text{Let } \Psi(\epsilon, \delta) = \frac{\epsilon^2}{\delta^2} (2 - \log \frac{\delta}{4}).$$

Lemma 2. *Let $P \in \mathcal{S}(\Sigma)$ and let S be a complete presentation of P . For any precision parameter ϵ , any confidence parameter δ , any $n \geq \Psi(\epsilon, \delta)$, with a probability greater than $1 - \delta$, $|P_n(w\Sigma^*) - P(w\Sigma^*)| \leq \epsilon$ for all $w \in \Sigma^*$.*

Proof. Use inequality (1). □

Check that for any α such that $-1/2 < \alpha < 0$ and any $\beta < -1$, if we define $\epsilon_k = k^\alpha$ and $\delta_k = k^\beta$, there exists K such that for all $k \geq K$, we have $k \geq \Psi(\epsilon_k, \delta_k)$. For such choices of α and β , we have $\lim_{k \rightarrow \infty} \epsilon_k = 0$ and $\sum_{k \geq 1} \delta_k < \infty$.

Lemma 3. *Let $P \in \mathcal{S}(\Sigma)$, $u_0, u_1, \dots, u_n \in \text{res}(P)$ and $\alpha_1, \dots, \alpha_n \in \mathbb{R}$ be such that $u_0^{-1}P = \sum_{i=1}^n \alpha_i u_i^{-1}P$. Then, with probability one, for any complete presentation S of P , there exists K s.t. $I(\{u_1, \dots, u_n\}, u_0, S_k, k^{-1/3})$ has a solution for every $k \geq K$.*

Proof. Let S be a complete presentation of P . Let $\alpha_0 = 1$ and let $R = \text{Max}\{|\alpha_i| : 0 \leq i \leq n\}$. With probability one, there exists K_1 s.t. $\forall k \geq K_1, \forall i = 0, \dots, n$, $|u_i^{-1}S_k| \geq \Psi([k^{1/3}(n+1)R]^{-1}, [(n+1)k^2]^{-1})$. Let $k \geq K_1$. For any $X \subseteq \Sigma^*$,

$$|u_0^{-1}P_{S_k}(X) - \sum_{i=1}^n \alpha_i u_i^{-1}P_{S_k}(X)| \leq |u_0^{-1}P_{S_k}(X) - u_0^{-1}P(X)| + \sum_{i=1}^n |\alpha_i| |u_i^{-1}P_{S_k}(X) - u_i^{-1}P(X)|.$$

From Lemma 2, with probability greater than $1 - 1/k^2$, for any $i = 0, \dots, n$ and any word w , $|u_i^{-1}P_{S_k}(w\Sigma^*) - u_i^{-1}P(w\Sigma^*)| \leq [k^{1/3}(n+1)R]^{-1}$ and therefore, $|u_0^{-1}P_{S_k}(w\Sigma^*) - \sum_{i=1}^n \alpha_i u_i^{-1}P_{S_k}(w\Sigma^*)| \leq k^{-1/3}$.

For any integer $k \geq K_1$, let A_k be the event: $|u_0^{-1}P_{S_k}(w\Sigma^*) - \sum_{i=1}^n \alpha_i u_i^{-1}P_{S_k}(w\Sigma^*)| > k^{-1/3}$. Since $\text{Pr}(A_k) < 1/k^2$, the probability that a finite number of A_k occurs is 1.

Therefore, with probability 1, there exists an integer K such that for any $k \geq K$, $I(\{u_1, \dots, u_n\}, u_0, S_k, k^{-1/3})$ has a solution. □

Lemma 4. *Let $P \in \mathcal{S}(\Sigma)$, let $u_0, u_1, \dots, u_n \in \text{res}(P)$ such that $\{u_1^{-1}P, \dots, u_n^{-1}P\}$ is linearly independent and let $\alpha_1, \dots, \alpha_n \in \mathbb{R}$ be such that $u_0^{-1}P = \sum_{i=1}^n \alpha_i u_i^{-1}P$. Then, with probability one, for any complete presentation S of P , there exists an integer K such that $\forall k \geq K$, any solution $\widehat{\alpha}_1, \dots, \widehat{\alpha}_n$ of $I(\{u_1, \dots, u_n\}, u_0, S_k, k^{-1/3})$ satisfies $|\widehat{\alpha}_i - \alpha_i| < O(k^{-1/3})$ for $1 \leq i \leq n$.*

Proof. Let $w_1, \dots, w_n \in \Sigma^*$ be such that the square matrix M defined by $M[i, j] = u_j^{-1}P(w_i\Sigma^*)$ for $1 \leq i, j \leq n$ is invertible. Let $A = (\alpha_1, \dots, \alpha_n)^t$, $U_0 = (u_0^{-1}P(w_1\Sigma^*), \dots, u_0^{-1}P(w_n\Sigma^*))^t$. We have $MA = U_0$. Let S be a complete presentation of P , let $k \in \mathbb{N}$ and let $\widehat{\alpha}_1, \dots, \widehat{\alpha}_n$ be a solution of $I(\{u_1, \dots, u_n\}, u_0, S_k, k^{-1/3})$. Let M_k be the square matrix defined by $M_k[i, j] = u_j^{-1}P_{S_k}(w_i\Sigma^*)$ for $1 \leq i, j \leq n$, let $A_k = (\widehat{\alpha}_1, \dots, \widehat{\alpha}_n)^t$ and $U_{0,k} = (u_0^{-1}P_{S_k}(w_1\Sigma^*), \dots, u_0^{-1}P_{S_k}(w_n\Sigma^*))^t$. We have

$$\|M_k A_k - U_{0,k}\|^2 = \sum_{i=1}^n [u_0^{-1}P_{S_k}(w_i\Sigma^*) - \sum_{j=1}^n \widehat{\alpha}_j u_j^{-1}P_{S_k}(w_i\Sigma^*)]^2 \leq nk^{-2/3}.$$

Check that

$$A - A_k = M^{-1}(MA - U_0 + U_0 - U_{0,k} + U_{0,k} - M_k A_k + M_k A_k - MA_k)$$

and therefore, for any $1 \leq i \leq n$

$$|\alpha_i - \widehat{\alpha}_i| \leq \|A - A_k\| \leq \|M^{-1}\|(\|U_0 - U_{0,k}\| + n^{1/2}k^{-1/3} + \|M_k - M\| \|A_k\|).$$

Now, by using Lemma 2 and Borel-Cantelli Lemma as in the proof of Lemma 3, with probability 1, there exists K such that for all $k \geq K$, $\|U_0 - U_{0,k}\| < O(k^{-1/3})$ and $\|M_k - M\| < O(k^{-1/3})$. Therefore, for all $k \geq K$, any solution $\widehat{\alpha}_1, \dots, \widehat{\alpha}_n$ of $I(\{u_1, \dots, u_n\}, u_0, S_k, k^{-1/3})$ satisfies $|\widehat{\alpha}_i - \alpha_i| < O(k^{-1/3})$ for $1 \leq i \leq n$. \square

Theorem 1. *Let $P \in \mathcal{S}_{\mathbb{R}}^{rat}(\Sigma)$ and A be the prefixial reduced representation of P . Then, with probability one, for any complete presentation S of P , there exists an integer K such that for any $k \geq K$, $DEES(S_k)$ returns a multiplicity automaton A_k whose support is the same as A 's. Moreover, there exists a constant C such that for any parameter α of A , the corresponding parameter α_k in A_k satisfies $|\alpha - \alpha_k| \leq Ck^{-1/3}$.*

Proof. Let Q_P be the set of states of A , i.e. the smallest prefixial subset of $res(P)$ such that $\{u^{-1}P : u \in Q_P\}$ spans the same vector space as $Res(P)$. Let $u \in Q_P$, let $Q_u = \{v \in Q_P | v < u\}$ and let $x \in \Sigma$.

- If $\{v^{-1}P | v \in Q_u \cup \{ux\}\}$ is linearly independent, from Lemma 1, with probability 1, there exists ϵ_{ux} and K_{ux} such that for any $k \geq K_{ux}$, $I(Q_u, ux, S_k, \epsilon_{ux})$ has no solution.
- If there exists $(\alpha_v)_{v \in Q_u}$ such that $(ux)^{-1}P = \sum_{v \in Q_u} \alpha_v v^{-1}P$, from Lemma 3, with probability 1, there exists an integer K_{ux} such that for any $k \geq K_{ux}$, $I(Q_u, ux, S_k, k^{-1/3})$ has a solution.

Therefore, with probability one, there exists an integer K such that for any $k \geq K$, $DEES(S_k)$ returns a multiplicity automaton A_k whose set of states is equal to Q_P . Use Lemmas 2 and 4 to check the last part of the proposition. \square

When the target is in $\mathcal{S}_{\mathbb{Q}}^{rat}(\Sigma)$, DEES can be used to exactly identify it. The proof is based on the representation of real numbers by continuous fraction. See [8] for a survey on continuous fraction and [6] for a similar application.

Let (ϵ_n) be a sequence of non negative real numbers which converges to 0, let $x \in \mathbb{Q}$, let (y_n) be a sequence of elements of \mathbb{Q} such that $|x - y_n| \leq \epsilon_n$ for all but finitely many n . It can be shown that there exists an integer N such that, for any $n \geq N$, x is the unique rational number $\frac{p}{q}$ which satisfies $|y_n - \frac{p}{q}| \leq \epsilon_n \leq \frac{1}{q^2}$. Moreover, the unique solution of these inequations can be computed from y_n .

Let $P \in \mathcal{S}_{\mathbb{Q}}^{rat}(\Sigma)$, let S be a complete presentation of P and let A_k the MA output by DEES on input S_k . Let \overline{A}_k be the MA derived from A_k by replacing every parameter α_k with a solution $\frac{p}{q}$ of $|\alpha - \frac{p}{q}| \leq k^{-1/4} \leq \frac{1}{q^2}$.

Theorem 2. *Let $P \in \mathcal{S}_{\mathbb{Q}}^{rat}(\Sigma)$ and A be the prefixial reduced representation of P . Then, with probability one, for any complete presentation S of P , there exists an integer K such that $\forall k \geq K$, $DEES(S_k)$ returns an MA A_k such that $\overline{A}_k = A$.*

Proof. From previous theorem, for every parameter α of A , the corresponding parameter α_k in A_k satisfies $|\alpha - \alpha_k| \leq Ck^{-1/3}$ for some constant C . Therefore, if k is sufficiently large, we have $|\alpha - \alpha_k| \leq k^{-1/4}$ and there exists an integer K such that $\alpha = p/q$ is the unique solution of $|\alpha - \frac{p}{q}| \leq k^{-1/4} \leq \frac{1}{q^2}$. \square

4 Learning rational stochastic languages

We have seen that $\mathcal{S}_{\mathbb{Q}}^{rat}(\Sigma)$ is identifiable in the limit. Moreover, DEES runs in polynomial time and aims at computing a representation of the target which is minimal and whose parameters depends only on the target to be learned. DEES computes estimates which are proved to converge reasonably fast to these parameters. That is, DEES compute functions which are likely to be close to the target. But these functions are not stochastic languages and it remains to study how they can be used in a grammatical inference perspective.

Any rational stochastic language P defines a vector subspace of $\mathbb{R}\langle\langle\Sigma\rangle\rangle$ in which the stochastic languages form a compact convex subset.

Proposition 2. *Let p_1, \dots, p_n be n independent stochastic languages. Then, $\Lambda = \{\vec{\alpha} = (\alpha_1, \dots, \alpha_n) \in \mathbb{R}^n : \sum_{i=1}^n \alpha_i p_i \in \mathcal{S}(\Sigma)\}$ is a compact convex subset of \mathbb{R}^n .*

Proof. First, check that for any $\vec{\alpha}, \vec{\beta} \in \Lambda$ and any $\gamma \in [0, 1]$, the series $\sum_{i=1}^n [\gamma \alpha_i + (1 - \gamma) \beta_i] p_i$ is a stochastic language. Hence, Λ is convex.

For every word w , the mapping $\vec{\alpha} \rightarrow \sum_{i=1}^n \alpha_i p_i(w)$ defined from \mathbb{R}^n into \mathbb{R} is linear; and so is the mapping $\vec{\alpha} \rightarrow \sum_{i=1}^n \alpha_i$. Λ is closed since these mappings are continuous and since

$$\Lambda = \left\{ \vec{\alpha} \in \mathbb{R}^n : \sum_{i=1}^n \alpha_i p_i(w) \geq 0 \text{ for every word } w \text{ and } \sum_{i=1}^n \alpha_i = 1 \right\}.$$

Now, let us show that Λ is bounded. Suppose that for any integer k , there exists $\vec{\alpha}_k \in \Lambda$ such that $\|\vec{\alpha}_k\| \geq k$. Since $\vec{\alpha}_k / \|\vec{\alpha}_k\|$ belongs to the unit sphere in \mathbb{R}^n , which is compact, there exists a subsequence $\vec{\alpha}_{\phi(k)}$ such that $\vec{\alpha}_{\phi(k)} / \|\vec{\alpha}_{\phi(k)}\|$ converges to some $\vec{\alpha}$ satisfying $\|\vec{\alpha}\| = 1$. Let $q_k = \sum_{i=1}^n \alpha_{i,k} p_i$ and $r = \sum_{i=1}^n \alpha_i p_i$.

For any $0 < \lambda \leq \|\vec{\alpha}_k\|$, $p_1 + \lambda \frac{q_k - p_1}{\|\vec{\alpha}_k\|} = (1 - \frac{\lambda}{\|\vec{\alpha}_k\|}) p_1 + \frac{\lambda}{\|\vec{\alpha}_k\|} q_k$ is a stochastic language since $\mathcal{S}(\Sigma)$ is convex; for every $\lambda > 0$, $p_1 + \lambda \frac{q_{\phi(k)} - p_1}{\|\vec{\alpha}_{\phi(k)}\|}$ converges to $p_1 + \lambda r$ when $l \rightarrow \infty$, which is a stochastic language since Λ is closed. Therefore, for any $\lambda > 0$, $p_1 + \lambda r$ is a stochastic language. Since $p_1(w) + \lambda r(w) \in [0, 1]$ for every word w , we must have $r = 0$, i.e. $\alpha_i = 0$ for any $1 \leq i \leq n$ since the languages p_1, \dots, p_n are independent, which is impossible since $\|\vec{\alpha}\| = 1$. Therefore, Λ is bounded. \square

The MA A output by DEES generally do not compute stochastic languages. However, we wish that the series r_A they compute share some properties with them. Next proposition gives sufficient conditions which guaranty that $\sum_{k \geq 0} r_A(\Sigma^k) = 1$.

Proposition 3. *Let $A = \langle \Sigma, Q = \{q_1, \dots, q_n\}, \varphi, \iota, \tau \rangle$ be an MA and let M be the square matrix defined by $M[i, j] = [\varphi(q_i, \Sigma, q_j)]_{1 \leq i, j \leq n}$. Suppose that the spectral radius of M satisfies $\rho(M) < 1$. Let $\vec{\iota} = (\iota(q_1), \dots, \iota(q_n))$ and $\vec{\tau} = (\tau(q_1), \dots, \tau(q_n))^t$.*

1. *Then, the matrix $(I - M)$ is invertible and $\sum_{k \geq 0} M^k$ converges to $(I - M)^{-1}$.*
2. *$\forall q_i \in Q, \forall K \geq 0, \sum_{k \geq K} r_{A, q_i}(\Sigma^k)$ converges to $M^K \sum_{j=1}^n (I - M)^{-1}[i, j] \tau(q_j)$ and $\sum_{k \geq K} r_A(\Sigma^k)$ converges to $\vec{\tau} M^K (I - M)^{-1} \vec{\iota}$.*
3. *If $\forall q \in Q, \tau(q) + \varphi(q, \Sigma, Q) = 1$, then $\forall q \in Q, r_{A, q}(\sum_{k \geq 0} \Sigma^k) = 1$. If moreover $\sum_{q \in Q} \iota(q) = 1$, then $r(\sum_{k \geq 0} \Sigma^k) = 1$.*

- Proof.* 1. Since $\rho(M) < 1$, 1 is not an eigen value of M and $I - M$ is invertible. From Gelfand's formula, $\lim_{k \rightarrow \infty} \|M^k\| = 0$. Since for any integer k , $(I - M)(I + M + \dots + M^k) = I - M^{k+1}$, the sum $\sum_{k \geq 0} M^k$ converges to $(I - M)^{-1}$.
2. Since $r_{A,q_i}(\Sigma^k) = \sum_{j=1}^n M^k[i, j] \tau(q_j)$, $\sum_{k \geq K} r_{A,q_i}(\Sigma^k) = M^K \sum_{j=1}^n (I - M)^{-1}[i, j] \tau(q_j)$ and $\sum_{k \geq K} r_A(\Sigma^k) = \sum_{i=1}^n \iota(q_i) r_{A,q_i}(\Sigma^{\geq K}) = \vec{\iota} M^K (I - M)^{-1} \vec{\tau}$.
3. Let $s_i = r_{A,q_i}(\Sigma^*)$ for $1 \leq i \leq n$ and $\vec{s} = (s_1, \dots, s_n)^t$. We have $(I - M) \vec{s} = \vec{\tau}$. Since $I - M$ is invertible, there exists one and only one s such that $(I - M) \vec{s} = \vec{\tau}$. But since $\tau(q) + \varphi(q, \Sigma, Q) = 1$ for any state q , the vector $(1, \dots, 1)^t$ is clearly a solution. Therefore, $s_i = 1$ for $1 \leq i \leq n$. If $\sum_{q \in Q} \iota(q) = 1$, then $r(\Sigma^*) = \sum_{q \in Q} \iota(q) r_{A,q}(\Sigma^*) = 1$. \square

Proposition 4. Let $A = \langle \Sigma, Q, \varphi, \iota, \tau \rangle$ be a reduced representation of a stochastic language P . Let $Q = \{q_1, \dots, q_n\}$ and let M be the square matrix defined by $M[i, j] = [\varphi(q_i, \Sigma, q_j)]_{1 \leq i, j \leq n}$. Then the spectral radius of M satisfies $\rho(M) < 1$.

Proof. From Prop. 2, let R be such that $\{\vec{\alpha} \in \mathbb{R}^n : \sum_{i=1}^n \alpha_i P_{A,q_i} \in \mathcal{S}(\Sigma)\} \subseteq B(0, R)$. For every $u \in \text{res}(P_A)$ and every $1 \leq i \leq n$, we have

$$u^{-1} P_{A,q_i} = \frac{\sum_{1 \leq j \leq n} \varphi(q_i, u, q_j) P_{A,q_j}}{P_{A,q_i}(u \Sigma^*)}.$$

Therefore, for every word u and every k , we have $|\varphi(q_i, u, q_j)| \leq R \cdot P_{A,q_i}(u \Sigma^*)$ and

$$|\varphi(q_i, \Sigma^k, q_j)| \leq \sum_{u \in \Sigma^k} |\varphi(q_i, u, q_j)| \leq R \cdot P_{A,q_i}(\Sigma^{\geq k}).$$

Now, let λ be an eigen value of M associated with the eigen vector v and let i be an index such that $|v_i| = \text{Max}\{|v_j| : j = 1, \dots, n\}$. For every integer k , we have

$$M^k v = \lambda^k v \text{ and } |\lambda^k v_i| = \left| \sum_{j=1}^n \varphi(q_i, \Sigma^k, q_j) v_j \right| \leq n R \cdot P_{A,q_i}(\Sigma^{\geq k}) |v_i|$$

which implies that $|\lambda| < 1$ since $P_{A,q_i}(\Sigma^{\geq k})$ converges to 0 when $k \rightarrow \infty$. \square

If the spectral radius of a matrix is < 1 , the power of M decrease exponentially fast.

Lemma 5. Let $M \in \mathbb{R}^{n \times n}$ be such that $\rho(M) < 1$. Then, there exists $C \in \mathbb{R}$ and $\rho \in [0, 1[$ such that for any integer $k \geq 0$, $\|M^k\| \leq C \rho^k$.

Proof. Let $\rho \in]\rho(M), 1[$. From Gelfand's formula, there exists an integer K such that for any $k \geq K$, $\|M^k\|^{1/k} \leq \rho$. Let $C = \text{Max}\{\|M^h\| / \rho^h : h < K\}$. Let $k \in \mathbb{N}$ and let $a, b \in \mathbb{N}$ be such that $k = aK + b$ and $b < K$. We have

$$\|M^k\| = \|M^{aK+b}\| \leq \|M^{aK}\| \|M^b\| \leq \rho^{aK} \|M^b\| \leq \rho^k \frac{\|M^b\|}{\rho^b} \leq C \rho^k.$$

Proposition 5. Let $P \in \mathcal{S}_{\mathbb{R}}^{\text{rat}}(\Sigma)$. There exists a constant C and $\rho \in [0, 1[$ such that for any integer k , $P(\Sigma^{\geq k}) \leq C \rho^k$.

Proof. Let $A = \langle \Sigma, Q, \varphi, \iota, \tau \rangle$ be a reduced representation of P and let M be the square matrix defined by $M[i, j] = [\varphi(q_i, \Sigma, q_j)]_{1 \leq i, j \leq n}$. From Prop. 4, the spectral radius of M is ≤ 1 . From Lemma 5, there exists C_1 and $\rho \in [0, 1[$ such that $\|M^k\| \leq C_1 \rho^k$ for every integer k . Let $\vec{\iota}_A = (\iota(q_1), \dots, \iota(q_n))$ and $\vec{\tau}_A = (\tau(q_1), \dots, \tau(q_n))^t$. We have

$$P(\Sigma^{\geq k}) \leq \|\iota_A\| \cdot \|M^k\| \cdot \|(I - M)^{-1}\| \cdot \|\vec{\tau}_A\| \leq C \rho^k$$

with $C = C_1 \|\vec{\iota}_A\| \cdot \|(I - M)^{-1}\| \cdot \|\vec{\tau}_A\|$. \square

It is not difficult to design an MA A which generates a stochastic language P and such that $\varphi(q, u, q')$ is unbounded when $u \in \Sigma^*$. However, the next proposition proves that this situation never happens when A is a reduced representation of P .

Proposition 6. *Let $P \in \mathcal{S}_{\mathbb{R}}^{rat}(\Sigma)$ and let $A = \langle \Sigma, Q, \varphi, \iota, \tau \rangle$ be a reduced representation of P . Then, there exists a constant C and $\rho \in [0, 1[$ such that for any integer k and any pair of states q, q' , $\sum_{u \in \Sigma^k} |\varphi(q, u, q')| \leq C \rho^k$.*

Proof. Let k be an integer and let $q, q' \in Q$. Let $P_k = \{u \in \Sigma^k : \varphi(q, u, q') \geq 0\}$ and $N_k = \Sigma^k \setminus P_k$.

$$P_k^{-1} P_{A,q} = \sum_{u \in P_k} \frac{P_{A,q}(u \Sigma^*)}{\sum_{u \in P_k} P_{A,q}(u \Sigma^*)} u^{-1} P_{A,q} = \sum_{q'' \in Q} \frac{\sum_{u \in P_k} \varphi(q, u, q'')}{\sum_{u \in P_k} P_{A,q}(u \Sigma^*)} P_{A,q''}$$

is a stochastic language which is a linear combination of the independent stochastic languages $P_{A,q''}$. From prop. 2, there exists a constant R which depends only on A s.t.

$$\left| \sum_{u \in P_k} \varphi(q, u, q') \right| = \sum_{u \in P_k} \varphi(q, u, q') \leq R \sum_{u \in P_k} P_{A,q}(u \Sigma^*).$$

Similarly, we have $|\sum_{u \in N_k} \varphi(q, u, q')| = \sum_{u \in N_k} |\varphi(q, u, q')| \leq R \sum_{u \in N_k} P_{A,q}(u \Sigma^*)$. Let C and $\rho \in]0, 1[$ be such that $P_{A,q}(\Sigma^{\geq k}) \leq C \rho^k$ for any state q and any integer k . We have

$$\sum_{u \in \Sigma^k} |\varphi(q, u, q')| \leq R \sum_{u \in \Sigma^k} P_{A,q}(u \Sigma^*) \leq RC \rho^k.$$

\square

MA representation of rational stochastic languages are unstable (see Fig. 3). Arbitrarily close to an MA A which generates a stochastic language, we can find an MA B such that the sum $\sum_{w \in \Sigma^*} r_B(w)$ converges to any real number or even diverges. However, the next theorem shows that when A is a reduced representation of a stochastic language, any MA B sufficiently close to A defines a series which is absolutely convergent. Moreover, simple syntactical conditions ensure that $r_B(\Sigma^*) = 1$.

Theorem 3. *Let $P \in \mathcal{S}_{\mathbb{R}}^{rat}(\Sigma)$ and let $A = \langle \Sigma, Q, \varphi_A, \iota_A, \tau_A \rangle$ be a reduced representation of P . Let C_A and $\rho_A \in]0, 1[$ be such that for any integer k and any pair of states q, q' , $\sum_{u \in \Sigma^k} |\varphi_A(q, u, q')| \leq C_A \rho_A^k$. Then, for any $\rho > \rho_A$, there exists C and $\alpha > 0$ such that for any MA $B = \langle \Sigma, Q, \varphi_B, \iota_B, \tau_B \rangle$ satisfying*

$$\forall q, q' \in Q, \forall x \in \Sigma, |\varphi_A(q, x, q') - \varphi_B(q, x, q')| < \alpha \quad (2)$$




Figure 3. These MA compute a series r_ϵ such that $\sum_{w \in \Sigma^*} r_\epsilon(w) = 1$ if $\epsilon \neq 0$ and $\sum_{w \in \Sigma^*} r_0(w) = 2/5$. Note that when $\epsilon = 0$, the series r_{0,q_1} and r_{0,q_2} are dependent.

we have $\sum_{u \in \Sigma^k} |\varphi_B(q, u, q')| \leq C\rho^k$ for any pair of states q, q' and any integer k . As a consequence, the series r_B is absolutely convergent. Moreover, if B satisfies also

$$\forall q \in Q, \tau_B(q) + \varphi_B(q, \Sigma, Q) = 1 \text{ and } \sum_{q \in Q} \iota_B(q) = 1 \quad (3)$$

then, α can be chosen such that (2) implies that $r_{B,q}(\Sigma^*) = 1$ for any state q and $r_B(\Sigma^*) = 1$.

Proof. Let k be such that $(2nC_A)^{1/k} \leq \rho/\rho_A$ where $n = |Q|$. There exists $\alpha > 0$ such that for any MA $B = \langle \Sigma, Q, \varphi_B, \iota_B, \tau_B \rangle$ satisfying (2), we have

$$\forall q, q' \in Q, \sum_{u \in \Sigma^k} |\varphi_B(q, u, q') - \varphi_A(q, u, q')| < C_A \rho_A^k.$$

Since $\sum_{u \in \Sigma^k} |\varphi_A(q, u, q')| \leq C_A \rho_A^k$, we must have also

$$\sum_{u \in \Sigma^k} |\varphi_B(q, u, q')| \leq 2C_A \rho_A^k \leq \frac{\rho^k}{n}.$$

Let $C_1 = \text{Max}\{\sum_{u \in \Sigma^{<k}} |\varphi_B(q, u, q')| : q, q' \in Q\}$. Let $l, a, b \in \mathbb{N}$ such that $l = ak + b$ and $b < k$. Let $u \in \Sigma^l$ and let $u = u_0 \dots u_a$ where $|u_i| = k$ for $0 \leq i < a$ and $|u_a| = b$. For any pair of states q_0, q_{a+1} , we have

$$\varphi_B(q_0, u, q_{a+1}) = \sum_{q_1, \dots, q_a \in Q} \prod_{i=0}^a \varphi_B(q_i, u_i, q_{i+1})$$

and

$$\begin{aligned} \sum_{u \in \Sigma^l} \varphi_B(q_0, u, q_{a+1}) &= \sum_{u_0, \dots, u_{a-1} \in \Sigma^k} \sum_{u_a \in \Sigma^b} \sum_{q_1, \dots, q_a \in Q} \prod_{i=0}^a \varphi_B(q_i, u_i, q_{i+1}) \\ &= \sum_{q_1, \dots, q_a \in Q} \sum_{u_0, \dots, u_{a-1} \in \Sigma^k} \sum_{u_a \in \Sigma^b} \prod_{i=0}^a \varphi_B(q_i, u_i, q_{i+1}) \\ &= \sum_{q_1, \dots, q_a \in Q} \prod_{i=0}^{a-1} \left(\sum_{u \in \Sigma^k} \varphi_B(q_i, u, q_{i+1}) \right) \left(\sum_{u \in \Sigma^b} \varphi_B(q_a, u, q_{a+1}) \right). \end{aligned}$$

Hence, $\sum_{u \in \Sigma^l} |\varphi_B(q_0, u, q_{m+1})| \leq n^a \cdot \left(\frac{\rho^k}{n}\right)^a \cdot C_1 \leq C\rho^l$ where $C = \frac{C_1}{\rho^{k-1}}$.

Now, let us prove that r_B is absolutely convergent.

$$\sum_{w \in \Sigma^*} |r_B(w)| \leq \sum_{k \in \mathbb{N}} \sum_{u \in \Sigma^k} \sum_{q, q' \in Q} \iota_B(q) \varphi_B(q, u, q') \tau_B(q') \leq C'$$

where $C' = Cn^2 \text{Max}\{|\iota_B(q)\tau_B(q')| : q, q' \in Q\}/(1 - \rho)$.

Lastly, let M_B be the square matrix defined by $M_B[i, j] = \varphi_B(q_i, \Sigma, q_j)$. Since the spectral radius of a matrix depends continuously on its coefficients and since A is a reduced representation of a stochastic language, any MA satisfying (2) with α sufficiently small must have a spectral radius $\neq 1$ (Prop. 4). Therefore, if B satisfies (3) and (2) with α sufficiently small, the Prop. 3 entails the conclusion. \square

It remains to show how a series which converges absolutely to 1 can be used to approximate a stochastic language.

Let r be a series over Σ such that $\sum_{w \in \Sigma^*} r(w)$ converges absolutely to 1. Therefore, $r(X) = \sum_{u \in X} r(u)$ is defined without ambiguity for every $X \subseteq \Sigma^*$ and $r(X)$ is bounded by $\bar{r} = \sum_{u \in \Sigma^*} |r(u)|$. Let S be the smallest subset of Σ^* such that

$$\varepsilon \in S \text{ and } \forall u \in \Sigma^*, \forall x \in \Sigma, u \in S \text{ and } r(ux\Sigma^*) > 0 \Rightarrow ux \in S.$$

S is a prefixial subset of Σ^* and $\forall u \in S, r(u\Sigma^*) > 0$. For every word $u \in S$, let us define $N(u) = \cup\{ux\Sigma^* : x \in \Sigma, r(ux\Sigma^*) \leq 0\} \cup \{u : \text{if } r(u) \leq 0\}$ and $N = \cup\{N(u) : u \in \Sigma^*\}$. Then, for every $u \in S$, let us define λ_u by:

$$\lambda_\varepsilon = (1 - r(N(\varepsilon)))^{-1} \text{ and } \lambda_{ux} = \lambda_u \frac{r(ux\Sigma^*)}{r(ux\Sigma^*) - r(N(ux))}.$$

Lemma 6. For every word $u \in S$, $e^{r(N)}/\bar{r} \leq \lambda_u \leq 1$.

Proof. First, check that $r(N(u)) \leq 0$ for every $u \in S$. Therefore, $\lambda_u \leq 1$. Now, check that if $u, uv \in S$ then $v = \varepsilon$ or $N(u) \cap N(uv) = \emptyset$. Let $u = x_1 \dots x_n \in \Sigma^*$ where $x_1, \dots, x_n \in \Sigma$ and let $u_0 = \varepsilon$ and $u_i = u_{i-1}x_i$ for $1 \leq i \leq n$. We have

$$\lambda_u = \prod_{i=0}^n \frac{r(u_i\Sigma^*)}{r(u_i\Sigma^*) - r(N(u_i))} = \prod_{i=0}^n \left(1 - \frac{r(N(u_i))}{r(u_i\Sigma^*)}\right)^{-1}$$

and

$$\log \lambda_u = - \sum_{i=0}^n \log \left(1 - \frac{r(N(u_i))}{r(u_i\Sigma^*)}\right) \geq \sum_{i=0}^n \frac{r(N(u_i))}{r(u_i\Sigma^*)}.$$

Since $r(u_i\Sigma^*) \leq \bar{r}$, $\log \lambda_u \geq \sum_{i=0}^n r(N(u_i))/\bar{r} = r(\cup_{i=0}^n N(u_i))/\bar{r} \geq r(N)/\bar{r}$. Therefore, $\lambda_u \geq e^{r(N)/\bar{r}}$. \square

Let p_r be the series defined by: $p_r(u) = 0$ if $u \in N$ and $p_r(u) = \lambda_u r(u)$ otherwise. We show that p_r is a stochastic language.

Lemma 7. $p_r(\varepsilon) + \lambda_\varepsilon \sum_{x \in S \cap \Sigma} r(x\Sigma^*) = 1$,

– For any $u \in \Sigma^*$ and any $x \in \Sigma$, if $ux \in S$ then

$$p_r(ux) + \lambda_{ux} \sum_{\{y \in \Sigma: uxy \in S\}} r(uxy\Sigma^*) = \lambda_u r(ux\Sigma^*).$$

Proof. First, check that for every $u \in S$,

$$p_r(u) + \lambda_u \sum_{x \in u^{-1}S \cap \Sigma} r(ux\Sigma^*) = \lambda_u (r(u\Sigma^*) - r(N(u))).$$

Then, $p_r(\varepsilon) + \lambda_\varepsilon \sum_{x \in S \cap \Sigma} r(x\Sigma^*) = \lambda_\varepsilon (1 - r(N(\varepsilon))) = 1$. Now, let $u \in \Sigma^*$ and $x \in \Sigma$ s.t. $ux \in S$, $p_r(ux) + \lambda_{ux} \sum_{\{y \in \Sigma: uxy \in S\}} r(uxy\Sigma^*) = \lambda_{ux} (r(ux\Sigma^*) - r(N(ux))) = \lambda_u r(ux\Sigma^*)$. \square

Lemma 8. Let Q be a prefixial finite subset of Σ^* and let $Q_s = (Q\Sigma \setminus Q) \cap S$. Then

$$p_r(Q) = 1 - \sum_{ux \in Q_s, x \in \Sigma} \lambda_u r(ux\Sigma^*).$$

Proof. By induction on Q . When $Q = \{\varepsilon\}$, the relation comes directly from Lemma 7. Now, suppose that the relation is true for a prefixial subset Q' , let $u_0 \in Q'$ and $x_0 \in \Sigma$ such that $u_0x_0 \notin Q'$ and let $Q = Q' \cup \{u_0x_0\}$. We have

$$p_r(Q) = p_r(Q') + p_r(u_0x_0) = 1 - \sum_{ux \in Q'_s, x \in \Sigma} \lambda_u r(ux\Sigma^*) + p_r(u_0x_0)$$

where $Q'_s = (Q'\Sigma \setminus Q') \cap S$, from inductive hypothesis.

If $u_0x_0 \notin S$, check that $p_r(u_0x_0) = 0$ and that $Q_s = Q'_s$. Therefore, $p_r(Q) = 1 - \sum_{ux \in Q'_s, x \in \Sigma} \lambda_u r(ux\Sigma^*)$.

If $u_0x_0 \in S$, then $Q_s = Q'_s \setminus \{u_0x_0\} \cup (u_0x_0\Sigma \cap S)$. Therefore,

$$\begin{aligned} p_r(Q) &= 1 - \sum_{ux \in Q'_s, x \in \Sigma} \lambda_u r(ux\Sigma^*) + p_r(u_0x_0) \\ &= 1 - \sum_{ux \in Q_s, x \in \Sigma} \lambda_u r(ux\Sigma^*) - \lambda_{u_0} r(u_0x_0\Sigma^*) \\ &\quad + \lambda_{u_0x_0} \sum_{u_0x_0x \in S, x \in \Sigma} r(u_0x_0x\Sigma^*) + p_r(u_0x_0) \\ &= 1 - \sum_{ux \in Q_s, x \in \Sigma} \lambda_u r(ux\Sigma^*) \text{ from Lemma 7.} \end{aligned} \quad \square$$

Proposition 7. Let r be a formal series over Σ such that $\sum_{w \in \Sigma^*} r(w)$ converges absolutely to 1. Then, p_r is a stochastic language such that for every $u \in \Sigma^* \setminus N$,

$$(1 + r(N)/\bar{r})r(u) \leq e^{r(N)/\bar{r}}r(u) \leq p_r(u) \leq r(u).$$

Proof. From Lemma 6, the only thing that remains to be proved is that p_r is a stochastic language. Clearly, $p_r(u) \in [0, 1]$ for every word u . From Lemma 8, for any integer k ,

$$|1 - p_r(\Sigma^{\leq k})| \leq \sum_{u \in \Sigma^{k+1} \cap S} r(u\Sigma^*) \leq r(\Sigma^{> k})$$

which tends to 0 since r is absolutely convergent. \square

To sum up, DEES computes MA A whose structure is equal to the structure of the target from some steps, and whose parameters tends reasonably fast to the true parameters. From some steps, they define absolutely rational series r_A which converge absolutely to 1. By using these MA, it is possible to efficiently compute $p_{r_A}(u)$ or $p_{r_A}(u\Sigma^*)$ for any word u . Moreover, since r_A converges absolutely and since A tends to the target, the weight $r_A(N)$ of the negative values tends to 0 and p_{r_A} converges to the target.

5 Conclusion

We have defined an inference algorithm DEES designed to learn rational stochastic languages which strictly contains the class of stochastic languages computable by PA (or HMM). We have shown that the class of rational stochastic languages over \mathbb{Q} is strongly identifiable in the limit. Moreover, DEES is an efficient inference algorithm which can be used in practical cases of grammatical inference. The experiments we have already carried out confirm the theoretical results of this paper: the fact that DEES aims at building a natural and minimal representation of the target provides a very significant improvement of the results obtained by classical probabilistic inference algorithms.

References

1. Amos Beimel, Francesco Bergadano, Nader H. Bshouty, Eyal Kushilevitz, and Stefano Varricchio. On the applications of multiplicity automata in learning. In *IEEE Symposium on Foundations of Computer Science*, pages 349–358, 1996.
2. Amos Beimel, Francesco Bergadano, Nader H. Bshouty, Eyal Kushilevitz, and Stefano Varricchio. Learning functions represented as multiplicity automata. *Journal of the ACM*, 47(3):506–530, 2000.
3. F. Bergadano and S. Varricchio. Learning behaviors of automata from multiplicity and equivalence queries. In *Italian Conf. on Algorithms and Complexity*, 1994.
4. J. Berstel and C. Reutenauer. *Les séries rationnelles et leurs langages*. Masson, 1984.
5. R.C. Carrasco and J. Oncina. Learning stochastic regular grammars by means of a state merging method. In *ICGI*, pages 139–152, Heidelberg, September 1994. Springer-Verlag.
6. F. Denis and Y. Esposito. Learning classes of probabilistic automata. In *COLT 2004*, number 3120 in LNAI, pages 124–139, 2004.
7. F. Denis and Y. Esposito. Rational stochastic languages. Technical report, LIF - Université de Provence, 2006.
8. G. H. Hardy and E. M. Wright. *An introduction to the theory of numbers*. Oxford University Press, 1979.
9. G. Lugosi. *Principles of Nonparametric Learning*, chapter Pattern classification and learning theory, pages 1–56. Springer, 2002.
10. Jacques Sakarovitch. *Éléments de théorie des automates*. Éditions Vuibert, 2003.
11. Arto Salomaa and M. Soittola. *Automata: Theoretic Aspects of Formal Power Series*. Springer-Verlag, 1978.
12. Franck Thollard, Pierre Dupont, and Colin de la Higuera. Probabilistic DFA inference using Kullback-Leibler divergence and minimality. In *Proc. 17th ICML*. KAUFM, 975–982.
13. V. N. Vapnik. *Statistical Learning Theory*. John Wiley, 1998.