

# A Classification of the Projective Lines over Small Rings

Metod Saniga,<sup>1–3</sup> Michel Planat,<sup>2</sup> Maurice R. Kibler<sup>3</sup>  
and Petr Pracna<sup>2–4</sup>

<sup>1</sup>Astronomical Institute, Slovak Academy of Sciences  
SK-05960 Tatranská Lomnica, Slovak Republic  
(msaniga@astro.sk)

<sup>2</sup>Institut FEMTO-ST, CNRS, Département LPMO, 32 Avenue de l'Observatoire  
F-25044 Besançon Cedex, France  
(planat@lpmo.edu)

<sup>3</sup>Institut de Physique Nucléaire de Lyon, IN2P3-CNRS/Université Claude Bernard Lyon 1  
43 Boulevard du 11 Novembre 1918, F-69622 Villeurbanne Cedex, France  
(kibler@ipnl.in2p3.fr)

<sup>4</sup>J. Heyrovský Institute of Physical Chemistry, Academy of Sciences of the Czech Republic  
Dolejškova 3, CZ-182 23 Prague 8, Czech Republic  
(pracna@jh-inst.cas.cz)

---

## Abstract

A compact classification of the projective lines defined over (commutative) rings (with unity) of all orders up to thirty-one is given. There are altogether sixty-five different types of them. For each type we introduce the total number of points on the line, the number of points represented by coordinates with at least one entry being a unit, the cardinality of the neighbourhood of a generic point of the line as well as those of the intersections between the neighbourhoods of two and three mutually distant points, the number of ‘Jacobson’ points per a neighbourhood, the maximum number of pairwise distant points and, finally, a list of representative/base rings. The classification is presented in form of a table in order to see readily not only the fine traits of the hierarchy, but also the changes in the structure of the lines as one goes from one type to the other. We hope this study will serve as an impetus to a search for possible applications of these remarkable geometries in physics, chemistry, biology and other natural sciences as well.

**Keywords:** Projective Ring Lines – Rings of Small Orders

---

## 1 Introduction

Recently, projective lines defined over rings instead of fields have become the subject of renewed interest not only for their own sake [1]–[4], but also in view of their interesting potential applications in (quantum) physics [5]–[7]. It was the latter fact that motivated our in-depth study of the structure of projective lines over a large number of distinct commutative rings with unity of order up to thirty-one — the study that yields, as far as we know, the first compact classification of these beautiful geometric configurations. The key element of our classification scheme is, of course, the neighbour (or parallel) relation [8],[9], which is a geometrical concept intimately related with the structure of and the connection between the maximal ideals of a ring under consideration. Being simply an identity relation for fields, this concept acquires a highly non-trivial character if the base ring features three and/or more maximal ideals and, as we shall see in detail, endows the corresponding line with a rich and quite involved intrinsic structure.

## 2 The Projective Line over a Ring with Unity

Given a ring  $R$  with unity (see, e.g., [10]–[12] and also [5] or [7] for a brief recollection of basic definitions, concepts and notations of ring theory) and  $GL(2, R)$ , the general linear group of invertible two-by-two

matrices with entries in  $R$ , a pair  $(\alpha, \beta) \in R^2$  is called *admissible* over  $R$  if there exist  $\gamma, \delta \in R$  such that [9]

$$\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \text{GL}(2, R). \quad (1)$$

The projective line over  $R$ , denoted as  $PR(1)$ , is defined as the set of classes of ordered pairs  $(\varrho\alpha, \varrho\beta)$ , where  $\varrho$  is a unit and  $(\alpha, \beta)$  is admissible [1],[2],[4],[9]. Such a line carries two non-trivial, mutually complementary relations of neighbour and distant. In particular, its two distinct points  $X := (\varrho\alpha, \varrho\beta)$  and  $Y := (\varrho\gamma, \varrho\delta)$  are called *neighbour* (or, *parallel*) if

$$\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \notin \text{GL}(2, R) \quad (2)$$

and *distant* otherwise, i. e., if condition (1) is met. The neighbour relation is reflexive (every point is obviously neighbour to itself) and symmetric (i. e., if  $X$  is neighbour to  $Y$  then  $Y$  is neighbour to  $X$  too), but, in general, not transitive (i. e.,  $X$  being neighbour to  $Y$  and  $Y$  being neighbour to  $Z$  does not necessarily mean that  $X$  is neighbour to  $Z$  — see, e. g., [4],[8],[9]). Given a point of  $PR(1)$ , the set of all neighbour points to it will be called its *neighbourhood*.<sup>1</sup> Obviously, if  $R$  is a field then ‘neighbour’ simply reduces to ‘identical’ and ‘distant’ to ‘different’. For a *finite commutative* ring  $R$ , Eq. (1) reads

$$\det \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in R^*, \quad (3)$$

and Eq. (2) reduces to

$$\det \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in R \setminus R^*, \quad (4)$$

where  $R^*$  denotes the set of *units* (invertible elements) and  $R \setminus R^*$  stands for the set of *zero-divisors* of  $R$  (including the trivial zero divisor, 0).

To illustrate the concept of a ring line, we shall examine in detail the structure of the projective line defined over the direct product ring  $R_\diamond \equiv Z_4 \otimes Z_4$ , with  $Z_4$  being the ring of integers modulo 4, i. e., the set  $\{0, 1, 2, 3\}$  endowed with the addition and multiplication properties as shown in Table 1.

Table 1: Addition (*left*) and multiplication (*right*) in  $Z_4$ .

$\oplus$	0	1	2	3
0	0	1	2	3
1	1	2	3	0
2	2	3	0	1
3	3	0	1	2

$\otimes$	0	1	2	3
0	0	0	0	0
1	0	1	2	3
2	0	2	0	2
3	0	3	2	1

The ring  $R_\diamond$  is, like  $Z_4$  itself, of characteristic four, and features the following sixteen elements

$$\begin{aligned} R_\diamond = \{ & a \equiv [0, 0], b \equiv [0, 1], c \equiv [0, 2], d \equiv [0, 3], \\ & e \equiv [1, 0], h \equiv [1, 1], i \equiv [1, 2], j \equiv [1, 3], \\ & f \equiv [2, 0], k \equiv [2, 1], l \equiv [2, 2], m \equiv [2, 3], \\ & g \equiv [3, 0], n \equiv [3, 1], p \equiv [3, 2], q \equiv [3, 3] \}. \end{aligned} \quad (5)$$

It contains two (proper) maximal ideals,

$$\mathcal{I}_1 = \{a, c, f, l, b, d, k, m\}, \quad (6)$$

$$\mathcal{I}_2 = \{a, c, f, l, e, g, i, p\}, \quad (7)$$

yielding a non-trivial Jacobson radical

$$\mathcal{J} = \mathcal{I}_1 \cap \mathcal{I}_2 = \{a, c, f, l\}, \quad (8)$$

as it can readily be ascertained from its addition and multiplication properties (Table 2).

---

<sup>1</sup>To avoid any confusion, the reader should be cautious that some authors (e. g. [1],[4]) use this term for the set of *distant* points instead.

Table 2: Addition (*top*) and multiplication (*bottom*) in  $R_\diamond$ .

$\oplus$	$a$	$b$	$c$	$d$	$e$	$f$	$g$	$h$	$i$	$j$	$k$	$l$	$m$	$n$	$p$	$q$
$a$	$a$	$b$	$c$	$d$	$e$	$f$	$g$	$h$	$i$	$j$	$k$	$l$	$m$	$n$	$p$	$q$
$b$	$b$	$c$	$d$	$a$	$h$	$k$	$n$	$i$	$j$	$e$	$l$	$m$	$f$	$p$	$q$	$g$
$c$	$c$	$d$	$a$	$b$	$i$	$l$	$p$	$j$	$e$	$h$	$m$	$f$	$k$	$q$	$g$	$n$
$d$	$d$	$a$	$b$	$c$	$j$	$m$	$q$	$e$	$h$	$i$	$f$	$k$	$l$	$g$	$n$	$p$
$e$	$e$	$h$	$i$	$j$	$f$	$g$	$a$	$k$	$l$	$m$	$n$	$p$	$q$	$b$	$c$	$d$
$f$	$f$	$k$	$l$	$m$	$g$	$a$	$e$	$n$	$p$	$q$	$b$	$c$	$d$	$h$	$i$	$j$
$g$	$g$	$n$	$p$	$q$	$a$	$e$	$f$	$b$	$c$	$d$	$h$	$i$	$j$	$k$	$l$	$m$
$h$	$h$	$i$	$j$	$e$	$k$	$n$	$b$	$l$	$m$	$f$	$p$	$q$	$g$	$c$	$d$	$a$
$i$	$i$	$j$	$e$	$h$	$l$	$p$	$c$	$m$	$f$	$k$	$q$	$g$	$n$	$d$	$a$	$b$
$j$	$j$	$e$	$h$	$i$	$m$	$q$	$d$	$f$	$k$	$l$	$g$	$n$	$p$	$a$	$b$	$c$
$k$	$k$	$l$	$m$	$f$	$n$	$b$	$h$	$p$	$q$	$g$	$c$	$d$	$a$	$i$	$j$	$e$
$l$	$l$	$m$	$f$	$k$	$p$	$c$	$i$	$q$	$g$	$n$	$d$	$a$	$b$	$j$	$e$	$h$
$m$	$m$	$f$	$k$	$l$	$q$	$d$	$j$	$g$	$n$	$p$	$a$	$b$	$c$	$e$	$h$	$i$
$n$	$n$	$p$	$q$	$g$	$b$	$h$	$k$	$c$	$d$	$a$	$i$	$j$	$e$	$l$	$m$	$f$
$p$	$p$	$q$	$g$	$n$	$c$	$i$	$l$	$d$	$a$	$b$	$j$	$e$	$h$	$m$	$f$	$k$
$q$	$q$	$g$	$n$	$p$	$d$	$j$	$m$	$a$	$b$	$c$	$e$	$h$	$i$	$f$	$k$	$l$

$\otimes$	$a$	$b$	$c$	$d$	$e$	$f$	$g$	$h$	$i$	$j$	$k$	$l$	$m$	$n$	$p$	$q$
$a$	$a$	$a$	$a$	$a$	$a$	$a$	$a$	$a$	$a$	$a$	$a$	$a$	$a$	$a$	$a$	$a$
$b$	$a$	$b$	$c$	$d$	$a$	$a$	$a$	$b$	$c$	$d$	$b$	$c$	$d$	$b$	$c$	$d$
$c$	$a$	$c$	$a$	$c$	$a$	$a$	$a$	$c$	$a$	$c$	$c$	$a$	$c$	$c$	$a$	$c$
$d$	$a$	$d$	$c$	$b$	$a$	$a$	$a$	$d$	$c$	$b$	$d$	$c$	$b$	$d$	$c$	$b$
$e$	$a$	$a$	$a$	$a$	$e$	$f$	$g$	$e$	$e$	$e$	$f$	$f$	$f$	$g$	$g$	$g$
$f$	$a$	$a$	$a$	$a$	$f$	$a$	$f$	$f$	$f$	$f$	$a$	$a$	$a$	$f$	$f$	$f$
$g$	$a$	$a$	$a$	$a$	$g$	$f$	$e$	$g$	$g$	$g$	$f$	$f$	$f$	$e$	$e$	$e$
$h$	$a$	$b$	$c$	$d$	$e$	$f$	$g$	$h$	$i$	$j$	$k$	$l$	$m$	$n$	$p$	$q$
$i$	$a$	$c$	$a$	$c$	$e$	$f$	$g$	$i$	$e$	$i$	$l$	$f$	$l$	$p$	$g$	$p$
$j$	$a$	$d$	$c$	$b$	$e$	$f$	$g$	$j$	$i$	$h$	$m$	$l$	$k$	$q$	$p$	$n$
$k$	$a$	$b$	$c$	$d$	$f$	$a$	$f$	$k$	$l$	$m$	$b$	$c$	$d$	$k$	$l$	$m$
$l$	$a$	$c$	$a$	$c$	$f$	$a$	$f$	$l$	$f$	$l$	$c$	$a$	$c$	$l$	$f$	$l$
$m$	$a$	$d$	$c$	$b$	$f$	$a$	$f$	$m$	$l$	$k$	$d$	$c$	$b$	$m$	$l$	$k$
$n$	$a$	$b$	$c$	$d$	$g$	$f$	$e$	$n$	$p$	$q$	$k$	$l$	$m$	$h$	$i$	$j$
$p$	$a$	$c$	$a$	$c$	$g$	$f$	$e$	$p$	$g$	$p$	$l$	$f$	$l$	$i$	$e$	$i$
$q$	$a$	$d$	$c$	$b$	$g$	$f$	$e$	$q$	$p$	$n$	$m$	$l$	$k$	$j$	$i$	$h$

From these tables it also follows that  $a$  and  $h$  are, respectively, the addition and multiplication identities ('0' and '1') of the ring and that

$$R_\diamond^* = \{h \equiv 1, j, n, q\} \tag{9}$$

and

$$R_\diamond \setminus R_\diamond^* = \{a \equiv 0, b, c, d, e, f, g, i, k, l, m, p\}. \tag{10}$$

Now we can employ the admissibility condition and find out that the projective line  $PR_\diamond(1)$  contains altogether thirty-six points, which can be partitioned into two distinct groups. The first group ('type I points') consists of the points represented by coordinates where at least one entry is a unit and there are the following twenty-eight points here

$$\begin{aligned} &(1, 0), (1, b), (1, c), (1, d), (1, e), (1, f), (1, g), (1, i), (1, k), (1, l), (1, m), (1, p), \\ &(0, 1), (b, 1), (c, 1), (d, 1), (e, 1), (f, 1), (g, 1), (i, 1), (k, 1), (l, 1), (m, 1), (p, 1), \\ &(1, 1), (1, j), (1, n), (1, q); \end{aligned} \tag{11}$$

it is easy to verify that for any finite commutative ring this number is always equal to the sum of the total elements of the ring and the number of its zero-divisors. The other group ('type II points') is composed of

those points whose representing coordinates are both zero-divisors and we find the following eight points ranked here

$$(e, b), (e, k), (i, b), (i, k), (b, e), (k, e), (b, i), (k, i); \quad (12)$$

these points, in general, exist only if the ring has two or more maximal ideals and their number depends on the properties and interconnection between these ideals. To reveal all the subtleties of the structure of the line, one has to make use of the neighbour/distant relation. The reasoning here is, without any loss of generality, much facilitated by considering three distinguished points of the line, viz.  $U := (1, 0)$ ,  $V := (0, 1)$ ,  $W := (1, 1)$ , which are obviously pairwise distant and have the following neighbourhoods

$$U : \quad (1, b), (\underline{1, c}), (1, d), (1, e), (\underline{1, f}), (1, g), (1, i), (1, k), (\underline{1, l}), (1, m), (1, p), \\ (e, b), (e, k), (i, b), (i, k), (b, e), (k, e), (b, i), (k, i) \quad (13)$$

$$V : \quad (b, 1), (\underline{c, 1}), (d, 1), (e, 1), (\underline{f, 1}), (g, 1), (i, 1), (k, 1), (\underline{l, 1}), (m, 1), (p, 1), \\ (e, b), (e, k), (i, b), (i, k), (b, e), (k, e), (b, i), (k, i) \quad (14)$$

$$W : \quad (1, b), (1, d), (1, e), (1, g), (1, i), (1, k), (1, m), (1, p), \\ (b, 1), (d, 1), (e, 1), (g, 1), (i, 1), (k, 1), (m, 1), (p, 1), \\ (\underline{1, j}), (\underline{1, n}), (\underline{1, q}). \quad (15)$$

We see that each neighbourhood features nineteen points and has three ‘Jacobson’ points (underlined), i.e., the points unique to the particular neighbourhood, that the neighbourhoods pairwise overlap in eight points and have no common element if considered altogether; moreover, one easily checks that there exists no point of the line that would be simultaneously distant to all the three distinguished points. Now, as the coordinate system on this line can *always* be chosen in such a way that the coordinates of *any* three mutually distant points are made identical to those of  $U$ ,  $V$  and  $W$ , we can conclude that the neighbourhood of any point of the line features nineteen distinct points, the neighbourhoods of any two distant points share eight points, the neighbourhoods of any three mutually distant points have no point in common and three is the maximum number of mutually distant points; a nice ‘conic’ representation of the line exhibiting all these properties is given in Fig. 1.




Figure 1: A schematic sketch of the structure of the projective line  $PR_{\diamond}(1)$ . The three double circles stand for three pairwise distant points, whilst the remaining points of the line are all located on the neighbourhoods of these three points (three sets of points located on three different ellipses centered at the points in question). Every bullet represents *two* distinct points of the line, while each of the three small circles represents *three* ‘Jacobson’ points.

### 3 Classifying Lines over Rings of Small Orders

Following the same strategy as in the previous section, we have examined the properties of the projective lines over commutative rings with unity of all orders up to thirty-one. The most relevant results/findings of our study are in a succinct, and in a number of respects also unique, form displayed in Table 3; note that

the line analysed in the preceding section is of type 16/12. The following abbreviations are used for the number of points: ‘Tot’ – all the points of the line; ‘TpI’ – points of type I; ‘1N’ – in the neighbourhood of a generic point (the point itself exclusive); ‘ $\cap 2N$ ’ – common to the neighbourhoods of two distant points; ‘ $\cap 3N$ ’ – shared by the neighbourhoods of three pairwise distant points; ‘Jcb’ – having ‘Jacobson’ property; ‘MD’ – the maximum set of pairwise distant ones. The type of a line is given in the form A/B, where A is the total number of elements of the associated ring and B is the number of its zero-divisors. In all cases except for 8/4, 16/8, 16/12, 24/16 and 27/9 the table gives all the representative rings; due to a lack of space here, a full list of rings for each of the five specified types is given in a separate table (Table 4), using the notation of [13] (orders  $\leq 16$ ) and [14] (orders  $>16$ ).

Table 3: The basic types of a projective line over small commutative rings with unity. For the types denoted by bold-facing there also exist “non-commutative” projective lines of the particular orders [15].

Line Type	Cardinalities of Points							Representative Rings
	Tot	TpI	1N	$\cap 2N$	$\cap 3N$	Jcb	MD	
31/1	32	32	0	0	0	0	32	$GF(31)$
30/22	72	52	41	20	6	7	3	$GF(2) \otimes GF(3) \otimes GF(5)$
29/1	30	30	0	0	0	0	30	$GF(29)$
28/10	40	38	11	2	0	3	5	$GF(4) \otimes GF(7)$
28/16	48	44	19	4	0	11	3	$GF(7) \otimes [Z_4 \text{ or } GF(2)[x]/\langle x^2 \rangle]$
28/22	72	50	43	22	6	5	3	$GF(2) \otimes GF(2) \otimes GF(7)$
27/1	28	28	0	0	0	0	28	$GF(27)$
27/9	36	36	8	0	0	8	4	$Z_{27}, GF(3)[x]/\langle x^3 \rangle, \dots$
27/11	40	38	12	2	0	6	4	$GF(3) \otimes GF(9)$
<b>27/15</b>	48	42	20	6	0	2	4	$GF(3) \otimes [Z_9 \text{ or } GF(3)[x]/\langle x^2 \rangle]$
27/19	64	46	36	18	6	0	4	$GF(3) \otimes GF(3) \otimes GF(3)$
26/14	42	40	15	2	0	11	3	$GF(2) \otimes GF(13)$
25/1	26	26	0	0	0	0	26	$GF(25)$
25/5	30	30	4	0	0	4	6	$Z_{25}, GF(5)[x]/\langle x^2 \rangle$
25/9	36	34	10	2	0	0	6	$GF(5) \otimes GF(5)$
24/10	36	34	11	2	0	5	4	$GF(3) \otimes GF(8)$
24/16	48	40	23	8	0	7	3	$GF(3) \otimes Z_8, GF(3) \otimes GF(2)[x]/\langle x^3 \rangle, \dots$
24/18	60	42	35	18	6	5	3	$GF(2) \otimes GF(3) \otimes GF(4)$
<b>24/20</b>	72	44	47	28	12	3	3	$GF(2) \otimes GF(3) \otimes [Z_4 \text{ or } GF(2)[x]/\langle x^2 \rangle]$
24/22	108	46	83	62	42	1	3	$GF(2) \otimes GF(2) \otimes GF(2) \otimes GF(3)$
23/1	24	24	0	0	0	0	24	$GF(23)$
22/12	36	34	13	2	0	9	3	$GF(2) \otimes GF(11)$
21/9	32	30	10	2	0	4	4	$GF(3) \otimes GF(7)$
20/8	30	28	9	2	0	1	5	$GF(5) \otimes GF(4)$
20/12	36	32	15	4	0	7	3	$GF(5) \otimes [Z_4 \text{ or } GF(2)[x]/\langle x^2 \rangle]$
20/16	54	36	33	18	6	3	3	$GF(5) \otimes GF(2) \otimes GF(2)$
19/1	20	20	0	0	0	0	20	$GF(19)$
18/10	30	28	11	2	0	7	3	$GF(2) \otimes GF(9)$
18/12	36	30	17	6	0	5	3	$GF(2) \otimes [Z_9 \text{ or } GF(3)[x]/\langle x^2 \rangle]$
18/14	48	32	29	16	6	3	3	$GF(2) \otimes GF(3) \otimes GF(3)$
17/1	18	18	0	0	0	0	18	$GF(17)$

Table 3: (Continued)

Line Type	Cardinalities of Points							Representative Rings
	Tot	TpI	1N	$\cap 2N$	$\cap 3N$	Jcb	MD	
16/1	17	17	0	0	0	0	17	$GF(16)$
<b>16/4</b>	20	20	3	0	0	3	5	$Z_4[x]/\langle x^2 + x + 1 \rangle, GF(4)[x]/\langle x^2 \rangle$
16/7	25	23	8	2	0	0	5	$GF(4) \otimes GF(4)$
<b>16/8</b>	24	24	7	0	0	7	3	$Z_{16}, Z_4[x]/\langle x^2 \rangle, GF(2)[x]/\langle x^4 \rangle, \dots$
16/9	27	25	10	2	0	6	3	$GF(2) \otimes GF(8)$
<b>16/10</b>	30	26	13	4	0	5	3	$GF(4) \otimes [Z_4 \text{ or } GF(2)[x]/\langle x^2 \rangle]$
<b>16/12</b>	36	28	19	8	0	3	3	$Z_4 \otimes Z_4, GF(2) \otimes Z_8, \dots$
16/13	45	29	28	16	6	2	3	$GF(2) \otimes GF(2) \otimes GF(4)$
<b>16/14</b>	54	30	37	24	12	1	3	$GF(2) \otimes GF(2) \otimes [Z_4 \text{ or } GF(2)[x]/\langle x^2 \rangle]$
16/15	81	31	64	50	36	0	3	$GF(2) \otimes GF(2) \otimes GF(2) \otimes GF(2)$
15/7	24	22	8	2	0	2	4	$GF(3) \otimes GF(5)$
14/8	24	22	9	2	0	5	3	$GF(2) \otimes GF(7)$
13/1	14	14	0	0	0	0	14	$GF(13)$
12/6	20	18	7	2	0	1	4	$GF(3) \otimes GF(4)$
12/8	24	20	11	4	0	3	3	$GF(3) \otimes [Z_4 \text{ or } GF(2)[x]/\langle x^2 \rangle]$
12/10	36	22	23	14	6	1	3	$GF(2) \otimes GF(2) \otimes GF(3)$
11/1	12	12	0	0	0	0	12	$GF(11)$
10/6	18	16	7	2	0	3	3	$GF(2) \otimes GF(5)$
9/1	10	10	0	0	0	0	10	$GF(9)$
9/3	12	12	2	0	0	2	4	$Z_9, GF(3)[x]/\langle x^2 \rangle$
9/5	16	14	6	2	0	0	4	$GF(3) \otimes GF(3)$
8/1	9	9	0	0	0	0	9	$GF(8)$
8/4	12	12	3	0	0	3	3	$Z_8, GF(2)[x]/\langle x^3 \rangle, \dots$
8/5	15	13	6	2	0	2	3	$GF(2) \otimes GF(4)$
<b>8/6</b>	18	14	9	4	0	1	3	$GF(2) \otimes [Z_4 \text{ or } GF(2)[x]/\langle x^2 \rangle]$
8/7	27	15	18	12	6	0	3	$GF(2) \otimes GF(2) \otimes GF(2)$
7/1	8	8	0	0	0	0	8	$GF(7)$
6/4	12	10	5	2	0	1	3	$GF(2) \otimes GF(3)$
5/1	6	6	0	0	0	0	6	$GF(5)$
4/1	5	5	0	0	0	0	5	$GF(4)$
4/2	6	6	1	0	0	1	3	$Z_4, GF(2)[x]/\langle x^2 \rangle$
4/3	9	7	4	2	0	0	3	$GF(2) \otimes GF(2)$
3/1	4	4	0	0	0	0	4	$GF(3)$
2/1	3	3	0	0	0	0	3	$GF(2)$

From Table 3 one can discern a number of interesting features of the structure of this hierarchy. The most marked one is a general increase of the total number of points, those of type II and the population of neighbourhoods with the increasing number of zero-divisors of the base ring. This is accompanied by strengthening of the ‘coupling,’ or ‘entanglement,’ between the neighbourhoods of mutually distant points, which is embodied in gradually increasing overlaps between the neighbourhoods of first two and then three such points; note, for example, that for lines of 16/15 type, the number of points in the intersection of the neighbourhoods of three pairwise distant points is greater than that of type I. One further observes that various types of a projective line form natural groups (Gr) differing from each other in the total number of elements of the associated base ring. Each such group can further be divided into classes (Cl) of the same maximum number of mutually distant points; thus, for example, Gr-16 consists of three classes Cl-17, Cl-5 and Cl-3, the last-mentioned being the most populated. Classes generated by the Galois fields,  $GF(q)$ , contain just a single entry and are regarded as trivial. A further subdivision within a given class is into cells (Ce) according as the neighbourhoods of mutually distant points are disjoint

Table 4: A comprehensive list of all commutative rings with unity which generate the projective lines of the type 8/4, 16/8, 16/12, 24/16 and 27/9.

Type of Line	Representative Rings
8/4	1.4, 2.19, 2.20, 3.14, 3.19
16/8	1.5, 2.28, 2.29, 3.40, 3.42, 3.61, 3.64, 4.118, 4.119, 4.165, 4.166, 4.168, 4.170, 5.99, 5.102, 5.109, 5.110, 5.111
16/12	2.19, 3.49, 4.81, 4.82, 4.167, 5.107, 5.114, 5.115
24/16	1.8, 2.39, 2.40, 3.28, 3.38
27/9	1.4, 2.17, 2.24, 2.25, 3.14, 3.23

( $Ce \cap 1$ ), overlaps for pairs ( $Ce \cap 2$ ) and/or triples ( $Ce \cap 3$ ) of them; so, for example, the class Cl-3 of Gr-16 is seen to comprise all the three kinds of a cell, of the cardinalities one, three and three, respectively, whilst the same class of Gr-2 features just a single cell,  $Ce \cap 1$ . A point that deserves particular attention here is a considerable change in the structure of the line under the transition between the classes within a given group. This change is likely to be found ever more pronounced with the increasing order of the ring and for lines within our scope it is best visible in Gr-16, on the boundary between its Cl-5 and Cl-3; one sees that drop, when moving from 16/7 to 16/8, in the maximum number of pairwise distant and the total number of points, as well as in the cardinality of a neighbourhood, is accompanied by reappearance of ‘Jacobson’ points and disappearance of the cell of type  $Ce \cap 2$ . Another noteworthy property is gradual decrease in the number of ‘Jacobson’ points within any (non-trivial) class. It is also to be noted that the neighbourhoods of mutually distant points are disjoint on the lines defined over local rings (types 4/2, 8/4, 9/3, 16/4, 16/8, 25/5 and 27/9), a fact which entails the transitivity of the neighbour relation.

## 4 Conclusion

All projective lines defined over commutative rings with unity of orders two to thirty-one have been classified. We have found altogether sixty-five different types of them. Each type is characterized by the following string of parameters: the total number of points on the line, the number of points represented by coordinates with at least one entry being a unit, the cardinality of the neighbourhood of a generic point of the line as well as those of the intersections between the neighbourhoods of two and three mutually distant points, the number of ‘Jacobson’ points per a neighbourhood, the maximum number of pairwise distant points and, finally, by a list of representative rings. The exposition of the ideas and the classification itself are presented in the way to stir the interest of physicists, chemists, biologists and scholars of other natural sciences to look for possible applications of these abstract finite geometries in their domains of research. As per physics, a couple of the above-introduced types of a projective ring line, namely the 4/2 and 8/6 ones, have already been successfully employed to account for some subtleties of the structure of *two-qubit* systems [5],[6]; our most recent study [16] indicates that it is also the line of type 16/15 whose structure is relevant for these quantum systems, with type II points (and, so, zero-divisors) playing a particular role here. The standard model of elementary particles and their interactions is another domain where the combinatorics of finite ring lines is likely to find its proper setting; in this respect, it should be emphasized that the line of type 4/2 could be of interest for the classification of the six quarks and six leptons.

## Acknowledgements

This work was partially supported by the Science and Technology Assistance Agency under the contract # APVT-51-012704, the VEGA project # 2/6070/26 (both from Slovak Republic), the trans-national ECO-NET project # 12651NJ “Geometries Over Finite Rings and the Properties of Mutually Unbiased Bases” (France) and by the project 1ET400400410 of the Academy of Sciences of the Czech Republic.

## References

- [1] Blunck A, Havlicek H. Projective representations I: Projective lines over a ring. *Abh Math Sem Univ Hamburg* 2000;70:287–99.
- [2] Blunck A, Havlicek H. Radical parallelism on projective lines and non-linear models of affine spaces. *Mathematica Pannonica* 2003;14:113–27.
- [3] Blunck A, Havlicek H. On distant-isomorphisms of projective lines. *Aequationes Mathematicae* 2005;69:146–163.
- [4] Havlicek H. Divisible designs, Laguerre geometry, and beyond. *Quaderni del Seminario Matematico di Brescia* 2006;11:1–63. A preprint available from <http://www.geometrie.tuwien.ac.at/havlicek/dd-laguerre.pdf>.
- [5] Saniga M, Planat M. The projective line over the finite quotient ring  $\text{GF}(2)[x]/\langle x^3 - x \rangle$  and quantum entanglement I. Theoretical background. *Theoretical and Mathematical Physics* 2006; submitted. Available from [quant-ph/0603051](http://arxiv.org/abs/quant-ph/0603051).
- [6] Saniga M, Planat M, Minarovjeh M. The projective line over the finite quotient ring  $\text{GF}(2)[x]/\langle x^3 - x \rangle$  and quantum entanglement II. The Mermin “Magic” Square/Pentagram. *Theoretical and Mathematical Physics* 2006; submitted. Available from [quant-ph/0603206](http://arxiv.org/abs/quant-ph/0603206).
- [7] Saniga M, Planat M. On the fine structure of the projective line over  $\text{GF}(2) \otimes \text{GF}(2) \otimes \text{GF}(2)$ . *Indagationes Mathematicae* 2006; submitted. Available from [math.AG/0604307](http://arxiv.org/abs/math.AG/0604307).
- [8] Veldkamp FD. Geometry over rings. In: Buekenhout F, editor. *Handbook of incidence geometry*. Amsterdam: Elsevier; 1995. p. 1033–84.
- [9] Herzer A. Chain geometries. In: Buekenhout F, editor. *Handbook of incidence geometry*. Amsterdam: Elsevier; 1995. p. 781–842.
- [10] Fraleigh JB. *A first course in abstract algebra* (5th edition). Reading (MA): Addison-Wesley; 1994. p. 273–362.
- [11] McDonald BR. *Finite rings with identity*. New York: Marcel Dekker; 1974.
- [12] Raghavendran R. Finite associative rings. *Comp Mathematica* 1969;21:195–229.
- [13] Nöbauer C. *The Book of the Rings — Part I*. 2000; Available from <http://www.algebra.uni-linz.ac.at/~noebasi/pub/rings.ps>.
- [14] Nöbauer C. *The Book of the Rings — Part II*. 2000; Available from <http://www.algebra.uni-linz.ac.at/~noebasi/pub/ringsII.ps>.
- [15] Saniga M, Planat M, Pracna P. A classification of the projective lines over small rings II. Non-commutative case. Available from [math.AG/0606500](http://arxiv.org/abs/math.AG/0606500).
- [16] Planat M, Saniga M, Kibler MR. Quantum entanglement and finite ring geometry. *SIGMA* 2006; accepted. Available from [quant-ph/0605239](http://arxiv.org/abs/quant-ph/0605239).