Synthesis and X-ray crystal structure of [2(phosphinomethyl)ferrocenyl]diphenyl-phosphine

Francis Labrue, Bénédicte Pons, Louis Ricard, Angela Marinetti

► To cite this version:

Francis Labrue, Bénédicte Pons, Louis Ricard, Angela Marinetti. Synthesis and X-ray crystal structure of [2(phosphinomethyl)ferrocenyl]diphenyl-phosphine. *Journal of Organometallic Chemistry*, 2005, 690, pp.2285-2290. 10.1016/j.jorganchem.2005.02.034 . hal-00114278

HAL Id: hal-00114278

<https://hal.science/hal-00114278>

Submitted on 17 Nov 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Synthesis and X-ray Crystal Structure of [2-(Phosphinomethyl)ferrocenyl]diphenylphosphine

Francis Labrue^b, Bénédicte Pons^b, Louis Ricard^c, Angela Marinetti^{a*}

^a Institut de Chimie des Substances Naturelles, C.N.R.S., Avenue de la Terrasse, 91198 Gif-sur-Yvette Cedex, France.

^b Laboratoire de Synthèse Sélective Organique et Produits Naturels – E.N.S.C.P. 11, rue Pierre et Marie Curie -75231 Paris Cedex 06

^c Laboratoire Hétéroéléments et Coordination, Ecole Polytechnique – 91128 Palaiseau Cedex

Abstract

[2-(phosphinomethyl)ferrocenyl]diphenylphosphine **2**, is an air stable primary phosphine bearing a 1,2-disubstituted ferrocene framework, which has been prepared by reduction of the corresponding phosphonate. Confirmation of its structure has been obtained by X-ray single-crystals diffraction analysis. Despite its high stability toward oxidation, phosphine **2** still displays a normal coordinative behaviour toward [(*p*-cymene)RuCl₂]₂. The expected (*p*-cymene)RuCl₂(phosphine) complex is formed by coordination of the primary phosphine function, while the conceivably competitive complexation of the PPh₂ group was not observed.

Keywords : ferrocenyl derivative, diphosphine, primary phosphine

1. Introduction

Primary phosphines are highly useful intermediates in organophosphorus chemistry, owing to the reactive character of the P-H bond. They are however also typically air-sensitive, often pyrophoric and, thus, difficult compounds to handle. Recent work from W. Henderson [1,2,3] pointed out the availability of especially ‘user-friendly’ primary phosphines which incorporates ferrocenyl groups, i.e. ferrocenylmethyl- and ferrocenylethylphosphines **1**. These primary phosphines are remarkably air stable, both in solution and in the solid state. Albeit the origin of the exceptionally high air-stability can’t be explained to date, the synthetic utility of **1** was easily anticipated.

Scheme 1

Based on the pioneering work above, we envisioned the synthesis of the analogous primary phosphine **2** which bears an additional diphenylphosphino function on the cyclopentadienyl moiety.

* Corresponding author. E-mail address: angela.marinetti@icsn.cnrs-gif.fr

If the ‘user-friendly’ physical properties of **1** would be retained in diphosphine **2**, this compound could represent a highly convenient synthon to produce new ferrocenyl-based diphosphines with planar chirality.

Ferrocenyl-based diphosphines with planar chirality constitute a well known, efficient class of chiral auxiliaries for transition metal catalysed reactions [4]. Togni’s Josiphos [5] is the lead-compound in this series, however a variety of other diphosphines bearing chiral, 1,2-disubstituted ferrocene moieties and, possibly, an additional chiral carbon centre, have been prepared [6] since the initial work of Hayashi and Kumada [7]. Several synthetic approaches have been applied to their synthesis, however, as far as we are aware, the use of primary phosphines, such as **2**, as synthetic intermediates has never been considered before.

We report hereafter on a synthetic approach to racemic and enantio-enriched **2**, as well as on its structural characterization by X-ray diffraction.

2. Results and discussion

For the synthesis of the target diphosphine **2**, the known oxide **4**⁸ and sulphide **5** [7] of [2-(N,N-dimethylaminomethyl)ferrocenyl]diphenylphosphine **3** [7,9], have been considered as suitable starting materials. Conversion of their nitrogen functions into trimethylammonium iodide functions should create leaving groups to be displaced then by a suitable phosphorus reagent. An analogous procedure had been applied to the synthesis of **1**, by using tris(hydroxymethyl)phosphine as the nucleophile [1].

Scheme 2

While nitrogen methylation of the trivalent phosphine **3**, is hampered by the potentially concurrent alkylation of phosphorus, it can be suitably performed on both the phosphine oxide and sulphide to

afford **6** and **7** respectively. Starting from **6**, the reaction with tris(hydroxymethyl)phosphine in refluxing methanol, according to the Henderson procedure [10,11], did not afford the expected bis(hydroxymethyl)phosphino derivative, mainly due to the low reactivity of the starting material. Different approaches were then considered, where the ammonium iodide function would serve as a leaving group in Michaelis-Becker or Arbuzov-like reactions with sodium diethylphosphite and trimethylphosphite respectively. The last reaction led successfully to the desired phosphonates **8** and **9** in 57% and 20% yields, respectively.

Scheme 3

The use of ammonium iodides in Arbuzov-like reactions [12] has been applied previously to the synthesis of diethyl ferrocenylmethylphosphonate [13], a ferrocene derivative closely related to **8** or **9**.

Reduction of both the phosphonate and phosphine oxide functions of **8** has been performed in a single step by using a trichlorosilane-triethylamine mixture, in benzene or toluene, by heating at 120°C in a sealed tube. The final diphosphine **2** [^{31}P NMR δ -23 and -127 ppm, $^1J_{\text{P-H}} = 196$ Hz] is a stable compound which has been purified by column chromatography and stored in air without noticeable oxidation.

Crystals of **2** have been grown from pentane, which were suitable for X-ray crystal structure determination. ORTEP drawing of **2** is reported in Figure 1. Selected bond angles and distances are given in Table 1.

Scheme 4

Compound **10** is an orange-red, air stable solid. ^{31}P NMR data are diagnostic of the exclusive complexation of the PH_2 fragment, with an almost unchanged chemical shift for the PPh_2 unit [δ -25 ppm] and a considerably large $\Delta\delta$ value [106 ppm] for the PH_2 group [δ -21 ppm], with respect to the uncomplexed diphosphine **2**. Complexation of phosphorus makes the two P-H bonds enantiotopic thus inducing two different $^1\text{J}_{\text{H-P}}$ couplings of 385 and 348 Hz, respectively, as well as separate ^1H NMR signals for the corresponding protons at δ 4.26 and 4.48 ppm, respectively.

Thus, the efficient synthetic approach to the primary phosphine **2** reported herein, affords a new ‘user-friendly’ ligand as well as a valuable precursor for ferrocene-based diphosphines.

Finally, we wished to check if the synthetic approach defined above would be suitable for the synthesis of enantiomerically pure **2** through stereospecific transformations of a chiral starting material. Thus, the known phosphine sulphides (*R*)- and (*S*)-**5** have been prepared separately through the reported resolution method, which implies fractional crystallisation with dibenzoyltartaric acid [7]. (*R*)-**5** was then converted into the corresponding oxide (*R*)-**4** by reaction with oxone [15].

Scheme 5

The next step, the Arbuzov-like reaction of the ammonium salt **6** with trimethylphosphite, causes partial racemisation and, consequently, phosphonate **8** was obtained in only 46% enantiomeric excess (measured by HPLC on Daicel OD-H column). Racemisation has been barely noticed for ferrocenes bearing planar chirality, including ferrocenylphosphines [16]. Racemisation, e.g. coordination of iron by the other face of the Cp, may take place by either an intermolecular exchange process *via* triple-decker sandwich intermediates, or *via* a dissociative process [*a*], or again *via* η^1 coordination of the Cp ring followed by 1,5-proton shift [*b*]. The last two mechanisms are shown in Scheme 6 hereafter.

Scheme 6. Intramolecular racemisation processes for chiral ferrocenes.

Both a dissociative process and slippage of the Cp ring from η^5 - to η^1 -coordination are plausibly promoted, in the experimental conditions of the Arbuzov reaction (Scheme 3), by temporary complexation of trimethylphosphite ($L = \text{P}(\text{OMe})_3$). An haptotropic shift of the Cp ligand of a ferrocene derivative, in the presence of trimethylphosphite, has been documented recently [17].

The observed partial racemisation process prevents application of the synthetic method above (Schemes 2 and 3) to the synthesis of an enantiomerically pure diphosphine **2**.

Other approaches to enantiomerically pure **2** are being considered. These include resolution of racemic **2** or even resolution of phosphonate **8**, given that reduction of **8** in the reported conditions, proved to be stereospecific. Also, the use of more suitable precursors, allowing the synthesis of **8** in milder conditions, is under investigation.

3. Experimental

The [2-(*N,N*-dimethylaminomethyl)ferrocenyl]diphenylphosphine **3**, [2-(*N,N*-dimethylaminomethyl)ferrocenyl]diphenylphosphine sulfide **5** have been prepared according to Hayashi et al. [7] All experiments were performed under an inert atmosphere. NMR spectra were recorded on either a Bruker AM 200 or an AM 400 spectrometer.

[2-(*N,N*-dimethylaminomethyl)ferrocenyl]diphenylphosphine oxide (**4**).

From 3. To a suspension of phosphine **3** (7.7g, 16 mmol) in methanol (130 mL) at 0°C, was added dropwise 2.1 mL of 30% hydrogen peroxide. After 30 min at room temperature, aqueous sodium sulfite was added to decompose the excess hydrogen peroxide. After evaporation of methanol, the

reaction mixture was extracted with dichloromethane. The extract was washed with water, dried over anhydrous magnesium sulphate and then evaporated. Purification of the residue by column chromatography on basic alumina (ether/methanol, 99:1) gave the corresponding oxide **4** in 91% yield (7.1g).

From (R)-5. (R)-[2-(N,N-dimethylaminomethyl)ferrocenyl]diphenylphosphine sulfide **5** [7] (2.4g, 5.2 mmol, $[\alpha]_D = -47$ (c = 0.5, CHCl₃), e.e. = 94%, based on the $[\alpha]_D$ value) was dissolved in a THF/MeOH 1:1 mixture (100 mL). Then 50 mL of a 0.2M buffered solution (pH 6-7) of oxone[®] were added slowly at 0°C. After 20 min stirring at room temperature, the mixture was hydrolysed with aqueous sodium thiosulfate and extracted with chloroform. The organic extracts were dried over MgSO₄ and solvents were removed to afford the crude oxide **4** which was purified as above. (R)-**4** was obtained in 80% yield (1.8 g); $[\alpha]_D = -104$ (c = 0.2, CHCl₃). A 10% amount of (2-formylferrocenyl)diphenylphosphine oxide [³¹P NMR (CDCl₃) δ 29.0; ¹H NMR (CDCl₃) δ 4.24 (br s, 1H), 4.41 (s, 5H, Cp), 4.78 (br s, 1H), 5.24 (br s, 1H), 7.3-7.9 (m, 10h, Ph), 10.4 (1H, CHO); ¹³C NMR (50 MHz) δ 194.3 (s, CHO) ppm] was formed as side-product when addition of oxone was performed at room temperature.

4: Orange-red solid. Spectral data are in agreement with the reported data [8]. ³¹P NMR (CDCl₃) δ 29.7; ¹H NMR (CDCl₃) δ 1.94 (s, 6H, NMe₂), 3.36 (AB, J=13.4 Hz, 1H, CH₂), 3.64 (AB, 1H, CH₂), 3.92 (br s, 1H), 4.20 (s, 5H, Cp), 4.33 (br s, 1H), 4.62 (br s, 1H), 7.3-7.9 (m, 10h, Ph) ppm. Anal. Calcd. for C₂₅H₂₆FeNP: C, 67.74; H, 5.91; N, 3.16. Found: C, 67.62, H, 5.96; N, 3.20.

Trimethyl[(2-diphenylphosphinyl)ferrocenylmethyl]ammonium iodide (6). To a solution of amine **4** (6.5 g, 14.6 mmol) in methanol (15 mL), an excess methyl iodide (2.7 mL, 44 mmol) was added at room temperature. The solution was heated for 10 min at about 60°C and then cooled to room temperature. 130 mL of ether were added with stirring. The ammonium salt **6** separates as an orange oil which crystallizes on standing (7.3g, 85% yield). ³¹P NMR (CDCl₃) δ 30.8; ¹H NMR (CDCl₃) δ 3.10 (s, 9H, NMe₃), 4.16 (s, 5H, Cp), 4.35 (br s, 1H), 4.72 (br s, 1H), 5.23 (AB, J=12.7 Hz, 1H, CH₂), 5.30 (br s, 1H), 5.45 (AB, 1H, CH₂), 7.4-7.8 (m, 10h, Ph) ppm. The crude ammonium salt was used without further purification.

Trimethyl[(2-diphenylthiophosphinyl)ferrocenylmethyl]ammonium iodide (7) was prepared by the same procedure as for **6**, in 70% yield. ³¹P NMR (CDCl₃) δ 41.4; ¹H NMR (CDCl₃) δ 3.06 (s, 9H, NMe₃), 4.18 (br s, 1H), 4.36 (s, 5H, Cp), 4.68 (br s, 1H), 5.33 (br s, 1H), 5.41 (AB, J=13.2 Hz, 1H, CH₂), 5.92 (AB, 1H, CH₂), 7.4-7.9 (m, 10h, Ph) ppm. The crude ammonium salt was used without further purification.

Dimethyl [2-(diphenylphosphinyl)ferrocenyl]methylphosphonate (8). A solution containing the ammonium salt **6** (4.0g, 6.8 mmol) and trimethylphosphite (4 mL) in xylene (10 mL) was heated at 145°C for 2h30. Solvent and excess P(OMe)₃ were removed by distillation under vacuum. The residue was taken up in dichloromethane and purified by chromatography on alumina with ether/methanol 95:5 as the eluent. 2.0g of **8** were obtained (57% yield). ³¹P NMR (CDCl₃) δ 29.6 and 30.9 ppm; ¹H NMR (CD₂Cl₂) 400 MHz, δ 3.06 (dd, J = 20.7 Hz, J = 15.7 Hz, 1H, CH₂), 3.30 (d, ³J_{H-P} = 10.9 Hz, 3H, OMe), 3.45 (d, ³J_{H-P} = 10.9 Hz, 3H, OMe), 3.74 (dd, J = 19.0 Hz, J = 15.7 Hz, 1H, CH₂), 4.00 (br m, 1H), 4.11 (s, 5H, Cp), 4.43 (q, J = 2.4 Hz, 1H), 4.72 (br s, 1H), 7.4-7.9 (m, 10h, Ph); ¹³C NMR (50 MHz, selected data) δ 24.8 (d, ¹J_{PC} = 138.6 Hz, PCH₂), 51.8 (d, ²J_{PC} = 6.8 Hz, OCH₃), 52.3 (d, ²J_{PC} = 6.8 Hz, OCH₃), 70.6 (s, Cp), 70.7 (dd, ¹J_{PC} = 114.0 Hz, ³J_{PC} = 6.0 Hz, PC), 71.9 (d, J_{PC} = 11.0 Hz, CH), 72.6 (d, J_{PC} = 14.4 Hz, CH), 73.4 (dd, J_{PC} = 9.6 Hz, J_{PC} = 2.4 Hz, CH), 82.9 (d, J_{PC} = 10.0 Hz, J_{PC} = 4.0 Hz, C), 134.9 (d, J_{PC} = 105.5 Hz, C_{ipso}), 133.8 (d, J_{PC} = 106.3 Hz, C_{ipso}) ppm. Mass spectrum: m/z 508 (M, 95%), 443 (M-Cp, 100%). Anal. Calcd. for C₂₅H₂₆FeO₄P₂: C, 59.08; H, 5.16. Found: C, 58.99; H, 5.20.

When starting from the ammonium iodide of (*R*)-**4** above, phosphonate (*R*)-**8** was obtained with an enantiomeric excess of 46%. The enantiomeric excess was determined by HPLC analysis: Chiracel OD-H, hexane/iPrOH 85:15, *t*₁ = 12.6 min (major), *t*₂ = 17.4 min. [α]_D = -34 (c = 0.2, CHCl₃).

Dimethyl [2-(diphenylthiophosphinyl)ferrocenyl]methylphosphonate (9). The same procedure as for **8**, afforded **9** in 20% yield. ³¹P NMR (CDCl₃) δ 29.7 and 42.5; ¹H NMR (CDCl₃) 400 MHz, δ 3.25 (t, J = 18 Hz, 1H, CH₂), 3.33 (d, ³J_{H-P} = 10.1 Hz, 3H, OMe), 3.51 (d, ³J_{H-P} = 10.2 Hz, 3H, OMe), 3.76 (br s, 1H), 4.02 (t, J = 17.3 Hz, 1H, CH₂), 4.24 (s, 5H, Cp), 4.34 (br s, 1H), 4.83 (br s, 1H), 7.3-7.8 (m, 10h, Ph); ¹³C NMR (100 MHz, selected data) δ 24.8 (d, ¹J_{PC} = 138 Hz, PCH₂), 51.9 (d, ²J_{PC} = 6 Hz, OCH₃), 52.5 (d, ²J_{PC} = 6 Hz, OCH₃), 69.8 (d, J_{PC} = 9.8 Hz, CH), 71.2 (s, Cp), 73.0 (dd, ¹J_{PC} = 89.5 Hz, ³J_{PC} = 6 Hz, PC), 74.1 (d, J_{PC} = 12 Hz, CH), 74.5 (d, J_{PC} = 2 Hz, CH), 82.9 (d, J_{PC} = 13 Hz, C), 133.4 (d, J_{PC} = 86 Hz, C_{ipso}), 134.7 (d, J_{PC} = 87 Hz, C_{ipso}) ppm. Mass spectrum: m/z 524 (M, 34%), 459 (M-Cp, 100%)

[2-(phosphinomethyl)ferrocenyl]diphenylphosphine (2).

A toluene solution of phosphonate **8** (1.0g, 2.0 mmol) was added to a solution of trichlorosilane (4.0 mL, 40 mmol) in toluene (20 mL) at 0°C. An excess triethylamine (4.5 mL, 32 mmol) was then added. The mixture was then heated in a sealed glass tube at 120°C for 5h. After cooling to room

temperature, hydrolysis was performed by slow addition of an aqueous NaOH solution (20% weight). After extraction with dichloromethane, the combined organic layer were washed with water and dried over magnesium sulphate. Purification was performed by column chromatography on alumina with cyclohexane-ether 94:6 as the eluent. Yield 70% (0.51g, orange-red solid). The enantiomeric excess was determined by HPLC analysis: Chiracel OJ, hexane/*i*-PrOH 98:2, $t_1 = 6.3$ min (major), $t_2 = 9.8$ min. E.e. = 45%, $[\alpha]_D = -54$ ($c = 0.2$, CHCl_3).

^{31}P NMR (CDCl_3) δ -127.5 ($J_{\text{H-P}} = 196$ Hz) and -22.9 ($J_{\text{P-P}} = 1.6$ Hz); ^1H NMR (CDCl_3) 400 MHz, δ 2.71 (dm, $J_{\text{H-P}} = 197.2$ Hz, 2H, PH_2), 2.7-2.9 (m, 2H, CH_2), 3.73 (br, 1H), 4.00 (s, 5H, Cp), 4.25 (t, $J = 2.4$ Hz, 1H), 4.18 (q, $J = 2.0$ Hz, 1H, CH), 7.2-7.6 (m, 10h, Ph); ^{13}C NMR (100 MHz, C_6D_6) δ 13.8 (dd, $J_{\text{PC}} = 12.0$ Hz, $J_{\text{PC}} = 10.2$ Hz, PCH_2), 69.3 (s, CH), 70.1 (s, Cp), 70.7 (t, $J_{\text{PC}} = 3.8$ Hz, CH), 71.0 (d, $J_{\text{PC}} = 3.9$ Hz, CH), 75.7 (dd, $J_{\text{PC}} = 7.6$ Hz, $J_{\text{PC}} = 1.5$ Hz, C), 95.0 (dd, $J_{\text{PC}} = 25.8$ Hz, $J_{\text{PC}} = 3.8$ Hz, PC), 138.4 (d, $J_{\text{PC}} = 9.5$ Hz, C_{ipso}), 140.6 (d, $J_{\text{PC}} = 10.6$ Hz, C_{ipso}) ppm. Mass spectrum (C.I.): m/z 417 ($\text{M}+1$). Anal. Calcd. for $\text{C}_{23}\text{H}_{22}\text{FeP}_2$: C, 66.37; H, 5.33. Found: C, 66.35; H, 5.65.

[(p-cymene)RuCl₂(2)] complex (10).

A solution containing diphosphine **2** (83mg, 0.2 mmol) and $[(p\text{-cymene})\text{RuCl}_2]_2$ (61 mg, 0.1 mmol) in CH_2Cl_2 (0.5 mL) – EtOH (1.5 mL) was stirred at room temperature for 2.5h. During this time a red solid precipitates which was recovered by filtration.

^{31}P NMR (CDCl_3) δ -25.5 ($J_{\text{H-P}} = 196$ Hz, PH_2) and -21.1; ^1H NMR (CDCl_3) 400 MHz, δ 1.00 (d, $J = 6.9$ Hz, 3H, CHMe), 1.04 (d, $J = 6.9$ Hz, 3H, CHMe), 2.07 (s, 3H, Me), 2.36 (m, 1H, CHMe_2) 3.33 (m, 2H, CH_2), 3.88 (m, 1H), 3.98 (s, 5H, Cp), 4.26 (dm, $J_{\text{H-P}} = 385$ Hz, 1H, PH_2), 4.35 (t, $J = 2.5$ Hz, 1H), 4.45 (br s, 1H, CH), 4.48 (dm, $J_{\text{H-P}} = 348$ Hz, 1H, PH_2), 5.09 (d, $J_{\text{H-P}} = 5.8$ Hz, 1H, CH), 5.20 (d, $J_{\text{H-P}} = 5.9$ Hz, 1H, CH), 5.25 (d, $J_{\text{H-P}} = 5.3$ Hz, 1H, CH), 5.28 (d, $J_{\text{H-P}} = 5.0$ Hz, 1H, CH), 7.3-7.6 (m, 10h, Ph) ppm.

Crystal data for 2.

Molecular formula	$\text{C}_{23}\text{H}_{22}\text{FeP}_2$
Molecular weight	416.2
Crystal habit	orange plate
Crystal size (mm)	0.22 x 0.18 x 0.12
Crystal system	monoclinic
Space group	$\text{P}2_1/\text{n}$
a (Å)	23.028(10)
b (Å)	8.4680(10)
c (Å)	23.0340(10)
α (°)	90.00
β (°)	116.4100(10)

$\gamma(^{\circ})$	90.00
$V(\text{\AA}^3)$	4022.9(5)
Z	8
$D(\text{g.cm}^{-3})$	1.374
$F(000)$	1728
$\mu(\text{cm}^{-1})$	0.912
Diffractionmeter	KappaCCD
X-ray source	MoK α
$\lambda(\text{\AA})$	0.71069
T (K)	150.0(1)
Reflections measured	14272
Unique data	8928
R _{int}	0.0247
Reflection used	7220
Refinement type	Fsqd
Hydrogen atoms	mixed
Parameters refined	482
Reflections/parameter	14
wR ₂	0.0831
R ₁	0.0358
GoF	0.976
Difference peak / hole (e $\text{\AA}^{-3}$)	0.728(0.060) / -0.422(0.060)

Supplementary material

Crystallographic data for the structural analysis of compound **2** have been deposited with the Cambridge Crystallographic Data Centre, CCDC No. 264508. Copies of this information may be obtained free of charge from: The Director, CCDC, 12 Union Road, Cambridge, CB2 1EZ UK. Fax. +44(1223)336-033 or Email: deposit@ccdc.cam.ac.uk or www: <http://www.ccdc.cam.ac.uk>.

References

- [1] N.J. Goodwin, W. Henderson, B.K. Nicholson, Chem. Commun. (1997) 31.
- [2] N.J. Goodwin, W. Henderson, B.K. Nicholson, J. Fawcett, D.R. Russell, J. Chem. Soc. Dalton Trans. (1999) 1785.
- [3] W. Henderson, S.R. Alley, J. Organomet. Chem. 636 (2002) 120
- [4] For a recent update of their catalytic applications see: T.J. Colacot, Chem. Rev. 103 (2003) 3101
- [5] A. Togni, C. Breutel, A. Schnyder, F. Spindler, H. Landert, A. Tijani, J. Am. Chem. Soc. 116 (1994) 4062
- [6] Representative examples: (a) M. Sawamura, A. Yamauchi, T. Takegawa, Y. Ito, J. Chem. Soc. Chem. Commun. (1991) 874. (b) M. Sawamura, H. Hamashima, Y. Ito, Tetrahedron : Asymmetry 2 (1991) 593. (c) Riant, G. Argouarch, D. Guillaneux, O. Samuel, H.B. Kagan, J. Org. Chem. 63 (1998) 3511. (d) J.J. Almerna Perea, A. Börner, P. Knochel, Tetrahedron Lett.

-
- 39 (1998) 8073. (e) T. Ireland, G. Grossheimann, C. Wieser-Jeunesse, P. Knochel, *Angew. Chem. Int. Ed.* 38 (1999) 3212. (f) W. Zhang, T. Shimanuki, T. Kida, Y. Nakatsuji, I. Ikeda, *J. Org. Chem.* 54 (1999) 6247. (g) T. Sturm, W. Weissensteiner, F. Spindler, *Adv. Synth. Catal.* 345 (2003) 160. (h) P. Barbaro, C. Bianchini, G. Giambastiani, D. Masi, S.L. Parisel, A. Togni, *Synthesis* (2004) 345.
- [7] T. Hayashi, T. Mise, M. Fukushima, M. Kagotani, N. Nagashima, Y. Hamada, A. Matsumoto, S. Kawakami, M. Konishi, K. Yamamoto, M. Kumada, *Bull. Chem. Soc. Jpn.* 53 (1980) 1138
- [8] Y. Miyake, A. Yamauchi, Y. Nishibayashi, S. Uemura, *Bull. Chem. Soc. Jpn* 76 (2003) 381.
- [9] G. Marr, T. Hunt, *J. Chem. Soc. (C)* (1969) 1070.
- [10] N.J. Goodwin, W. Henderson, J.K. Sarfo, *Chem. Commun.* (1996) 1551.
- [11] N.J. Goodwin, W. Henderson, B.K. Nicholson, J.K. Sarfo, J. Fawcett, D.R. Russel, *J. Chem. Soc. Dalton Trans.* (1997) 4377.
- [12] Selected examples: (a) J. Szmuszkovicz, *J. Am. Chem. Soc.* 80 (1958) 3782. (b) S. Hirai, R.G. Harvey, E.V. Jensen, *Tetrahedron* 22 (1966) 1625. (c) H. Gross, H. Seibt, I. Keitel, *J. Prakt. Chem.* 317 (1979) 890. (d) M. Mikolajczyk, P. Graczyk, M.W. Wieczorek, G. Bujacz, *Tetrahedron Lett.* 29 (1988) 6801. (e) J-W. Weener, J.P.G. Versleijen, A. Meetsma, W. Ten Hoeve, A.M. Van Leusen, *Eur. J. Org. Chem.* (1998) 1511.
- [13] V.I. Boev, *Zh. Obshch. Khim.* 48 (1978) 1594.
- [14] For representative examples of ruthenium complexes of primary phosphines see ref. 2 and J.F. van der Maelen Uria, S. Garcia-Granda, J.A. Cabeza, I. de Rio, *Acta Crystallogr. Sect. C* 50 (1994) 1064.
- [15] L.A. Wozniak, W.J. Stec, *Tetrahedron Lett.* 40 (1999) 2537.
- [16] (a) M. Tsukazaki, M. Tinkl, A. Roglans, B.J. Chapell, N.J. Taylor, V. Snieckus, *J. Am. Chem. Soc.* 1996, 118, 685. (b) R.S. Laufer, U. Veith, N.J. Taylor, V. Snieckus, *Org. Lett.* 2000, 2, 629. (c) O.J. Curnow, G.M. Fern, *Organometallics* 21 (2002) 2827. (d) O.J. Curnow, G.M. Fern, M.L. Hamilton, A. Zahl, R. van Eldik, *Organometallics* 23 (2004) 906 and references therein.
- [17] T. Mizuta, Y. Imamura, K. Miyoshi, *J. Am. Chem. Soc.* 125 (2003) 2068.