

# BSDEs with stochastic Lipschitz condition and quadratic PDEs in Hilbert spaces

Philippe Briand

*IRMAR, Université Rennes 1, 35042 Rennes Cedex, FRANCE*  
 philippe.briand@univ-rennes1.fr

Fulvia Confortola

*Dipartimento di Matematica, Politecnico di Milano*  
*piazza Leonardo da Vinci 32, 20133 Milano, Italy*  
 fulvia.confortola@unimib.it

January 29, 2006

## Abstract

This paper is devoted to the study of the differentiability of solutions to real-valued backward stochastic differential equations (BSDEs for short) with quadratic generators driven by a cylindrical Wiener process. The main novelty of this problem consists in the fact that the gradient equation of a quadratic BSDE has generators which satisfy stochastic Lipschitz conditions involving BMO martingales. We show some applications to the nonlinear Kolmogorov equations.

**Key words.** BMO-martingales, backward stochastic differential equations, Kolmogorov equations.

**MSC classification.** 60H10, 35K55.

## 1 Introduction

In this paper we are concerned with a real valued BSDE

$$Y_\tau = \Phi(X_T) + \int_\tau^T F(r, X_r, Y_r, Z_r) dr - \int_\tau^T Z_r dW_r, \quad \tau \in [t, T],$$

where  $W$  is a cylindrical Wiener process in some infinite dimensional Hilbert space  $\Xi$  and the generator  $F$  has quadratic growth with respect to the variable  $z$ . Quadratic BSDEs has been intensively studied by Kobylanski [13], and then by Lepeltier and San Martin in [14] and more recently by Briand and Hu in [3]. The process  $X$ , appearing in the generator and in the terminal value of the BSDE, takes its values in an Hilbert space  $H$  and it is solution of the following forward equation

$$\begin{cases} dX_\tau = AX_\tau d\tau + b(\tau, X_\tau) d\tau + \sigma(\tau, X_\tau) dW_\tau, & \tau \in [t, T], \\ X_t = x \in H. \end{cases}$$

$A$  is the generator of a strongly continuous semigroup of bounded linear operators  $\{e^{tA}\}$  in  $H$ ,  $b$  and  $\sigma$  are functions with values in  $H$  and  $L_2(\Xi, H)$  – the space of Hilbert-Schmidt operators from  $\Xi$  to  $H$  – respectively. Under suitable assumptions on the coefficients, there exists a unique adapted process  $(X, Y, Z)$  in the space  $H \times \mathbf{R} \times L_2(\Xi, \mathbf{R})$  solution to this forward-backward system. The processes  $X, Y, Z$  depend on the values of  $x$  and  $t$  occurring as initial conditions in the forward equation: we may denote them by  $X^{t,x}$ ,  $Y^{t,x}$  and  $Z^{t,x}$ .

Nonlinear BSDEs were first introduced by Pardoux and Peng [19] and, since then, have been studied with great interest in finite and infinite dimensions: we refer the reader to [8], [6] and [18] for an exposition of this subject and to [15] for coupled forward-backward systems. The interest in BSDEs comes from their connections with different mathematical fields, such as finance, stochastic control and partial differential equations. In this paper, we are concerned with the relation between BSDEs and nonlinear PDEs known as the nonlinear Feynman-Kac formula. More precisely, let us consider the following nonlinear PDE

$$\partial_t u(t, x) + \mathcal{L}_t[u(t, \cdot)](x) + F(t, x, u(t, x), \sigma(t, x)^* \nabla_x u(t, x)) = 0, \quad u(T, x) = \Phi(x),$$

where  $\mathcal{L}_t$  is the infinitesimal generator of the diffusion  $X$ . Then the solution  $u$  is given by the formula  $u(t, x) = Y_t^{t,x}$  which generalizes the Feynman-Kac formula to a nonlinear setting.

Numerous results (for instance [21, 20, 17, 18, 13]) show the connections between BSDEs set from a forward-backward system and solutions of a large class of quasilinear parabolic and elliptic PDEs. In the finite dimensional case, solutions to PDEs are usually understood in the viscosity sense. Here we work in infinite dimensional spaces and consider solutions in the so called mild sense (see e.g. [9]), which are intermediate between classical and viscosity solutions. This notion of solution seems natural in infinite dimensional framework: to have a mild solution its enough to prove that it is Gâteaux differentiable. Hence we don't have to impose heavy assumptions on the coefficients as for the classical solutions. However a mild solution is Gâteaux differentiable and thus more regular than a viscosity solution. For the probabilistic approach, this means that, in the infinite dimensional case, one has to study the regularity of  $X^{t,x}$ ,  $Y^{t,x}$  and  $Z^{t,x}$  with respect to  $t$  and  $x$  in order to solve the PDE.

This problem of regular dependence of the solution of a stochastic forward-backward system has been studied in finite dimension by Pardoux, Peng [20] and by El Karoui, Peng and Quenez [8], and, in infinite dimension, by Fuhrman and Tessitore in [9], [10]. In both cases,  $F$  is assumed to be Lipschitz continuous with respect to  $y$  and  $z$ . In [1], in infinite dimension, the generator  $F$  is assumed to be only Lipschitz continuous only with respect to  $z$  and monotone with respect to  $y$  in the spirit of the works [21], [17] and more recently [2].

In this work, we want to achieve this program when  $F$  is quadratic with respect to  $z$  meaning that the PDE is quadratic in the gradient. We will only consider the case of a bounded function  $\Phi$ . The study of the differentiability of the process  $Y$  with respect to  $x$  in this quadratic framework open an interesting problem of solvability of linear BSDEs with stochastic Lipschitz condition. Let us show with an example what happens in order to motivate the assumptions we will work with.

Let  $(Y^x, Z^x)$  be the solution to the BSDE – all processes are real in this example –

$$Y_t^x = \Phi(x + W_t) + \frac{1}{2} \int_t^T |Z_s^x|^2 ds - \int_t^T Z_s^x dW_s$$

where  $\Phi$  is bounded and  $\mathcal{C}^1$ . If  $(G^x, H^x)$  stands for the gradient with respect to  $x$  of  $(Y^x, Z^x)$

then we have, at least formally,

$$G_t^x = \Phi'(x + W_t) + \int_t^T Z_s^x H_s^x ds - \int_t^T H_s^x dW_s.$$

In this linear equation, of course, the process  $Z^x$  is not bounded in general so the usual Lipschitz assumption is not satisfied. It is only known that the process  $Z^x$  is such that  $\int_0^t Z_s^x dW_s$  is a BMO-martingale: this fact was used in [11] to prove a uniqueness result. BSDEs under stochastic Lipschitz condition have already been studied in [7] and more recently in [4]. However, the results in these papers do not fit our BMO-framework. This is the starting point of this paper.

The plan of the paper is as follows: Section 2 is devoted to notations. In Section 3 we recall some known results about BMO-martingales and we state a result of existence and uniqueness for BSDEs with generators satisfying a stochastic Lipschitz condition with BMO feature. In section 4 we apply the previous result to the study the regularity of the map  $(t, x) \mapsto (Y^{t,x}, Z^{t,x})$  solution of the forward-backward system. The last section contain the applications to nonlinear Kolmogorov PDEs.

## 2 Notations

### 2.1 Vector spaces and stochastic processes

In the following, all stochastic processes will be defined on subsets of a fixed time interval  $[0, T]$ .

The letters  $\Xi$ ,  $H$  and  $K$  will always denote Hilbert spaces. Scalar product is denoted  $\langle \cdot, \cdot \rangle$ , with a subscript to specify the space if necessary. All Hilbert spaces are assumed to be real and separable.  $L_2(\Xi, K)$  is the space of Hilbert-Schmidt operators from  $\Xi$  to  $K$  endowed with the Hilbert-Schmidt norm. We observe that if  $K = \mathbf{R}$  the space  $L_2(\Xi, \mathbf{R})$  is the space  $L(\Xi, \mathbf{R})$  of bounded linear operators from  $\Xi$  to  $\mathbf{R}$ . By the Riesz isometry the dual space  $\Xi^* = L(\Xi, \mathbf{R})$  can be identified with  $\Xi$ .

$W = \{W_t\}_{t \geq 0}$  is a cylindrical Wiener process with values in the infinite dimensional Hilbert space  $\Xi$ , defined on a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ ; this means that a family  $W(t)$ ,  $t \geq 0$ , is a family of linear mappings from  $\Xi$  to  $L^2(\Omega)$  such that

- (i) for every  $u \in \Xi$ ,  $\{W(t)u, t \geq 0\}$  is a real (continuous) Wiener process;
- (ii) for every  $u, v \in \Xi$  and  $t \geq 0$ ,  $\mathbb{E}(W(t)u \cdot W(t)v) = \langle u, v \rangle_{\Xi}$ .

$\{\mathcal{F}_t\}_{t \in [0, T]}$  will denote the natural filtration of  $W$ , augmented with the family  $\mathcal{N}$  of  $\mathbb{P}$ -null sets of  $\mathcal{F}_T$ :

$$\mathcal{F}_t = \sigma(W(s) : s \in [0, t]) \vee \mathcal{N}.$$

The filtration  $\{\mathcal{F}_t\}_{t \in [0, T]}$  satisfies the usual conditions. All the concepts of measurability for stochastic processes (e.g. predictability etc.) refer to this filtration. By  $\mathcal{P}$  we denote the predictable  $\sigma$ -algebra on  $\Omega \times [0, T]$  and by  $\mathcal{B}(\Lambda)$  the Borel  $\sigma$ -algebra of any topological space  $\Lambda$ .

Next we define several classes of stochastic processes which we use in the sequel. For any real  $p > 0$ ,  $\mathcal{S}^p(K)$ , or  $\mathcal{S}^p$  when no confusion is possible, denotes the set of  $K$ -valued, adapted and càdlàg processes  $\{Y_t\}_{t \in [0, T]}$  such that

$$\|Y\|_{\mathcal{S}^p} := \mathbb{E} \left[ \sup_{t \in [0, T]} |Y_t|^p \right]^{1 \wedge 1/p} < +\infty.$$

If  $p \geq 1$ ,  $\|\cdot\|_{\mathcal{S}^p}$  is a norm on  $\mathcal{S}^p$  and if  $p \in (0, 1)$ ,  $(X, X') \mapsto \|X - X'\|_{\mathcal{S}^p}$  defines a distance on  $\mathcal{S}^p$ . Under this metric,  $\mathcal{S}^p$  is complete.  $M^p$  ( $M^p(L_2(\Xi, K))$ ) denotes the set of (equivalent classes of) predictable processes  $\{Z_t\}_{t \in [0, T]}$  with values in  $L_2(\Xi, K)$  such that

$$\|Z\|_{M^p} := \mathbb{E} \left[ \left( \int_0^T |Z_s|^2 ds \right)^{p/2} \right]^{1 \wedge 1/p} < +\infty.$$

For  $p \geq 1$ ,  $M^p$  is a Banach space endowed with this norm and for  $p \in (0, 1)$ ,  $M^p$  is a complete metric space with the resulting distance. We set  $\mathcal{S} = \cup_{p > 1} \mathcal{S}^p$ ,  $M = \cup_{p > 1} M^p$  and  $\mathcal{S}^\infty$  stands for the set of predictable bounded processes.

Given an element  $\Psi$  of  $L^2_{\mathcal{P}}(\Omega \times [0, T]; L_2(\Xi, K))$ , one can define the Itô stochastic integral  $\int_0^t \Psi(\sigma) dW_\sigma$ ,  $t \in [0, T]$ ; it is a  $K$ -valued martingale with continuous path such that

$$\mathbb{E} \left[ \sup_{t \in [0, T]} \left| \int_0^t \Psi(\sigma) dW_\sigma \right|^2 \right]^{1/2} < +\infty.$$

The previous definitions have obvious extensions to processes defined on subintervals of  $[0, T]$ .

## 2.2 The class $\mathcal{G}$

$F : X \rightarrow V$ , where  $X$  and  $V$  are two Banach spaces, has a directional derivative at point  $x \in X$  in the direction  $h \in X$  when

$$\nabla F(x; h) = \lim_{s \rightarrow 0} \frac{F(x + sh) - F(x)}{s},$$

exists in the topology of  $V$ .  $F$  is said to be Gâteaux differentiable at point  $x$  if  $\nabla F(x; h)$  exists for every  $h$  and there exists an element of  $L(X, V)$ , denoted  $\nabla F(x)$  and called Gâteaux derivative, such that  $\nabla F(x; h) = \nabla F(x)h$  for every  $h \in X$ .

**Definition 2.1.**  $F : X \rightarrow V$  belongs to the class  $\mathcal{G}^1(X; V)$  if it is continuous, Gâteaux differentiable on  $X$ , and  $\nabla F : X \rightarrow L(X, V)$  is strongly continuous.

In particular, for every  $h \in X$  the map  $\nabla F(\cdot)h : X \rightarrow V$  is continuous. Let us recall some features of the class  $\mathcal{G}^1(X, V)$  proved in [9].

**Lemma 2.2.** Suppose  $F \in \mathcal{G}^1(X, V)$ . Then

- (i)  $(x, h) \mapsto \nabla F(x)h$  is continuous from  $X \times X$  to  $V$ ;
- (ii) If  $G \in \mathcal{G}^1(V, Z)$  then  $G(F) \in \mathcal{G}^1(X, Z)$  and  $\nabla(G(F))(x) = \nabla G(F(x))\nabla F(x)$ .

**Lemma 2.3.** A map  $F : X \rightarrow V$  belongs to  $\mathcal{G}^1(X, V)$  provided the following conditions hold:

- (i) the directional derivatives  $\nabla F(x; h)$  exist at every point  $x \in X$  and in every direction  $h \in X$ ;
- (ii) for every  $h$ , the mapping  $\nabla F(\cdot; h) : X \rightarrow V$  is continuous;
- (iii) for every  $x$ , the mapping  $h \mapsto \nabla F(x; h)$  is continuous from  $X$  to  $V$ .

These definitions can be generalized to functions depending on several variables. For instance, if  $F$  is a function from  $X \times Y$  into  $V$ , the partial directional and Gâteaux derivatives with respect to the first argument, at point  $(x, y)$  and in the direction  $h \in X$ , are denoted  $\nabla_x F(x, y; h)$  and  $\nabla_x F(x, y)$  respectively.

**Definition 2.4.**  $F : X \times Y \rightarrow V$  belongs to the class  $\mathcal{G}^{1,0}(X \times Y; V)$  if it is continuous, Gâteaux differentiable with respect to  $x$  on  $X \times Y$ , and  $\nabla_x F : X \times Y \rightarrow L(X, V)$  is strongly continuous.

As in Lemma 2.2, the map  $(x, y, h) \mapsto \nabla_x F(x, y)h$  is continuous from  $X \times Y \times X$  to  $V$ , and the chain rules hold. One can also extend Lemma 2.3 in the following way.

**Lemma 2.5.** *A continuous map  $F : X \times Y \rightarrow V$  belongs to  $\mathcal{G}^{1,0}(X \times Y, V)$  provided the following conditions hold:*

- (i) *the directional derivatives  $\nabla_x F(x, y; h)$  exist at every point  $(x, y) \in X \times Y$  and in every direction  $h \in X$ ;*
- (ii) *for every  $h$ , the mapping  $\nabla F(\cdot, \cdot; h) : X \times Y \rightarrow V$  is continuous;*
- (iii) *for every  $(x, y)$ , the mapping  $h \mapsto \nabla_x F(x, y; h)$  is continuous from  $X$  to  $V$ .*

When  $F$  depends on additional arguments, the previous definitions and properties have obvious generalizations. For instance, we say that  $F : X \times Y \times Z \rightarrow V$  belongs to  $\mathcal{G}^{1,1,0}(X \times Y \times Z; V)$  if it is continuous, Gâteaux differentiable with respect to  $x$  and  $y$  on  $X \times Y \times Z$ , and  $\nabla_x F : X \times Y \times Z \rightarrow L(X, V)$  and  $\nabla_y F : X \times Y \times Z \rightarrow L(Y, V)$  are strongly continuous.

### 3 BSDEs with random Lipschitz condition

In this section, we want to study the BSDE

$$Y_t = \xi + \int_t^T f(s, Y_s, Z_s) ds - \int_t^T Z_s dW_s \quad (1)$$

when the generator  $f$  is Lipschitz but with random Lipschitz constants. This kind of BSDEs were also considered in [7] and more recently in [4]. However our framework is different from the setting of the results obtained in these papers. Let us recall that a generator is a random function  $f : [0, T] \times \Omega \times \mathbf{R} \times L_2(\Xi, \mathbf{R}) \rightarrow \mathbf{R}$  which is measurable with respect to  $\mathcal{P} \otimes \mathcal{B}(\mathbf{R}) \otimes \mathcal{B}(\Xi)$  and a terminal condition is simply a real  $\mathcal{F}_T$ -measurable random variable. From now on, we deal only with generators such that,  $\mathbb{P}$ -a.s., for each  $t \in [0, T]$ ,  $(y, z) \rightarrow f(t, y, z)$  is continuous.

By a solution to the BSDE (1) we mean a pair  $(Y, Z) = \{(Y_t, Z_t)\}_{t \in [0, T]}$  of predictable processes with values in  $\mathbf{R} \times L_2(\Xi, \mathbf{R})$  such that  $\mathbb{P}$ -a.s.,  $t \mapsto Y_t$  is continuous,  $t \mapsto Z_t$  belongs to  $L^2(0, T)$ ,  $t \mapsto f(t, Y_t, Z_t)$  belongs to  $L^1(0, T)$  and  $\mathbb{P}$ -a.s.

$$Y_t = \xi + \int_t^T f(s, Y_s, Z_s) ds - \int_t^T Z_s dW_s, \quad 0 \leq t \leq T.$$

We will work with the following assumption on the generator.

**Assumption A1.** There exist a real process  $K$  and a constant  $\alpha \in (0, 1)$  such that  $\mathbb{P}$ -a.s.:

- for each  $t \in [0, T]$ ,  $(y, z) \rightarrow f(t, y, z)$  is continuous ;
- for each  $(t, z) \in [0, T] \times L_2(\Xi, \mathbf{R})$ ,

$$\forall y, p \in \mathbf{R}, \quad (y - p)(f(t, y, z) - f(t, p, z)) \leq K_t^{2\alpha} |y - p|^2$$

- for each  $(t, y) \in [0, T] \times \mathbf{R}$ ,

$$\forall (z, q) \in L_2(\Xi, \mathbf{R}) \times L_2(\Xi, \mathbf{R}), \quad |f(t, y, z) - f(t, y, q)| \leq K_t |z - q|_{L_2(\Xi, \mathbf{R})}.$$

In the classical theory, the process  $K$  is constant but for the application we have in mind we will only assume the following.

**Assumption A2.**  $\{K_s\}_{s \in [0, T]}$  is a predictable real process bounded from below by 1 such that there is a constant  $C$  such that, for any stopping time  $\tau \leq T$ ,

$$\mathbb{E} \left( \int_{\tau}^T |K_s|^2 ds \mid \mathcal{F}_{\tau} \right) \leq C^2.$$

$N$  denotes the smallest constant  $C$  for which the previous statement is true.

This assumption says that, for any  $u \in L_2(\Xi, \mathbf{R})$  such that  $\|u\|_{L_2(\Xi, \mathbf{R})} = 1$  the martingale

$$M_t = \int_0^t K_s u dW_s, \quad 0 \leq t \leq T$$

is a BMO-martingale with  $\|M\|_{BMO_2} = N$ . We refer to [12] for the theory of BMO-martingales and we just recall the properties we will use in the sequel. It follows from the inequality ([12, p. 26]),

$$\forall n \in \mathbf{N}^*, \quad \mathbb{E} [\langle M \rangle_T^n] = \mathbb{E} \left[ \left( \int_0^T |K_s|^2 ds \right)^n \right] \leq n! N^{2n}$$

that  $M$  belongs to  $\mathbb{H}^p$  for all  $p \geq 1$  and moreover

$$\forall \alpha \in (0, 1), \quad \forall p \geq 1, \quad \eta(p)^p := \mathbb{E} \left[ \exp \left( p \int_0^T |K_s|^{2\alpha} ds \right) \right] < +\infty. \quad (2)$$

The very important feature of BMO-martingales is the following: the exponential martingale

$$\mathcal{E}(M)_t = \mathcal{E}_t = \exp \left( \int_0^t K_s u \cdot dW_s - \frac{1}{2} \int_0^t |K_s|^2 ds \right)$$

is a uniformly integrable martingale. More precisely,  $\{\mathcal{E}_t\}_{0 \leq t \leq T}$  satisfies a reverse Hölder inequality. Let  $\Phi$  be the function defined on  $(1, +\infty)$  by

$$\Phi(p) = \left( 1 + \frac{1}{p^2} \log \frac{2p-1}{2(p-1)} \right)^{1/2} - 1;$$

$\Phi$  is nonincreasing with  $\lim_{p \rightarrow 1} \Phi(p) = +\infty$ ,  $\lim_{p \rightarrow +\infty} \Phi(p) = 0$ . Let  $q_*$  be such that  $\Phi(q_*) = N$ . Then, for each  $1 < q < q_*$  and for all stopping time  $\tau \leq T$ ,

$$\mathbb{E} (\mathcal{E}(M)_T^q \mid \mathcal{F}_{\tau}) \leq K(q, N) \mathcal{E}(M)_{\tau}^q \quad (3)$$

where the constant  $K(q, N)$  can be chosen depending only on  $q$  and  $N = \|M\|_{BMO_2}$  e.g.

$$K(q, N) = \frac{2}{1 - 2(q-1)(2q-1)^{-1} \exp(q^2(N^2 + 2N))}.$$

*Remark 3.1.* If we denote  $\mathbb{P}^*$  the probability measure on  $(\Omega, \mathcal{F}_T)$  whose density with respect to  $\mathbb{P}$  is given by  $\mathcal{E}_T$  then  $\mathbb{P}$  and  $\mathbb{P}^*$  are equivalent.

Moreover, it follows from (3) and Hölder's inequality that, if  $X$  belongs to  $L^p(\mathbb{P})$  then  $X$  belongs to  $L^s(\mathbb{P}^*)$  for all  $s < p/p_*$  where  $p_*$  is the conjugate exponent of  $q^*$ .

We assume also some integrability conditions on the data. For this, let  $p_*$  be the conjugate exponent of  $q_*$ .

**Assumption A3.** There exists  $p^* > p_*$  such that

$$\mathbb{E} \left[ |\xi|^{p^*} + \left( \int_0^T |f(s, 0, 0)| ds \right)^{p^*} \right] < +\infty.$$

As usual for BSDEs, we begin with some apriori estimate. The first one shows that, one can control the process  $Y$  as soon as the process  $Z$  has some integrability property. The following lemma relies heavily on the reverse Hölder's inequality.

**Lemma 3.2.** *Let the assumptions A1, A2 and A3 hold. If  $(Y, Z)$  is a solution to (1) such that, for some  $r > p_*$ ,  $Z \in M^r$ , then, for each  $p \in (p_*, p^*)$ ,  $Y \in \mathcal{S}^p$  and*

$$\|Y\|_{\mathcal{S}^p} \leq C \left\| |\xi| + \int_0^T |f(s, 0, 0)| ds \right\|_{p^*},$$

for a suitable constant  $C$  depending on  $p$ ,  $p^*$ ,  $p_*$  and  $N$ .

*Proof.* The starting point to obtain this estimate is a linearization of the generator of the BSDE (1). Let us set

$$a_s = \frac{f(s, Y_s, Z_s) - f(s, 0, Z_s)}{Y_s}, \quad b_s = \frac{f(s, 0, Z_s) - f(s, 0, 0)}{|Z_s|_{L_2(\Xi, \mathbf{R})}^2} Z_s.$$

Then,  $(Y, Z)$  solves the linear BSDE

$$Y_t = \xi + \int_t^T (f(s, 0, 0) + a_s Y_s + \langle b_s, Z_s \rangle_{L_2(\Xi, \mathbf{R})}) ds - \int_t^T Z_s dW_s.$$

As usual, let us set  $e_t = e^{\int_0^t a_s ds}$ . We have,

$$e_t Y_t = e_T \xi + \int_t^T e_s f(s, 0, 0) ds - \int_t^T e_s Z_s \cdot dW_s^*,$$

where we have set  $W_s^* = W_s - \int_0^s b_r dr$ . Of course, we want to take the conditional expectation of the previous equality with respect to the probability  $\mathbb{P}^*$  whose density is

$$\mathcal{E}(I(b))_T = \exp \left( \int_0^T b_s dW_s - \frac{1}{2} \int_0^T |b_s|_{L_2(\Xi, \mathbf{R})}^2 ds \right)$$

under which  $B^*$  is a Brownian motion. To do this, let us observe that  $|b_s|_{L_2(\Xi, \mathbf{R})} \leq K_s$  so that  $\|I(b)\|_{BMO_2} \leq \|M\|_{BMO_2}$  and  $\mathcal{E}(I(b))$  satisfies the reverse Hölder inequality (3) for all  $q < q_*$  (with the same constant).

Moreover, it follows from A1 that  $a_s \leq K_s^{2\alpha}$  and, in particular, (2) says that the process  $e$  belongs to all  $\mathcal{S}^p$  spaces. Thus  $e_T \xi$  belongs to  $L^p$  for all  $p < p_*$  and the same is true for  $\int_0^T e_s |f(s, 0, 0)| ds$ . In the same way, we have, for all  $\rho < r$ ,

$$\mathbb{E} \left[ \left( \int_0^T e_s^2 |Z_s|^2 ds \right)^{\rho/2} \right] \leq \mathbb{E} \left[ \sup e_t^\rho \left( \int_0^T |Z_s|^2 ds \right)^{\rho/2} \right] < +\infty.$$

Using Lemma 3.1, we deduce that  $e_T \xi$  and  $\int_0^T e_s |f(s, 0, 0)| ds$  belongs to  $L^p(\mathbb{P}^*)$  for all  $p < p^*/p_*$  and  $\left( \int_0^T |Z_s|^2 ds \right)^{1/2}$  belongs to  $L^s$  for all  $s < r/p_*$ .

Thus we can take the conditional expectation to obtain

$$e_t Y_t = \mathbb{E}^* \left( e_T \xi + \int_t^T e_s f(s, 0, 0) ds \mid \mathcal{F}_t \right),$$

and, as a byproduct of this equality, we get

$$|Y_t| \leq (\mathcal{E}_t)^{-1} \mathbb{E} \left( \mathcal{E}_T \left( |\xi| e_T / e_t + \int_t^T |f(s, 0, 0)| e_s / e_t ds \right) \mid \mathcal{F}_t \right).$$

Taking into account A1, we have  $a_s \leq K_s^{2\alpha}$  and, for all  $s > t$ ,

$$e_s / e_t \leq \exp \left( \int_t^s K_r^{2\alpha} dr \right) \leq \exp \left( \int_0^T K_r^{2\alpha} dr \right),$$

from which we deduce the inequality

$$|Y_t| \leq (\mathcal{E}_t)^{-1} \mathbb{E} (\mathcal{E}_T \Gamma_T X \mid \mathcal{F}_t),$$

where we have set

$$\Gamma_T = \exp \left( \int_0^T K_r^{2\alpha} dr \right), \quad \text{and} \quad X = \left( |\xi| + \int_0^T |f(s, 0, 0)| ds \right).$$

Using the reverse Hölder inequality, for each  $r > p_*$ , we have,  $q = r/(r-1) < q_*$  and

$$|Y_t| \leq (\mathcal{E}_t)^{-1} \mathbb{E} (\mathcal{E}_T^q \mid \mathcal{F}_t)^{1/q} \mathbb{E} (\Gamma_T^r X^r \mid \mathcal{F}_t)^{1/r} \leq K(q, N)^{1/q} \mathbb{E} (\Gamma_T^r X^r \mid \mathcal{F}_t)^{1/r}$$

Doob's inequality gives for all  $p_* < r < p$ ,

$$\mathbb{E} \left[ \sup_{t \in [0, T]} |Y_t|^p \right] \leq K(q, N)^{p/q} \left( \frac{p}{p-r} \right)^{p/r} \mathbb{E} [\Gamma_T^p X^p].$$

Now, let  $p \in (p_*, p^*)$ , from Hölder inequality, we have, for each  $p_* < r < p$ ,

$$\mathbb{E} \left[ \sup_{t \in [0, T]} |Y_t|^p \right] \leq K(q, N)^{p/q} \left( \frac{p}{p-r} \right)^{p/r} \eta (pp^*/(p^* - p))^p \mathbb{E} [X^{p^*}]^{p/p^*}.$$

It follows that, for  $p_* < r < p < p^*$ ,

$$\|Y\|_{\mathcal{S}^p} \leq K \left( \frac{r}{r-1}, N \right)^{(r-1)/r} \left( \frac{p}{p-r} \right)^{1/r} \eta \left( \frac{pp^*}{p^* - p} \right) \left\| |\xi| + \int_0^T |f(s, 0, 0)| ds \right\|_{p^*},$$

which gives the result taking  $r = (p + p_*)/2$ . □

We keep on by showing that one can obtain an estimate for the process  $Z$  in terms of the norm of  $Y$ . This kind of results is quite classical see e.g. [2]. We give the proof in our framework for the ease of the reader.

**Lemma 3.3.** *Let us assume that*

$$y \cdot f(t, y, z) \leq |y|f_t + K_t^{2\alpha}|y|^2 + K_t|y||z|$$

for nonnegative processes  $f$  and  $K$ .

If  $(Y, Z)$  solves the BSDE (1), with  $Y \in \mathcal{S}^q$  then, for each  $p < q$ ,  $Z \in \mathbb{M}^p$  and

$$\|Z\|_{\mathbb{M}^p} \leq C \left( \|Y\|_{\mathcal{S}^p} + \left\| \int_0^T f_s ds \right\|_p + \|Y\|_{\mathcal{S}^q} \left\| \left( \int_0^T (K_s^{2\alpha} + K_s^2) ds \right)^{1/2} \right\|_{pq/(q-p)} \right),$$

where  $C$  depends only on  $p$  and  $q$ .

*Proof.* We follow [2]. For each integer  $n \geq 1$ , let us introduce the stopping time

$$\tau_n = \inf \left\{ t \in [0, T], \int_0^t |Z_r|^2 dr \geq n \right\} \wedge T.$$

Itô's formula gives us,

$$|Y_0|^2 + \int_0^{\tau_n} |Z_r|^2 dr = |Y_{\tau_n}|^2 + 2 \int_0^{\tau_n} \langle Y_r, f(r, Y_r, Z_r) \rangle dr - 2 \int_0^{\tau_n} \langle Y_r, Z_r dW_r \rangle.$$

But, from the assumption on  $f$ , we have,

$$2y \cdot f(r, y, z) \leq 2|y|f_r + 2K_r^{2\alpha}|y|^2 + 2K_r^2|y|^2 + |z|^2/2.$$

Thus, since  $\tau_n \leq T$ , we deduce that

$$\frac{1}{2} \int_0^{\tau_n} |Z_r|^2 dr \leq Y_*^2 + 2Y_* \int_0^T f_r dr + 2Y_*^2 \int_0^T (K_r^{2\alpha} + K_r^2) dr + 2 \left| \int_0^{\tau_n} \langle Y_r, Z_r dW_r \rangle \right|.$$

It follows that

$$\int_0^{\tau_n} |Z_r|^2 dr \leq 4 \left( Y_*^2 + \left( \int_0^T f_r dr \right)^2 + Y_*^2 \int_0^T (K_r^{2\alpha} + K_r^2) dr + \left| \int_0^{\tau_n} \langle Y_r, Z_r dW_r \rangle \right| \right)$$

and thus that

$$\begin{aligned} & \left( \int_0^{\tau_n} |Z_r|^2 dr \right)^{p/2} \\ & \leq c_p \left( Y_*^p + \left( \int_0^T f_r dr \right)^p + Y_*^p \left( \int_0^T (K_r^{2\alpha} + K_r^2) dr \right)^{p/2} + \left| \int_0^{\tau_n} \langle Y_r, Z_r dW_r \rangle \right|^{p/2} \right). \end{aligned} \quad (4)$$

But by the BDG inequality, we get

$$c_p \mathbb{E} \left[ \left| \int_0^{\tau_n} \langle Y_r, Z_r dW_r \rangle \right|^{p/2} \right] \leq d_p \mathbb{E} \left[ \left( \int_0^{\tau_n} |Y_r|^2 |Z_r|^2 dr \right)^{p/4} \right] \leq d_p \mathbb{E} \left[ Y_*^{p/2} \left( \int_0^{\tau_n} |Z_r|^2 dr \right)^{p/4} \right],$$

and thus

$$c_p \mathbb{E} \left[ \left| \int_0^{\tau_n} \langle Y_r, Z_r dW_r \rangle \right|^{p/2} \right] \leq \frac{d_p^2}{2} \mathbb{E} [Y_*^p] + \frac{1}{2} \mathbb{E} \left[ \left( \int_0^{\tau_n} |Z_r|^2 dr \right)^{p/2} \right].$$

Coming back to the estimate (4), we get, for each  $n \geq 1$ ,

$$\mathbb{E} \left[ \left( \int_0^{\tau_n} |Z_r|^2 dr \right)^{p/2} \right] \leq C_p \mathbb{E} \left[ Y_*^p + \left( \int_0^T f_r dr \right)^p + Y_*^p \left( \int_0^T (K_s^{2\alpha} + K_s^2) ds \right)^{p/2} \right]$$

and, Fatou's lemma implies that

$$\mathbb{E} \left[ \left( \int_0^T |Z_r|^2 dr \right)^{p/2} \right] \leq C_p \mathbb{E} \left[ Y_*^p + \left( \int_0^T f_r dr \right)^p + Y_*^p \left( \int_0^T (K_s^{2\alpha} + K_s^2) ds \right)^{p/2} \right].$$

The result follows from Hölder's inequality.  $\square$

The previous two lemmas lead the following result.

**Corollary 3.4.** *Let the assumptions A1, A2 and A3 hold. If  $(Y, Z)$  is a solution to (1) such that, for some  $r > p_*$ ,  $Y \in \mathcal{S}^r$ , then, for each  $p \in (p_*, p^*)$ ,  $(Y, Z) \in \mathcal{S}^p \times \mathcal{M}^p$  and*

$$\|Y\|_{\mathcal{S}^p} + \|Z\|_{\mathcal{M}^p} \leq C \left\| \|\xi\| + \int_0^T |f(s, 0, 0)| ds \right\|_{p^*} \left( 1 + \left\| \left( \int_0^T (K_s^{2\alpha} + K_s^2) ds \right)^{1/2} \right\|_{p(p^*+p)/(p^*-p)} \right)$$

where  $C$  depends on  $p$ ,  $p_*$ ,  $p^*$  and  $N$ .

*Proof.* Since  $Y$  belongs to  $\mathcal{S}^p$  for some  $p > p_*$ , there exists by Lemma 3.3  $r \in (p_*, p^*)$  such that  $Z$  belongs to  $\mathcal{M}^r$ . It follows from Lemma 3.2 that  $Y$  belongs to  $\mathcal{S}^p$  for all  $p < p^*$  and then by Lemma 3.3  $Z \in \mathcal{M}^p$  for all  $p < p^*$ .

The inequality comes from the choice  $q = (p+p^*)/2$  in Lemma 3.3 together with the estimate of Lemma 3.2.  $\square$

**Assumption A4.** There exists a nonnegative predictable process  $f$  such that,

$$\mathbb{E} \left[ \left( \int_0^T f(s) ds \right)^{p^*} \right] < +\infty$$

and  $\mathbb{P}$ -a.s.

$$\forall (t, y, z) \in [0, T] \times \mathbf{R} \times L_2(\Xi, \mathbf{R}), \quad |f(t, y, z)| \leq f(t) + K_t^{2\alpha} |y| + K_t |z|.$$

**Theorem 3.5.** *Let the assumptions A1, A2, A3 and A4 hold. Then BSDE (1) has a unique solution  $(Y, Z)$  which belongs to  $\mathcal{S}^p \times \mathcal{M}^p$  for all  $p < p^*$ .*

*Proof.* Let us prove first uniqueness. Let  $(Y^1, Z^1)$  and  $(Y^2, Z^2)$  be solutions to (1) such that  $Y^1$  and  $Y^2$  belongs to  $\mathcal{S}^p$  for  $p > p_*$ . The by Corollary 3.4,  $(Y^1, Z^1)$  and  $(Y^2, Z^2)$  belongs to  $\mathcal{S}^p \times \mathcal{M}^p$  for all  $p < p_*$ . Moreover,  $U = Y^1 - Y^2$  and  $V = Z^1 - Z^2$  solves the BSDE

$$U_t = \int_t^T F(s, U_s, V_s) ds - \int_t^T V_s \cdot dW_s,$$

where  $F(t, u, v) = f(t, Y_t^2 + u, Z_t^2 + v) - f(t, Y_t^2, Z_t^2)$ . We have  $F(t, 0, 0) = 0$  and  $F$  satisfies A1 with the same process  $K$ . It follows from Corollary 3.4 that  $(U, V) \equiv (0, 0)$ .

Let us turn to existence. For each integer  $n \geq 1$ , let  $\tau_n$  be the following stopping time:

$$\tau_n = \inf \left\{ t \in [0, T] : \int_0^t (f(s) + K_s^2) ds \geq n \right\} \wedge T.$$

Let  $\xi^n = \xi \mathbf{1}_{|\xi| \leq n}$  and  $(Y^n, Z^n)$  be the solution to the BSDE

$$Y_t^n = \xi^n + \int_t^T \mathbf{1}_{s \leq \tau_n} f(s, Y_s^n, Z_s^n) ds - \int_t^T Z_s^n dW_s.$$

The existence of the solution  $(Y^n, Z^n)$  to the previous equation comes from [16]. Indeed, we have, setting  $f^n(t, y, z) = \mathbf{1}_{t \leq \tau_n} f(t, y, z)$ ,

$$|f^n(t, y, z)| \leq \mathbf{1}_{t \leq \tau_n} (f(t) + K_t^{2\alpha} + K_t^2/2) (1 + |y|) + |z|^2/2,$$

and,  $\mathbb{P}$ -a.s.

$$\int_0^T \mathbf{1}_{t \leq \tau_n} (f(t) + K_t^{2\alpha} + K_t^2/2) dt \leq 5n/2.$$

Since  $\xi^n$  is bounded by  $n$ , the previous BSDE has a unique solution  $(Y^n, Z^n)$  such that  $Y^n$  is a bounded process and  $Z^n \in M^2$ . Since

$$\int_0^T |f^n(t, 0, 0)| dt \leq n,$$

we know, from Corollary 3.4, that  $(Y^n, Z^n) \in \mathcal{S}^p \times M^p$  for all  $p$ .

Moreover, still by Corollary 3.4, the sequence  $((Y^n, Z^n))_{n \geq 1}$  is bounded in  $\mathcal{K}^p := \mathcal{S}^p \times M^p$  for all  $p < p^*$ .

Let us show that  $((Y^n, Z^n))_{n \geq 1}$  is a Cauchy sequence in  $\mathcal{K}^p := \mathcal{S}^p \times M^p$  for all  $p < p^*$ . Let  $m > n \geq 1$  and let us set as before  $U = Y^m - Y^n$ ,  $V = Z^m - Z^n$ . Then  $(U, V)$  solves the BSDE

$$U_t = \xi^m - \xi^n + \int_t^T F(s, U_s, V_s) ds - \int_t^T V_s dW_s$$

where

$$F(t, u, v) = \mathbf{1}_{t \leq \tau_m} (f(t, u + Y_t^n, v + Z_t^n) - f(t, Y_t^n, Z_t^n)) - \mathbf{1}_{\tau_n < t \leq \tau_m} f(t, Y_t^n, Z_t^n).$$

$F$  satisfies A1 and  $F(t, 0, 0) = -\mathbf{1}_{\tau_n < t \leq \tau_m} f(t, Y_t^n, Z_t^n)$  belongs to  $L^p$  for all  $p \geq 1$ .

Since  $\xi \in L^{p^*}$ ,  $\|\xi^m - \xi^n\|_{p^*} \rightarrow 0$  if  $n \rightarrow \infty$ . Moreover, we have, from A4 and Hölder inequality,

$$\int_0^T |F(t, 0, 0)| dt \leq \int_{\tau_n}^T f(t) dt + \sup_t |Y_t^n| \int_{\tau_n}^T K_t^{2\alpha} dt + \left( \int_{\tau_n}^T K_t^2 dt \right)^{1/2} \left( \int_0^T |Z_t^n|^2 dt \right)^{1/2}.$$

Let  $p < p^*$ . We choose  $p < q < r < p^*$ . It follows from the previous inequality, using Hölder inequality, that

$$\left\| \int_0^T |F(t, 0, 0)| dt \right\|_q \leq \left\| \int_{\tau_n}^T f(t) dt \right\|_q + \|Y^n\|_{\mathcal{S}^r} \left\| \int_{\tau_n}^T K_t^{2\alpha} dt \right\|_{\frac{qr}{r-q}} + \|Z^n\|_{M^r} \left\| \left( \int_{\tau_n}^T K_t^2 dt \right)^{1/2} \right\|_{\frac{qr}{r-q}}.$$

Let us recall that  $\tau_n \rightarrow T$   $\mathbb{P}$ -a.s and that the sequence  $((Y^n, Z^n))_{n \geq 1}$  is bounded in  $\mathcal{K}^r$ . Since  $\int_0^T f(t) dt$  belongs to  $L^{p^*}$ ,  $\int_0^T K_t^{2\alpha} dt$  and  $\int_0^T K_t^2 dt$  has moments of all order, the right hand side of the previous inequality tends to 0 as  $n$  tends to infinity.

It follows from Corollary 3.4 – applied with  $q$  instead of  $p^*$  – that  $((Y^n, Z^n))_{n \geq 1}$  is a Cauchy sequence in  $\mathcal{K}^p$  and this is valid as soon as  $p < p^*$ .

It is easy to check that the limit of this sequence is a solution to BSDE (1) □

## 4 The forward-backward system

In this section, we apply the previous results on BSDEs to study the differentiability of the solution to the following quadratic BSDE

$$Y_\tau^{t,x} = \Phi(X_T^{t,x}) + \int_\tau^T F(r, X_r^{t,x}, Y_r^{t,x}, Z_r^{t,x}) dr - \int_\tau^T Z_r^{t,x} dW_r, \quad 0 \leq \tau \leq T, \quad (5)$$

where  $\{X_\tau^{t,x}\}_{0 \leq \tau \leq T}$  is the solution to

$$X_\tau^{t,x} = e^{(\tau-t)A}x + \int_t^\tau e^{(\tau-r)A}b(r, X_r^{t,x}) dr + \int_t^\tau e^{(\tau-r)A}\sigma(r, X_r^{t,x}) dW_r, \quad t \leq \tau \leq T. \quad (6)$$

As usual, we have set  $X_\tau^{t,x} = x$  for  $\tau < t$ . Of course, from Itô's formula, we have

$$dX_\tau^{t,x} = AX_\tau^{t,x} d\tau + b(\tau, X_\tau^{t,x}) d\tau + \sigma(\tau, X_\tau^{t,x}) dW_\tau, \quad \tau \in [t, T], \quad X_\tau^{t,x} = x \in H, \quad \tau \leq t.$$

But a solution of this equation is always understood as an  $(\mathcal{F}_t)$ -predictable continuous process  $X$  solving (6).

We will work under the following assumption on the diffusion coefficients.

**Assumption A5. (i)** The operator  $A$  is the generator of a strongly continuous semigroup  $e^{tA}$ ,  $t \geq 0$ , in the Hilbert space  $H$ .

**(ii)** The mapping  $b : [0, T] \times H \rightarrow H$  is measurable and satisfies, for some constant  $L > 0$ ,

$$\begin{aligned} |b(t, x) - b(t, y)| &\leq L|x - y|, & t \in [0, T], \quad x, y \in H, \\ |b(t, x)| &\leq L(1 + |x|), & t \in [0, T], \quad x \in H. \end{aligned}$$

**(iii)**  $\sigma : [0, T] \times H \rightarrow L(\Xi, H)$  is such that, for every  $v \in \Xi$ , the map  $\sigma v : [0, T] \times H \rightarrow H$  is measurable,  $e^{sA}\sigma(t, x) \in L_2(\Xi, H)$  for every  $s > 0$ ,  $t \in [0, T]$  and  $x \in H$ , and

$$\begin{aligned} |e^{sA}\sigma(t, x)|_{L_2(\Xi, H)} &\leq L s^{-\gamma}(1 + |x|), \\ |e^{sA}\sigma(t, x) - e^{sA}\sigma(t, y)|_{L_2(\Xi, H)} &\leq L s^{-\gamma}|x - y|, \\ |\sigma(t, x)|_{L(\Xi, H)} &\leq L(1 + |x|), \end{aligned}$$

for some constants  $L > 0$  and  $\gamma \in [0, 1/2)$ .

**(iv)** For every  $s > 0$ ,  $t \in [0, T]$ ,

$$b(t, \cdot) \in \mathcal{G}^1(H, H), \quad e^{sA}\sigma(t, \cdot) \in \mathcal{G}^1(H, L_2(\Xi, H)).$$

A consequence of the previous assumptions is that, for every  $s > 0$ ,  $t \in [0, T]$ ,  $x, h \in H$ ,

$$|\nabla_x b(t, x)h| \leq L|h|, \quad |\nabla_x(e^{sA}\sigma(t, x))h|_{L_2(\Xi, H)} \leq L s^{-\gamma}|h|.$$

The following results are proved by Fuhrman and Tessitore in [9].

**Proposition 4.1.** *Let A5 hold. Then, for each  $(t, x) \in [0, T] \times H$ , (6) has a unique solution  $\{X_\tau^{t,x}\}_{0 \leq \tau \leq T}$ . Moreover, for every  $p > 1$ ,*

(i)  $X^{t,x}$  belongs to  $\mathcal{S}^p(H)$  and there exists a constant  $C$  such that

$$\mathbb{E} \left[ \sup_{\tau \in [0, T]} |X_\tau^{t,x}|^p \right] \leq C(1 + |x|)^p, \quad (7)$$

(ii) The map  $(t, x) \mapsto X^{t,x}$  belongs to  $\mathcal{G}^{0,1}([0, T] \times H, \mathcal{S}^p(H))$ .

(iii) For every  $h \in H$ , the directional derivative process  $\nabla_x X_\tau^{t,x} h$ ,  $\tau \in [0, T]$ , solves the equation:

$$\begin{cases} \nabla_x X_\tau^{t,x} h &= e^{(\tau-t)A} h + \int_t^\tau e^{(\tau-r)A} \nabla_x b(r, X_r^{t,x}) \nabla_x X_r^{t,x} h \, dr \\ &+ \int_t^\tau \nabla_x (e^{(\tau-r)A} \sigma(r, X_r^{t,x})) \nabla_x X_r^{t,x} h \, dW_r, \quad \tau \in [t, T], \\ \nabla_x X_\tau^{t,x} h &= h, \quad \tau \in [0, t). \end{cases}$$

(iii) Finally  $\left\| \nabla_x X_\tau^{t,x} h \right\|_{\mathcal{S}^p} \leq c|h|$  for some constant  $c$ .

We assume that  $F : [0, T] \times H \times \mathbf{R} \times L_2(\Xi, \mathbf{R}) \rightarrow \mathbf{R}$  and  $\Phi : H \rightarrow \mathbf{R}$  are measurable functions such that

**Assumption A6.** There exists  $C \geq 0$  and  $\alpha \in (0, 1)$  such that

- $|F(t, x, y, z)| \leq C(1 + |y| + |z|^2)$  and  $\Phi$  is bounded ;
- $F(s, \cdot, \cdot, \cdot)$  is  $\mathcal{G}^{1,1,1}(H \times \mathbf{R} \times L_2(\Xi, \mathbf{R}); \mathbf{R})$  and  $\Phi$  is  $\mathcal{G}^1(H; \mathbf{R})$  ;
- $|\nabla_x \Phi(x)| \leq C(1 + |x|^n)$  ;
- $|\nabla_x F(s, x, y, z)| \leq C(1 + |x|^n + |z|^2)$  ;
- $|\nabla_z F(s, x, y, z)| \leq C(1 + |z|)$  ;
- $|\nabla_y F(s, x, y, z)| \leq C(1 + |z|)^{2\alpha}$  ;

We know from results of [13, 14] (these results can be easily generalised to the case of a cylindrical Wiener process) that under A6 the BSDE (5) has a unique bounded solution and that there exists a constant  $C$  such that, for each  $(t, x)$ ,

$$\left\| \sup_{u \in [0, T]} |Y_u^{t,x}| \right\|_\infty + \left\| \int_0^\cdot Z_s^{t,x} \cdot dW_s \right\|_{BMO_2} \leq C. \quad (8)$$

For the existence and the bound for the process  $Y$  we refer to [14, Corollary 1], uniqueness follows from [13, Theorem 2.6] and finally the estimate for the BMO-norm of  $Z$  comes from a direct computation starting from Itô's formula applied to  $\varphi(x) = (e^{2Cx} - 2Cx - 1)/(2C^2)$ . In particular, for each  $p \geq 1$ ,

$$\left\| \left( \int_0^T |Z_s^{t,x}|^2 \, ds \right)^{1/2} \right\|_p \leq C_p. \quad (9)$$

**Proposition 4.2.** *Let the assumption A6 hold.*

The map  $(t, x) \mapsto (Y^{t,x}, Z^{t,x})$  belongs to  $\mathcal{G}^{0,1}([0, T] \times H; \mathcal{S}^p \times \mathcal{M}^p)$  for each  $p > 1$ . Moreover, for every  $x \in H$  and  $h \in H$ , the directional derivative process  $\left\{ \nabla_x Y_u^{t,x} h, \nabla_x Z_u^{t,x} h \right\}_{u \in [0, T]}$  solves the BSDE: for  $\tau \in [0, T]$ ,

$$\begin{aligned} \nabla_x Y_u^{t,x} h &= \nabla_x \Phi(X_T^{t,x}) \nabla_x X_T^{t,x} h + \int_u^T \nabla_x F(s, X_s^{t,x}, Y_s^{t,x}, Z_s^{t,x}) \nabla_x X_s^{t,x} h ds \\ &+ \int_u^T (\nabla_y F(s, X_s^{t,x}, Y_s^{t,x}, Z_s^{t,x}) \nabla_x Y_s^{t,x} h + \nabla_z F(s, X_s^{t,x}, Y_s^{t,x}, Z_s^{t,x}) \nabla_x Z_s^{t,x} h) ds \\ &- \int_u^T \nabla_x Z_s^{t,x} h dW_s \end{aligned} \quad (10)$$

and there exists  $C_p$  such that

$$\|\nabla_x Y^{t,x} h\|_{\mathcal{S}^p} + \|\nabla_x Z^{t,x} h\|_{\mathcal{M}^p} \leq C_p(1 + |x|^n) |h|.$$

*Proof.* The continuity of the map  $(t, x) \mapsto (Y^{t,x}, Z^{t,x})$  follows from a mere extension of Kobylanski's stability result [13, Theorem 2.8].

For the differentiability, let us remark that, in view of A6 and (9), for all  $p > 1$ ,

$$\left\| \left| \nabla_x \Phi(X_T^{t,x}) \nabla_x X_T^{t,x} h \right| + \int_0^T |\nabla_x F(s, X_s^{t,x}, Y_s^{t,x}, Z_s^{t,x}) \nabla_x X_s^{t,x} h| ds \right\|_p \leq C_p(1 + |x|^n) |h|.$$

It follows from Theorem 3.5, that the BSDE (10) has a unique solution which belongs to  $\mathcal{S}^p \times \mathcal{M}^p$  for all  $p \geq 1$ . And moreover, for  $p > 1$ , it follows from Corollary 3.4 and (9), that

$$\|\nabla_x Y^{t,x} h\|_{\mathcal{S}^p} + \|\nabla_x Z^{t,x} h\|_{\mathcal{M}^p} \leq C(1 + |x|^n) |h|.$$

Let us fix  $(t, x) \in [0, T] \times H$ . We remove the parameters  $t$  and  $x$  for notational simplicity. For  $\varepsilon > 0$ , we set  $X^\varepsilon = X^{t,x+\varepsilon h}$ , where  $h$  is some vector in  $H$ , and we consider  $(Y^\varepsilon, Z^\varepsilon)$  the solution in  $\mathcal{S}^p \times \mathcal{M}^p$  to the BSDE

$$Y_t^\varepsilon = \Phi(X_T^{t,\varepsilon}) + \int_t^T F(s, X_s^\varepsilon, Y_s^\varepsilon, Z_s^\varepsilon) ds - \int_t^T Z_s^\varepsilon dW_s.$$

When  $\varepsilon \rightarrow 0$ ,  $(X^\varepsilon, Y^\varepsilon, Z^\varepsilon) \rightarrow (X, Y, Z)$  in  $\mathcal{S}^p \times \mathcal{S}^p \times \mathcal{M}^p$  for all  $p > 1$ . We also denote  $(G, N)$  the solution to the BSDE (10) and it remains to prove that the directional derivative of the map  $(t, x) \mapsto (Y^{t,x}, Z^{t,x})$  in the direction  $h \in H$  is given by  $(G, N)$ .

Let us consider  $U^\varepsilon = \varepsilon^{-1}(Y^\varepsilon - Y) - G$ ,  $V^\varepsilon = \varepsilon^{-1}(Z^\varepsilon - Z) - N$ . We have,

$$\begin{aligned} U_t^\varepsilon &= \frac{1}{\varepsilon} (\Phi(X_T^\varepsilon) - \Phi(X_T)) - \nabla_x \Phi(X_T) \nabla_x X_T h + \\ &+ \frac{1}{\varepsilon} \int_t^T (F(s, X_s^\varepsilon, Y_s^\varepsilon, Z_s^\varepsilon) - F(s, X_s, Y_s, Z_s)) ds - \int_t^T V_s^\varepsilon dW_s \\ &- \int_t^T \nabla_x F(s, X_s, Y_s, Z_s) \nabla_x X_s h ds - \int_t^T \nabla_y F(s, X_s, Y_s, Z_s) G_s ds \\ &- \int_t^T \nabla_z F(s, X_s, Y_s, Z_s) N_s ds. \end{aligned}$$

Using the fact that  $\psi(s, \cdot, \cdot, \cdot)$  belongs to  $\mathcal{G}^{1,1,1}$ , we can write

$$\begin{aligned} \frac{1}{\varepsilon} (F(s, X_s^\varepsilon, Y_s^\varepsilon, Z_s^\varepsilon) - F(s, X_s, Y_s, Z_s)) &= \frac{1}{\varepsilon} (F(s, X_s^\varepsilon, Y_s, Z_s) - F(s, X_s, Y_s, Z_s)) + \\ &+ A_s^\varepsilon \frac{Y_s^\varepsilon - Y_s}{\varepsilon} + B_s^\varepsilon \frac{Z_s^\varepsilon - Z_s}{\varepsilon} \end{aligned}$$

where  $A_s^\varepsilon \in L(\mathbf{R}, \mathbf{R})$  and  $B_s^\varepsilon \in L(L_2(\Xi, \mathbf{R}), \mathbf{R})$  are defined by

$$\begin{aligned} \forall y \in \mathbf{R}, \quad A_s^\varepsilon y &= \int_0^1 \nabla_y F(s, X_s^\varepsilon, Y_s + \alpha(Y_s^\varepsilon - Y_s), Z_s) y \, d\alpha, \\ \forall z \in L_2(\Xi, \mathbf{R}), \quad B_s^\varepsilon z &= \int_0^1 \nabla_z F(s, X_s^\varepsilon, Y_s^\varepsilon, Z_s + \alpha(Z_s^\varepsilon - Z_s)) z \, d\alpha. \end{aligned}$$

Then  $(U^\varepsilon, V^\varepsilon)$  solves the following BSDE

$$U_t^\varepsilon = \zeta^\varepsilon + \int_t^T (A_s^\varepsilon U_s^\varepsilon + B_s^\varepsilon V_s^\varepsilon) \, ds + \int_t^T (P^\varepsilon(s) + Q^\varepsilon(s) + R^\varepsilon(s)) \, ds - \int_t^T V_s^\varepsilon \, dW_s$$

where we have set

$$\begin{aligned} P^\varepsilon(s) &= (A_s^\varepsilon - \nabla_y F(s, X_s, Y_s, Z_s)) G_s, \quad Q^\varepsilon(s) = (B_s^\varepsilon - \nabla_z F(s, X_s, Y_s, Z_s)) N_s, \\ R^\varepsilon(s) &= \varepsilon^{-1} (F(s, X_s^\varepsilon, Y_s, Z_s) - F(s, X_s, Y_s, Z_s)) - \nabla_x F(s, X_s, Y_s, Z_s) \nabla_x X_s h, \\ \zeta^\varepsilon &= \varepsilon^{-1} (\Phi(X_T^\varepsilon) - \Phi(X_T)) - \nabla_x \Phi(X_T) \nabla_x X_T h. \end{aligned}$$

It follows from A6 that

$$A_s^\varepsilon \leq C(1 + |Z_s| + |Z_s^\varepsilon|)^{2\alpha}, \quad |B_s^\varepsilon| \leq C(1 + |Z_s| + |Z_s^\varepsilon|),$$

and

$$|P^\varepsilon(s)| \leq C(1 + |Z_s| + |Z_s^\varepsilon|)^{2\alpha} |G_s|, \quad |Q^\varepsilon(s)| \leq C(1 + |Z_s| + |Z_s^\varepsilon|) |H_s|$$

For  $p$  large enough, we have from Corollary 3.4 taking into account (8) and (9),

$$\|U^\varepsilon\|_{S^p} + \|V^\varepsilon\|_{M^p} \leq C \left\| \left| \zeta^\varepsilon \right| + \int_0^T (|P^\varepsilon(s)| + |Q^\varepsilon(s)| + |R^\varepsilon(s)|) \, ds \right\|_{p+1}.$$

The right hand side of the previous inequality tends to 0 as  $\varepsilon \rightarrow 0$  in view of the regularity and the growth of  $F$  and  $\Phi$  (see A6).

The proof that the maps  $x \mapsto (\nabla_x Y^{t,x} h, \nabla_x Z^{t,x} h)$  and  $h \mapsto (\nabla_x Y^{t,x} h, \nabla_x Z^{t,x} h)$  are continuous (for every  $h$  and  $x$  respectively) comes once again of Corollary 3.4.  $\square$

*Remark 4.3.* Since  $\sup_{t,x} \|\sup_u |Y(u, t, x)|\|_\infty < \infty$ , one can change  $C$  by  $C(|y|)$  in the assumptions on the gradient on  $F$  in A6.

## 5 Application to nonlinear PDEs

In this section we are interested in finding a probabilistic representation in our framework for the solution to

$$\begin{cases} \partial_t u(t, x) + \mathcal{L}_t[u(t, \cdot)](x) + F(t, x, u(t, x), \sigma(t, x)^* \nabla_x u(t, x)) = 0, & t \in [0, T], x \in H, \\ u(T, x) = \Phi(x), \end{cases} \quad (11)$$

where  $\mathcal{L}_t$  is the operator:

$$\mathcal{L}_t[\phi](x) = \frac{1}{2} \text{Trace} (\sigma(t, x) \sigma(t, x)^* \nabla^2 \phi(x)) + \langle Ax + b(t, x), \nabla \phi(x) \rangle,$$

where  $\nabla \phi$  and  $\nabla^2 \phi$  are the first and the second Gâteaux derivatives of  $\phi$  (identified with elements of  $H$  and  $L(H)$  respectively). This definition is formal, since the domain of  $\mathcal{L}_t$  is not specified.

We will refer to this equation as the nonlinear Kolmogorov equation. In this equation,  $F : [0, T] \times H \times \mathbf{R} \times \Xi \rightarrow \mathbf{R}$  is a given function verifying A6 and  $\nabla_x u(t, x)$  is the Gâteaux derivative of  $u(t, x)$  with respect to  $x$ : it is identified with an element of  $H$ , so that  $\sigma(t, x)^* \nabla_x u(t, x) \in \Xi$ .

Under the assumption A5, we can define a transition semigroup  $P_{t, \tau}$  with the help of  $X^{t, x}$  solution to (6) by the formula

$$P_{t, \tau}[\phi](x) = \mathbb{E} [\phi(X_\tau^{t, x})], \quad x \in H.$$

The estimate (7) shows that  $P_{t, \tau}$  is well defined as a linear operator from  $\mathcal{B}_p(H)$ , the set of measurable functions from  $H$  to  $\mathbf{R}$  with polynomial growth, into itself; the semigroup property  $P_{t, s} P_{s, \tau} = P_{t, \tau}$ ,  $t \leq s \leq \tau$ , is well known.

When  $\phi$  is sufficiently regular, the function  $v(t, x) = P_{t, T}[\phi](x)$ , is a classical solution of the backward Kolmogorov equation (11) with  $F \equiv 0$ ; we refer to [5] and [22] for a detailed exposition. When  $\phi$  is not regular, the function  $v$  defined by the formula  $v(t, x) = P_{t, T}[\phi](x)$  can be considered as a generalized solution of this equation.

For the nonlinear case, we consider the variation of constants formula for (11):

$$u(t, x) = \int_t^T P_{t, \tau} [F(\tau, \cdot, u(\tau, \cdot), \sigma(\tau, \cdot)^* \nabla_x u(\tau, \cdot))](x) d\tau + P_{t, T}[\Phi](x), \quad t \in [0, T], x \in H, \quad (12)$$

and we notice that this formula is meaningful, provided  $F(t, \cdot, \cdot, \cdot)$ ,  $u(t, \cdot)$  and  $\nabla_x u(t, \cdot)$  have polynomial growth. We use this formula as a definition for the solution of (11):

**Definition 5.1.** We say that a function  $u : [0, T] \times H \rightarrow \mathbf{R}$  is a mild solution of the nonlinear Kolmogorov equation (11) if the following conditions hold:

- (i)  $u \in \mathcal{G}^{0,1}([0, T] \times H, \mathbf{R})$ ;
- (ii) there exists  $C > 0$  and  $d \in \mathbf{N}$  such that  $|\nabla_x u(t, x)h| \leq C|h|(1 + |x|^d)$  for all  $t \in [0, T]$ ,  $x \in H$ ,  $h \in H$ ;
- (iii) equality (12) holds.

*Remark 5.2.* We obtain an equivalent formulation of (11) and (12) by considering the Gâteaux derivative  $\nabla_x u(t, x)$  as an element of  $\Xi^* = L(\Xi, \mathbf{R}) = L_2(\Xi, \mathbf{R})$ . In this case, we take a function  $F : [0, T] \times H \times \mathbf{R} \times L_2(\Xi, \mathbf{R}) \rightarrow \mathbf{R}$  and we write the equation in the form

$$\partial_t u(t, x) + \mathcal{L}_t[u(t, \cdot)](x) + F(t, x, u(t, x), \nabla_x u(t, x) \sigma(t, x)) = 0.$$

The two forms are equivalent provided we identify  $\Xi^* = L_2(\Xi, \mathbf{R})$  with  $\Xi$  by the Riesz isometry.

We are now ready to state the main result of this section.

**Theorem 5.3.** *Let the assumptions A5 and A6 hold.*

*The nonlinear Kolmogorov equation (11) has a unique mild solution  $u$  given by the formula*

$$u(t, x) = Y_t^{t,x}, \quad (t, x) \in [0, T] \times H$$

where  $(Y^{t,x}, Z^{t,x})$  is the solution to the BSDE (5) and  $X^{t,x}$  the solution to (6). Moreover, we have,  $\mathbb{P}$ -a.s.

$$Y_s^{t,x} = u(s, X_s^{t,x}), \quad Z_s^{t,x} \sigma(s, X_s^{t,x})^* \nabla_x u(s, t, X_s^{t,x}).$$

*Proof.* Let us first recall a result of [9, Lemma 6.3]. Let  $\{e_i\}$  be a basis of  $\Xi$  and let us consider the standard real Wiener process  $W_\tau^i = \int_0^\tau \langle e_i, dW_\sigma \rangle$ ,  $\tau \geq 0$ .

If  $v \in \mathcal{G}^{0,1}([0, T] \times H, \mathbf{R})$ , for every  $i$ , the quadratic variation of  $v(s, X_s^{t,x})$  and  $W_s^i$  is given by

$$[v(\cdot, X^{t,x}), W^i]_s = \int_t^s \nabla_x v(\tau, X_\tau^{t,x}) G(\tau, X_\tau^{t,x}) e_i d\tau, \quad s \in [t, T]. \quad (13)$$

(a) *Existence.* Let us recall that for  $s \in [t, T]$ ,  $Y_s^{t,x}$  is measurable with respect to  $\mathcal{F}_{[t,s]}$  and  $\mathcal{F}_s$ ; it follows that  $Y_t^{t,x}$  is deterministic (see also [6]). Moreover, as a byproduct of Proposition 4.2, the function  $u$  defined by the formula  $u(t, x) = Y_t^{t,x}$  has the regularity properties stated in Definition 5.1. It remains to verify that equality (12) holds true for  $u$ .

To this purpose we first fix  $t \in [0, T]$  and  $x \in H$ . Since  $(Y^{t,x}, Z^{t,x})$  solves the BSDE (5), we have, for  $s \in [t, T]$ ,

$$Y_s^{t,x} + \int_s^T Z_\tau^{t,x} dW_\tau = \Phi(X_T^{t,x}) + \int_s^T F(\tau, X_\tau^{t,x}, Y_\tau^{t,x}, Z_\tau^{t,x}) d\tau,$$

and, taking expectation for  $s = t$  we obtain, coming back to the definition of  $u$  and  $P_{t,T}$ ,

$$u(t, x) = P_{t,T}[\Phi](x) + \mathbb{E} \left[ \int_t^T F(\tau, X_\tau^{t,x}, Y_\tau^{t,x}, Z_\tau^{t,x}) d\tau \right]. \quad (14)$$

Moreover, we have, for each  $i$ ,

$$[Y^{t,x}, W^i]_s = \int_t^s \langle Z_\tau, e_i \rangle d\tau, \quad s \in [t, T].$$

Now let us observe that the processes  $Y$  and  $Z$  satisfy the Markov property: for  $t \leq s \leq T$ ,  $\mathbb{P}$ -a.s.

$$\begin{aligned} Y_\tau^{s, X_s^{t,x}} &= Y_\tau^{t,x} \quad \text{for } \tau \in [s, T] \\ Z_\tau^{s, X_s^{t,x}} &= Z_\tau^{t,x} \quad \text{for a.e. } \tau \in [s, T]. \end{aligned}$$

In fact the solution of the backward equation is uniquely determined on an interval  $[s, T]$  by the values of the process  $X$  on the same interval. The process  $X$  is the unique solution of the forward equation (6) and satisfies the Markov property.

As consequence we have,  $\mathbb{P}$ -a.s.,

$$u(\tau, X_\tau^{t,x}) = Y_\tau^{t,x}, \quad \tau \in [t, T].$$

It follows from (13) that, for each  $i$ ,

$$[Y^{t,x}, W^i]_s = \int_t^s \nabla_x u(\tau, X_\tau^{t,x}) \sigma(\tau, X_\tau^{t,x}) e_i d\tau, \quad s \in [t, T].$$

Therefore, for a.a.  $\tau \in [t, T]$ , we have  $\mathbb{P}$ -a.s.

$$\nabla_x u(\tau, X_\tau^{t,x}) \sigma(\tau, X_\tau^{t,x}) e_i = \langle Z_\tau^{t,x}, e_i \rangle,$$

for every  $i$ . Identifying  $\nabla_x u(t, x)$  with an element of  $\Xi$ , we conclude that for a.a.  $\tau \in [t, T]$ ,

$$\sigma(\tau, X_\tau^{t,x})^* \nabla_x u(\tau, t, X_\tau^{t,x}) = Z_\tau^{t,x}.$$

Thus,  $F(\tau, X_\tau^{t,x}, Y_\tau^{t,x}, Z_\tau^{t,x})$  can be rewritten as

$$F(\tau, X_\tau^{t,x}, u(\tau, X_\tau^{t,x}), \sigma(\tau, X_\tau^{t,x})^* \nabla_x u(\tau, X_\tau^{t,x}))$$

and (14) leads to

$$u(t, x) = P_{t,T}[\phi](x) + \int_t^T P_{t,\tau}[F(\tau, \cdot, u(\tau, \cdot), \sigma(\tau, \cdot)^* \nabla_x u(\tau, \cdot))](x) d\tau$$

which is (12).

(b) *Uniqueness.* Let  $u$  be a mild solution. We look for a convenient expression for the process  $u(s, X_s^{t,x})$ ,  $s \in [t, T]$ . By (12) and the definition of  $P_{t,\tau}$ , for every  $s \in [t, T]$  and  $x \in H$ ,

$$\begin{aligned} u(s, x) &= \mathbb{E}[\Phi(X_T^{s,x})] \\ &\quad + \mathbb{E}\left[\int_s^T F(\tau, X_\tau^{t,x}, u(\tau, X_\tau^{t,x}), \sigma(\tau, X_\tau^{t,x})^* \nabla_x u(\tau, X_\tau^{t,x})) d\tau\right]. \end{aligned}$$

Since  $X_\tau^{t,x}$  is independent of  $\mathcal{F}_s$ , we can replace the expectation by the conditional expectation given  $\mathcal{F}_s$ :

$$\begin{aligned} u(s, x) &= \mathbb{E}^{\mathcal{F}_s}[\Phi(X_T^{s,x})] \\ &\quad + \mathbb{E}^{\mathcal{F}_s}\left[\int_s^T F(\tau, X_\tau^{t,x}, u(\tau, X_\tau^{t,x}), \sigma(\tau, X_\tau^{t,x})^* \nabla_x u(\tau, X_\tau^{t,x})) d\tau\right]. \end{aligned}$$

Taking into account the Markov property of  $X$ ,  $\mathbb{P}$ -a.s.

$$X_\tau^{s, X_s^{t,x}} = X_\tau^{t,x}, \quad \tau \in [s, T],$$

we have

$$\begin{aligned} u(s, X_s^{t,x}) &= \mathbb{E}^{\mathcal{F}_s}[\Phi(X_T^{t,x})] \\ &\quad + \mathbb{E}^{\mathcal{F}_s}\left[\int_s^T F(\tau, X_\tau^{t,x}, u(\tau, X_\tau^{t,x}), \sigma(\tau, X_\tau^{t,x})^* \nabla_x u(\tau, X_\tau^{t,x})) d\tau\right]. \end{aligned}$$

If we set

$$\xi = \Phi(X_T^{t,x}) + \int_t^T F(\tau, X_\tau^{t,x}, u(\tau, X_\tau^{t,x}), \sigma(\tau, X_\tau^{t,x})^* \nabla_x u(\tau, X_\tau^{t,x})) d\tau$$

the previous equality leads to

$$\begin{aligned} & u(s, X_s^{t,x}) \\ &= \mathbb{E}^{\mathcal{F}_s} [\xi] - \int_t^s F(\tau, X_\tau^{t,x}, u(\tau, X_\tau^{t,x}), \sigma(\tau, X_\tau^{t,x})^* \nabla_x u(\tau, X_\tau^{t,x})) d\tau. \end{aligned}$$

Let us observe that  $\mathbb{E}^{\mathcal{F}_t} [\xi] = u(t, x)$ . Since  $\xi \in L^2(\Omega; \mathbf{R})$  is  $\mathcal{F}_{[t,T]}$ -measurable, by the representation theorem, there exists  $\tilde{Z} \in L^2_{\mathcal{P}}(\Omega \times [t, T]; L_2(\Xi, \mathbf{R}))$  such that

$$\mathbb{E}^{\mathcal{F}_s} [\xi] = u(t, x) + \int_t^s \tilde{Z}_\tau dW_\tau, \quad s \in [t, T].$$

We conclude that the process  $u(s, X_s^{t,x})$ ,  $s \in [t, T]$  is a (real) continuous semimartingale with canonical decomposition

$$\begin{aligned} u(s, X_s^{t,x}) &= u(t, x) + \int_t^s \tilde{Z}_\tau dW_\tau \\ &\quad - \int_t^s F(\tau, X_\tau^{t,x}, u(\tau, X_\tau^{t,x}), \sigma(\tau, X_\tau^{t,x})^* \nabla_x u(\tau, X_\tau^{t,x})) d\tau. \end{aligned} \tag{15}$$

Using (13) and arguing as in the proof of existence, we deduce that for a.a.  $\tau \in [t, T]$ ,  $\mathbb{P}$ -a.s.

$$\sigma(\tau, X_\tau^{t,x})^* \nabla_x u(\tau, X_\tau^{t,x}) = \tilde{Z}_\tau.$$

Substituting into (15) we obtain

$$\begin{aligned} u(s, X_s^{t,x}) &= u(t, x) + \int_t^s \sigma(\tau, X_\tau^{t,x})^* \nabla_x u(\tau, X_\tau^{t,x}) dW_\tau \\ &\quad - \int_t^s F(\tau, X_\tau^{t,x}, u(\tau, X_\tau^{t,x}), \sigma(\tau, X_\tau^{t,x})^* \nabla_x u(\tau, X_\tau^{t,x})) d\tau, \end{aligned}$$

for  $s \in [t, T]$ . Since  $u(T, X_T^{t,x}) = \Phi(X_T^{t,x})$ , we deduce that

$$\left\{ (u(s, X_s^{t,x}), \sigma(\tau, X_\tau^{t,x})^* \nabla_x u(\tau, X_\tau^{t,x})) \right\}_{s \in [t, T]}$$

solves the backward equation (5). By uniqueness, we have  $Y_s^{t,x} = u(s, X_s^{t,x})$ , for each  $s \in [t, T]$  and in particular, for  $s = t$ ,  $u(t, x) = Y_t^{t,x}$ . □

## References

- [1] Ph. Briand and F. Confortola. Differentiability of backward stochastic differential equations in Hilbert spaces with monotone generators. *Appl. Math. Optim.* to appear, <http://fr.arxiv.org/abs/math.PR/0603428>, 2006.
- [2] Ph. Briand, B. Delyon, Y. Hu, É. Pardoux, and L. Stoica.  $L^p$  solutions of backward stochastic differential equations. *Stochastic Process. Appl.*, 108(1):109–129, 2003.
- [3] Ph. Briand and Y. Hu. BSDE with quadratic growth and unbounded terminal value. *Probab. Theory Related Fields*, 136(4):604–618, 2006.

- [4] Q. Chen, Q. Ran, and J. Wang.  $L^p$  solutions of BSDEs with stochastic Lipschitz condition. Preprint, 2006.
- [5] G. Da Prato and J. Zabczyk. *Stochastic equations in infinite dimensions*, volume 44 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1992.
- [6] N. El Karoui. Backward stochastic differential equations: a general introduction. In *Backward stochastic differential equations (Paris, 1995–1996)*, volume 364 of *Pitman Res. Notes Math. Ser.*, pages 7–26. Longman, Harlow, 1997.
- [7] N. El Karoui and S.-J. Huang. A general result of existence and uniqueness of backward stochastic differential equations. In N. El Karoui and L. Mazliak, editors, *Backward stochastic differential equations*, volume 364 of *Pitman Res. Notes Math. Ser.*, pages 27–36. Longman, Harlow, 1997.
- [8] N. El Karoui, S. Peng, and M.-C. Quenez. Backward stochastic differential equations in finance. *Math. Finance*, 7(1):1–71, 1997.
- [9] M. Fuhrman and G. Tessitore. Nonlinear Kolmogorov equations in infinite dimensional spaces: the backward stochastic differential equations approach and applications to optimal control. *Ann. Probab.*, 30(3):1397–1465, 2002.
- [10] M. Fuhrman and G. Tessitore. Infinite horizon backward stochastic differential equations and elliptic equations in Hilbert spaces. *Ann. Probab.*, 32(1B):607–660, 2004.
- [11] Y. Hu, P. Imkeller, and M. Müller. Utility maximization in incomplete markets. *Ann. Appl. Probab.*, 15(3):1691–1712, 2005.
- [12] N. Kazamaki. *Continuous exponential martingales and BMO*, volume 1579 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1994.
- [13] M. Kobylanski. Backward stochastic differential equations and partial differential equations with quadratic growth. *Ann. Probab.*, 28(2):558–602, 2000.
- [14] J.-P. Lepeltier and J. San Martin. Existence for BSDE with superlinear-quadratic coefficient. *Stochastics Stochastics Rep.*, 63(3-4):227–240, 1998.
- [15] J. Ma and J. Yong. *Forward-backward stochastic differential equations and their applications*, volume 1702 of *Lecture Notes in Math*. Springer-Verlag, Berlin, 1999.
- [16] M.-A. Morlais. Quadratic BSDEs driven by a continuous martingale and application to the utility maximization problem. Preprint 06-12, IRMAR, 2006.
- [17] E. Pardoux. Backward stochastic differential equations and viscosity solutions of systems of semilinear parabolic and elliptic PDEs of second order. In L. Decreusefond, J. Gjerde, B. Øksendal, and A. S. Üstünel, editors, *Stochastic analysis and related topics VI (The Geilo Workshop, 1996)*, volume 42 of *Progr. Probab.*, pages 79–127. Birkhäuser Boston, Boston, MA, 1998.
- [18] E. Pardoux. BSDEs, weak convergence and homogenization of semilinear PDEs. In *Non-linear analysis, differential equations and control (Montreal, QC, 1998)*, pages 503–549. Kluwer Acad. Publ., Dordrecht, 1999.

- [19] E. Pardoux and S. Peng. Adapted solution of a backward stochastic differential equation. *Systems Control Lett.*, 14(1):55–61, 1990.
- [20] E. Pardoux and S. Peng. Backward stochastic differential equations and quasilinear parabolic partial differential equations. In B. L. Rozovskii and R. B. Sowers, editors, *Stochastic partial differential equations and their applications (Charlotte, NC, 1991)*, volume 176 of *Lecture Notes in Control and Inform. Sci.*, pages 200–217. Springer, Berlin, 1992.
- [21] S. Peng. Probabilistic interpretation for systems of quasilinear parabolic partial differential equations. *Stochastics Stochastics Rep.*, 37(1-2):61–74, 1991.
- [22] J. Zabczyk. Parabolic equations on Hilbert spaces. In *Stochastic PDE's and Kolmogorov equations in infinite dimensions (Cetraro, 1998)*, volume 1715 of *Lecture Notes in Math.*, pages 117–213. Springer, Berlin, 1999.