

REMARKS ON NON-LINEAR SCHRÖDINGER EQUATION WITH MAGNETIC FIELDS

LAURENT MICHEL

ABSTRACT. We study the non-linear Schrödinger equation with time depending magnetic field without smallness assumption at infinity. We obtain some results on the Cauchy problem, WKB asymptotics and instability.

1. INTRODUCTION

We consider the non-linear Schrödinger equation with magnetic field on \mathbb{R}^n

$$(1.1) \quad i\partial_t u = H_{A(t)} u - b^\gamma f(x, u)$$

with initial condition

$$(1.2) \quad u|_{t=t_0} = \varphi.$$

Here

$$H_{A(t)} = \sum_{j=1}^n (i\partial_{x_j} - bA_j(t, x))^2, t \in \mathbb{R}, x \in \mathbb{R}^n$$

is the time-dependent Schrödinger operator associated to the magnetic potential $A(t, x) = (A_1(t, x), \dots, A_n(t, x))$, $b \in]0, +\infty[$ is a parameter quantizing the strength of the magnetic field and $\gamma \geq 0$. We sometimes omit the space dependence and write $A(t)$ instead of $A(t, x)$. The aim of this note is to show that recent improvement in the analysis of non-linear Schrödinger equations can be adapted to the case with magnetic field. As an important preliminary, we study the local Cauchy problem for (1.1) in energetic space. Let us begin with the general framework of our study.

We suppose that the magnetic potential is a smooth function $A \in C^\infty(\mathbb{R}_t \times \mathbb{R}_x^n, \mathbb{R}^n)$ and that it satisfies the following assumption.

Assumption 1. (1) $\forall \alpha \in \mathbb{N}^n \quad \sup_{(t,x) \in \mathbb{R} \times \mathbb{R}^n} |\partial_x^\alpha \partial_t A| \leq C_\alpha.$

(2) $\forall |\alpha| \geq 1, \quad \sup_{(t,x) \in \mathbb{R} \times \mathbb{R}^n} |\partial_x^\alpha A| \leq C_\alpha.$

(3) $\exists \epsilon > 0, \forall |\alpha| \geq 1, \quad \sup_{(t,x) \in \mathbb{R} \times \mathbb{R}^n} |\partial_x^\alpha B| \leq C_\alpha \langle x \rangle^{-1-\epsilon}$

where $B(t, x)$ is the matrix defined by $B_{jk} = \partial_{x_j} A_k - \partial_{x_k} A_j$.

Remark that compactly supported perturbations of linear (with respect to x) magnetic potentials satisfy the above hypothesis.

Under Assumption 1, the domain $D(H_{A(t)}) = \{u \in L^2(\mathbb{R}_x^n), H_{A(t)} u \in L^2(\mathbb{R}_x^n)\}$ does not depend on t . Indeed, for $t, t' \in \mathbb{R}$ one has

$$(1.3) \quad H_{A(t')} = H_{A(t)} + bW(t, t')(i\nabla_x - bA(t)) + b(i\nabla_x - bA(t))W(t, t') + b^2W(t, t')^2$$

Key words and phrases. Schrödinger equation, Magnetic fields, Strichartz estimate.

with $x \mapsto W(t, t', x) = \int_t^{t'} \partial_s A(s, x) ds$ bounded as well as its x -derivatives uniformly with respect to t, t' in any compact set. In fact, the above identity shows that the space

$$H_{mg}^\beta(\mathbb{R}^n) = \{u \in L^2(\mathbb{R}^n), (1 + H_{A(t)})^{\beta/2} u \in L^2(\mathbb{R}^n)\}$$

does not depend on $t \in \mathbb{R}$. As $D(H_{A(t)}) = H_{mg}^2(\mathbb{R}^n)$, the above statement is straightforward. Moreover, the natural norms on this space are equivalent and this equivalence is uniform with respect to the parameter b for close times. More precisely, denoting $m_A = \sup_{(t,x) \in \mathbb{R} \times \mathbb{R}^n} |\partial_t A(t, x)|$, we have the following

Proposition 1.1. *Suppose that Assumption 1 is satisfied and let $\beta > 0$ and $T > 0$. Then, for all $t, t' \in \mathbb{R}$ such that $|t - t'| \leq b^{-1}T$ and all $u \in H_{mg}^\beta$ we have*

$$\|(H_{A(t')} + 1)^\beta u\|_{L^2} \leq (1 + 2m_A T + m_A^2 T^2)^\beta \|(H_{A(t)} + 1)^\beta u\|_{L^2}.$$

Proof. It is a straightforward consequence of equation (1.3), Assumption 1 and the fact that $(i\nabla_x - bA(t))(H_{A(t)} + 1)^{-1}$ is bounded by 1 in L^2 . \square

For $\beta \in \mathbb{N}$ we set

$$(1.4) \quad \|u\|_{H_{A(t)}^\beta} = \|(i\nabla_x - bA(t))^\beta u\|_{L^2} + \|u\|_{L^2}.$$

This norm is clearly equivalent (uniformly with respect to b) to $\|(1 + H_{A(t)})^{\beta/2} u\|_{L^2}$. In regard of Proposition 1.1 we define the magnetic Sobolev norm by

$$\|u\|_{H_{mg}^\beta} = \|u\|_{H_{A(t_0)}^\beta}.$$

Under Assumption 1 it is well-known (see [15], Th 4.6, p143 or [18]) that for $\varphi \in H_{mg}^1$, the linear Schrödinger equation

$$(1.5) \quad i\partial_t u = H_{A(t)} u, \quad u|_{t=s} = \varphi$$

has a solution $U_0(t, s)\varphi$. The operator $U_0(t, s)$ maps H_{mg}^1 into itself, is continuous from L^2 into L^2 and from H_{mg}^1 into H_{mg}^1 . Moreover, $U_0(t, s)\varphi$ is the unique H_{mg}^1 valued solution of (1.5) and $U_0(t, s)$ is unitary.

The first aim of this paper is to solve the Cauchy problem for the non-linear equation in the most appropriate space. We state the assumptions on the non-linearity f . We suppose that $f : \mathbb{R}^n \times \mathbb{C} \rightarrow \mathbb{C}$ is a measurable function such that

Assumption 2. (1) $f(x, 0) = 0$ almost every where.

(2) $\exists M \geq 0, \alpha \in [0, \frac{4}{n-2}[$ ($\alpha \in [0, \infty[$ if $n = 1, 2$) such that

$$|f(x, z_1) - f(x, z_2)| \leq M(1 + |z_1|^\alpha + |z_2|^\alpha)|z_1 - z_2|$$

for almost all $x \in \mathbb{R}^n$ and all $z_1, z_2 \in \mathbb{C}$.

(3) $\forall z \in \mathbb{C}, f(x, z) = (z/|z|)f(x, |z|)$

Remark that these assumptions are often used in the case $A = 0$. More precisely, in the case $A = 0$, the second property of the above assumption corresponds to a subcritical non-linearity with respect to H^1 .

Let us introduce some energy functional associated to these non-linearities. We define

$$F(x, z) = \int_0^{|z|} f(x, s) ds, \quad G(u) = \int_{\mathbb{R}^n} F(x, u(x)) dx$$

and for $t \in \mathbb{R}$ and $u \in H_{mg}^1$ we define the energy

$$E(b, t, u) = \int_{\mathbb{R}^n} \frac{1}{2} |(i\nabla_x - bA(t, x))u(x)|^2 dx - b^\gamma G(u).$$

Formally, it is not hard to see that any sufficiently regular solution of (1.1), (1.2), enjoys the following energy evolution law:

$$E(b, t, u) = E(b, 0, \varphi) - \operatorname{Re} \int_0^t \langle \partial_s A(s)u(x), (i\nabla - A(s))u(s) \rangle_{L^2} ds.$$

Therefore, the natural space to solve (1.1), (1.2) seems to be H_{mg}^1 .

Now we are in position to state our first result.

Theorem 1. *Suppose that Assumptions 1 and 2 are satisfied and let $\varphi \in H_{mg}^1$. Then, there exists $T_b, T^b > 0$ and a unique $u \in C([- T_b, T^b], H_{mg}^1) \cap C^1([- T_b, T^b], H_{mg}^{-1})$ solution of (1.1). Moreover, either $T_b = \infty$ (resp. $T^b = \infty$), or $\lim_{t \rightarrow -T_b} \|u(t)\|_{H_{mg}^1} = \infty$ (resp. $\lim_{t \rightarrow T^b} \|u(t)\|_{H_{mg}^1} = \infty$) and*

$$(1.6) \quad \|u(t)\|_{L^2} = \|\varphi\|_{L^2},$$

$$(1.7) \quad E(b, t, u) = E(b, 0, \varphi) - \operatorname{Re} \int_0^t \langle \partial_s A(s)u(x), (i\nabla - A(s))u(s) \rangle_{L^2} ds,$$

for all $t \in] - T_b, T^b[$. Additionally, there exists $\epsilon > 0$ such that, for all $b > 0$ and $\varphi \in H_{mg}^1$ such that $\|\varphi\|_{H_{mg}^1} \leq Cb$, we have $T_b, T^b \geq \epsilon b^{-\delta}$ with $\delta = \max(1, 2\gamma, \frac{2\gamma}{\alpha})$.

Let us make a few remarks on this result. The Cauchy problem for non-linear Schrödinger equation has a long story. In absence of magnetic field there are numerous results; see for instance [9, 10], [5].

In presence of magnetic field, the behavior of A when $|x|$ becomes large plays an important role. In the case where the magnetic potential A is bounded, the spaces H_{mg}^1 and H^1 coincide and the Cauchy problem can be solved in H^1 using usual techniques. If the magnetic field is unbounded, it is not possible to solve the Cauchy problem in H^1 as multiplication by A is not bounded on L^2 .

To avoid this difficulty some authors work in the weighted Sobolev space $\Sigma = \{u \in H^1(\mathbb{R}^n), (1 + |x|)u \in L^2, \}$ (see for instance [7], [14]). In particular, they require some decay of the initial data at infinity.

In the case of [7], this decay is required because the author use dispersive properties for the Laplacian instead of $H_{\Lambda(t)}$. In [14] the author use magnetic Strichartz estimates but their method based on fixed-point theorem is not adapted to the magnetic context and requires decay of the solution at infinity.

On the other hand, there exists also of a result of Cazenave and Esteban [4] dealing with the special case where the magnetic field B is constant (and hence, A is linear with respect to x). In a way, this paper is more satisfactory as they need only u_0 to belong to the energy space. Nevertheless, their result applies only to constant magnetic field.

Our theorem is, then a generalization of the above results. Before going further, let us remark that for unbounded A , the spaces H^1 , H_{mg}^1 and Σ are different. First, it is evident that Σ is contained in $H^1 \cap H_{mg}^1$. Let us give an example where Σ is strictly contained in H_{mg}^1 . For this purpose, we restrict ourselves to the case where the dimension $n = 2$ and consider the magnetic potential $A(x, y) = (y, x)$.

Let $g \in H^1(\mathbb{R}^2)$ be such that $|x|g \notin L^2$, then a simple calculus shows that $f(x, y) = g(x, y)e^{-ixy}$ belongs to $H_{mg}^1 \setminus \Sigma$.

In the case of defocusing non-linearities the energy law implies the following result.

Corollary 1.2. *Suppose that $F(x, z) \leq 0$ for all x, z , then $T_b, T^b = +\infty$.*

Proof. For $F \leq 0$, we deduce from (1.7) and Cauchy-Schwarz inequality, that

$$\|(i\nabla - A(t))u(t)\|_{L^2} \leq C_1 + C_2 \int_0^t \|(i\nabla - A(s))u(s)\|_{L^2} ds$$

for some fixed constant $C_1, C_2 > 0$. Hence, Gronwall Lemma shows that $\|(i\nabla - A(t))u(t)\|_{L^2}$ remains bounded on any bounded time-interval. Using (1.6) and the characterization of T_b , we obtain the result. \square

The next section contains the proof of Theorem 1. In section 3 we give some qualitative results on the solution of (1.1) in the limit $b \rightarrow \infty$. More precisely, we can construct WKB solutions and prove instability results with respect to initial data and parameter b .

2. CAUCHY PROBLEM IN THE ENERGY SPACE

The proof of theorem 1 relies on the Strichartz estimates proved in [18] for the problem

$$(2.1) \quad i\partial_t u = H_{A(t)} u + g(t), \quad u|_{t=s} = \varphi$$

Theorem 2. (Yajima) *Let I be a finite real interval, (q, r) and $(\gamma_j, \rho_j), j = 1, 2$ be such that $r, \rho_j \in [2, \frac{2n}{n-2}[$, $\frac{2}{q} = n(\frac{1}{2} - \frac{1}{r})$ and $\frac{2}{\gamma_j} = n(\frac{1}{2} - \frac{1}{\rho_j})$. Let $g_j \in L^{\gamma_j}(I, L^{\rho_j}(\mathbb{R}_x^n)), j = 1, 2$, where γ_j', ρ_j' are the conjugate exponent of γ_j, ρ_j . Then the solution u to (2.1) with $g = g_1 + g_2$ satisfies*

$$(2.2) \quad \|u\|_{L^q(I, L^r(\mathbb{R}_x^n))} \leq C(\|g_1\|_{L^{\gamma_1}(I, L^{\rho_1}(\mathbb{R}_x^n))} + \|g_2\|_{L^{\gamma_2}(I, L^{\rho_2}(\mathbb{R}_x^n))} + \|\varphi\|_{L^2(\mathbb{R}^n)})$$

where the constant C depends only on the length of I and the constant C_α of Assumption 1.

Proof. In the case $g = 0$ it is exactly Theorem 1 of [18]. In the general case it suffices to work as in the proof of Proposition 2.15 of [2] using a celebrated result of Christ and Kiselev [6]. The fact that the constant C depends only on the C_α is a direct consequence of the construction of Yajima [18]. \square

Remark 2.1. *In the case where the magnetic potential is not regular, there are some recent results of A. Stefanov [16] and Georgiev-Tarulli [8] which provide Strichartz estimates under smallness assumption on the magnetic fields. This should lead to the corresponding existence and uniqueness result for NLS in the case of small magnetic field. This could also have consequences on the well-posedness of the Schrödinger-Maxwell system (see [12], [13], [17] for results on this topics).*

It is important to notice that Theorem 1 is not a straightforward consequence of the above Strichartz estimate. Indeed, if we try to apply a fixed point method to equation (1.1), a problem occurs when we try to control the norm of the non-linearity in the H_{mg}^1 norm. Consider for instance the case $f(u) = |u|^2 u$, then

$$(i\nabla_x - bA(t))(|u|^2 u) = |u|^2(i\nabla_x - bA(t))(u) + ui\nabla_x(|u|^2).$$

The first term of the right hand side of this equality will be controlled by $\|u\|_{H_{mg}^1}$, whereas in the second term, as $A(t, x)$ is not bounded with respect to x , there is no chance to control $i\nabla_x(|u^2|)$ by $(i\nabla_x - bA(t))(|u^2|)$. For the same reason it does not seem easy to solve the Cauchy problem in magnetic Sobolev spaces of high degree.

To overcome this difficulty, we work as in [5], [4] and approximate the solution of (1.1) by solution of a non-linear Schrödinger equation with non-linearity linearized at infinity. In the work of Cazenave and Weissler, the main tool to justify the approximation is an energy conservation. In our case, the Hamiltonian depends on time, so that the energy is not conserved. Nevertheless, the error term is controlled by the H_{mg}^1 -norm so that it is possible to implement the same strategy. Another difference involved by the dependance with respect to time of the Hamiltonian is that usual techniques to solve the Cauchy problem with regular initial data and nice non-linearities can not apply in our context. Therefore, additionnaly to the approximation of the non-linearity, we have to introduce an approximation of the magnetic field itself and justify the convergence to our initial problem.

Let us introduce the approximated nonlinearities used in the sequel. Following [5], we decompose $f = \tilde{f}_1 + \tilde{f}_2$ with

$$(2.3) \quad \tilde{f}_1(x, z) = 1_{\{|z| \leq 1\}} f(x, z) + 1_{\{|z| \geq 1\}} f(x, 1)z$$

and

$$(2.4) \quad \tilde{f}_2(x, z) = 1_{\{|z| \geq 1\}} (f(x, z) - f(x, 1)z).$$

Next we define $f_m = \tilde{f}_1 + \tilde{f}_{2,m}$ where

$$(2.5) \quad \tilde{f}_{2,m}(x, z) = 1_{\{|z| \leq m\}} \tilde{f}_2(x, z) + 1_{\{|z| \geq m\}} \tilde{f}_2(x, m) \frac{z}{m}$$

Remark that these functions satisfy Assumption 2. We consider also the energy functional associated to these approximated non-linearities. We define

$$(2.6) \quad F_m(x, z) = \int_0^{|z|} f_m(x, s) ds, \quad G_m(u) = \int_{\mathbb{R}^n} F_m(x, u(x)) dx$$

and for $t \in \mathbb{R}$ and $u \in H_{mg}^1$ we set

$$(2.7) \quad E_m(b, t, u) = \int_{\mathbb{R}^n} \frac{1}{2} |(i\nabla_x - bA(t, x))u(x)|^2 dx - G_m(u).$$

Finally, we remark that replacing the magnetic potential $A(t, x)$ by $A(t + t_0, x)$ it suffices to prove Theorem 1 for $t_0 = 0$.

On the other hand, to enlight the notations we prove the theorem in the particular case $b = 1$. To get the general case it suffices to keep track of b along the proof. We will also restrict our study to $t \geq 0$, the other case being treated by reversing time in the equation.

2.1. Preliminary results. In the sequel, we will need Sobolev embeddings in the magnetic context. In this subsection, A is a magnetic potential satisfying Assumption 1.

Lemma 2.2. *Let $0 < s < \frac{n}{2}$ and $p_s = \frac{2n}{n-2s}$, then H_A^s is continuously embedded in $L^p(\mathbb{R}^n)$ for all $p \in [2, p_s]$ and there exists $C > 0$ independent of A such that*

$$(2.8) \quad \|u\|_{L^p} \leq C \|u\|_{H_A^s}$$

Proof. From the diamagnetic inequality (see [1]), we know that almost everywhere we have

$$|u| = |(\mathbf{H}_A + 1)^{-\frac{\alpha}{2}}(\mathbf{H}_A + 1)^{\frac{\alpha}{2}}u| \leq (-\Delta + 1)^{-\frac{\alpha}{2}}|(\mathbf{H}_A + 1)^{\frac{\alpha}{2}}u|.$$

Taking the L^p norm, the result follows from standard Sobolev inequalities. \square

Next we prove a technical result on the non-linearity.

Proposition 2.3. *Let $M > 0$, $r_1 = \rho_1 = 2$ and $r_2 = \rho_2 = \alpha + 2$ then*

- (1) *the sequence $(\tilde{f}_{2,m}(\cdot, u))_{m \in \mathbb{N}^*}$ converges to $\tilde{f}_2(\cdot, u)$ in $L^{\rho_2'}(\mathbb{R}^n)$ uniformly with respect to $u \in H_A^1$ such that $\|u\|_{H_A^1} \leq M$.*
- (2) *there exists $C(M) > 0$ independent of A such that for all $m \in \mathbb{N}^*$ and for all $u, v \in H_A^1$ with $\max(\|u\|_{H_A^1}, \|v\|_{H_A^1}) \leq M$ we have*

$$\|\tilde{f}_1(\cdot, u) - \tilde{f}_1(\cdot, v)\|_{L^{\rho_1'}(\mathbb{R}^n)} \leq C(M)\|u - v\|_{L^{r_1}}$$

$$\|\tilde{f}_{2,m}(\cdot, u) - \tilde{f}_{2,m}(\cdot, v)\|_{L^{\rho_2'}(\mathbb{R}^n)} + \|\tilde{f}_2(\cdot, u) - \tilde{f}_2(\cdot, v)\|_{L^{\rho_2'}(\mathbb{R}^n)} \leq C(M)\|u - v\|_{L^{r_2}(\mathbb{R}^n)}$$

Proof. We follow the method of Example 3 in [5]. Taking χ the characteristic function of the set $\{x \in \mathbb{R}^n \mid |u(x)| > m\}$ and using Assumption 2, we have

$$(2.9) \quad \|\tilde{f}_2(u) - \tilde{f}_{2,m}(u)\|_{L^{\rho_2'}(\mathbb{R}^n)} \leq 2\|\chi|u|^{\alpha+1}\|_{L^{\rho_2'}} = 2\|\chi u\|_{L^{\alpha+2}}^{\alpha+1}.$$

On the other hand, using Hölder inequality and Lemma 2.2 we get for $p = \frac{2n}{n-2}$,

$$(2.10) \quad \|u\|_{H_A^1} \geq C\|\chi u\|_{L^p} \geq Cm^{1-\frac{\alpha+2}{p}}\|\chi u\|_{L^{\alpha+2}}^{\frac{\alpha+2}{p}}.$$

As $\alpha < \frac{4}{n-2}$ then $1 - \frac{\alpha}{p+2} > 0$. Combining equations (2.9) and (2.9), we obtain the first point of the proposition.

The second assertion follows, as in example 3 in [5], from Hölder inequality, Assumption 2 and Lemma 2.2. The fact that the constant $C(M)$ is independent of the magnetic fields follows from the uniformity of the constant in Lemma 2.2. \square

Lemma 2.4. *Let $T > 0$ and γ_k , $k = 1, 2$ be defined by $\frac{2}{\gamma_k} = n(\frac{1}{2} - \frac{1}{\rho_k})$. For $M > 0$ there exists a constant $C(M)$ independent of A , such that for all $u, v \in H_A^1$ with $\|u\|_{H_A^1} \leq M$ and $\|v\|_{H_A^1} \leq M$ we have*

$$|G(u) - G(v)| + |G_m(u) - G_m(v)| \leq C(M)(\|v - u\|_{L^2} + \|v - u\|_{L^2}^{\nu}),$$

with $\frac{2}{\nu} = \frac{n}{2} - \frac{n}{\alpha+2}$ and for all $u, v \in L^\infty([0, T]H_A^1)$,

$$\begin{aligned} \|\tilde{f}_1(\cdot, u) - \tilde{f}_1(\cdot, v)\|_{L^{\gamma_1'}([0, T], L^{\rho_1'}(\mathbb{R}^n))} &\leq C(M)T\|u - v\|_{L^{\gamma_1}([0, T], L^{r_1}(\mathbb{R}^n))}. \\ \|\tilde{f}_{2,m}(\cdot, u) - \tilde{f}_{2,m}(\cdot, v)\|_{L^{\gamma_2'}([0, T], L^{\rho_2'}(\mathbb{R}^n))} &+ \|\tilde{f}_2(\cdot, u) - \tilde{f}_2(\cdot, v)\|_{L^{\gamma_2'}([0, T], L^{\rho_2'}(\mathbb{R}^n))} \\ &\leq C(M)T^{\frac{\gamma_2-1}{\gamma_2}}\|u - v\|_{L^{\gamma_2}([0, T], L^{r_2}(\mathbb{R}^n))} \end{aligned}$$

Moreover, $G_m \rightarrow G$ as $m \rightarrow \infty$ uniformly on bounded sets of H_A^1 .

Proof. Remark that $G(u) = \int_0^1 \langle f(x, su), u \rangle_{L^2} ds$ and $G_m(u) = \int_0^1 \langle f_m(x, su), u \rangle_{L^2} ds$ and copy the proof of Lemma 3.3 in [5], replacing classical Sobolev inequalities by Lemma 2.2 and using Proposition 2.3. \square

We are now in position to prove the uniqueness part of Theorem 1.

Proposition 2.5. *Let $T > 0$ and $u, v \in C([0, T], H_{mg}^1) \cap C^1([0, T], H_{mg}^{-1})$ be solution of (1.1). Then $u = v$.*

Proof. Let $u, v \in C([0, T[, H_{mg}^1) \cap C^1([0, T[, H_{mg}^{-1})$ be solution of (1.1), and set $w = v - u$. Then $w(0) = 0$ and

$$i\partial_t w - H_{A(t)} w = \tilde{f}_1(u) - \tilde{f}_1(v) + \tilde{f}_2(u) - \tilde{f}_2(v).$$

Let $r \in [2, \frac{2}{n-2}]$ and $q > 2$ such that $\frac{2}{q} = n(\frac{1}{2} - \frac{1}{r})$. Apply Theorem 2 together with Lemma 2.4, we get

$$\|w\|_{L^q([0, T[, L^r)} \leq C(T + T^{\gamma_2})(\|w\|_{L^\infty([0, T[, L^2)} + \|w\|_{L^{\gamma_2}([0, T[, L^{\rho_2})})$$

where $\gamma_2 = \frac{\alpha+1}{\alpha+2}$ and $\frac{2}{\gamma_2} = n(\frac{1}{2} - \frac{1}{\rho_2})$. As we can alternatively take (q, r) to be equal to $(2, \infty)$ and (γ_2, ρ_2) , we get the announced result by summing the obtained inequalities and making $T > 0$ small enough. \square

2.2. Autonomous case. In this section we explain briefly how to solve the Cauchy problem in H_{mg}^1 when the magnetic field $A(t, x) = A(x)$ is time independent. In this context, the functional E does not depend on time and formally we have the following conservation of energy. Suppose that u is solution of (1.1) then

$$E(b, u(t)) = E(b, \varphi), \quad \forall t.$$

More precisely, we prove the following

Proposition 2.6. *Let $M > 0$ and C_α , $\alpha \in \mathbb{N}^n$ a family of finite positive numbers. There exists $T > 0$ depending only on M and the C_α such that for all A satisfying $\partial_t A = 0$ and Assumption 1 with C_α and for all $\varphi \in H_A^1$ such that $\|\varphi\|_{H_A^1} \leq M$, there exists a unique $u \in C^0([0, T[, H_A^1) \cap C^1([0, T[, H_A^{-1})$ maximal solution of*

$$i\partial_t u = H_A u + f(x, u)$$

with initial condition $u|_{t=0} = \varphi$. Moreover, for all $t \in [0, T[$ we have

$$E(b, u(t)) = E(b, \varphi),$$

and if $T < \infty$ then $\lim_{t \rightarrow T} \|u\|_{H_A^1} = \infty$.

The proof is slight adaption of [5], [4] to our context. We need also to investigate the dependence of the existence time with respect to the magnetic field. However, the scheme of proof is the same and consists to consider an approximate problem and justify convergence on fixed time intervals. Let us give the main steps of the proof.

Step 1. Let f_m be defined by (2.3), (2.4), (2.5) and let A be a magnetic field satisfying the above hypotheses. Consider the problem

$$(2.11) \quad i\partial_t u = H_A u + f_m(x, u), \quad u_{t=0} = \varphi$$

with $\varphi \in H_A^1$. We have the following

Lemma 2.7. *Let $\varphi \in H_A^1$, then there exists $\tau_{m,A} > 0$ such that there exists $u_m \in C([0, \tau_{m,A}[, H_A^1) \cap C^1([0, \tau_{m,A}[, H_A^{-1})$ solution of (2.11). Moreover we have for all $t \in [0, \tau_{m,A}[$,*

$$(2.12) \quad E_m(u_m) = E_m(\varphi)$$

and

$$(2.13) \quad \|u_m(t)\|_{L^2} = \|\varphi\|_{L^2}.$$

Proof. The proof is the same as that of Lemma 3.5 of [5], replacing usual derivatives by magnetic derivatives. \square

Step 2. We show that the existence time $\tau_{m,A}$ can be bounded from below uniformly with respect to $m \in \mathbb{N}$ and A satisfying Assumptions of the above proposition.

Lemma 2.8. *Let $M > 0$. There exists $T_1 = T_1(M) > 0$ such that for all $m \in \mathbb{N}$, all A satisfying Assumption 1 and all $\varphi \in H_A^1$ with $\|\varphi\|_{H_A^1} \leq M$ we have*

$$\|u_m\|_{L^\infty([0,T_1],H_A^1)} \leq 2\|\varphi\|_{H_A^1}.$$

Proof. The proof is exactly the same as in Lemma 3.6 of [5], making use of Lemma 2.7 (in particular, we use strongly the conservation of energy) and Proposition 2.3 to get uniformity with respect to A . \square

Step 3. The final step is to prove convergence of the u_m to solution of the initial problem. First we prove convergence in L^2 .

Lemma 2.9. *Let $M > 0$ and C_α , $\alpha \in \mathbb{N}^n$ a family of finite positive numbers. There exists $T_2 > 0$ depending only on M and the C_α such that for all A satisfying Assumption 1 with C_α and for all $\varphi \in H_A^1$ such that $\|\varphi\|_{H_A^1} \leq M$, such that $(u_m)_{m \in \mathbb{N}}$ is a Cauchy sequence in $C([0, T_2], L^2)$.*

Proof. The proof is the same as in [5], making use of Theorem 2, Lemma 2.4, Proposition 2.3 and Lemma 2.7. \square

Now, we can complete the proof of Theorem 1. We denote u the limit of u_m in $C([0, T_2], L^2)$. From Lemma 2.8, it follows that $u \in L^\infty([0, T_2], H_A^1)$ and by Lemma 2.2, u_m converges to u in $C([0, T_2], L^r)$ for all $r \geq 2n/(n-2)$. Hence, it follows from Proposition 2.3 that $f_m(u_m)$ converges to $f(u)$ in $C([0, T_2], H_A^{-1})$ and u solves 1.1 in $L^\infty([0, T_2], H_A^{-1})$. Moreover, combining Lemma 2.4 and 2.7 we prove that

$$E(b, t, u) = E(b, 0, \varphi).$$

This shows that $u \in C([0, T_2], H_{mg}^1)$ and hence $u \in C^1([0, T_2], H_A^{-1})$.

2.3. Cauchy problem in the time-depending case. We suppose now that $A(t, x)$ satisfies Assumption 1. The strategy of proof is the same as in autonomous case and we first consider the problem

$$(2.14) \quad i\partial_t u = H_{A(t)} u + f_m(x, u), \quad u_{t=0} = \varphi$$

At least formally, we can see that the energy of the solution of this equation satisfies the following rule

$$(2.15) \quad E(t, u) = E(0, \varphi) - \operatorname{Re} \int_0^t \langle \partial_s A(s) u(s), (i\nabla_x - A(s)) u(s) \rangle ds.$$

This will replace the energy conservation in our approach. On the other hand another problem occurs if we try to apply the proof of [5]. Indeed, the first step should be to obtain a generalization of Lemma 2.7 in the time depending framework. Following the proof of Lemma 3.5 in [5], we should regularize the initial data and solve the Cauchy problem in H_{mg}^2 . The issue is that contrary to the autonomous case, the existence of smooth solution is not easy to prove. Indeed, the key point in the approach of [5] is that for any $g \in C([0, T], H^1)$ Lipschitz continuous with respect to time, the function $v(t) = \int_0^t U_0(t, s) g(s) ds$ is also Lipschitz continuous with respect to time. Such a result is easily proved in the autonomous case as the

identity $U_0(t+h, s) = U_0(t, s-h)$ permits to use the assumption on g . This fails to be true in the time-depending case. For this reason, we prove the existence in H_{mg}^1 in a direct way.

2.3.1. Existence of solution for approximated problem. In the case where the magnetic potential depends on time, we can not use the method of [5] to prove existence of solution on (2.14) in H_{mg}^1 . However we can prove the following.

Proposition 2.10. *Let $\varphi \in H_{mg}^1$, then there exists $\tilde{T} > 0$ such that there exists $u_m \in C([0, \tilde{T}[, H_{mg}^1) \cap C^1([0, \tilde{T}[, H_{mg}^{-1})$ solution of (2.14). Moreover we have for all $t \in [0, \tilde{T}]$,*

$$(2.16) \quad E_m(t, u_m) = E_m(t, \varphi) - \operatorname{Re} \int_0^t \langle \partial_s A(s)u(s), (i\nabla_x - A(s))u(s) \rangle ds.$$

and

$$(2.17) \quad \|u_m(t)\|_{L^2} = \|\varphi\|_{L^2}.$$

Proof. The method consists in approximating the magnetic potential $A(t, x)$ by potentials which are piecewise constant with respect to time. More precisely, remark that thanks to Assumption 1 and Proposition 2.6 there exists $T_2 = T_2(M) > 0$ such that for all $t_0 \in [0, T_2]$ the Cauchy problem

$$i\partial_t u = H_{A(t_0)} u(t) + f_m(u(t)), \quad u|_{t=t_0} = \varphi$$

can be solved in $C([t_0, t_0 + T_2], H_{A(t_0)}^1)$ for all initial data such that $\|\varphi\|_{H_{A(t_0)}^1} \leq M$.

Let $T \in]0, T_2[$ and for $n \in \mathbb{N}^*$, $k \in \{0, \dots, n-1\}$ define $t_n^k = \frac{kT}{n}$. We set $A_n(t, x) = A(t_n^k, x)$, $\forall t \in [t_n^k, t_n^{k+1}[$ and $A_n(T, x) = A(T, x)$. Next, we define the Hamiltonian $H_n = (i\nabla_x - A_n)^2$ and we look for solutions $u_{n,m}$ of

$$(2.18) \quad i\partial_t u = H_n u + f_m(u), \quad u|_{t=0} = \varphi.$$

From uniqueness in the autonomous case, such a function is given by

$$(2.19) \quad u_{n,m}(t, x) = \sum_{k=0}^{n-1} 1_{[t_n^k, t_n^{k+1}[}(t) v_{k,n,m}(t, x)$$

where $v_{k,n,m}(t, x)$ is defined as follows. We choose $v_{0,n,m}$ to be solution of

$$(2.20) \quad \begin{cases} i\partial_t v_{0,n,m} = (i\nabla_x - A(t_n^0, x))^2 v_{0,n,m} + f_m(v_{0,n,m}) \\ v_{0,n,m}(t_n^0, x) = \varphi(x) \end{cases}$$

and for $k \geq 1$, $v_{k,n,m}(t, x)$ is the solution of

$$(2.21) \quad \begin{cases} i\partial_t v_{k,n,m} = (i\nabla_x - A(t_n^k, x))^2 v_{k,n,m} + f_m(v_{k,n,m}) \\ v_{k,n,m}(t_n^k, x) = v_{k-1,n,m}(t_n^k, x). \end{cases}$$

Thanks to Proposition 2.6, the function $v_{k,n,m}$ are well defined and belong to $C^0([t_n^k, t_n^k + T_2], H_{mg}^1)$ and satisfy the following conservation equations

$$E_{n,m}(t, v_{k,n,m}(t)) = E_{n,m}(t_n^k, v_{k,n,m}(t_n^k))$$

for all $k = 0, \dots, n-1$, $t \in [t_n^k, t_n^{k+1}[$ and where for all $w \in H_{mg}^1(\mathbb{R}^n)$,

$$E_{n,m}(t, w) = \frac{1}{2} \int_{\mathbb{R}^n} |(i\nabla_x - A_n(t, x))w(x)|^2 dx - G_m(w).$$

Let us write $A(t_n^k, x) = A(t_n^{k-1}, x) + W_{n,k}(x)$ with $W_{n,k}(x) = \frac{1}{2} \int_{t_n^{k-1}}^{t_n^k} \partial_t A(t, x) dt$ and use $v_{k,n,m}(t_n^k, x) = v_{k-1,n,m}(t_n^k, x)$, then

$$(2.22) \quad \begin{aligned} E_{n,m}(t_n^k, u_{n,m}(t_n^k)) &= E_{n,m}(t_n^{k-1}, u_{n,m}(t_n^{k-1})) \\ &\quad - \int_{t_n^{k-1}}^{t_n^k} \operatorname{Re} \langle (i\nabla_x - A(t_n^{k-1}))u_{n,m}(t_n^{k-1}), \partial_t A(t, x)u_{n,m}(t_n^{k-1}) \rangle dt \\ &\quad + \|W_{n,k}u_{n,m}(t_n^{k-1})\|_{L^2}^2. \end{aligned}$$

Thanks to Assumption 1 and conservation of mass, we have $\|W_{n,k}u_{n,m}(t_n^k)\|_{L^2}^2 = O(\frac{\|\varphi\|_{L^2}^2}{n^2})$ uniformly with respect to k, n, m .

Hence, taking the sum of equations (2.22) for $k = 1, \dots, k_0$ with $k_0 = [\frac{nt}{T}]$, and using the fact that the energy is constant on $[t_n^{k_0}, t_n^{k_0+1}[$ we get for $t \in [t_n^{k_0}, t_n^{k_0+1}[$

$$(2.23) \quad \begin{aligned} E_{n,m}(t, u_{n,m}(t)) &= E_{n,m}(0, \varphi) \\ &\quad - \sum_{k=1}^{k_0} \int_{t_n^{k-1}}^{t_n^k} \operatorname{Re} \langle (i\nabla_x - A(t_n^{k-1}))u_{n,m}(t_n^{k-1}), \partial_t A(t, x)u_{n,m}(t_n^{k-1}) \rangle dt \\ &\quad + O(\frac{t}{n} \|\varphi\|_{L^2}^2). \end{aligned}$$

With this equation we can show that the sequence $(u_{n,m})_{(n,m) \in \mathbb{N} \times \mathbb{N}}$ is bounded in H_{mg}^1 . The proof is a discretization of the proof of Lemma 3.6 in [5]. Let $M = 2\|\varphi\|_{H_{mg}^1}$ and let $T_{n,m} > 0$ the maximal time such that $1 + 2m_A T_{n,m} + m_A^2 T_{n,m}^2 \leq \frac{5}{4}$ and for $t \in [0, T_{n,m}[$,

$$\|u_{n,m}\|_{H_{mg}^1} \leq M.$$

Thanks to Propositions 1.1, 2.3 and Lemma 2.2 there exists $K(M) > 0$ independent of $n, m \in \mathbb{N}$, such that

$$\|\partial_t u_{n,m}\|_{H_{mg}^{-1}} \leq K(M), \quad \forall n, m \in \mathbb{N}, \forall t \in [0, T_{n,m}[$$

and consequently,

$$(2.24) \quad \|u_{n,m} - \varphi\|_{L^2} \leq 2MK(M)t, \quad \forall t \in [0, T_{n,m}[.$$

On the other hand, it follows from (2.23) that

$$(2.25) \quad \begin{aligned} \frac{1}{2} \|(i\nabla_x - A_n(t))u_{n,m}(t)\|_{L^2}^2 &\leq \frac{1}{2} \|(i\nabla_x - A(0))\varphi\|_{L^2}^2 + G_m(u_{n,m}) - G_m(\varphi) \\ &\quad - \sum_{k=1}^n \int_{t_n^{k-1}}^{t_n^k} \operatorname{Re} \langle (i\nabla_x - A(t_n^k))u_{n,m}(t_n^k), \partial_s A(s, x)u_{n,m}(t_n^{k-1}) \rangle ds \\ &\quad + O(\frac{t}{n} \|\varphi\|_{L^2}^2). \end{aligned}$$

As $\partial_t A$ is bounded, the fourth term of the right hand side of (2.25) is bounded by CtM^2 . Moreover it follows from Lemma 2.4 and estimate (2.24) that

$$|G_m(u_{n,m}) - G_m(\varphi)| \leq C(M)(t + t^\nu).$$

Combining these equations with Proposition 1.1 we get

$$\|u_{n,m}\|_{H_{mg}^1}^2 \leq \frac{25}{16} \frac{M^2}{2} + C(M)(T_{n,m} + T_{n,m}^\nu).$$

Taking $0 < T_{n,m} < T$ with T sufficiently small independently on n, m , this proves that

$$(2.26) \quad \|u_{n,m}(t)\|_{L^\infty([0,T], H_{mg}^1)} \leq M, \quad \forall n, m \in \mathbb{N}$$

Let now $p, q \in \mathbb{N}$, then

$$i\partial_t(u_{p,m} - u_{q,m})(t) = H_p(u_{p,m} - u_{q,m})(t) + R_{p,q,m}(t) + g_m(u_{p,m}(t)) - g_m(u_{q,m}(t))$$

and $(u_{p,m} - u_{q,m})|_{t=0} = 0$, where

$$R_{p,q,m}(t) = ((A_q - A_p)(i\nabla - A(0)) + (i\nabla - A(0))(A_q - A_p)(t) + (A_p^2 - A_q^2)(t) + 2A(0)(A_q - A_p)(t))u_{q,m}(t).$$

Thanks to Theorem 2, we have for $\tilde{T} \in]0, T[$, $r \in [\frac{2n}{n-2} [$ and $\frac{2}{q} = n(\frac{1}{2} - \frac{1}{r})$,

$$\begin{aligned} \|u_{p,m} - u_{q,m}\|_{L^q([0,\tilde{T}], L^r(\mathbb{R}^n))} &\leq \|R_{p,q,m}\|_{L^\infty([0,\tilde{T}], L^2(\mathbb{R}^n))} \\ &+ C(M)(T\|u - v\|_{L^\infty([0,\tilde{T}], L^2(\mathbb{R}^n))} + \tilde{T}^{\frac{\gamma_2-1}{\gamma_2}}\|u - v\|_{L^{\gamma_2}([0,\tilde{T}], L^{r_2}(\mathbb{R}^n))}). \end{aligned}$$

On the other hand, $\epsilon > 0$ being fixed, for p, q large enough we have

$$\sup_{(t,x) \in \mathbb{R} \times \mathbb{R}^n} |A_p - A_q| \leq \epsilon.$$

Hence,

$$\|R_{p,q,m}\|_{L^\infty([0,\tilde{T}], L^2(\mathbb{R}^n))} \leq 2\epsilon\|u_{q,m}\|_{H_{mg}^1} + C\epsilon\|u_{q,m}\|_{L^2} \leq CM\epsilon,$$

and for p, q large enough we get

$$\begin{aligned} \|u_{p,m} - u_{q,m}\|_{L^q([0,\tilde{T}], L^r(\mathbb{R}^n))} &\leq \epsilon + C(M)\tilde{T}\|u - v\|_{L^\infty([0,\tilde{T}], L^2(\mathbb{R}^n))} \\ &+ C(M)\tilde{T}^{\frac{\gamma_2-1}{\gamma_2}}\|u - v\|_{L^{\gamma_2}([0,\tilde{T}], L^{r_2}(\mathbb{R}^n))}. \end{aligned}$$

This estimate is available, both for $(q, r) = (\infty, 2)$ and $(q, r) = (\gamma_2, \rho_2)$. Summing the two inequalities obtained and making $\tilde{T} > 0$ small enough, we get

$$\|u_{p,m} - u_{q,m}\|_{L^q([0,\tilde{T}], L^r(\mathbb{R}^n))} \leq 2\epsilon.$$

Therefore, the sequence $(u_{n,m})_{n \in \mathbb{N}}$ converges, as n goes to infinity, to a limit $u_m \in L^2$ which is solution of (2.14). Moreover, as $(u_{n,m})_{n \in \mathbb{N}}$ is bounded in H_{mg}^1 we can suppose that it converges weakly to u_m in H_{mg}^1 .

Now let's go back to equation (2.23). Using the fact that $u_{n,m}$ converges in L^2 and converges weakly in H_{mg}^1 it is no hard to see that $E_{n,m}(t, u_{n,m}) - E_{n,m}(0, \varphi)$ converges as $n \rightarrow \infty$, to $\text{Re} \int_0^t \langle \partial_s A(s)u_m(s), (i\nabla_x - A(s))u_m(s) \rangle ds$. From Proposition 2.3 and weak lower semicontinuity of the magnetic Sobolev norm $\|(i\nabla - A(t)) \cdot\|_{L^2}$ it follows that

$$E_m(t, u_m) \leq E_m(0, \varphi) - \text{Re} \int_0^t \langle \partial_s A(s)u(s), (i\nabla_x - A(s))u(s) \rangle ds.$$

Finally, $t > 0$ being fixed, consider $v_{n,m}(s) = u_{n,m}(t - s)$, which is solution of

$$i\partial_s v_{n,m} = -H_{A(t-s)} v_{n,m} - g_m(v_{n,m})$$

with initial data $v_{n,m}(s = 0) = u_{n,m}(t)$. Then we can do the same computations as above to get the converse inequality and hence (2.16) is proved. \square

2.3.2. *Convergence to the initial problem.* In this section, we show that the sequence u_m converges to a solution of (1.1) when m goes to infinity.

Lemma 2.11. *There exists $\tilde{T}_2 > 0$ depending only on $\|\varphi\|_{H_{mg}^1}$ such that $(u_m)_{m \in \mathbb{N}}$ is a Cauchy sequence in $C([0, \tilde{T}_2], L^2)$.*

Proof. The proof is the same as in [5], making use of Theorem 2, Lemma 2.4, Proposition 2.3 and Proposition 2.10. \square

Now, we can complete the proof of Theorem 1. This is the same as in [5] and we recall it for reader convenience. We denote u the limit of u_m in $C([0, \tilde{T}_2], L^2)$. From estimate (2.26), it follows that $u \in L^\infty([0, \tilde{T}_2], H_{mg}^1)$ and by Lemma 2.2, u_m converges to u in $C([0, \tilde{T}_2], L^r)$ for all $r \geq 2n/(n-2)$. Hence, it follows from Proposition 2.3 that $f_m(u_m)$ converges to $f(u)$ in $C([0, \tilde{T}_2], H_{mg}^{-1})$ and u solves (1.1) in $L^\infty([0, \tilde{T}_2], H_{mg}^{-1})$. Moreover, combining Lemma 2.4 and Proposition 2.10 we prove that

$$E(t, u) = E(0, \varphi) - \operatorname{Re} \int_0^t \langle \partial_s A(s)u(s), (i\nabla_x - A(s))u(s) \rangle ds.$$

This shows that $u \in C([0, \tilde{T}_2], H_{mg}^1)$ and hence $u \in C^1([0, \tilde{T}_2], H_{mg}^{-1})$.

3. WKB APPROXIMATION

In this section we justify WKB approximation for solution of (1.1) when the strength of the magnetic field b goes to infinity and obtain instability results. We stress our attention on the case where the magnetic field and the non-linearity have the same strength; that is we consider the case $\gamma = 2$ and search approximate solution for

$$(3.1) \quad \begin{cases} i\partial_s u = H_{A(s)} u + b^2 u g(|u|^2) \\ u|_{s=0} = a_0(x) e^{ibS(x)} \end{cases}$$

where g does not depend on x . Remark that with the previous notations, $f = ug(|u|^2)$. In this section we still ask f to satisfy Assumption 2 and we require additionally

Assumption 3. $g \in C^\infty(\mathbb{R}_+, \mathbb{R})$ with $g' > 0$.

Remark that if we suppose that $a_0 \in H^1$ and $\nabla S + A(0) \in L^2$ then the initial data satisfies $\|a_0(x) e^{ibS(x)}\|_{H_{mg}^1} = O(b)$. Therefore, under Assumptions 1, 2 and 3 it follows from Theorem 1 that there exists a unique solution of (3.1) in $C(-T_b, T_b], H_{mg}^1)$ with $T_b, T^b \geq Cb^{-\delta}$, $\delta = \max(2, \frac{2}{\alpha})$. In fact this solution takes a particular form.

Theorem 3. *Let $\sigma > \frac{n}{2} + 2$ and suppose that Assumptions 1, 2 and 3 are satisfied. Assume additionally that $\partial_t A$ belongs to $H^{\sigma-1}(\mathbb{R}^n)$ for all $t \in \mathbb{R}$ and take a_0 in $H^\sigma(\mathbb{R}^n)$ and S such that $\nabla S + A(t=0)$ belongs to $H^{\sigma-1}(\mathbb{R}^n)$. Then, there exists $T > 0$ and α_b, ϕ_b in $C([0, T], H^\sigma(\mathbb{R}^n)) \cap C^1([0, T], H^{\sigma-1}(\mathbb{R}^n))$ such that $u(t, x) = \alpha_b(bt, x) e^{ib(S(x) + \phi_b(bt, x))}$ is solution of (3.1) on $[0, b^{-1}T]$.*

Proof. We start the proof by a time rescaling leading to a semiclassical feature. We denote $h = b^{-1} > 0$ and set $u(s) = v(bs)$. Then equation (3.1) is equivalent to

$$(3.2) \quad \begin{cases} ih\partial_t v = (ih\nabla_x - A(ht))^2 v + vg(|v(t)|^2) \\ v|_{t=0} = a_0(x) e^{ih^{-1}S(x)} \end{cases}$$

We follow the general method initiated by Grenier [11] for the semiclassical Schrödinger equation and look for a phase and an amplitude depending on the parameter h . Putting the ansatz $v(t, x) = \alpha_h(t, x)e^{ih^{-1}\phi_h(t, x)}$ in the equations (3.2) we get

$$(3.3) \quad \begin{cases} \partial_t \phi_h + |\nabla_A \phi_h|^2 + g(|\alpha_h|)^2 = 0 \\ \partial_t \alpha_h + \nabla_A \phi_h \cdot \nabla \alpha_h + \operatorname{div}(\nabla_A \phi_h) \alpha_h = ih \Delta \alpha_h \end{cases}$$

where $\nabla_A \phi = (\nabla_x \phi + A(ht))$. Next we set $\varphi_h(t, x) = \nabla_A \phi_h(t, x) \in \mathbb{R}^n$ and differentiate the above eikonal equation with respect to x . We obtain

$$(3.4) \quad \begin{cases} \partial_t \varphi_h + 2\varphi_h \cdot \nabla \varphi_h + 2g'(|\alpha_h|^2) \operatorname{Re}(\overline{\alpha_h} \nabla \alpha_h) = h \partial_t A(ht, x) \\ \partial_t \alpha_h + \varphi_h \cdot \nabla \alpha_h + \operatorname{div}(\varphi_h) \alpha_h = ih \Delta \alpha_h \end{cases}$$

Separating real and imaginary parts of $\alpha_h = \alpha_{1,h} + i\alpha_{2,h}$, (3.4) becomes

$$(3.5) \quad \partial_t w_h + \sum_{j=1}^n A_j(w_h) \partial_{x_j} w_h = hLw_h + \nu_h$$

with

$$(3.6) \quad w_h = \begin{pmatrix} \alpha_{1,h} \\ \alpha_{2,h} \\ \varphi_{1,h} \\ \vdots \\ \varphi_{n,h} \end{pmatrix}, \nu_h = \begin{pmatrix} 0 \\ 0 \\ h \partial_t A_1(ht, x) \\ \vdots \\ h \partial_t A_n(ht, x) \end{pmatrix}$$

$$(3.7) \quad L = \begin{pmatrix} 0 & -\Delta & 0 \\ \Delta & 0 & 0 \\ 0 & 0 & 0_{n \times n} \end{pmatrix}$$

and

$$(3.8) \quad A_j(w) = \begin{pmatrix} \varphi_{j,h} & 0 & \alpha_1 & \dots & \alpha_1 \\ 0 & \varphi_{j,h} & \alpha_2 & \dots & \alpha_2 \\ 2g'\alpha_1 & 2g'\alpha_2 & v_j & 0 & 0 \\ \vdots & \vdots & 0 & \ddots & 0 \\ 2g'\alpha_1 & 2g'\alpha_2 & 0 & 0 & \varphi_{j,h} \end{pmatrix}$$

This system has the same form as in [11], [3] with the exception of the source term ν_h in right hand side of (3.5) and the initial data. Thanks to the assumptions, ν_h belongs to $H^{\sigma-1}(R^n)$, whereas the initial condition in (3.2) yields

$$(3.9) \quad w_h(t=0) = \begin{pmatrix} \operatorname{Re} a_o \\ \operatorname{Im} a_o \\ \partial_{x_1} S + A_1(0) \\ \vdots \\ \partial_{x_n} S + A_n(0) \end{pmatrix}$$

which belongs to $H^{\sigma-1}(\mathbb{R}^n)$.

On the other hand, thanks to the assumption on g' , the system (3.5) can be symmetrized by

$$(3.10) \quad S = \begin{pmatrix} I_2 & 0 \\ 0 & \frac{1}{g'} I_n \end{pmatrix}$$

which is symmetric and positive. It follows from general theory of hyperbolic systems that the problem (3.5) together with initial condition (3.9) has a unique solution $w_h \in L^\infty([0, T_h], H^{\sigma-1})$ for some $T_h > 0$.

Hence, we have to bound T_h from below by a constant independent of h . This is done by computing classical energies estimates as in [11], [3], and using the fact that $\partial_t A$ as well as $\nabla_x S + A(0)$ belong to $H^{\sigma-1}$.

Finally we define α_h and ϕ_h by $\alpha_h = w_{1,h} + iw_{2,h}$ and

$$\phi_h = S(x) - \int_0^t |\varphi_h|^2 + f(|\alpha_h|^2) ds.$$

By construction, ϕ_h belongs to L^2 . Moreover, a simple calculus shows that $\nabla_x \phi_h = \varphi_h - A(ht)$ belongs to $H^{\sigma-1}$ so that ϕ_h is in fact in H^σ . Going back to the equation on α_h and making energies estimates we show that $\alpha_h \in H^\sigma$. Finally, it a direct calculus shows that (α_h, ϕ_h) defined above solves (3.3) □

Remark 3.1. *The above solution belongs to the magnetic sobolev space H_{mg}^1 . Indeed,*

$$(i\nabla_x - bA)(\alpha_b e^{ib\phi_b}) = (i\nabla \alpha_b - b(\nabla \phi_b + A)\alpha_b) e^{ib\phi_b}$$

belongs to L^2 . Therefore the solution built in Theorem 3 coincide with the one of Theorem 1.

With Theorem 3 in hand it is easy to prove instability results.

Proposition 3.2. *Let $\sigma > \frac{n}{2} + 2$ and let A satisfy the assumptions of Theorem 3. Suppose that S is such that $\nabla S + A(t=0)$ belongs to $H^{\sigma-1}(\mathbb{R}^n)$. Then, there exists a_0 and $\tilde{a}_{0,b}$ in $H^\sigma(\mathbb{R}^n)$ and $0 < t_b < Cb^{-1}$ such that*

$$\|a_0 - \tilde{a}_{0,b}\|_{L^2} \rightarrow 0 \text{ as } b \rightarrow \infty$$

and the solutions u_b (resp. \tilde{u}_b) associated to (3.1) with initial data $a_0 e^{ibS(x)}$ (resp. $\tilde{a}_0 e^{ibS(x)}$) satisfy

$$\|u_b - \tilde{u}_b\|_{L^\infty([0, t_b], L^2)} \geq 1.$$

Proof. It is a straightforward consequence of Theorem 3 and the methods of [3]. □

REFERENCES

1. J. Avron, I. Herbst, and B. Simon, *Schrödinger operators with magnetic fields. I. General interactions*, Duke Math. J. **45** (1978), no. 4, 847–883.
2. N. Burq, P. Gérard, and N. Tzvetkov, *On nonlinear Schrödinger equations in exterior domains*, Ann. Inst. H. Poincaré Anal. Non Linéaire **21** (2004), no. 3, 295–318.
3. R. Carles, *Geometric optics and instability for semi-classical schrödinger equations*, Arch. Ration. Mech. Anal. (to appear).
4. T. Cazenave and M. J. Esteban, *On the stability of stationary states for nonlinear Schrödinger equations with an external magnetic field*, Mat. Apl. Comput. **7** (1988), no. 3, 155–168.
5. T. Cazenave and F. B. Weissler, *The Cauchy problem for the nonlinear Schrödinger equation in H^1* , Manuscripta Math. **61** (1988), no. 4, 477–494.
6. M. Christ and A. Kiselev, *Maximal functions associated to filtrations*, J. Funct. Anal. **179** (2001), no. 2, 409–425.
7. A. De Bouard, *Nonlinear Schroedinger equations with magnetic fields*, Differential Integral Equations **4** (1991), no. 1, 73–88.
8. V. Georgiev and M. Tarulli, *Scale invariant energy smoothing estimates for the Schrödinger equation with small magnetic potential*, Asymptot. Anal. **47** (2006), no. 1-2, 107–138.

9. J. Ginibre and G. Velo, *The classical field limit of scattering theory for nonrelativistic many-boson systems. II*, Comm. Math. Phys. **68** (1979), no. 1, 45–68.
10. ———, *The global Cauchy problem for the nonlinear Schrödinger equation revisited*, Ann. Inst. H. Poincaré Anal. Non Linéaire **2** (1985), no. 4, 309–327.
11. E. Grenier, *Semiclassical limit of the nonlinear Schrödinger equation in small time*, Proc. Amer. Math. Soc. **126** (1998), no. 2, 523–530.
12. Ginibre J. and Velo G., *Long range scattering for some schrödinger related nonlinear systems*, preprint.
13. M. Nakamura and T. Wada, *Local well-posedness for the Maxwell-Schrödinger equation*, Math. Ann. **332** (2005), no. 3, 565–604.
14. Y. Nakamura and A. Shimomura, *Local well-posedness and smoothing effects of strong solutions for nonlinear Schrödinger equations with potentials and magnetic fields*, Hokkaido Math. J. **34** (2005), no. 1, 37–63.
15. A. Pazy, *Semigroups of linear operators and applications to partial differential equations*, Applied Mathematical Sciences, vol. 44, Springer-Verlag, New York, 1983.
16. A. Stefanov, *Strichartz estimates for the magnetic schrödinger equation*, <http://arxiv.org/abs/math.AP/0509314> (2005).
17. Y. Tsutsumi, *Global existence and asymptotic behavior of solutions for the Maxwell-Schrödinger equations in three space dimensions*, Comm. Math. Phys. **151** (1993), no. 3, 543–576.
18. K. Yajima, *Schrödinger evolution equations with magnetic fields*, J. Analyse Math. **56** (1991), 29–76.

(L. Michel) LABORATOIRE J. A. DIEUDONNÉ, UNIVERSITÉ DE NICE SOPHIA-ANTIPOLIS, PARC VALROSE, 06108 NICE CEDEX 02, FRANCE
E-mail address, L. Michel: lmichel@math.unice.fr