



HAL
open science

A storage model with random release rate for modeling exposure to food contaminants

Patrice Bertail, Stéphan Cléménçon, Jessica Tressou

► To cite this version:

Patrice Bertail, Stéphan Cléménçon, Jessica Tressou. A storage model with random release rate for modeling exposure to food contaminants. 2007. hal-00138279v2

HAL Id: hal-00138279

<https://hal.science/hal-00138279v2>

Preprint submitted on 4 Apr 2007 (v2), last revised 3 Oct 2007 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A storage model with random release rate for modeling exposure to food contaminants

Patrice Bertail

CREST (INSEE) - Laboratoire de Statistiques

Stéphan Cléménçon*

MODALX - Université Paris X & Unité Mét@risk - INRA & LPMA - UMR CNRS 7599

Jessica Tressou

Unité Mét@risk - INRA & Hong Kong University of Sciences and Technology

April 4, 2007

Abstract

This paper is devoted to present and study a specific continuous-time piecewise-deterministic Markov process for describing the temporal evolution of exposure to a given food contaminant. The quantity X of food contaminant present in the body evolves through its accumulation after repeated dietary intakes on the one hand and the pharmacokinetics behavior of the chemical on the other hand. In the dynamic modeling considered here, the accumulation phenomenon is modeled by a simple marked point process with positive i.i.d. marks and elimination in between intakes occurs at a random linear rate θX , randomness of the coefficient θ accounting for the variability of the elimination process due to metabolic factors. Via embedded chain analysis, ergodic properties of this extension of the standard compound Poisson dam with (deterministic) linear release rate are investigated, the latter being of crucial importance for describing the long-term behavior of the exposure process $(X_t)_{t \geq 0}$ and assessing values of quantities such as the proportion of time the body burden in contaminant is over a certain threshold. We also highlight the fact that the exposure process is generally not directly observable in practice and establish a validity framework for simulation-based statistical methods by coupling analysis. Eventually, applications to methyl mercury contamination data are considered.

Key words and phrases: *pharmacokinetics model, dietary contamination, food safety, piecewise-deterministic Markov process, simulation method, stability analysis.*

AMS 2000 Mathematics Subject Classification: 60J25, 62P10, 92B15.

*Address of corresponding author: Stéphan Cléménçon Unité Met@risk - AgroParisTech - 16, rue Claude Bernard - 75231 Paris Cedex 05 - Email: Stephan.Clemencon@agroparistech.fr - Tel: +33 1 44 08 18 18 - Fax: +33 1 44 08 72 76

1 Introduction

Certain foods may contain varying amounts of chemicals such as methyl mercury (present in sea food), dioxins (in poultry, meat) or mycotoxins (in cereals, dried fruits, etc.), which may cause major health problems when accumulating inside the body in excessive doses. Food safety is now a crucial stake as regards public health in many countries (as an example, it is a thematic top priority of the 7th European Research Framework program, see <http://ec.europa.eu/research/fp7/>). This topic naturally interfaces with various disciplines, such as biology, nutritional medicine, toxicology and of course applied mathematics with the aim to develop rigorous methods for quantitative risk assessment. A scientific literature devoted to probabilistic and statistical methods for the study of dietary exposure to food contaminants is progressively carving out a place in applied probability and statistics journals (see [53], [21], [27] or [8] for instance).

Static viewpoints for the probabilistic modeling of the quantity X of a given food contaminant ingested on a short period have been considered in recent works, mainly focusing on the tail behavior of X and allowing for computation of the probability that X exceeds a maximum tolerable dose (see [7], [52]). However, as highlighted in [57], such approaches for food risk analysis do not take into account the accumulating and eliminating processes occurring in the body, which naturally requires to introduce time as a crucial description parameter of a comprehensive model (see also the discussion in [24]).

This paper aims at proposing a dynamic modeling of exposure to a certain food contaminant, incorporating important features of the phenomenon, in particular in a way that the model may account for the contaminant pharmacokinetics in man following intakes. The case of methyl mercury food contamination shall serve as a running illustration of the concepts and methods studied in this article: mathematical modeling of the pharmacokinetics behavior in man of methyl mercury (essentially present in sea food products) has received increasing attention in the toxicology literature (see [40], [51] [50], [1] or [23]) and dose-response relationships have been extensively investigated for this contaminant, establishing clearly its negative impact on human health (refer to [16]). In our modeling the amount of contaminant present in the body evolves through its accumulation after repeated intakes (food consumption) and according to the pharmacokinetics governing its elimination/excretion, so that its temporal evolution is described by a piecewise-deterministic Markov process (*PDM process* in abbreviated form): the accumulation process is modeled by a marked point process in a standard fashion, while the elimination phenomenon is described by a differential equation with random coefficients, randomness accounting for the variability of the rate at which the total contaminant body burden decreases in between intakes due to metabolic factors. Such a process slightly extends storage models with general release rules widely used in operations research and engineering for dealing with problems such as water storage in dams, in that one allows here the (content-dependent) release rate to be random, as strongly advocated by biological modeling, and inter-intake times are not required to be exponentially distributed (the choice of a memoryless distribution being totally inadequate in the dietary context). Having practical use of the proposed

exposure model for public health guidance in view (see [57]), we also discuss its relation to available data in the present paper: as sample paths of the exposure process cannot be observed in general, we set theoretical grounds for practical inference techniques based on intensive computer simulation methods. A thorough statistical analysis of toxicological and intake data based on the concepts and results developed in this article is carried out in a forthcoming companion paper (see [6]).

The outline of the paper is as follows. In section 2 a class of stochastic models with a reasonably simple (markovian) structure for describing the evolution through time of food contaminant exposure is introduced. In the important case when the (random) elimination rate is linear (such a feature being strongly motivated by previous works on kinetics modeling), theoretical properties of the *exposure process* are thoroughly investigated in section 3. Turning to the problem of estimating the steady-state or time-dependent features of the model that are relevant from the toxicology viewpoint and taking account of the fact that the exposure process is not observable in practice, statistical procedures relying on simulation methods are presented and studied in section 4. Finally, empirical studies related to methyl mercury food contamination are carried out in section 5, with the aim to illustrate the relevance of the modeling and the statistical methods studied in this paper. Technical proofs are postponed to the Appendix.

2 Modeling the exposure to a food contaminant

Suppose that an exhaustive list of P types of food, indexed by $p = 1, \dots, P$, involved in the alimentation of a given population and possibly contaminated by a certain chemical, is drawn up. Each type of food $p \in \{1, \dots, P\}$ is contaminated in random ratio $K^{(p)}$, with probability distribution $F_{\mathcal{K}^{(p)}}$, regarding to the chemical of interest. Concerning this specific contaminant exposure, a meal may be viewed as a realization of a r.v. $Q = (Q^{(1)}, \dots, Q^{(P)})$ indicating the quantity of food of each type consumed, normalized by the body weight. For a meal Q drawn from a distribution F_Q on $(\mathbb{R}_+^P, \mathcal{B}_{\mathbb{R}_+^P})$, cooked from foods of which toxicity is described by a contamination ratio vector $K = (K^{(1)}, \dots, K^{(P)})$ drawn from $F_{\mathcal{K}} = \otimes_{p=1}^P F_{\mathcal{K}^{(p)}}$, the global contaminant intake is

$$U = \langle K, Q \rangle, \tag{1}$$

denoting by $\langle \cdot, \cdot \rangle$ the standard inner product on \mathbb{R}^P . Its probability distribution F_U is the image of $F_{\mathcal{K}} \otimes F_Q$ by the inner product $\langle \cdot, \cdot \rangle$, assuming that the quantities of food consumed are independent from the contamination levels. Here and throughout, we suppose that the contaminant intake distribution F_U has a density f_U with respect to λ , the Lebesgue measure on \mathbb{R}_+ .

By convention, $T_0 = 0$ is chosen as time origin. The food contamination phenomenon through time for an individual of the population of interest may be classically modeled by a marked point process $\{(T_n, Q_n, K_n)\}_{n \geq 1}$ on $\mathbb{R}_+ \times \mathbb{R}_+^P \times \mathbb{R}_+^P$, the T_n 's being the successive times at which the individual consumes foods among the list $\{1, \dots, P\}$ and the marks (Q_n, K_n)

being respectively the vector of food quantities and the vector of contamination ratios related to the meal had at time T_n , $n \geq 1$. The process $\{(T_n, Q_n)\}_{n \geq 1}$ describing dietary behavior is assumed independent from the sequence $(K_n)_{n \geq 1}$ of chemical contamination ratios. Although the modeling of dietary behaviors could certainly give rise to a huge variety of models, depending on the dependence structure between (T_n, Q_n) and past values $\{(T_m, Q_m)\}_{m < n}$ that one stipulates, we make here the simplifying assumption that the marks Q_n , $n \geq 1$, form an i.i.d. sequence with common distribution F_Q , independent from the location times $(T_n)_{n \geq 1}$. This assumption is acceptable for chemicals present in a few types of food, such as methyl mercury, our running example, but certainly not for all contaminants. For chemicals present in many foods of everyday consumption such as Ochratoxin A (present in cereals, coffee, *etc.*), it would be necessary to introduce additional autoregressive structure in the model for capturing important features of any realistic diet (the consumption of certain food being typically alternated for reasons related to taste or nutritional aspects). Such a modeling task is beyond the scope of the present paper and is left for further investigation. Finally, we suppose that the inter-intake times $\Delta T_{n+1} = T_{n+1} - T_n$, $n \geq 1$, form a sequence of i.i.d. r.v.'s with common probability distribution $G(dt) = g(t)dt$ and finite expectation $m_G = \int_{t=0}^{\infty} tG(dt) < \infty$, the sequence $(T_n)_{n \geq 1}$ of intake times being thus a pure renewal process.

Contamination sources other than dietary are neglected in the present study and we denote by $X(t)$ the total body burden in contaminant at time $t \geq 0$. In between intakes, we assume that the *contamination exposure process* $X(t)$ is governed by the differential equation

$$\dot{x}(t) = -r(x(t), \theta), \quad (2)$$

θ being a random parameter, taking its values in a set $\Theta \subset \mathbb{R}^d$ with $d \geq 1$ say, and accounting in the modeling for fluctuations of the elimination rate due to metabolic factors at the intake times (the successive values θ_n , $n \in \mathbb{N}$, of θ are thus fixed at times T_0, T_1, \dots). And the function $r(x, \theta)$ is assumed to be strictly positive and continuous on $\mathbb{R}_+ \times \Theta$, such that for all $\theta \in \Theta$, $r(0, \theta) = 0$ (so that when $X(t)$ eventually reaches the level 0, the process stays at this level until the next intake) and for all $(\epsilon, \theta) \in (0, 1) \times \Theta$:

$$\inf_{\epsilon < x < \epsilon^{-1}} r(x, \theta) > 0 \text{ and } \sup_{0 < x < \epsilon^{-1}} r(x, \theta) < \infty. \quad (3)$$

Under these conditions, for any initial value $x(0) \geq 0$ and metabolic parameter value $\theta \in \Theta$, Eq. (2) has clearly a unique solution.

Other approaches may be naturally adopted for describing the elimination phenomenon occurring in between intakes. For instance, toxicokinetic models based on stochastic differential equations or decreasing jump processes (as in inventory modeling) could be pertinently considered for this purpose.

Remark 1 (ON PHARMACOKINETICS MODELING) In toxicology, Eq. (2) is widely used with $r(x, \theta) = \theta x$ for modeling the kinetics in man of certain contaminants following intakes. As shown by many pharmacokinetics studies, there is considerable empirical

evidence that it properly describes the way the elimination rate depends on the total body burden of the chemical in numerous cases (see [11], [49] or [26] for further details on *linear pharmacokinetics models*). In this context, the release parameter $\log 2/\theta$ is known as the *biological half-life* of the contaminant (the time required for X to decrease by half in the organism in absence of new contaminant intake). For methyl mercury (MeHg), our running example in this paper, the half-life is known to fluctuate around six weeks (see [51] and the references therein). For such dietary contaminants, of which biological half-life is measured in weeks rather than days, it is naturally essential to take account of the kinetics in man for successful modeling of the exposure phenomenon.

We assume that $(\theta_n)_{n \in \mathbb{N}}$ is an i.i.d. sequence with common distribution $H(d\theta)$. For a given value of the metabolic parameter $\theta \in \Theta$, the time necessary for the body burden (without further intake) to decrease from $x_0 > 0$ to $x \in (0, x_0)$ is given by

$$\tau_\theta(x_0, x) = \int_x^{x_0} \frac{1}{r(y, \theta)} dy.$$

Under the assumptions stated above, we clearly have that $H(\{\tau_\theta(x_0, x) < \infty\}) = 1$ for all $0 < x \leq x_0$. The contaminant may be thus entirely eliminated from the body (the amount x reaching then the level 0) with probability one in the sole case when the following condition holds.

Condition (\mathcal{C}_1): $H(\{\tau_\theta(x_0, 0) < \infty\}) = 1$ for some $x_0 > 0$.

In such a case we would also have $H(\{\tau_\theta(x, 0) < \infty\}) = 1$ for all $x \geq 0$. In this respect, it is noteworthy that, in the linear case mentioned in Remark 2, $\tau_\theta(x, 0) = \infty$ for all $\theta > 0$ and $x > 0$.

Hence, in between intake times and given the current value of the metabolic parameter θ , the exposure process moves in a deterministic fashion according to (2), and has the same (upward) jumps as the process of cumulative intakes

$$B(t) = \sum_{n=1}^{N(t)} U_n,$$

with $U_n = \langle K_n, Q_n \rangle$, $n \in \mathbb{N}$, and denoting by $N(t) = \sum_{n \in \mathbb{N}} \mathbb{I}\{T_n \leq t\}$ the number of intakes until time t . The exposure process X is piecewise-deterministic with càd-làg trajectories (see a typical sample path in Fig.1) and satisfies the equation

$$X(t) = X(0) + B(t) - \sum_{n=1}^{N(t)+1} \int_{T_{n-1}}^{T_n \wedge t} r(X(s), \theta_n) ds, \quad (4)$$

$X(0)$ denoting the total body burden in contaminant at initial time $T_0 = 0$ and with $a \wedge b = \inf\{a, b\}$ for all $(a, b) \in \mathbb{R}^2$. For an account of such piecewise deterministic processes, one may refer to [20] (see also [19] and ergodic results may be found in [15]).




Figure 1: Sample path of the exposure process X , modeling the evolution of the total body burden of a given dietary contaminant through time.

For the continuous-time process thus defined to be Markovian, one has to record the current value $\theta(t) = \sum_{n \in \mathbb{N}} \theta_n \mathbb{I}_{\{t \in [T_n, T_{n+1}]\}}$ of the metabolic parameter as well as the backward recurrence time $A(t) = t - T_{N(t)}$ (the time since the last intake). By construction, the process $(X(t), \theta(t), A(t))_{t \geq 0}$ is strongly Markovian with generator

$$\begin{aligned} \mathcal{G}\phi(x, \theta, t) = & \zeta(t) \int_{u=0}^{\infty} \int_{\theta' \in \Theta} \{\phi(x+u, \theta', 0) - \phi(x, \theta, t)\} F_U(du) H(d\theta') \\ & - r(x, \theta) \partial_x \phi(x, \theta, t) + \partial_t \phi(x, \theta, t), \quad (5) \end{aligned}$$

denoting by $\zeta(t) = g(t) / \int_{s=t}^{\infty} g(s) ds$ the hazard rate of the inter-intake times and provided that $\phi(\cdot, \theta, \cdot) : (x, t) \mapsto \phi(x, \theta, t)$ is a bounded function with bounded continuous first derivatives in x and t for all $\theta \in \Theta$.

In the above setting, the time origin $T_0 = 0$ does not necessarily correspond to an intake time. Given the time $A(0) = a$ since the last intake at time $t = 0$, we let ΔT_1 have the density $g_a(t) = g(a+t) / \int_{s=a}^{\infty} g(s) ds$, making the renewal process $(\Delta T_n)_{n \in \mathbb{N}}$ possibly delayed, except naturally in the case when the inter-intake distribution G is exponential. However, the choice of such a memoryless distribution in the dietary context is clearly not pertinent, distributions with increasing hazard rate being much more adequate (see section 5). Here and throughout we denote by $\mathbb{P}_{x,a}$ the probability measure on the underlying space such that $(X(0), A(0)) = (x, a)$ and $\theta(0) \sim H$, and by $\mathbb{E}_{x,a}[\cdot]$ the $\mathbb{P}_{x,a}$ -expectation for all $x \geq 0$ and $a \in \text{supp}(G)$.

In the case when one neglects variability in the elimination process (*i.e.* when H is a Dirac measure) and under the additional assumption that the renewal times are exponentially distributed (making the process X itself Markovian, which facilitates much the study but is not relevant to our application as emphasized above), this modeling boils down to a standard *storage model with a general release rate* (see [13] and [12] for instance). We refer to Chapter XIV in [3] for an account of such processes, widely used in operations research for modeling queuing/storage systems. Basic communication properties of the stochastic

process $X = (X(t))_{t \geq 0}$ may be established in a fashion very similar to the ones of the latter processes. They are summarized in the next result (of which proof is omitted since it is a slight modification of the proof of Proposition 1.2 in chap. XIV of [3]).

Theorem 1 *Suppose that $G(dx) = g(x)dx$ has infinite tail. Assume further that either $g(x) > 0$ on $]0, \epsilon]$ for some $\epsilon > 0$ or else that F_U has infinite tail. Then X reaches any state $x > 0$ in finite time with positive probability whatever the starting point, i.e. for all $x_0 \geq 0$, $a \in \text{supp}(G)$, it holds*

$$\mathbb{P}_{x_0, a}(\tau_x < \infty) > 0, \quad (6)$$

with $\tau_x = \inf\{t \geq 0 : X_t = x\}$. Furthermore, if condition (C_1) is fulfilled, then (6) still holds for $x = 0$. Besides, either X "heads to infinity" with probability one, i.e. is such that $\mathbb{P}_{x_0, a}(\{X(t) \rightarrow \infty, \text{ as } t \rightarrow \infty\}) = 1$ for all $x_0 \geq 0$, or else X reaches any state $x > 0$ in finite time with probability one whatever the starting point, i.e. for all $x_0 \geq 0$, $a \in \text{supp}(G)$,

$$\mathbb{P}_{x_0, a}(\tau_x < \infty) = 1. \quad (7)$$

If (C_1) is satisfied, then (7) also holds for $x = 0$.

An important task is to find conditions ensuring that the limiting behavior of the exposure process X is represented by a stationary probability measure μ describing the equilibrium state to which the process settles as time goes to infinity. In particular, time averages over long periods, such as the mean time spent by the exposure process X over a possibly critical threshold $u > 0$, $T^{-1} \int_0^T \mathbb{I}_{\{X_t \geq u\}} dt$, for instance, are then asymptotically described by the distribution μ . And computing/estimating steady-state quantities would be then relevant for summarizing the exposure phenomenon in the long run and assessing the long-term toxicological risk. Beyond stochastic stability properties, determining the tail behavior of the steady-state distribution and evaluating the rate at which the exposure process converges to the stationary state is also of critical importance in practice. These questions are thoroughly investigated for linear pharmacokinetics models in the next section.

3 Probabilistic study in the 'linear kinetics' case

We now focus on the ergodicity properties of the exposure process $X(t)$ in the specific case when for a given metabolic state described by a real parameter θ , the elimination rate is proportional to the total body burden in contaminant, i.e. $r(x, \theta) = \theta x$.

Here we suppose that Θ is a subset of \mathbb{R}_+ , ensuring that (3) is satisfied. As mentioned before, the linear case is of crucial importance in toxicology, insofar as it suitably models the pharmacokinetics behavior in man of numerous chemicals (see [26]). We shall show that studying the long-term behavior of X boils down to investigating the properties of the embedded Markov chain $\tilde{X} = (X_n)_{n \geq 1}$ of which values correspond to the ones taken by the

exposure process just after intake times : $X_n = X(T_n)$ for all $n \geq 1$. By construction, the chain \tilde{X} satisfies the following autoregressive equation with random coefficients

$$X_{n+1} = e^{-\theta_n \Delta T_{n+1}} X_n + U_{n+1}, \text{ for all } n \geq 1, \quad (8)$$

and has transition probability $\Pi(x, dy) = \pi(x, y) dy$ with transition density

$$\pi(x, y) = \int_{\theta \in \Theta} \int_{t = \frac{1}{\theta} \log(1 \vee \frac{x}{y})}^{\infty} f_U(y - xe^{-\theta t}) G(dt) H(d\theta), \quad (9)$$

for all $(x, y) \in \mathbb{R}_+^{*2}$, where $a \vee b = \max(a, b)$. Ergodicity of such real-valued Markov chains Y , defined through stochastic recurrence equations of the form $Y_{n+1} = \alpha_n Y_n + \beta_n$, where $\{(\alpha_n, \beta_n)\}_{n \in \mathbb{N}}$ is a sequence of i.i.d. pairs of positive r.v.'s, has been extensively studied in the literature, such models being widely used in financial or insurance mathematics (see section 8.4 in [22] for instance). Specialized to our setting, well known results related to such processes enable to show that the embedded chains \tilde{X} is positive recurrent under the assumption that $\log(1 \vee U_1)$ has finite expectation, as stated in the next theorem, and then to specify the tail behavior of the limiting probability distribution. Furthermore, the simple autoregressive form of Eq. (8) makes Foster-Lyapunov conditions easily verifiable for such Markov chains, in order to refine their stochastic stability analysis.

Theorem 2 *Under the assumptions of Theorem 1, the chain \tilde{X} is λ -irreducible. Moreover, suppose that the following condition holds.*

(H1) $\mathbb{E}[\log(1 \vee U_1)] < \infty$.

Then \tilde{X} is recurrent positive with stationary probability distribution $\tilde{\mu}$.

If one assume further that f_U is continuous and strictly positive on \mathbb{R}_+ and:

(H2) *there exists some $\gamma \geq 1$ such that $\mathbb{E}[U_1^\gamma] < \infty$,*

then \tilde{X} is geometrically ergodic, $\tilde{\mu}$ has finite moment of order γ and there exist constants $R < \infty$ and $r > 1$ such that, for all $n \geq 1, x > 0$,

$$\sup_{\{\psi, |\psi(z)| \leq 1 + z^\gamma\}} \left| \int_{y=0}^{\infty} \psi(y) \Pi^n(x, dy) - \tilde{\mu}(\psi) \right| \leq R(1 + x^\gamma)r^{-n}, \quad (10)$$

denoting by Π^n the n -th iterate of Π and with $\tilde{\mu}(\psi) = \int_{y=0}^{\infty} \psi(y) \tilde{\mu}(dy)$ for any $\tilde{\mu}$ -integrable function ψ .

Suppose finally that the condition (H1) and the next one simultaneously hold,

(H3) *The r.v. U_1 is regularly varying with index $\kappa > 0$ (i.e. for all $t > 0, (1 - F_U(tx))/(1 - F_U(x)) \sim t^{-\kappa}$ as $x \rightarrow \infty$).*

Then the stationary law $\tilde{\mu}$ has regularly varying tail with index κ .

Remark 2 (ON THE TAIL BEHAVIOR ASSUMPTION FOR THE INTAKE DISTRIBUTION) The relevance of the regular variation assumption for modeling the tail behavior of dietary contaminant intakes related to certain chemicals is strongly supported in [52] and [7]. In these works, various estimation strategies for tail distribution features such as the Pareto index κ involved in (H3) are also proposed and implemented on several food contamination and consumption data sets. We refer to [22] for an excellent account of such notions arising in extreme values theory and techniques for modeling extremal events.

As pointed out in [38], stochastic stability analysis based on drift criteria in the continuous-time setting is not as straightforward as in the discrete-time case, generally due to the complex form of the generator and of candidate test functions. Fortunately, given the explicit relationship between X and the embedded discrete-time chain \tilde{X} in our specific case, ergodicity of the continuous-time model and tail properties of its limiting distribution may be investigated based on the results established above for \tilde{X} under mild conditions. However, under more restrictive moment conditions for the inter-intake distribution, as the one stated below, a simple test function for which the generator (5) is shown to satisfy a geometric drift condition, may be nevertheless exhibited, so as to establish geometric ergodicity for the Markov process $\{(X(t), \theta(t), A(t))\}_{t \geq 0}$.

(H4) There exists $\eta > 0$ such that $\mathbb{E}[\exp(\eta \Delta T_2)] < \infty$.

Theorem 3 *Under the assumptions of Theorem 1 and supposing that (H1) is fulfilled, $X(t)$ has an absolutely continuous limiting probability distribution μ given by*

$$\mu([u, \infty[) = m_G^{-1} \int_{x=u}^{\infty} \int_{t=0}^{\infty} \int_{\theta \in \Theta} t \wedge \frac{\log(x/u)}{\theta} \tilde{\mu}(dx) G(dt) H(d\theta), \quad (11)$$

in the sense that $\Gamma^{-1} \int_0^\Gamma \mathbb{I}_{\{X_t \leq u\}} dt \rightarrow \mu([0, u])$, $\mathbb{P}_{x_0, \alpha}$ -a.s., as $t \rightarrow \infty$ for all $x_0 \geq 0$ and $\alpha \in \text{supp}(G)$. Furthermore,

- *if (H3) holds and the set Θ is bounded, then μ is regularly varying with the same index as F_U ,*
- *and if (H2) and (H4) hold, then $\{(X(t), \theta(t), A(t))\}_{t \geq 0}$ is geometrically recurrent. In particular, μ has finite moment of order γ and for all $(x, \alpha) \in \mathbb{R}_+^* \times \text{supp}(G)$ there exist constants $\beta \in]0, 1[$, $B_\alpha < \infty$ such that*

$$\sup_{\psi(z) \leq 1+z^\gamma} |\mathbb{E}_{x, \alpha}[\psi(X_t)] - \mu(\psi)| \leq B_\alpha (1 + x^\gamma) \beta^t. \quad (12)$$

Remark 3 (ON THE TAIL BEHAVIOR OF THE STATIONARY DISTRIBUTION) When the U_n 's are heavy-tailed, and under the assumption that the ΔT_n 's are exponentially distributed (making $B(t)$ a time-homogeneous Lévy process), the fact that the stationary distribution μ inherits its tail behavior from F_U has been established in [2] for deterministic release rates. Besides, when assuming G exponential and θ fixed, one may exactly

identify the limit distribution μ in some specific cases (see section 8 in [12] or section 2 in Chap. XIV of [3]) using basic level crossing arguments (X being itself markovian in this case). If F_U is also exponential for instance, μ is a Gamma distribution.

Remark 4 (ON PRACTICAL RELEVANCE OF STEADY-STATE FEATURES) Now that it is established that the exposure process settles to an equilibrium regime after a 'certain time', the question of specifying precisely what is meant by 'certain time' naturally arises. It would be pertinent to describe the long run behavior of the exposure process and assess the long-term toxicological risk by computing steady-state characteristics solely if the time necessary to reach the equilibrium state approximately may be considered as a reasonable horizon at the human scale. As an illustration, the amount of time needed to be roughly in steady-state from a collection of datasets related to dietary MeHg contamination is evaluated through simulation in Section 5.

In order to exhibit connections between the exposure process $X = (X(t))_{t \geq 0}$ and possible negative effects of the chemical on human health, it is appropriate to consider simple characteristics of the process X , easily interpretable from an epidemiology viewpoint. In this respect, the mean exposure over a long time period $T^{-1} \int_{t=0}^T X(t) dt$ is one of the most relevant features. Its asymptotic behavior is refined in the next result.

Proposition 4 *Under the assumptions of Theorem 1 and supposing that (H2) is fulfilled for $\gamma = 1$, we have for all $(x_0, \mathbf{a}) \in \mathbb{R}_+ \times \text{supp}(\mathbf{G})$*

$$\bar{X}_T = \frac{1}{T} \int_{t=0}^T X(t) dt \rightarrow m_\mu, \mathbb{P}_{x_0, \mathbf{a}}\text{-a.s.}, \quad (13)$$

as $T \rightarrow \infty$ with $m_\mu = \int_{x=0}^{\infty} x \mu(dx)$. Moreover, if (H2) is fulfilled with $\gamma \geq 2$, then there exists a constant $0 < \sigma^2 < \infty$ s.t. for all $(x_0, \mathbf{a}) \in \mathbb{R}_+ \times \text{supp}(\mathbf{G})$ we have the following convergence in $\mathbb{P}_{x_0, \mathbf{a}}$ -distribution

$$\sqrt{T}(\bar{X}_T - m_\mu) \Rightarrow \mathcal{N}(0, \sigma^2) \text{ as } T \rightarrow \infty. \quad (14)$$

Remark 5 (ON THE LIMITING VARIANCE OF THE SAMPLE MEAN EXPOSURE VALUE) As will be shown in the proof below, the asymptotic variance σ^2 in (14) may be related to the limiting behavior of a certain additive functional of the Markov chain $((X_n, \theta_n, \Delta T_{n+1}))_{n \geq 1}$. In [4] (see also [5]), an estimator of the asymptotic variance of such functionals based on pseudo-renewal properties of the underlying chain (namely, on renewal properties of a Nummelin extension of the chain) has been proposed and a detailed study of its asymptotic properties has been carried out.

Beyond the asymptotic exposure mean or the asymptotic mean time spent by X above a certain threshold, other summary characteristics of the exposure process could be pertinently considered from an epidemiology viewpoint, among which the asymptotic tail conditional expectation $\mathbb{E}_\mu[X \mid X > u]$, denoting by $\mathbb{E}_\mu[\cdot]$ the expectation with respect to μ , after the fashion of risk evaluation in mathematical finance or insurance (see also [57]).

4 Simulation-based statistical inference

We now consider the statistical issues one faces when attempting to estimate certain features of linear rate exposure models. The main difficulty lies in the fact that the exposure process X is generally unobservable. Food consumption data (quantities of consumed food and consumption times) related to a single individual over long time periods are scarcely available in practice. And performing measurements at all consumption times so as to record the food contamination levels appears as not easily realizable. Instead, practitioners have at their disposal some massive databases, in which information related to the dietary habits of large population samples over short periods of time is gathered. Besides, some contamination data concerning certain chemicals and types of food are stored in data warehouses and available for statistical purposes. Finally, experiments for assessing models accounting for the pharmacokinetics behavior in man of various chemicals have been carried out. And data permitting to fit values or probability distributions on the parameters of these models are consequently available.

Estimation of steady-state or time-dependent features of the law \mathcal{L}_X of the process X given the starting point $(X(0), A(0)) = (x_0, \mathbf{a}) \in \mathbb{R}_+ \times \text{supp}(G)$ could thus be based on preliminary computation of consistent estimates \hat{G} , \hat{F}_U and \hat{H} of the unknown distribution functions G , F_U and H . Hence, when no closed form analytic expression for the quantity of interest $\mathcal{Q}(X)$ is available from (G, F_U, H) , ruling out the possibility of computing plug-in estimates, a feasible method could consist in simulating sample paths starting from (x_0, \mathbf{a}) of the approximate process \hat{X} with law $\mathcal{L}_{\hat{X}}$ corresponding to the estimated distribution functions $(\hat{G}, \hat{F}_U, \hat{H})$ and construct estimators based on the trajectories thus obtained.

Beyond stochastic modeling of the exposure phenomenon, the main goal of this paper is to provide theoretical grounds for the application of such statistical methods in toxicological risk evaluation. This leads up to investigate the *stability* of the stochastic model described in section 2 with respect to G , F_U and H (*stability analysis* may be viewed as the counterpart of *sensitivity analysis* in a probabilistic framework), and consider the *continuity problem* consisting in evaluating a measure of closeness between \mathcal{L}_X and $\mathcal{L}_{\hat{X}}$ making the mapping $\mathcal{L}_X \mapsto \mathcal{Q}(X)$ continuous for the functional of interest \mathcal{Q} (refer to [41] for an account on this topic). Hence, convergence preservation results may be obtained via the *continuous-mapping approach* as described in [58], where it is applied to establish stochastic-process limits for queuing systems. For simplicity's sake, we take $\mathbf{a} = 0$ in the following study and do not consider the stability issue related to the approximation of the starting point $(X(0), A(0))$, straightforward modifications of the argument below permitting to deal with the latter problem. For notational convenience, we omit to index by $(x_0, 0)$ the probabilities and expectations considered in the sequel.

Let $0 < T < \infty$. Considering the geometry of the (càd-làg) sample paths of the exposure process X (see Fig. 1), we use the \mathcal{M}_2 topology on the Skorohod's space $\mathbb{D}_T = \mathbb{D}([0, T], \mathbb{R})$ induced by the Hausdorff distance on the space of completed graphs (the completed graph of $x \in \mathbb{D}_T$ being obtained by connecting $(t, x(t))$ to $(t, x(t-))$ with a line segment for all discontinuity points), allowing trajectories to be eventually close even if their jumps do not

exactly match (the \mathcal{J}_2 topology would be actually sufficient for our purpose, refer to [31] or [58] for an account on topological concepts for sets of stochastic processes). In order to evaluate how close the approximating and true laws are, we shall establish an upper bound for the L_1 -Wasserstein Kantorovich distance between the distributions $\mathcal{L}_{X^{(T)}}$ and $\mathcal{L}_{\hat{X}^{(T)}}$ of $X^{(T)} = (X(t))_{t \in [0, T]}$ and $\hat{X}^{(T)} = (\hat{X}(t))_{t \in [0, T]}$, which metric on the space of probability laws on \mathbb{D}_T is defined as follows (refer to [42], [9]):

$$\mathcal{W}_1^{(T)}(\mathcal{L}, \mathcal{L}') = \inf_{Z' \sim \mathcal{L}', Z \sim \mathcal{L}} \mathbb{E}[\mathfrak{m}_{\mathcal{M}_2}^{(T)}(Z', Z)],$$

where the infimum is taken over all pairs (Z', Z) with marginals \mathcal{L}' and \mathcal{L} and $\mathfrak{m}_{\mathcal{M}_2}^{(T)}(Z', Z) = \mathfrak{m}_{\mathcal{H}}^{(T)}(\Gamma_{Z'}, \Gamma_Z)$, denoting by $\Gamma_{Z'}$ and Γ_Z the completed graphs of Z' and Z and by $\mathfrak{m}_{\mathcal{H}}^{(T)}$ the Hausdorff metric on the set of all compact subsets of $[0, T] \times \mathbb{R}$ related to the distance $\mathfrak{m}((t_1, x_1), (t_2, x_2)) = |t_1 - t_2| + |x_1 - x_2|$ on $[0, T] \times \mathbb{R}$. It is well-known that this metric implies weak convergence (see [9]).

As claimed in the next theorem, the law $\mathcal{L}_{\hat{X}^{(T)}}$ gets closer and closer to $\mathcal{L}_{X^{(T)}}$ as the distribution functions \hat{G} , \hat{F}_U and \hat{H} respectively tend to G , F_U and H in the Mallows sense. For $p \in [1, \infty)$, we denote by $M_p(F_1, F_2) = (\int_0^1 |F_1^{-1}(t) - F_2^{-1}(t)|^p dt)^{1/p}$ the L_p -Mallows distance between two distribution functions F_1 and F_2 on the real line.

Theorem 5 *Let (G, F_U, H) (respectively, $(\hat{G}^{(n)}, \hat{F}_U^{(n)}, \hat{H}^{(n)})$ for $n \in \mathbb{N}$) be a triplet of distribution functions on \mathbb{R}_+ defining a linear exposure process X (respectively, $\hat{X}_{(n)}$) starting from $x_0 \geq 0$ and fulfilling Theorem 1's assumptions and (H2) with $\gamma = 1$. Suppose that $(\hat{G}^{(n)}, \hat{F}_U^{(n)}, \hat{H}^{(n)})$ tends to (G, F_U, H) in the L_1 -Mallows distance as $n \rightarrow \infty$. And assume further that G (respectively, $\hat{G}^{(n)}$) has finite variance σ_G^2 (resp., $\sigma_{\hat{G}^{(n)}}^2$) and H (resp., $\hat{H}^{(n)}$) has finite mean. If $\sigma_{\hat{G}^{(n)}}^2$ remains bounded, then:*

$$\sup_{T > 0} T^{-2} \mathcal{W}_1^{(T)}(\mathcal{L}_{X^{(T)}}, \mathcal{L}_{\hat{X}^{(T)}}) \rightarrow 0, \text{ as } n \rightarrow \infty. \quad (15)$$

And for all $T > 0$ we have the weak convergence:

$$\hat{X}_{(n)}^{(T)} \Rightarrow X^{(T)} \text{ in } \mathbb{D}_T, \text{ as } n \rightarrow \infty. \quad (16)$$

Remark 6 Before showing how this theoretical result applies to the problem of approximating/estimating general functionals of the exposure process, a few remarks are in order.

- We point out that similar results hold for the L_p -Wasserstein Kantorovich distance with $p \in [1, \infty)$ under suitable moment conditions.
- It may also be convenient to consider the function space $\mathbb{D}_\infty = \mathbb{D}([0, \infty), \mathbb{R})$ in which X has its sample paths and on which the metric

$$\mathfrak{m}_{\mathcal{M}_2}^{(\infty)}(x, x') = \int_{t=0}^{\infty} 2^{-t} \mathfrak{m}_{\mathcal{M}_2}^{(t)}(\{x(s)\}_{s \in [0, t]}, \{x'(s)\}_{s \in [0, t]}) dt$$

for $(x, x') \in \mathbb{D}_\infty^2$ may be considered. It is noteworthy that (15) also immediately provides a control of the L_1 -Wasserstein distance $\mathcal{W}_1^{(\infty)}$ corresponding to that metric between \mathcal{L}_X and $\mathcal{L}_{\hat{X}}$.

- In statistical applications, one is led to consider random estimates $\hat{G}^{(n)}, \hat{F}_U^{(n)}, \hat{H}^{(n)}$. Clearly, if both the convergence $(\hat{G}^{(n)}, \hat{F}_U^{(n)}, \hat{H}^{(n)}) \rightarrow (G, F_U, H)$ (i.e. 'L₁-consistency' of the distribution estimates) and the boundedness of $\sigma_{\hat{G}^{(n)}}^2$ hold almost surely, then the results of the preceding theorem (and those stated in the next corollary) also hold almost surely.

By showing that good approximations/estimations of the distributions H , G and F_U also induces good approximation/estimation of general functionals of the exposure process, the next result establishes the asymptotic validity of simulation estimators under general conditions.

Corollary 6 *Let (G, F_U, H) (respectively $(\hat{G}^{(n)}, \hat{F}_U^{(n)}, \hat{H}^{(n)})$ for $n \in \mathbb{N}$) be a triplet of distribution functions on \mathbb{R}_+ defining a linear exposure process X (respectively $\hat{X}_{(n)}$) starting from $x_0 \geq 0$ and fulfilling the assumptions of Theorem 5. Let $0 < T \leq \infty$.*

- (i) *Let \mathcal{Q} be a measurable function mapping \mathbb{D}_T into some metric space $(\mathcal{S}, \mathcal{D})$ with $\text{Disc}(\mathcal{Q})$ as set of discontinuity points and such that $\mathbb{P}(X^{(T)} \in \text{Disc}(\mathcal{Q})) = 0$. Then we have the convergence in distribution*

$$\mathcal{Q}(\hat{X}_{(n)}^{(T)}) \Rightarrow \mathcal{Q}(X^{(T)}) \text{ in } (\mathcal{S}, \mathcal{D}).$$

- (ii) *For any Lipschitz function $\phi : (\mathbb{D}_T, \mathfrak{m}_{\mathcal{M}_2}^{(T)}) \rightarrow \mathbb{R}$, we have*

$$\mathbb{E} \left[\phi(\hat{X}_{(n)}^{(T)}) \right] \rightarrow \mathbb{E} \left[\phi(X^{(T)}) \right] \text{ as } n \rightarrow \infty.$$

PROOF. The first assertion derives from Theorem 5 and the convergence (in distribution) preservation result stated in Theorem 3.4.3 of [58], while the second one is an immediate consequence of the first assertion of Theorem 5 (see also [9]). ■

We conclude this section by giving several examples, illustrating how the results above apply to certain functionals of the exposure process in practice. Among the *time-dependent* features of the exposure process, the following quantities are of considerable importance to practitioners in the field of risk assessment of chemicals in food and diet (see [45] and the references therein).

Maximum exposure value. The mapping that assigns to any finite length trajectory $\{x(t)\}_{0 \leq t \leq T} \in \mathbb{D}_T$ its maximum value $\sup_{0 \leq t \leq T} x(t)$ is Lipschitz with respect to the Hausdorff distance $\mathfrak{m}_{\mathcal{M}_2}^{(T)}$ (see Theorem 13.4.1 in [58]). Under the assumptions of

Theorem 5, the expected supremum is finite and, given consistent estimates $\hat{G}^{(n)}$, $\hat{F}_U^{(n)}$ and $\hat{H}^{(n)}$ of G , F_U and H , one may thus construct a consistent estimate of $\mathbb{E}[\sup_{0 \leq t \leq T} X(t)]$ by implementing a standard Monte-Carlo procedure for approximating the expectation $\mathbb{E}[\sup_{0 \leq t \leq T} \hat{X}_{(n)}(t) dt]$.

Mean exposure value. The function $\{x(t)\}_{t \in [0, T]} \in \mathbb{D}_T \mapsto T^{-1} \int_{t=0}^T x(t) dt$ being continuous with respect to \mathcal{M}_2 -topology, we have $T^{-1} \int_{t=0}^T \hat{X}_{(n)}(t) dt \Rightarrow T^{-1} \int_{t=0}^T X(t) dt$ as soon as $\hat{X}_{(n)}^{(T)} \Rightarrow X^{(T)}$ in \mathbb{D}_T . By straightforward uniform integrability arguments, it may be seen that convergence in mean also holds, so that the mean exposure value $\mathbb{E}[T^{-1} \int_{t=0}^T X(t) dt]$ may be consistently estimated by Monte-Carlo simulations.

Average time spent over a critical threshold. Let $u > 0$ be some critical level. In a very similar fashion, it follows from the continuity of $\{x(t)\}_{t \in [0, T]} \in \mathbb{D}_T \mapsto T^{-1} \int_{t=0}^T \mathbb{I}_{\{x(t) \geq u\}} dt$, that a Monte-Carlo procedure also allows to estimate the expectation of the average time spent by the exposure process above the threshold value u , namely $\mathbb{E}[T^{-1} \int_{t=0}^T \mathbb{I}_{\{X(t) \geq u\}} dt]$.

First passage times. Given the starting point x_0 of the exposure process X , the distribution of the first passage time beyond a certain (possibly critical) level $x \geq 0$, *i.e.* the hitting time $\tau_x^+ = \inf\{t \geq 0, X(t) \geq x\}$, is also a characteristic of crucial interest for toxicologists. The mapping $X \in D((0, \infty), \mathbb{R}) \mapsto \tau_x^+$ being continuous w.r.t. the \mathcal{M}_2 -topology (refer to Theorem 13.6.4 in [58]), we have $\hat{\tau}_x^+ = \inf\{t \geq 0, \hat{X}(t) \geq x\} \Rightarrow \tau_x^+$ as soon as $\hat{X} \Rightarrow X$.

In practice, one is also concerned with *steady-state* characteristics, describing the long term behavior of the exposure process. For instance, the steady-state mean exposure m_μ or the limiting time average spent above a given critical value $u > 0$, $\mu([u, \infty[) = \lim_{T \rightarrow \infty} T^{-1} \int_{t=0}^T \mathbb{I}_{\{X(t) \geq u\}} dt$, can be pertinently used as quantitative indicators for chronic risk characterization (see also [57]). As shall be seen below, such quantities may be consistently estimated in a specific asymptotic framework stipulating that both T and n tend to infinity. As a matter of fact, one may naturally write

$$\begin{aligned} \mathbb{E} \left[T^{-1} \int_{t=0}^T \hat{X}_{(n)}(t) dt \right] - m_\mu &= \left\{ \mathbb{E} \left[T^{-1} \int_{t=0}^T \hat{X}_{(n)}(t) dt \right] - \mathbb{E} \left[T^{-1} \int_{t=0}^T X(t) dt \right] \right\} \\ &+ \left\{ \mathbb{E} \left[T^{-1} \int_{t=0}^T X(t) dt \right] - m_\mu \right\}. \end{aligned} \quad (17)$$

And the term between brackets on the left hand side of (17) tends to 0 as $T \rightarrow \infty$ by virtue of Theorem 3, while it follows from the coupling argument of Theorem 5 (see **A4** in the Appendix) that the term on the right hand side is less than $T \times M_1(\hat{F}_U^{(n)}, F_U) + T^2 \times (M_1(\hat{G}^{(n)}, G) + M_1(\hat{H}^{(n)}, H))$ up to a multiplicative constant. Hence, if T and n simultaneously tend to infinity in a way that the latter quantity converges to 0, consistency of $\mathbb{E} \left[T^{-1} \int_{t=0}^T \hat{X}_{(n)}(t) dt \right]$ as an estimator of m_μ is clearly established.

Besides, with regard to statistical applications, Theorem 5 paves the way for studying the asymptotic validity of *bootstrap* procedures in order to construct accurate confidence intervals (based on sample paths simulated from bootstrapped versions of the estimates $\hat{G}^{(n)}$, $\hat{F}_U^{(n)}$ and $\hat{H}^{(n)}$). This is beyond the scope of the present paper but will be the subject of further investigation.

5 Application to methyl mercury data

As an illustration of the mathematical toxicological model analyzed above, some numerical results related to dietary methyl mercury (MeHg) contamination are now exhibited. As previously mentioned, this chemical is present in sea food quasi-solely and a clear indication of its adverse effects on human health has been given by observational epidemiological studies (see [55], [18] and [28] and references therein), leading recently regulatory authorities to develop sea food standards for protecting the safety of the consumer.

At present, dietary risk assessment is conducted from a static viewpoint, comparing the weekly intakes to a reference dose called *Provisional Tolerable Weekly Intake* (PTWI), which is considered to represent the contaminant dose an individual can ingest per week over all his life without appreciable risk. For methyl mercury, the PTWI has been set to 1.6 micrograms per kilogram of body weight per week ($\mu\text{g}/\text{kgbw}/\text{w}$ in abbreviated form) by an international expert committee of FAO/WHO (see [25]). Another reference dose of $0.7 \mu\text{g}/\text{kgbw}/\text{w}$, established from a previous evaluation by the (U.S.) National Research Council [56], is sometimes used from a more conservative perspective. Hence, in a dynamic approach, a deterministic exposure process of reference for risk assessment could be built by considering weekly intakes exactly equal to one of these static reference doses, that is $X_{\text{ref},d} = d/(1 - 2^{-1/\text{HL}})$, where d is the static dose of reference and HL is the biological half-life of the contaminant expressed in weeks. This yields $X_{\text{ref},0.7} = 6.42 \mu\text{g}/\text{kgbw}$ and $X_{\text{ref},1.6} = 14.67 \mu\text{g}/\text{kgbw}$.

Datasets and empirical estimates of distributions F_U , G and H . Food contamination data related to fish and other seafoods available on the French market were generated by accredited laboratories from official national surveys performed between 1994 and 2003 by the French Ministry of Agriculture and Fisheries [35] and the French Research Institute for Exploitation of the Sea [30]. Our dataset comprises of 2832 analytical data.

The national individual consumption survey INCA (see [17]) provides the quantity consumed of an extensive list of foods over a week, among which fish and sea foods, as well as the time when consumption occurred with at least the information about the type of meal (breakfast, lunch, dinner or snack). The survey is composed of two samples: 1985 adults aged 15 years or over and 1018 children aged between 3 to 14 years. Here we consider three subpopulations: men, women, and children under 15.

For simplicity, MeHg intakes are computed through a deterministic procedure currently used in national and international risk assessments. From the INCA food list, 92 different fish or seafood species are determined and a mean level of contamination is computed from

the contamination data, as in [18, 54]. Intakes are then obtained by applying relation (1). For comparability sake, all consumptions are divided by the associated individual body weight, also provided in the INCA survey. After the work of [52], the intake distribution F_U is modeled as a Log-Gamma distribution (*i.e.* $\ln(1 + U)$ is drawn from a Gamma distribution). It is noteworthy that this is a heavy-tailed probability distribution (*i.e.* it fulfills assumption H3, see also Remark 3). It is fitted by means of standard maximum likelihood techniques.

The times of consumption available from the INCA survey allow us to compute the inter-intake times or at least produce right censored values (providing then the information that some durations between successive intakes are larger than a certain time). A Gamma distribution (which has increasing hazard rate) is retained for modeling the inter-intake distribution G : its parameters are fixed for each subpopulation using a right censored maximum likelihood procedure.

The pharmacokinetics in man of MeHg has been thoroughly investigated in several studies (see [44], [50] and [51] for instance), almost all coming to the conclusion that the *half-life* of methyl mercury in man (see Remark 1) fluctuates around six weeks. As we could not dispose of any raw MeHg half-life data for modeling its variability, we choose a Dirac distribution for H putting mass 1 on a *biological half-life* $\log(2)/\theta$ of 6 weeks in the present article.

Table 1 sums up the characteristics of the intake and inter-intake distributions for the three subpopulations, with the convention that a Gamma distribution with parameter α and β has density proportional to $x^{\alpha-1} \exp(-x/\beta)$ and mean $\alpha\beta$. The tail index of the (Log-Gamma) intake distribution is given by $\kappa = 1/\beta$ (see [22], page 153). Observe here that, although the inter-intake times tend to be larger in the children subpopulation than in the adult subpopulation, children are more exposed to extreme risks, insofar as the tail of the corresponding intake distribution (respectively, of the corresponding steady-state distribution) is the heaviest (see Theorem 3). This fact was already mentioned in a former static approach, see [54].


Table 1: Parameters of the intake and inter-intake distributions for each of the three subpopulations fitted by Maximum Likelihood.

		Women	Men	Children	
	sample size	n	1961	1605	1612
Intake distribution (Log-Gamma parameters)	α	0.90	0.85	1.20	
	β	4.88	5.00	3.50	
Mean intake (Unit: $\mu\text{g}/\text{kgbw}$)		m_{F_U}	0.23	0.21	0.50
	sample size	n	2144	1769	1816
Inter-intake distribution (Gamma parameters)	α	1.10	1.03	1.06	
	β	100.88	111.34	148.36	
Mean inter-intake (Unit: hours)		m_G	111	114	157

Time to steady-state. In order to evaluate the time to steady-state, Monte-Carlo simulations have been carried out: using the instrumental distributions described above, $M = 1000$ trajectories have been started from different initial values x_0 . For each path, the (temporal) mean exposure over the time interval $[0, T]$ is computed and individual results have been averaged. In Figure 2(a), the resulting estimates for the women subpopulation are displayed, as time T grows: as expected, all empirical mean exposures converge to the same quantity (namely m_μ) and the relative error is lower than 10% after 24 half-lives (two years and ten months approximately), whatever the starting value x_0 .

The same procedure is used for the mean time spent beyond the reference threshold $u = X_{\text{ref},0.7}$, see Figure 2(b), in the case of the women subpopulation again. The relative error gets now lower than 10% after 58 half-lives (about 6 years and 10 months). Running the same simulations for the men subpopulation, for any of the initial values x_0 considered, the relative errors are less than 10% after only 28 half-lives only, for both the mean exposure and the probability to exceed $X_{\text{ref},0.7}$ in steady-state, while for the children dietary behavior, 28 half-lives are required for the mean exposure and 77 for the probability to exceed $X_{\text{ref},0.7}$ (nearly 9 years).

Naturally, the time to steady-state strongly depends on the initial value x_0 and of the functional of interest. However, on the basis of these simulation results, it may be stated that, for realistic initial values, the time to steady-state for basic quantities as the ones considered here fluctuates between 3 and 10 years, which is, anyhow, a reasonable horizon at the human scale.



(a) Mean exposure of the women subpopulation versus time ($x_0 = 0, 1, \dots, 5$).

(b) Average time spent over u in the women subpopulation versus time ($x_0 = 0, 1, \dots, 5$, $u = X_{\text{ref},0.7} = 6.42 \mu\text{g}/\text{kgbw}$).

Figure 2: Convergence to steady-state

Remark 7 (CONVERGENCE RATE BOUNDS) The problem of determining computable bounds on the convergence rate of ergodic Markov processes has recently received much attention in the applied probability literature (see [37], [48], [34], [46] or [47]). Using suitably calibrated parameters of the drift and minorization conditions (Equations (18), (19), (23), and (22)) established along the proofs of theorems 2 and 3 in the Appendix, rough numerical estimates of the constants involved in rate bounds (10) and (12) can be computed from Theorem 2.2 in [47] (in the discrete-time case) and Theorem 4.1 in [47] (in the continuous-time setup). Explicit computations based on these theoretical results shall be carried out in a forthcoming paper (see [6]). However, as such computable bounds are quite loose in general and are, furthermore, related to the total variation norm (whereas one rather focuses on specific functional of interest in practice), the problem is handled here by exploiting the simulation approach.

Estimation by computer simulation. We now focus on estimating certain relevant time-dependent and steady-state quantities among those enumerated in the previous section via the simulation approach studied in Section 4 from our MeHg datasets. Except for the mean time required for exceeding a given threshold, all quantities of interest are estimated by averaging over $M = 1000$ replicated trajectories on $[0, T]$, taking T equal to 5 years and a starting value x_0 close to the estimated steady-state mean exposure (namely, 2.99 for the women subpopulation, 2.64 for the men subpopulation and 4.61 for the children subpopulation, refer to the preceding paragraph). Numerical results related to the estimation of the steady-state mean exposure (m_μ), the probability to exceed the dynamic reference doses in steady-state ($\mu([X_{\text{ref},0.7}, \infty[)$ and $\mu([X_{\text{ref},1.6}, \infty[)$, the mean time to run over the lowest reference dose ($\mathbb{E}[\tau_{X_{\text{ref},0.7}}]$) and the expected maximum exposure over 5 and 10 years ($\mathbb{E}[\max_{t \leq 5/10 \text{ years}} X(t)]$) are displayed in Table 2 for each of the three subpopulations.

Table 2: Estimated features of the exposure processes - men, women and children

Estimates	(Unit)	Women	Men	Children
x_0	($\mu\text{g}/\text{kgbw}$)	2.99	2.64	4.61
m_μ	($\mu\text{g}/\text{kgbw}$)	3.00	2.66	4.60
$\mu([X_{\text{ref},0.7}, \infty[)$	(%)	0.45%	0.22%	13.36%
$\mu([X_{\text{ref},1.6}, \infty[)$	(%)	0.00%	0.00%	0.15%
$\mathbb{E}[\max_{t \leq 5 \text{ years}} X(t)]$	($\mu\text{g}/\text{kgbw}$)	6.51	5.97	11.99
$\mathbb{E}[\max_{t \leq 10 \text{ years}} X(t)]$	($\mu\text{g}/\text{kgbw}$)	7.24	6.47	13.66
$\mathbb{E}[\tau_{X_{\text{ref},0.7}}]$	(years)	8.12	18.26	0.55

We observe that children constitute the riskiest subpopulation (they actually tend to eat more proportionally to their body weight, even though their sea food consumption frequency is the lowest among the three groups). Figure 3 gives the Monte-Carlo distribution of the mean time to run beyond the threshold $u = X_{\text{ref},0.7} = 6.42 \mu\text{g}/\text{kgbw}$ for the men

and women subpopulations. This clearly illustrates the fact that women may reach the safe reference faster than men (8 years versus 18 years).




Figure 3: Monte-Carlo distribution for the time to run over u for the men and women subpopulations (women in red color, men in blue color, $u = X_{\text{ref},0.7} = 6.42 \mu\text{g}/\text{kgbw}$)

Remark 8 (SENSITIVITY TO THE CHOICE OF THE INSTRUMENTAL DISTRIBUTIONS) The distribution models used here for the governing probability measures F_U and G have been chosen because they provide a good overall fit to the data (and may be easily seen to satisfy all the assumptions required in the ergodicity and stability analyses). According to our own practical experience, the numerical results displayed here would not have been significantly different, if one had chosen to model G by a Weibull distribution for instance. However, in a future study (see [6]) special attention shall be given to the statistical issues of validating the mathematical toxicological model, by investigating how sensitive the latter is to changes with respect to the estimation method chosen (considering also the use of nonparametric approaches).

Remark 9 (NAIVE MONTE-CARLO SIMULATION AND HIGH THRESHOLD) From the perspective of public health guidance practice, it is of prime importance to evaluate the probability of occurrence of rare (extremely risky) events, the probability to exceed a large threshold u such as $u = X_{\text{ref},1.6}$ for instance. In this respect, we point out that the naive Monte-Carlo simulation proposed here leads to estimate this probability by zero (see Table 2), so seldom this threshold is reached on a time interval $[0, T]$ of reasonable length. Treading in the steps of [14], it is shown in [6] that one may remedy to this problem by implementing a suitable *particle filtering algorithm*.

A Appendix - Technical proofs

A.1 Proof of Theorem 2

From conditions required by Theorem 1, aperiodicity and irreducibility properties are immediately established for the discrete-time chain \tilde{X} . Besides, under mild irreducibility conditions, the stability of the *random coefficients autoregressive model* on \mathbb{R}^d

$$Y_{n+1} = \alpha_n Y_n + \beta_n,$$

where (α_n, β_n) , $n = 1, \dots$ are i.i.d. r.v.'s on $\mathbb{R}_+^* \times \mathbb{R}^d$, has been investigated in detail since the seminal contribution of [32] (see [43] and the references therein). Under the assumption that $\mathbb{E}[\log(1 \vee \|\beta_1\|)] < \infty$ and $\mathbb{E}[\log(1 \vee \alpha_1)] < \infty$, it is well known that a sufficient and necessary condition for the chain X to have a (unique) probability measure is that $\mathbb{E}[\log(\alpha_1)] < 0$ (see Corollary 2.7 in [10] for instance). Based on this result, it is then straightforward that, under the assumptions of Theorem 1 and (H1), the chain \tilde{X} is positive recurrent with absolutely continuous stationary probability distribution $\tilde{\mu}(dx) = \tilde{f}(x)dx$. In the discrete-time context, analysis of the stability of Markov models $(Y_n)_{n \in \mathbb{N}}$ may be carried out by establishing suitable conditions for the 'drift' $\Delta V(y) = \mathbb{E}[V(Y_1) | Y_0 = y] - V(y)$ for appropriate non-negative test functions $V(y)$. Such 'Foster-Lyapunov' criteria stipulate the existence of a 'small set' S (*i.e.* an accessible set S to which the chain returns in a given number of steps with positive probability, uniformly bounded by below, see section 5.2 in [39]) toward which the chain drifts in the sense that:

$$\Delta V(x) \leq -f(x) + b\mathbb{I}_{\{x \in S\}}, \quad (18)$$

for some 'norm-like' function $f(x) \geq 1$ and $b < \infty$. Now for the chain \tilde{X} , any compact interval $[0, s]$ with $s > 0$ is small. Indeed, it follows from (9) that for all $x \in [0, s]$, the minorization condition below holds:

$$\Pi(x, \cdot) \geq \delta_s \cdot \mathbb{U}_s(\cdot), \quad (19)$$

with $\delta_s = s \times \inf_{y \in [0, s]} f_{\mathbb{U}}(y) > 0$ and denoting by $\mathbb{U}_s(\cdot)$ the uniform probability distribution on $[0, s]$. When $\gamma = 1$ for instance, take $V(x) = 1 + x$. The affine drift related to \tilde{X} is given by

$$\Delta V(x) = -cx + \mathbb{E}[\mathbb{U}_1],$$

with $c = 1 - \mathbb{E}[e^{-\theta_1 \Delta T_2}] > 0$. Choosing $S = [0, s]$ with $s \geq 1 + 2\mathbb{E}[\mathbb{U}_1]/c$, (18) is fulfilled with $f(x) = cV(x)/2$ and $b = \mathbb{E}[\mathbb{U}_1] + c/2$. Applying Theorem 15.0.1 in [39], we thus get that \tilde{X} is geometrically ergodic with invariant probability measure $\tilde{\mu}$ such that $\tilde{\mu}(V) = \int_{x=0}^{\infty} V(x)\tilde{\mu}(dx) < \infty$. In particular, $\tilde{\mu}$ has finite expectation and there exist constants $r > 1$, $R < \infty$ such that for all $x > 0$:

$$\sum_{n=0}^{\infty} r^n \|\Pi^n(x, \cdot) - \tilde{\mu}\|_V \leq RV(x), \quad (20)$$

with $\|\nu\|_V = \sup_{\psi: |\psi| \leq V} \left| \int \psi(x) \nu(dx) \right|$ for all bounded measure ν on the real line. When $V \equiv 1$, $\|\cdot\|_V$ coincides with the total variation norm $\|\cdot\|_{TV}$. For $\gamma > 1$, the results is proved in a similar fashion by taking $V(x) = 1 + x^\gamma$.

Finally, the last assertion of Theorem 2 immediately derives from Theorem 1 in [29].

A.2 Proof of Theorem 3

Set $X_0 = X(0)$. Observe that for all $t > 0$, $X(t) = X_{N(t)} e^{-\theta_{N(t)} A(t)}$, so that $X(t) \leq X_{N(t)}$. Hence we naturally have $\{X(t) \rightarrow \infty\} \subset \{X_n \rightarrow \infty\}$. Therefore, under (H1), we know that X is positive recurrent with stationary distribution $\tilde{\mu}$, so that in particular $\mathbb{P}(X_n \rightarrow \infty) = 0$. Furthermore, observe that for all $t > 0$, $u \geq 0$:

$$\int_{s=0}^t \mathbb{I}_{\{X(s) \geq u\}} ds = \sum_{k=1}^{N(t)} \int_{s=T_{k-1}}^{T_k} \mathbb{I}_{\{X(s) \geq u\}} ds + \int_{s=T_{N(t)}}^t \mathbb{I}_{\{X(s) \geq u\}} ds.$$

Therefore, for all $k \in \mathbb{N}$, $\int_{s=T_k}^{T_{k+1}} \mathbb{I}_{\{X(s) \geq u\}} ds = \mathbb{I}_{\{X_k \geq u\}} \cdot \Delta T_{k+1} \wedge \frac{\log(X_k/u)}{\theta_k}$.

Now, applying the strong law of large number (SLLN) to the positive recurrent chain $((X_n, \theta_n, \Delta T_{n+1}))_{n \in \mathbb{N}}$ with invariant probability distribution $\tilde{\mu}(dx) \otimes H(d\theta) \otimes G(dt)$, we get that

$$n^{-1} \sum_{k=1}^n \int_{s=T_k}^{T_{k+1}} \mathbb{I}_{\{X(s) \geq u\}} ds \rightarrow \int_{x=u}^{\infty} \int_{\theta \in \Theta} \int_{t=0}^{\infty} t \wedge \frac{\log(x/u)}{\theta} \tilde{\mu}(dx) H(d\theta) G(dt). \quad (21)$$

As we assumed $m_G = \mathbb{E}(\Delta T_k) < \infty$ for $k \geq 2$, we have the following convergence for the delayed renewal process: $N(t)/t \rightarrow m_G^{-1}$ as $t \rightarrow \infty$. Combined with (21), this yields $t^{-1} \int_{s=0}^t \mathbb{I}_{\{X(s) \geq u\}} ds \rightarrow \mu([u, \infty])$ as $t \rightarrow \infty$, with μ given by (11).

We thus proved that $X(t)$ has a limiting probability distribution μ , which has density $f(y)$ given by

$$f(y) = m_G^{-1} \int_{\theta \in \Theta} \int_{t=0}^{\infty} \tilde{f}(y e^{\theta t}) e^{\theta t} \bar{G}(t) dt H(d\theta),$$

denoting by $\bar{G} = 1 - G$ the inter-intake survival function.

Besides, if $\sup \Theta < \infty$, from (11), we immediately have that, for all $u > 0$, $t > 0$,

$$\frac{t \wedge \log 2}{m_G \sup \Theta} \bar{G}(t) \tilde{\mu}([2u, \infty]) \leq \mu([u, \infty]) \leq \tilde{\mu}([u, \infty]).$$

The distributions μ and $\tilde{\mu}$ have thus exactly the same right tail behavior.

Assuming now that (H2) and (H4) are both fulfilled, we turn to the study of the trivariate process $\{(X(t), \theta(t), A(t))\}_{t \geq 0}$. It may be easily seen as $\lambda \otimes H \otimes G$ -irreducible and any compact set $[0, s] \times [0, \bar{\theta}] \times [0, a]$, with $s > 0$, $\bar{\theta} > 0$, $a > 0$ is a 'petite set' for the latter (see [38] for an account of stochastic stability concepts related to continuous-time Markov

processes). Indeed, denote by $Q_t(\cdot | (x_0, \theta_0, a_0))$ the distribution of $(X(t), \theta(t), A(t))$ conditioned upon $(X(0), \theta(0), A(0)) = (x_0, \theta_0, a_0)$. One may easily check that its trace on the event $\{N(t) = 1\}$ has density $f_U(e^{\theta a} x - x_0 e^{-\theta_0(t-a)}) e^{\theta a} \bar{G}(a) g_{a_0}(t-a)$ with respect to $\lambda(dx) \otimes H(d\theta) \otimes \lambda(da)$. Hence, for all $(x_0, \theta_0, a_0) \in [0, s] \times [0, \bar{\theta}] \times [0, a]$, we have the minorization condition:

$$Q_t(\cdot | (x_0, \theta_0, a_0)) \geq \delta(s, \bar{\theta}, a) \cdot \mathbb{U}_s \otimes H(\cdot \cap [0, \bar{\theta}]) \otimes \mathbb{U}_a, \quad (22)$$

with $\delta(s, \bar{\theta}, a) = (s \times H([0, \bar{\theta}]) \times a) \cdot \inf_{x \in [0, se^{\bar{\theta}a}]} f_U(x) \cdot \inf_{v \in [t-a, t]} g(v) \bar{G}(a)$.

Following [33] (see Theorem 4.1 therein), let η, δ be such that $0 < \eta < \delta$ and consider the Lyapounov function $V(x, \theta, t) = (1 + x^\gamma)(1 + \theta)W(t)$ on \mathbb{R}_+^3 with $W(t) = 1 + G(t)e^{-\eta t} \left[1 + \int_{x=t}^{\infty} e^{\delta x} \frac{\bar{G}(x)}{\bar{G}(t)} dx \right]$ (notice that, under (H4), $W(t) \rightarrow \infty$ as $t \rightarrow \infty$). It may be easily seen that the test function V belongs to the domain of the infinitesimal generator \mathcal{G} (see Eq. (5)) and that $\mathcal{G}V(x, \theta, t) = -\eta V(x, \theta, t) + b(x, \theta, t)$, where

$$\begin{aligned} b(x, \theta, t) &= (1 + x^\gamma)(1 + \theta) [z(t) - \theta \gamma x^\gamma W(t)] \\ &\quad - \zeta(t) [(1 + x^\gamma)(1 + \theta) - (1 + \mathbb{E}[\theta])(1 + \mathbb{E}[(x + U)^\gamma])], \\ z(t) &= \eta + \frac{g(t)}{\bar{G}(t)} e^{-\eta t} \left(1 - 2G(t) + \int_{u=t}^{\infty} e^{\delta u} \bar{G}(u) du \right) - G(t) e^{(\delta - \eta)t}. \end{aligned}$$

Observe that $b(x, \theta, t) \rightarrow -\infty$ as (x, θ, t) tends to infinity. Hence, there exist $s > 0, \bar{\theta} > 0$ and $a > 0$ large enough and $b < \infty$ such that the following drift condition holds:

$$\mathcal{G}V(x, \theta, t) \leq -\eta V(x, \theta, t) + b \mathbb{I}_{\{(x, \theta, t) \in [0, s] \times [0, \bar{\theta}] \times [0, a]\}}. \quad (23)$$

Then, (12) directly follows from Theorem 5.3 in [36].

A.3 Proof of Proposition 4

Given $(X(0), A(0)) = (x_0, a)$, we have for all $T > 0$,

$$\bar{X}_T = T^{-1} \int_{t=0}^{T_1} X(t) dt + T^{-1} \sum_{k=1}^{N(T)-1} \int_{t=T_k}^{T_{k+1}} X(t) dt + T^{-1} \int_{T_{N(T)}}^T X(t) dt. \quad (24)$$

The first term in the right-hand side of (24) being bounded by $x_0 T_1 / T$, it almost surely converges to 0 as $T \rightarrow \infty$. Besides we have for all $k \geq 1$,

$$\int_{t=T_k}^{T_{k+1}} X(t) dt = \frac{X_k}{\theta_k} (1 - e^{-\theta_k \Delta T_{k+1}}).$$

Furthermore, by virtue of Theorem 2, assumption (H2) with $\gamma = 1$ ensures that $m_{\bar{\mu}} = \int_{x=0}^{\infty} x \bar{\mu}(dx) < \infty$ and consequently that

$$\tilde{m} = \int_{x=0}^{\infty} \int_{t=0}^{\infty} \int_{\theta \in \Theta} \frac{x(1 - e^{-\theta t})}{\theta} \bar{\mu}(dx) H(d\theta) G(dt) < \infty,$$

making the SLLN for the positive recurrent chain $((X_n, \theta_n, \Delta T_{n+1}))_{n \geq 1}$ applicable to $\sum_{n \geq 1} (1 - \exp(\theta_n \Delta T_{n+1})) X_n / \theta_n$ (refer to Theorem 17.3.2 in [39] for instance). We thus have that

$$N^{-1} \sum_{k=1}^N \frac{X_k}{\theta_k} (1 - e^{-\theta_k \Delta T_{k+1}}) \rightarrow m_{\tilde{\mu}} \int_{t=0}^{\infty} \int_{\theta \in \Theta} \frac{1 - e^{-\theta t}}{\theta} H(d\theta) G(dt) \text{ a.s.}, \quad (25)$$

as $N \rightarrow \infty$. Combining (25) with $N(T)/T \rightarrow m_G^{-1}$ a.s. as $T \rightarrow \infty$, this entails that the third term in (24) tends to 0 as $T \rightarrow \infty$ and establishes (13). Notice that $m_{\mu} = \int_{t=0}^{\infty} \int_{\theta \in \Theta} (1 - \exp(-\theta t)) / \theta H(d\theta) G(dt) m_{\tilde{\mu}} / m_G$.

We now turn to the proof of the Central Limit Theorem (CLT). Using again Theorem 2, we have that $\int x^2 \tilde{\mu}(dx) < \infty$ when (H2) holds for some $\gamma \geq 2$, so that

$$\int_{x=0}^{\infty} \int_{t=0}^{\infty} \int_{\theta \in \Theta} \frac{x^2 (1 - e^{-\theta t})^2}{\theta^2} \tilde{\mu}(dx) H(d\theta) G(dt) < \infty.$$

By virtue of the CLT for positive recurrent chains (see Theorem 17.0.1 in [39]), we have that $N^{-1/2} \sum_{k=1}^N \{(1 - e^{-\theta_k \Delta T_{k+1}}) X_k / \theta_k - \tilde{m}\}$ converges in distribution to $\mathcal{N}(0, \tilde{\sigma}^2)$ as $N \rightarrow \infty$, with $\tilde{\sigma}^2 = \mathbb{E}_{\mu}[(\frac{X_1(1 - e^{-\theta_1 \Delta T_2})}{\theta_1} - \tilde{m})^2] + 2 \sum_{k=2}^{\infty} \mathbb{E}_{\mu}[(\frac{X_1(1 - e^{-\theta_1 \Delta T_2})}{\theta_1} - \tilde{m})(\frac{X_k(1 - e^{-\theta_k \Delta T_{k+1}})}{\theta_k} - \tilde{m})]$.

One may then easily deduce (14) from (24) with $\sigma^2 = \tilde{\sigma}^2 / m_G$.

A.4 Proof of Theorem 5

Observe first that (16) immediately follows from (15) by virtue of standard properties of Wasserstein metrics. In order to prove (15), we construct a specific coupling of the laws $\mathcal{L}_{\hat{X}(T)}$ and $\mathcal{L}_{X(T)}$. Let $(V_k)_{k \in \mathbb{N}}$, $(V'_k)_{k \in \mathbb{N}}$ and $(V''_k)_{k \in \mathbb{N}}$ be three independent sequences of i.i.d. r.v.'s, uniformly distributed on $[0, 1]$. For all $(n, k) \in \mathbb{N}^2$, set

$$\begin{aligned} \Delta T_k &= G^{-1}(V_k), U_k = F_U^{-1}(V'_k), \theta_k = H^{-1}(V''_k), \\ \Delta \hat{T}_k^{(n)} &= \hat{G}^{(n)-1}(V_k), \hat{U}_k^{(n)} = \hat{F}_U^{(n)-1}(V'_k), \hat{\theta}_k^{(n)} = \hat{H}^{(n)-1}(V''_k), \end{aligned}$$

and define recursively for $k \in \mathbb{N}$, $X_{k+1} = X_k e^{-\theta_k \Delta T_{k+1}} + U_{k+1}$ and $\hat{X}_{k+1}^{(n)} = \hat{X}_k^{(n)} e^{-\hat{\theta}_k^{(n)} \Delta \hat{T}_{k+1}^{(n)}} + \hat{U}_{k+1}^{(n)}$ with $X_0 = \hat{X}_0^{(n)} = x_0$, as well as $T_{k+1} = \Delta T_{k+1} + T_k$ and $\hat{T}_{k+1}^{(n)} = \Delta \hat{T}_{k+1}^{(n)} + \hat{T}_k^{(n)}$ with $T_0 = \hat{T}_0^{(n)} = 0$. For notational convenience, the superscript (n) is omitted in the sequel. Using in particular the fact that $x \geq 0 \mapsto e^{-x}$ is 1-Lipschitz, straightforward computations yield

$$|\hat{X}_k - X_k| \leq x_0 \left\{ \sum_{i=1}^k \theta_i |\Delta T_{i+1} - \Delta \hat{T}_{i+1}| + \sum_{i=1}^k \Delta \hat{T}_{i+1} |\theta_i - \hat{\theta}_i| \right\}$$

$$+ \sum_{i=1}^k \mathbf{u}_i \left(\sum_{j=i}^{k-1} \theta_j |\Delta T_{j+1} - \Delta \hat{T}_{j+1}| + \sum_{j=i}^{k-1} \Delta \hat{T}_{j+1} |\theta_j - \hat{\theta}_j| \right) + \sum_{i=1}^k |\hat{\mathbf{u}}_i - \mathbf{u}_i| \quad (26)$$

Turning now to the coupling construction in continuous time, define $\mathbf{N}(t) = \sum_{k \geq 1} \mathbb{I}_{\{T_k \leq t\}}$ and $\hat{\mathbf{N}}(t) = \sum_{k \geq 1} \mathbb{I}_{\{\hat{T}_k \leq t\}}$, as well as $\mathbf{X}(t) = X_{\mathbf{N}(t)} \exp(-\theta_{\mathbf{N}(t)}(t - T_{\mathbf{N}(t)}))$ and $\hat{\mathbf{X}}(t) = \hat{X}_{\hat{\mathbf{N}}(t)} \exp(-\hat{\theta}_{\hat{\mathbf{N}}(t)}(t - \hat{T}_{\hat{\mathbf{N}}(t)}))$ for $t \geq 0$. Set also $T_k^+ = T_k \vee \hat{T}_k$ and $T_k^- = T_k \wedge \hat{T}_k$ for all $k \in \mathbb{N}$ and observe that

$$\mathfrak{m}_{\mathcal{H}}(\Gamma_{\hat{\mathbf{X}}(T)}, \Gamma_{\mathbf{X}(T)}) \leq \max_{0 \leq k \leq \mathbf{N}(T) \vee \hat{\mathbf{N}}(T)} M_k, \quad (27)$$

where

$$\begin{aligned} M_k &= \sup_{T_k^+ \leq t < T_{k+1}^-} |X(t - (\hat{T}_k - T_k)_+) - \hat{X}(t - (T_k - \hat{T}_k)_+)| + |T_k - \hat{T}_k| \\ &+ \sup_{T_{k+1}^- \leq t < T_{k+1}^+} |X(t - (t - T_{k+1})_+) - \hat{X}(t - (t - \hat{T}_{k+1})_+)| + |T_{k+1} - \hat{T}_{k+1}|, \end{aligned}$$

denoting by $x_+ = 0 \vee x$ the positive part of any $x \in \mathbb{R}$. And it follows from easy calculations that

$$\begin{aligned} M_k &\leq |X_k - \hat{X}_k| + X_k (\Delta T_{k+1} \wedge \Delta \hat{T}_{k+1}) |\theta_k - \hat{\theta}_k| + \sum_{i=1}^k |\Delta T_i - \Delta \hat{T}_i| \\ &+ |X_{k+1} - \hat{X}_{k+1}| + |\mathbf{u}_{k+1} - \hat{\mathbf{u}}_{k+1}| + \sum_{i=1}^{k+1} |\Delta T_i - \Delta \hat{T}_i|. \end{aligned}$$

By taking the expectation in (27) and then using the bounds $X_k \leq x_0 + \sum_{1 \leq i \leq k} \mathbf{u}_i$ and (26) combined with Wald's lemma, straightforward computations yield

$$\begin{aligned} \mathbb{E}[\mathfrak{m}_{\mathcal{M}_2}^{(T)}(\hat{\mathbf{X}}(T), \mathbf{X}(T))] &\leq (1 + \mathbb{E}[\mathbf{N}(T) \vee \hat{\mathbf{N}}(T)]) \{ 2x_0 (\mathfrak{m}_{\mathbf{H}} M_1(\mathbf{G}, \hat{\mathbf{G}}) + \mathfrak{m}_{\hat{\mathbf{G}}} M_1(\mathbf{H}, \hat{\mathbf{H}})) \\ &+ 3M_1(\mathbf{F}_{\mathbf{U}}, \hat{\mathbf{F}}_{\mathbf{U}}) + 2M_1(\mathbf{G}, \hat{\mathbf{G}}) + (x_0 + \mathfrak{m}_{\mathbf{F}_{\mathbf{U}}})(T + \mathfrak{m}_{\mathbf{G}} + \mathfrak{m}_{\hat{\mathbf{G}}}) \\ &\times M_1(\mathbf{H}, \hat{\mathbf{H}}) \} + \mathbb{E}[(1 + \mathbf{N}(T) \vee \hat{\mathbf{N}}(T))^2] \mathfrak{m}_{\mathbf{F}_{\mathbf{U}}}(\mathfrak{m}_{\mathbf{H}} M_1(\mathbf{G}, \hat{\mathbf{G}}) \\ &+ \mathfrak{m}_{\hat{\mathbf{G}}} M_1(\mathbf{H}, \hat{\mathbf{H}})), \end{aligned}$$

denoting by $\mathfrak{m}_{\mathbf{F}}$ (resp. $\mathfrak{m}_{\hat{\mathbf{F}}}$) the mean of the distribution function \mathbf{F} (resp. of the estimate $\hat{\mathbf{F}}$), \mathbf{F} being any of the distribution functions \mathbf{G} , $\mathbf{F}_{\mathbf{U}}$ or \mathbf{H} (notice that $\mathfrak{m}_{\hat{\mathbf{F}}} \leq \nu_{\mathbf{F}} + \mathfrak{m}_{\mathbf{F}}$). Besides, there exist constants $C, C' < \infty$ s.t. $\mathbb{E}(\mathbf{N}(T)) \vee \mathbb{E}(\hat{\mathbf{N}}(T)) \leq CT$ and $\mathbb{E}(\mathbf{N}(T)^2) \vee \mathbb{E}(\hat{\mathbf{N}}(T)^2) \leq C'T^2$ (refer to Propositions 6.1 and 6.3 of chap. V in [3] for instance). Observe that the constants C and C' may be chosen independent from the integer n indexing the sequence $\hat{\mathbf{G}}$, since by assumption the sequences $\mathfrak{m}_{\hat{\mathbf{G}}}$ and $\sigma_{\hat{\mathbf{G}}}^2$ are bounded. This establishes the desired result (15).

References

- [1] B. Aberg, L. Ekman, R. Falk, U. Greitz, G. Persson, and J.O. Snihs. Metabolism of methyl mercury compounds in man. *Arch. Environ. Health*, 19(4):478–484, 1969.
- [2] S. Asmussen. Subexponential asymptotics for stochastic processes: Extremal behavior, stationary distributions and first passage probabilities. *Adv. Appl. Probab.*, 8(2):354–374, 1998.
- [3] S. Asmussen. *Applied Probability and Queues*. Springer-Verlag, New York, 2003. Second Edition.
- [4] P. Bertail and S. Cléménçon. Edgeworth expansions for suitably normalized sample mean statistics of atomic Markov chains. *Prob. Th. Rel. Fields*, 130(3):388–414, 2004.
- [5] P. Bertail and S. Cléménçon. Regenerative-block bootstrap for Markov chains. *Bernoulli*, 12(4), 2005.
- [6] P. Bertail, S. Cléménçon, and J. Tressou. Statistical analysis of a dynamic model for food contaminant exposure with applications to dietary methylmercury contamination. 2007. <http://metarisk.inra.fr/>.
- [7] P. Bertail, M. Feinberg, J. Tressou, and P. Verger (Coordinateurs). *Analyse des Risques Alimentaires*. TEC&DOC, Lavoisier, Paris, 2006.
- [8] P. Bertail and J. Tressou. Incomplete generalized U-Statistics for food risk assessment. *Biometrics*, 62(1):66–74, 2006.
- [9] P.J. Bickel and D.A. Freedman. Some asymptotic theory for the bootstrap. *Ann. Statist.*, 9:1196–1217, 1981.
- [10] P. Bougerol and N. Picard. Strict stationarity of generalized autoregressive processes. *Ann. Probab.*, 20(4):1714–1730, 1992.
- [11] M. Brett, H.J. Weimann, W. Cawello, H. Zimmermann, G. Pabst, B. Sierakowski, R. Giesche, and A. Baumann. *Parameters for Compartment-free Pharmacokinetics: Standardisation of Study Design, Data Analysis and Reporting*. Shaker Verlag, Aachen, Germany, 1999.
- [12] P.J. Brockwell, S.I. Resnick, and R.L. Tweedie. Storage processes with general release rule and additive inputs. *Adv. Appl. Probab.*, 14(2):392–433, 1982.
- [13] S. Browne and K. Sigman. Work-modulated queues with applications to storage processes. *J. Appl. Probab.*, 29(3):699–712, 1992.
- [14] F. Cérou, P. Del Moral, F. LeGland, and P. Lezaud. Genetic genealogical models in rare event analysis. *Alea*, 1:183–196, 2006.
- [15] O.L.V. Costa. Stationary distributions for piecewise-deterministic Markov processes. *J. Appl. Probab.*, 27(1):60–73, 1990.
- [16] C. Cox, T.W. Clarkson, D.O. Marsh, L. Amin-Zaki, S. Tikriti, and G.G. Myers. Dose-response analysis of infants prenatally exposed to methylmercury: an application of a single compartment model to stringle-stranded hair analysis. *Environ. Res.*, 49:318–332, 1989.
- [17] CREDOC-AFSSA-DGAL. *Enquête INCA (individuelle et nationale sur les consommations alimentaires)*. TEC&DOC, Lavoisier, Paris, 1999. (Coordinateur : J.L. Volatier).
- [18] A. Crépet, J. Tressou, P. Verger, and J.Ch Leblanc. Management options to reduce exposure to methyl mercury through the consumption of fish and fishery products by the French population. *Regul. Toxicol. Pharmacol.*, 42:179–189, 2005.
- [19] M.H.A. Davis. Piecewise-deterministic Markov processes: A general class of non-diffusion stochastic models. *J. R. Statist. Soc.*, 46(3):353–388, 1984.
- [20] M.H.A. Davis. *Applied Stochastic Analysis*. Stochastics Monographs. Taylor & Francis, 1991.
- [21] L. Edler, K. Poirier, M. Dourson, J. Kleiner, B. Mileson, H. Nordmann, A. Renwick, W. Slob, K. Walton, and G. Würtzen. Mathematical modelling and quantitative methods. *Food Chem. Toxicol.*, 40:283–326, 2002.

- [22] P. Embrechts, C. Klüppelberg, and T. Mikosch. *Modelling Extremal Events for Insurance and Finance*. Applications of Mathematics. Springer-Verlag, 1997.
- [23] R. Falk, J.O. Snihs, L. Ekman, U. Greitz, and B. Aberg. Whole-body measurements on the distribution of mercury-203 in humans after oral-intake of methylradiomercury nitrate. *Acta Radiol.*, 9:55–72, 1970.
- [24] FAO/WHO. Evaluation of certain food additives and contaminants. Fifty-seventh report of the Joint FAO/WHO Expert Committee on Food Additives 909, 2002.
- [25] FAO/WHO. Evaluation of certain food additives and contaminants for methylmercury. Sixty first report of the Joint FAO/WHO Expert Committee on Food Additives, Technical Report Series, WHO, Geneva, Switzerland, 2003.
- [26] M. Gibaldi and D. Perrier. *Pharmacokinetics*. Drugs and the Pharmaceutical Sciences: a Series of Textbooks and Monographs. Marcel Dekker, New York, 1982. Second edition.
- [27] M.J. Gibney and H. van der Voet. Introduction to the Monte Carlo project and the approach to the validation of probabilistic models of dietary exposure to selected food chemicals. *Food Addit. Contam.*, 20(Suppl. 1):S1–S7, 2003.
- [28] P. Grandjean, P. Weihe, R. White, F. Debes, S. Araki, K. Yokoyama, K. Murata, N. Sorensen, R. Dahl, and P. Jorgensen. Cognitive deficit in 7-year-old children with prenatal exposure to methylmercury. *Neurotoxicology Teratology*, 19:417–428, 1997.
- [29] D.R. Grey. Regular variation in the tail behaviour of solutions of random difference equations. *Adv. Appl. Probab.*, 4(1):169–183, 1994.
- [30] IFREMER. Résultat du réseau national d’observation de la qualité du milieu marin pour les mollusques (RNO), 1994-1998.
- [31] J. Jacod and A.N. Shiryaev. *Limit Theorems for Stochastic Processes*. Springer, New York, 1987.
- [32] H. Kesten. Random difference equations and renewal theory for products of random matrices. *Acta Math.*, 131:207–248, 1973.
- [33] T. Konstantopoulos and G. Last. On the use of Lyapunov function methods in renewal theory. *Stoch. Proc. Appl.*, 79:165–178, 1999.
- [34] R. Lund, S.P. Meyn, and R.L. Tweedie. Rates of convergence of stochastically monotone Markov processes. *Ann. Appl. Probab.*, 6:218–237, 1996.
- [35] MAAPAR. Résultats des plans de surveillance pour les produits de la mer. Ministère de l’Agriculture, de l’Alimentation, de la Pêche et des Affaires Rurales, 1998-2002.
- [36] S.P. Meyn and R.L. Tweedie. Stability of Markovian processes III: Foster-Lyapunov criteria for continuous time processes. *Adv. Appl. Probab.*, 25(3):518–548, 1993.
- [37] S.P. Meyn and R.L. Tweedie. Computable bounds for geometric convergence rates of Markov chains. *Ann. Appl. Probab.*, 4(4):981–1011, 1994.
- [38] S.P. Meyn and R.L. Tweedie. A survey of Foster-Lyapunov conditions for general state space Markov processes. In *Proceedings of the Workshop on Stochastic Stability and Stochastic Stabilization*, Metz, France, 1994. Springer-Verlag.
- [39] S.P. Meyn and R.L. Tweedie. *Markov Chains and Stochastic Stability*. Springer-Verlag, 1996.
- [40] J.K. Miettinen. *Mercury, mercurials, and mercaptans*, chapter Absorption and elimination of dietary mercury and methyl mercury in man, pages 223–240. C.C. Thomas, Springfield, IL., M.W. Miller & T.W. Clarkson edition, 1973.
- [41] S.T. Rachev. *Probability metrics and the stability of stochastic models*. Wiley, New York, 1991.
- [42] S.T. Rachev and G. Rüschendorf. *Mass Transportation Problems I: Theory*. Springer, 1998.
- [43] S.T. Rachev and G. Samorodnitsky. Limit laws for a stochastic process and random recursion arising in probabilistic modelling. *Adv. Appl. Probab.*, 27(1):185–202, 1995.

- [44] T. Rahola, R.K. Aaron, and J. K. Miettinen. Half time studies of mercury and cadmium by whole body counting. In *Assessment of Radioactive Contamination in Man*, pages 553–562. Proceedings IAEA-SM-150-13, 1972.
- [45] A. G. Renwick, S. M. Barlow, I. Hertz-Picciotto, A. R. Boobis, E. Dybing, L. Edler, G. Eisenbrand, J. B. Greig, J. Kleiner, J. Lambe, D.J.G. Mueller, S. Tuijelaars and P.A. Van den Brandt, R. Walker, and R. Kroes. Risk characterisation of chemicals in food and diet. *Food Chem. Toxicol.*, 41(9):1211–1271, 2003.
- [46] G.O. Roberts and R.L. Tweedie. Bounds on regeneration times and convergence rates of Markov chains. *Stoch. Proc. Appl.*, 80:211–229, 1999.
- [47] G.O. Roberts and R.L. Tweedie. Rates of convergence of stochastically monotone and continuous time Markov models. *J. Appl. Probab.*, 37(2):359–373, 2000.
- [48] J.S. Rosenthal. Minorization conditions and convergence rates for Markov chain Monte Carlo. *J. Amer. Stat. Assoc.*, 90:558–566, 1995.
- [49] M. Rowland and T.N. Tozer. *Clinical Pharmacokinetics: Concepts and Applications*. 1995.
- [50] J.C. Smith, P.V. Allen, M.D. Turner, B. Most, H.L. Fisher, and L.L. Hall. The kinetics of intravenously administered methyl mercury in man. *Toxicol. Appl. Pharmacol.*, 128:251–256, 1994.
- [51] J.C. Smith and F.F. Farris. Methyl mercury pharmacokinetics in man: A reevaluation. *Toxicol. Appl. Pharmacol.*, 137:245–252, 1996.
- [52] J. Tressou. *Méthodes statistiques pour l'évaluation du risque alimentaire*. PhD thesis, Université Paris X, 2005. <http://tel.archives-ouvertes.fr/tel-00139909>.
- [53] J. Tressou. Non parametric modelling of the left censorship of analytical data in food risk exposure assessment. *J. Amer. Stat. Assoc.*, 101(476):1377–1386, 2006.
- [54] J. Tressou, A. Crépet, P. Bertail, M. H. Feinberg, and J. C. Leblanc. Probabilistic exposure assessment to food chemicals based on extreme value theory. application to heavy metals from fish and sea products. *Food Chem. Toxicol.*, 42(8):1349–1358, 2004.
- [55] T. Tsubaki and K. Irukayama. *Minamata Disease: Methyl-mercury Poisoning in Minamata and Niigata, Japan*. ElsevierScientific, New York, 1977.
- [56] NRC (National Research Council). Committee on the toxicological effects of methylmercury: Toxicological effects of methylmercury. Technical report, Washington DC, 2000.
- [57] P. Verger, J. Tressou, and S. Cléménçon. Integration of time as a description parameter in risk characterisation: application to methylmercury. *To appear in Regulatory Toxicology and Pharmacology*, 2007. Available at <http://metarisk.inapg.inra.fr/UserFiles/File/KDEM.pdf>.
- [58] W. Whitt. *Stochastic-Process Limits. An Introduction to Stochastic-Process Limits and their Application to Queues*. Springer, 2002.