



HAL
open science

In the Heat of the Summer: Lessons from the Heat Waves in Paris.

Emmanuelle Cadot, Victor G. Rodwin, Alfred Spira

► **To cite this version:**

Emmanuelle Cadot, Victor G. Rodwin, Alfred Spira. In the Heat of the Summer: Lessons from the Heat Waves in Paris.. *Journal of Urban Health*, 2007, [Epub ahead of print]. 10.1007/s11524-007-9161-y . inserm-00141162

HAL Id: inserm-00141162

<https://inserm.hal.science/inserm-00141162>

Submitted on 12 Apr 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

IN THE HEAT OF THE SUMMER
Lessons from the heat waves in Paris

E. Cadot¹, V.G. Rodwin² and A. Spira¹

¹ Atelier Parisien de Santé Publique, Service de Santé Publique et d'Epidémiologie, Hôpital de Bicêtre, 82, rue du Général Leclerc, 94276 Le Kremlin Bicêtre

² Wagner School, New York University and World Cities Project, International Longevity Center-USA

Climate change and human health are intertwined¹. The heat waves in Chicago, in 1995, and in Paris, in 2003, followed by Hurricane Katrina's destruction of New Orleans, raised awareness of the risks faced by vulnerable older people. Many cities have responded by announcing emergency preparedness plans; some of these plans have already been tested. Last summer, from July 27 to August 5th, New York City suffered a mild heat wave with temperatures reaching 100°F. Paris, as well, was hit by another heat wave from July 17th to July 29th with maximum temperatures reaching 104°F which was considerably milder than in 2003 when they often exceeded 110°F. In New York, there were 100 "excess deaths," an increase of 8% over past trends². In Paris, the number of excess deaths in 2006 (42), also an increase of 8%, was considerably lower than the 1,294 deaths registered in 2003 -- an increase of 190% compared to the preceding three-year average.

Given existing surveillance capacity, it is impossible to know whether the reduction in excess deaths, in Paris, was due, partly, to its enhanced preparedness or whether it reflects no more than the effects of a far milder heat wave. Nevertheless, the milder heat wave of 2006 does provide an opportunity to examine the actual implementation of the heat wave preparedness plan. In light of on-going efforts to develop such plans in cities, worldwide, and completed studies on the effects of the 2003 heat wave in Paris, what may be learned to promote urban health and improve understanding of the factors that put vulnerable older people at greatest risk?

The Paris heat wave preparedness plan. Approximately 9 months following the catastrophic heat wave of 2003, the Paris Mayor's Office developed a plan to target interventions for those who considered themselves to be most vulnerable. The Mayor wrote a letter to all older Parisians, as well as to all those suffering from severe handicaps and therefore eligible for special services. He invited these 400,000 Parisians to register with the City's Social Services Agency and thereby benefit from direct phone contact and offers of special assistance in the event of another heat wave. In June 2006, about 13,000 people, mostly those 65 years and over, had registered. On July 17th, the plan was activated when the National Weather Institute issued a heat wave warning. Until the 28th of July, the Social Services Agency phoned all registered individuals every other day, to check on them. When those who registered required assistance, based on a preliminary

screening, they were evaluated, once again, by physicians, to determine, their health status. Nearly 800 older people were evaluated by a medical team over these 11 days; 200 were called back. Most of the City's interventions were limited to providing information on what to do and where to go during the heat wave. About 30 people were transported to air conditioned adult day centers and 18 benefited from urgent medical attention.

Since the 2006 heat wave was milder in duration and most meteorological characteristics than the one in 2003, it is difficult to attribute the lower number of excess deaths only to the city's preparedness plan. It seems plausible, however, that the plan increased the number of daily contacts for some socially isolated older people and restored for them – in some small way -- a social network. Based on a synthesis of studies about the 2003 heat wave, this is precisely what must be done.

The August 2003 heat wave. Between August 1-20th, 2003, a deadly European heat wave resulted in some 14,800 excess deaths in France, an increase of 55%³. Paris was disproportionately affected (an increase of 190%) In contrast to the rest of France where 65% of excess mortality afflicted institutionalized older people, in Paris 74% of excess deaths occurred among those who were living at home.

It is well-known that Paris has a higher share of people 80 years and over than the rest of France. Also 66% of Parisian women in this age cohort live alone, which far exceeds the average for France (50%). But neither age, nor gender density, nor temperature levels, are sufficient to explain the concentration of excess deaths in Paris compared to the rest of France. Do they reflect the individual characteristics of Parisians or the character of the health care system and the neighborhoods in which they lived? The health system seems an implausible suspect, in light of its high performance along many dimensions⁴. But since this catastrophe took French physicians and the public health establishment by surprise, it is important to review at least partial answers to these questions, particularly since the evidence has not been widely available in English.

Findings to date. The most important risk factors for dying during the heat wave of 2003 were being a woman 75 years and over, and living alone at home⁵. Parisians who were unmarried (and more frequently live alone) were also at higher risk of death than with those who lived as couples. Higher risk associated with gender (being a woman), however, did not apply to foreign nationals. Older women, particularly those from Africa

and Asia – in contrast to French women – live, more often, in multi-generational families with stronger mutual support and social networks; hence a lower risk of social isolation, as Klinenberg found in his “autopsy” of the 1995 Chicago heat wave⁶. Moreover, mortality rates were highest in the south-eastern parts of the city, indicating a shift in the geographic mortality distribution from poorer to more well-to do neighborhoods, in comparison to the usual mortality gradient across Paris neighborhoods.

These findings could reflect temperature variations, as well as socio-economic and demographic differences among the north-east and south-west arrondissements of Paris.

Among a sample of vulnerable older people who received a personal autonomy allowance (PAA), based on their level of disability, living alone was the only significant risk factor for excess mortality. Whether PAA beneficiaries were community-dwelling or institutionalized, their disability levels were not significant risk factors. Among older nursing-home residents, Holstein et. al. followed a cohort of 4,403 residents from May through December of 2003 and found that women in better health were at higher risk of death than their more vulnerable counterparts. During the heat wave, there was an increased risk of death for all residents, but the mortality increase was significantly lower for the most dependent residents. Analysis of the weekly distribution of deaths reveals no “harvesting effect” – no compensating reduction in mortality after the heat wave, neither in the final days of August, nor in the weeks that followed. Thus, the deceased were not at risk of dying from medical, or any other causes, in the aftermath of the heat wave.

Lessons and directions for heat-wave preparedness: Perhaps the most important lesson of the Paris heat waves is to recognize that preparedness efforts and promotion of urban health must focus not only on individual, but also on neighborhood risk factors. The essential components of the heat wave preparedness plan focused on individual risk factors, e.g. encouraging older persons to drink more, offering special services to those who perceived themselves to be in need of assistance, and increasing medical surveillance of these individuals. In the future, more attention should focus on targeting some of the neighborhoods that suffered disproportionate excess mortality.

Notes

¹ Epstein, P. Climate Change and Human Health. *N Engl J Med.* 2005; 353: 14.

² *Deaths Associated with Heat Waves in 2006.* New York: NYCDHMH, November, 2006 (See internet site: <http://www.nyc.gov/html/doh/downloads/pdf/survey/survey-2006heatdeaths.pdf>).

³ A. Fouillet A, Rey G, Laurent F et al. Excess mortality related to the August 2003 heat wave in France. *Int Arch Occup Environ Health.* 2006; 80:16-24.

⁴ Gusmano MK, Rodwin VG, Weisz D. A new way to compare health systems: avoidable hospital conditions in Manhattan and Paris. *Health Aff.* 2006; 25:510-520.

⁵ Canouï-Poitaine F, Cadot E, Spira A. Excess deaths during the August 2003 heat wave in Paris, France. *Rev Epidemiol Sante Publique.* 2006; 54: 127-35.

⁶ Klinenberg E. *Heat Wave. A social autopsy of disaster in Chicago.* Chicago : The University of Chicago Press; 2002.