

A note on the entropy production formula

Vojkan Jakšić¹ and Claude-Alain Pillet^{2,3}

¹Department of Mathematics and Statistics
McGill University
805 Sherbrooke Street West
Montreal, QC, H3A 2K6, Canada

²PHYMAT, Université de Toulon
B.P. 132, 83957 La Garde Cedex, France

³FRUMAM
CPT-CNRS Luminy, Case 907
13288 Marseille Cedex 9, France

September 11, 2002

Abstract

We give an elementary derivation of the entropy production formula of [JP1]. Using this derivation we show that the entropy production of any normal, stationary state is zero.

1 Introduction

Let \mathcal{O} be a C^* - algebra, $E(\mathcal{O})$ the set of all states on \mathcal{O} and $\omega \in E(\mathcal{O})$. We assume that there exists a reference C^* - dynamics σ_ω^t on \mathcal{O} such that ω is a $(\sigma_\omega, -1)$ -KMS state. We denote by δ_ω the generator of σ_ω^t (i.e. $\sigma_\omega^t = e^{t\delta_\omega}$) and by $\mathcal{D}(\delta_\omega)$ its domain. Let $(\mathcal{H}_\omega, \pi_\omega, \Omega_\omega)$ be the GNS-representation of the algebra \mathcal{O} associated to the state ω .

A state $\eta \in E(\mathcal{O})$ is called ω - normal if there exists a density matrix ρ_η on \mathcal{H}_ω such that, for all $A \in \mathcal{O}$, $\eta(A) = \text{Tr}(\rho_\eta \pi_\omega(A))$. Let \mathcal{N}_ω be the set of all ω - normal states on \mathcal{O} .

For $\eta \in \mathcal{N}_\omega$, we denote by $\text{Ent}(\eta|\omega)$ the relative entropy of Araki [Ar1, Ar2]. (We use the notational convention for relative entropy of [BR, Don].) If $\eta \notin \mathcal{N}_\omega$, we set $\text{Ent}(\eta|\omega) = -\infty$. For unitary $U \in \mathcal{O}$ and $\eta \in E(\mathcal{O})$, we denote by η_U the state $\eta_U(A) \equiv \eta(U^*AU)$. The main result of this note is:

Theorem 1.1 *For any unitary $U \in \mathcal{O} \cap \mathcal{D}(\delta_\omega)$ and any $\eta \in E(\mathcal{O})$,*

$$\text{Ent}(\eta_U|\omega) = \text{Ent}(\eta|\omega) - i\eta(U^*\delta_\omega(U)). \quad (1.1)$$

As we shall explain below, Theorem 1.1 is a natural generalization of the entropy production formula derived in [JP1, JP2]. The method of proof we will use in this note, however, is quite different from the one in [JP1]. We will reduce the proof of Theorem 1.1 to a fairly elementary application of some well known identities in Araki's theory of perturbation of KMS structure. The proof in [JP1], based on Araki-Connes cocycles, was technically more involved and restricted to *faithful* states $\eta \in \mathcal{N}_\omega$.

We now relate Equ. (1.1) to the entropy production formula of [JP1, JP2]. Assume that there exists a C^* -dynamics τ^t on \mathcal{O} and that ω is τ -invariant. Let $V(t)$ be a time-dependent local perturbation, that is, $V(t)$ is norm-continuous, self-adjoint, \mathcal{O} -valued function on \mathbb{R} (the time-independent case of [JP1] of course follows by setting $V(t) \equiv V$). The perturbed time evolution is the strongly continuous family of $*$ -automorphisms of \mathcal{O} given by the formula

$$\tau_V^t(A) \equiv \tau^t(A) + \sum_{n \geq 1} i^n \int_0^t dt_1 \int_0^{t_1} dt_2 \cdots \int_0^{t_{n-1}} dt_n [\tau^{t_n}(V(t_n)), [\cdots, [\tau^{t_1}(V(t_1)), \tau^t(A)]]].$$

In the interaction representation, τ_V^t is given by

$$\tau_V^t(A) = \Gamma_V^t \tau^t(A) \Gamma_V^{t*},$$

where $\Gamma_V^t \in \mathcal{O}$ is a family of unitaries satisfying the differential equation

$$\frac{d}{dt} \Gamma_V^t = i \Gamma_V^t \tau^t(V(t)), \quad \Gamma_V^0 = \mathbf{1}.$$

Theorem 1.1 then has the following immediate corollary (see also Theorem 4.8 in [JP2]):

Corollary 1.2 *Assume that ω is τ -invariant and that $\Gamma_V^t \in \mathcal{D}(\delta_\omega)$. Then, for any $\eta \in E(\mathcal{O})$,*

$$\text{Ent}(\eta \circ \tau_V^t|\omega) = \text{Ent}(\eta|\omega) - i\eta(\Gamma_V^t \delta_\omega(\Gamma_V^{t*})). \quad (1.2)$$

From now on we will consider the time-independent case $V(t) \equiv V$. If $V \in \mathcal{D}(\delta_\omega)$, then $\Gamma_V^t \in \mathcal{D}(\delta_\omega)$ and

$$\frac{d}{dt} \Gamma_V^t \delta_\omega(\Gamma_V^{t*}) = -i \tau_V^t(\delta_\omega(V)). \quad (1.3)$$

Hence, (1.2) reduces to the entropy production formula of [JP1]:

$$\text{Ent}(\eta \circ \tau_V^t | \omega) = \text{Ent}(\eta | \omega) - \int_0^t \eta \circ \tau_V^s(\delta_\omega(V)) ds. \quad (1.4)$$

We emphasize that the above derivation of (1.4) allows for non-faithful η .

The entropy production of a state $\eta \in E(\mathcal{O})$ was defined in [JP1, JP2] by $\text{Ep}_V(\eta) \equiv \eta(\delta_\omega(V))$, see also [OHI, O1, O2, Ru, Sp]. On physical grounds, it is natural to conjecture that if η is ω -normal and τ_V -invariant, then $\text{Ep}_V(\eta) = 0$. For faithful η this was proven in [JP1]. Here, we establish this result in full generality.

Theorem 1.3 *Assume that ω is τ -invariant, that $V \in \mathcal{D}(\delta_\omega)$ and that η is τ_V -invariant and ω -normal. Then,*

$$\text{Ep}_V(\eta) = 0.$$

Remark. If $\text{Ent}(\eta | \omega) > -\infty$, then Theorem 1.3 is an immediate consequence of Equ. (1.4). The case $\text{Ent}(\eta | \omega) = -\infty$ requires a separate and somewhat delicate argument.

The results of this note were announced in the recent review [JP2] where the interested reader may find additional information and references about entropy production and its role in non-equilibrium quantum statistical mechanics.

Acknowledgment. The research of the first author was partly supported by NSERC. A part of this work has been done during the visit of the second author to the McGill University which was supported by NSERC.

2 Proof of Theorem 1.1

We assume that the reader is familiar with basic results of Tomita-Takesaki modular theory as discussed, for example, in [BR, DJP, Don, OP].

Let $\mathfrak{M}_\omega \equiv \pi_\omega(\mathcal{O})''$ be the enveloping von Neumann algebra. Since ω is $(\sigma_\omega, -1)$ -KMS state, the vector Ω_ω is separating for \mathfrak{M}_ω , and we denote by \mathcal{P} , J , Δ_ω the corresponding natural cone, modular conjugation and modular operator. We recall that $\Delta_\omega = e^{L_\omega}$, where L_ω is the unique self-adjoint operator on \mathcal{H}_ω such that

$$\pi_\omega(\sigma_\omega^t(A)) = e^{itL_\omega} \pi_\omega(A) e^{-itL_\omega}, \quad L_\omega \Omega_\omega = 0.$$

In particular, σ_ω^t extends naturally to a W^* -dynamics on \mathfrak{M}_ω which we again denote by σ_ω^t . In this context σ_ω^t is called modular dynamics.

Any state $\eta \in \mathcal{N}_\omega$ has a unique normal extension to \mathfrak{M}_ω which we denote by the same letter. Obviously, η is ω -normal iff η_U is ω -normal for all unitaries $U \in \mathcal{O}$ and so, in the proof of Theorem 1.1, we may restrict ourselves to ω -normal η 's.

We will use the fact that if $\gamma : \mathfrak{M}_\omega \mapsto \mathfrak{M}_\omega$ is a $*$ -automorphism, then

$$\text{Ent}(\eta \circ \gamma | \omega \circ \gamma) = \text{Ent}(\eta | \omega).$$

In particular,

$$\text{Ent}(\eta_U | \omega) = \text{Ent}(\eta | \omega_{U^*}).$$

Let Ψ_{U^*} be the unique vector representative of the state ω_{U^*} in the cone \mathcal{P} . A simple computation shows that

$$\Psi_{U^*} = \pi_\omega(U^*) J \pi_\omega(U^*) \Omega_\omega.$$

We will consider $P \equiv \pi_\omega(-iU^* \delta_\omega(U))$ as a local perturbation of the modular group σ_ω^t . Let α^t be the locally perturbed W^* -dynamics,

$$\alpha^t(A) \equiv e^{it(L_\omega + P)} A e^{-it(L_\omega + P)} = \Theta_P^t \sigma_\omega^t(A) \Theta_P^{t*},$$

where $e^{it(L_\omega + P)} e^{-itL_\omega} \equiv \Theta_P^t \in \mathfrak{M}_\omega$ is a family of unitaries satisfying

$$\frac{d}{dt} \Theta_P^t = i \Theta_P^t \sigma_\omega^t(P), \quad \Theta_P^0 = \mathbf{1}. \quad (2.5)$$

Let ψ be the unique $(\alpha, -1)$ -KMS state on \mathfrak{M}_ω . By the Araki theory, $\Omega_\omega \in \mathcal{D}(e^{(L_\omega + P)/2})$ and the unique vector representative of ψ in the natural cone \mathcal{P} is

$$\Psi = \frac{e^{(L_\omega + P)/2} \Omega_\omega}{\|e^{(L_\omega + P)/2} \Omega_\omega\|}.$$

Another fundamental result of Araki's theory is the relation

$$\text{Ent}(\eta | \psi) = \text{Ent}(\eta | \omega) + \eta(P) - \log \|e^{(L_\omega + P)/2} \Omega_\omega\|^2, \quad (2.6)$$

which holds for all ω -normal states η . (For η faithful, this relation was proven in [Ar1, Ar2], see also [BR]. Its extension to general η was obtained in [Don], see also the next section). Hence, to finish the proof it suffices to show that $e^{(L_\omega + P)/2} \Omega_\omega = \Psi_{U^*}$.

We set $T^t \equiv U^* \sigma_\omega^t(U)$ and observe that

$$\frac{d}{dt} T^t = iT^t \sigma_\omega^t(-iU^* \delta_\omega(U)), \quad T^0 = \mathbf{1}.$$

Comparison with Equ. (2.5) immediately leads to $\pi_\omega(T^t) = \Theta_P^t$ and therefore

$$\begin{aligned} e^{it(L_\omega+P)}\Omega_\omega &= \pi_\omega(T^t)e^{itL_\omega}\Omega_\omega \\ &= \pi_\omega(U^*)e^{itL_\omega}\pi_\omega(U)\Omega_\omega. \end{aligned} \tag{2.7}$$

Since the vector-valued function $z \mapsto e^{iz(L_\omega+P)}\Omega_\omega$ is analytic inside the strip $-1/2 < \text{Im } z < 0$ and strongly continuous on its closure, analytic continuation of the identity (2.7) to $z = -i/2$, yields

$$\begin{aligned} e^{(L_\omega+P)/2}\Omega_\omega &= \pi_\omega(U^*)\Delta_\omega^{1/2}\pi_\omega(U)\Omega_\omega \\ &= \pi_\omega(U^*)J\pi_\omega(U^*)\Omega_\omega \\ &= \Psi_{U^*}, \end{aligned}$$

which is the desired relation.

3 Proof of Theorem 1.3

Let Ω_η be the vector representative of η in the natural cone \mathcal{P} . The standard Liouvillean associated to the dynamics τ_V^t is $L_V = L + \pi_\omega(V) - J\pi_\omega(V)J$, where L is the standard Liouvillean associated to τ^t . We recall that L and L_V are uniquely specified by

$$\pi_\omega(\tau^t(A)) = e^{itL}\pi_\omega(A)e^{-itL}, \quad L\Omega_\omega = 0,$$

and

$$\pi_\omega(\tau_V^t(A)) = e^{itL_V}\pi_\omega(A)e^{-itL_V}, \quad L_V\Omega_\eta = 0.$$

We denote by s_η the support of the state η and set $s'_\eta = Js_\eta J$. Obviously

$$s_\eta\Omega_\eta = s'_\eta\Omega_\eta = \Omega_\eta,$$

and since η is τ_V -invariant

$$e^{itL_V}s_\eta = s_\eta e^{itL_V}, \quad e^{itL_V}s'_\eta = s'_\eta e^{itL_V}.$$

Let $\Delta_{\omega|\eta}$ be the relative modular operator. We recall that $\text{Ker } \Delta_{\omega|\eta} = \text{Ker } s'_\eta$,

$$J\Delta_{\omega|\eta}^{1/2}A\Omega_\eta = s'_\eta A^*\Omega_\omega,$$

for all $A \in \mathfrak{M}_\omega$ and that $\Delta_{\omega|\eta}$ is essentially self-adjoint on $\mathfrak{M}_\omega\Omega_\eta + (\mathbf{1} - s'_\eta)\mathcal{H}_\omega$. Hence

$$\Delta_{\omega \circ \tau_V^t | \eta \circ \tau_V^t} = e^{-itL_V}\Delta_{\omega|\eta}e^{itL_V},$$

and since η is τ_V -invariant,

$$e^{itL_V} \Delta_{\omega|\eta} e^{-itL_V} = \Delta_{\omega \circ \tau_V^{-t}|\eta} = \Delta_{\omega_{U^*}|\eta},$$

where $U^* \equiv \Gamma_V^t$.

As in the proof of Theorem 1.1, we set $P \equiv \pi_\omega(-iU^* \delta_\omega(U))$ and denote by α the perturbation of the modular dynamics σ_ω by P . It follows that ω_{U^*} is the unique $(\alpha, -1)$ -KMS state. Since also $\|e^{(L_\omega+P)/2} \Omega_\omega\| = 1$, the basic perturbation formula of Araki-Donald (see Lemma 5.7 in [Don]) yields

$$s'_\eta \log \Delta_{\omega_{U^*}|\eta} = s'_\eta \log \Delta_{\omega|\eta} - s'_\eta P.$$

Hence,

$$e^{itL_V} s'_\eta \log \Delta_{\omega|\eta} e^{-itL_V} = s'_\eta \log \Delta_{\omega|\eta} - s'_\eta P,$$

and we conclude that for any real number $\lambda \neq 0$,

$$e^{itL_V} (s'_\eta \log \Delta_{\omega|\eta} + i\lambda)^{-1} e^{-itL_V} = (s'_\eta \log \Delta_{\omega|\eta} - s'_\eta P + i\lambda)^{-1}.$$

Since $e^{-itL_V} \Omega_\eta = \Omega_\eta$, the second resolvent identity yields that for all real $\lambda \neq 0$,

$$(\Omega_\eta, (s'_\eta \log \Delta_{\omega|\eta} + i\lambda)^{-1} s'_\eta P (s'_\eta \log \Delta_{\omega|\eta} - s'_\eta P + i\lambda)^{-1} \Omega_\eta) = 0.$$

Since

$$s - \lim_{\lambda \rightarrow \infty} i\lambda (s'_\eta \log \Delta_{\omega|\eta} + i\lambda)^{-1} = \mathbf{1},$$

and

$$s - \lim_{\lambda \rightarrow \infty} i\lambda (s'_\eta \log \Delta_{\omega|\eta} - s'_\eta P + i\lambda)^{-1} = \mathbf{1},$$

we derive that

$$(\Omega_\eta, P \Omega_\eta) = (\Omega_\eta, s'_\eta P \Omega_\eta) = 0.$$

On the other hand, using Equ. (1.3), we get

$$P = \pi_\omega(-iU^* \delta_\omega(U)) = - \int_0^t \pi_\omega(\tau_V^s(\delta_\omega(V))) ds,$$

and since η is τ_V -invariant we conclude that

$$0 = (\Omega_\eta, P \Omega_\eta) = - \int_0^t \eta \circ \tau_V^s(\delta_\omega(V)) ds = -t\eta(\delta_\omega(V)),$$

for all t . This yields the statement.

References

- [Ar1] Araki, H.: Relative entropy of states of von Neumann algebras. *Publ. Res. Inst. Math. Sci. Kyoto Univ.* **11**, 809 (1975/76).
- [Ar2] Araki, H.: Relative entropy of states of von Neumann algebras II. *Publ. Res. Inst. Math. Sci. Kyoto Univ.* **13**, 173 (1977/78).
- [BR] Bratelli, O., Robinson, D.W.: *Operator Algebras and Quantum Statistical Mechanics 2*. Springer-Verlag, Berlin (1987).
- [DJP] Dereziński, J., Jakšić, V., Pillet, C.-A.: Perturbation theory of W^* -dynamics, Liouvillean and KMS-states. Submitted, mp-arc 01-440.
- [Don] Donald, M.J.: Relative Hamiltonians which are not bounded from above. *J. Func. Anal.* **91**, 143 (1990).
- [JP1] Jakšić, V., Pillet, C.-A.: On entropy production in quantum statistical mechanics. *Commun. Math. Phys.* **217**, 285 (2001).
- [JP2] Jakšić, V., Pillet, C.-A.: Mathematical theory of non-equilibrium quantum statistical mechanics. *J. Stat. Phys.* **108**, 787 (2002).
- [OHI] Ojima, I., Hasegawa, H., Ichiyang, M.: Entropy production and its positivity in non-linear response theory of quantum dynamical systems. *J. Stat. Phys.* **50**, 633 (1988).
- [O1] Ojima, I.: Entropy production and non-equilibrium stationarity in quantum dynamical systems: physical meaning of Van Hove limit. *J. Stat. Phys.* **56**, 203 (1989).
- [O2] Ojima, I.: Entropy production and non-equilibrium stationarity in quantum dynamical systems, in *Proceedings of international workshop on quantum aspects of optical communications*. *Lecture Notes in Physics* **378**, 164, Springer-Verlag, Berlin (1991).
- [OP] Ohya, M., Petz, D.: *Quantum Entropy and its Use*. Springer-Verlag, Berlin (1993).
- [Ru] Ruelle, D.: Entropy production in quantum spin systems. *Commun. Math. Phys.* **224**, 3 (2001).
- [Sp] Spohn, H.: Entropy production for quantum dynamical semigroups. *J. Math. Phys.* **19**, 227 (1978).