



« The concept of quantum state: new views on old phenomena », in Ashtekar, Abhay, Cohen, Robert S., Howard, Don, Renn, Jürgen, Sarkar, Sahotra & Shimony, Abner (eds.), *Revisiting the Foundations of Relativistic Physics : Festschrift in Honor of John Stachel, Boston Studies in the Philosophy and History of Science*, Dordrecht: Kluwer Academic Publishers, 451-478. , 2003

The concept of quantum state : new views on old phenomena

par

Michel PATY*

ABSTRACT.

Recent developments in the area of the knowledge of quantum systems have led to consider as physical facts statements that appeared formerly to be more related to interpretation, with free options. Of such a nature are the quantum behaviors of individual particles (diffraction, etc.), neutrinos oscillations, distant quantum correlations (local non-separability), Bose-Einstein condensation, cooling isolation of atoms and, recently, decoherence of quantum superposition states interacting with environment, that allows a better understanding of the transition from the quantum domain to the classical-macroscopic one. The debate on the interpretation of quantum mechanics has imperceptibly changed its nature through these developments, giving higher weight to a «physical interpretation» more clearly distinct from the philosophical one than in the old days of quantum mechanics. In particular, the concept of quantum state has undoubtedly acquired a direct physical meaning, in terms of *properties of a physical system* that is fully represented by a linear superposition of eigenstates, and able to propagate as such in space and time. The price for this new situation is an extension of meaning of the concepts of *physical magnitude* and *physical state* towards ones that do not correspond directly with numerical values.

* Equipe REHSEIS (UMR 7596, Centre National de la Recherche Scientifique (CNRS) et Université Paris 7-Denis Diderot), 37 rue Jacob, F-75006 Paris, France. E-mail : paty@paris7.jussieu.fr

“I think (...) that a theory cannot be produced from results of observations, but only from an invention”.

“I am not ashamed to put the concept of «real state of a physical system» [“existing objectively, independently of any observation or measure, and that can in principle be described through the means of expression of physics”] at the very centre of my meditation”.

Albert Einstein¹.

“(...) It is interesting to speculate on the possibility that a future theory will not be *intrinsically* ambiguous and approximate. Such a theory could not be fundamentally about «measurements», for that would again imply incompleteness of the system and unanalyzed interventions from outside. Rather, it should again become possible to say of a system not that such and such may be *observed* to be so, but that such and such *be* so. The theory would not be about *observables*, but about «beables»”.

John S. Bell²

¹ Respectively : Einstein, letter to Karl Popper, 11.9.1935, published as an appendix in Popper [1959] ; Einstein [1953], p. 6-7.

² Bell [1973, 1984].

1

INTRODUCTION. STATE FUNCTION AND «DIRECT REPRESENTATION» OF A QUANTUM SYSTEM OR STATE

Quantum physics is aimed at the deep structure of matter in general, from bodies of our environment and molecular associations of atoms up to atomic nuclei and to elementary particles actually or «virtually» contained in the latter, and up as well to cosmic objects and to the primordial phases of cosmology. It ensures the unity of matter in the variety of its organization patterns, the means of the theoretical understanding of this domain of physics being constituted around *quantum mechanics*. This last one is, in turn, applied to particular (atomic, nuclear) theoretical models, and enlarged, from a more fundamental point of view, to a *quantum field theory*, from quantum electrodynamics (QED)³ to electroweak and chromodynamic gauge field theories⁴. These recent developments have been performed within the conceptual frame of quantum mechanics and, as an effect, have confirmed its heuristic power and permitted at the same time physicists to get used to work with this tool-for-thought that is indispensable to explore quantum phenomena.

Quantum mechanics, as a theoretical scheme, is *practised* successfully, and today physicists, while applying strictly, in the working process of their physical thought on phenomena under study, the rules that govern the use of quantum magnitudes, no longer do worry very much about the «difficulties of the interpretation» that had heavily preoccupied the founding fathers and their immediate successors. As for interpretation, if they had to propose any, this would be for most of them the following : “what is important is that it works”. And, indeed, this might be a mark of unconcern or the expression of an immediately pragmatic philosophy that would remain blind to deeper reasons. This attitude comprises, in all events, a part of truth, of the kind of that we walk before knowing how : they have the theoretical (and even conceptual) tool and know to handle it before knowing exactly its nature, and worrying about it.

However, as soon as they ask themselves questions concerning the intelligibility of physical phenomena in the quantum area, they find again the terms of the old debate. But, differently from their elders, they meet these «from outside», so to speak, in this sense that these questions appear to them as posed only to a «second order» of the understanding : i.e., when they question themselves *not about this understanding*, that is itself provided by theory, a theory they know so well that it has become a «second nature» to their thought, but *about the reasons of it*.

To understand «to the first degree», that is at the level of their work in physics itself, is got at through handling the concepts and magnitudes that represent, reproduce or create the phenomena of interest. When physicists speak,

³ On the history of quantum electrodynamics, see Schweber [1994].

⁴ Cf., f. ex., Bimbot & Paty [1996].

today, of an «elementary particle» (for example, a proton), they mean, indeed, that it is described by «quantum numbers» or quantities that are «eigenvalues» of operators representing the adequate physical magnitudes and they have abandoned the classical image of a directly visible corpuscle, that no longer belongs to their referential background. *Quarks* themselves are quantum particles considered in this sense. Such *entities* or *physical «systems»* or *quantons*⁵, conceived in the specific way of quantum physics, are implicitly supposed to constitute in a way or another, according to the modalities of their description, objective elements of the real world, that manifest themselves to knowledge. Symmetries of quantum particles and fields allow us to *understand* in that way their properties and their arrangements.

Philosophical and epistemological difficulties arise only when one intends to understand *the nature of this understanding* : this is what I mean by «intelligibility to the second degree». This difference of degree with the founding debates comes from that the theoretical tool, «formalism», in the usual expression, is now already justified as a representation by its success. Physics builds its tools in an abstract manner and contents itself with these being well conceived, without trying to naturalize their origin ; admittedly they are abstract, symbolic, mathematical, and elaborated by thought from necessities laid by phenomena (physics has indeed established rules for that purpose, such as for example statements in the form of principles).

From this one sees clearly, undoubtedly better than in the past, when it was necessary also to construct these tools, that the problem of interpretation is twofold, but separated : physical and philosophical. The physical interpretation deals essentially, as it was traditionally the case since the birth of modern physics (in XVIIIth century), with the relationships between mathematically expressed magnitudes and the corresponding physical contents. The difference, from the physical point of view, between theoretical and conceptual elaborations aimed at quantum phenomena and systems, and those dealing with classical ones, is that the quantum are farther than the classical from the processes of observation by which these phenomena come to our senses.

Niels Bohr had rightly emphasized this difference of nature between the quantum and the classical. But he had formulated it in a manner that changed abruptly a simply physical state of things into a philosophical problematic about knowledge. There was, according to him, a barrier between the quantum and the classical, that was due to measurement, made necessary to know from perception. It resulted from there, in his view, that the knowledge of quantum phenomena can not grasp these directly as they stand in themselves, but has always to refer to classical representations.

The disjunction between the representation of quantum phenomena and that of classical ones can be pointed at in a philosophically more neutral manner : while classical physical phenomena and systems are *homogeneous* to the processes of their observation, quantum phenomena and systems are not, since observations and measurements relative to them belong ultimately to the domain of classical physics. But that does not entail any impossibility to represent

⁵ This term has been introduced by Mario Bunge (Bunge [1973]).

«directly», that is to say in terms of *properties* and *objects*, quantum phenomena and systems, at least if one refers *intelligibility* not to *perception* but to *understanding*, as it seems logical. If one can conceive objects of a quantum area or «world» in this sense, the questions of *physical interpretation* will be therefore largely independent from those bearing on more general considerations on knowledge, i.e. *philosophical interpretations*. One would then have shifted, so to speak, from a concern for *interpretation in general* to a more precise interest for the *physical meaning of quantum magnitudes* provided inside quantum theory itself.

The philosophical aspects of interpretation would then be set at a different level, granting a large autonomy to quantum physical thought, sensibly the same as for the other areas of physics : the former would no longer have to sacrifice to a so-called «foundational» need of being based on a peculiar philosophical interpretation, as in the first times of quantum mechanics⁶. And one would have then satisfied, up to some point, the realist demand of the theoretical physicist asking, with John Bell, for a theory that is not fundamentally about measurements, that considers physical systems in their inner completeness, and of which one could tell “not that such and such *may be observed* to be so, but that such and such *be so*”. In other words, a theory that “would not be on *observables*” [magnitudes able to be *observed*], “but about *beables*” [magnitudes able to *be*]⁷. This being eventually obtained with the argument presented in what follows, without modifying in any way the form of standard quantum theory, and only by understanding it differently (to an intermediate degree of understanding, involving some kind of *physical* interpretation).

It remains to know how to «interpret physically» the theoretical, conceptual or factual states of things that were problematic in a physico-philosophical mode for the «orthodox» or the «complementarity» «interpretation». We shall restrict ourselves, in what follows, to revisiting some characteristic and relatively simple quantum phenomena, renewed by recent results from high precision experiments, in the light of the proposed perspective on the physical interpretation of magnitudes and of theoretical formalism referred to the description of a world of properly quantum objects or systems. We shall see that they invite us directly to conceive quantum magnitudes in this manner, which entails the need to enlarge the meaning generally given to the concept of physical magnitude, and especially to that of state function representative of a physical system.

These phenomena served usually, since the beginnings of quantum mechanics, to illustrate the problems of interpretation. By a fair reversal of things, it is today possible to make arise directly from them the physical interpretation they are calling for. These phenomena are, first, *local non-separability*, whose epistemological status has undergone changes from a formal feature with optional physical meaning to an established physical fact, corroborated by experiments with distant correlated systems. Then, *diffraction of quantum particles*, performed not any more with many particle beams for statistical results, but with individual

⁶ See Paty [2000a].

⁷ Bell [1973, 198?]. See the complete quotation in the epigraph.

quantum systems for probabilities of individual events. Also, *indistinguishability of identical particles*, initially postulated or conceived as a formal property, and thereafter demonstrated by direct physical effects such as Bose-Einstein condensation, where a great number of identical atoms are accumulated in the same fundamental state up to a quasi-macroscopic level. Finally, recent *experiments of «decoherence»* have permitted to visualize superpositions of states in relation with mesoscopic systems in a situation of measurement by a classical device, in a tiny time interval before the dissipative loss of information occurred from interaction with the environment.

All these results converge towards a specification of the physical meaning of quantum concepts and magnitudes implied by the corresponding phenomena, obliging to associate factual evidences and physical contents conceived in terms of *properties of systems*, with «formal» properties whose interpretation remained until then optional or problematical. We will analyze some aspects of this new situation, trying to make out in which way they may contribute to deepen, to modify, or to base our theoretical comprehension of quantum features, by reducing the latitude of arbitrary choice in the interpretation and by adapting the norms of our intelligibility.

2

LOCAL NON-SEPARABILITY AS A FACT AND AS A PRINCIPLE

The objection opposed by Einstein in 1935 to the claim that quantum mechanics is a fundamental theory that will serve as the *basis* for any further progress in physics, known as «EPR argument», and afterwards reformulated and refined by its main author, raised several questions that overlap with interpretation⁸. The problem set was to know whether the theory (quantum mechanics) describes, or not, *real individual physical systems*, and if it describes them *completely*, that is to say adequately to all aspects rightfully attached to their *individuality* and in a one-to-one manner. The orthodox interpretation (in the philosophical sense) challenged the legitimacy of speaking of elements of reality independently of their conditions of observation, and Bohr's reply to Einstein's argument was exactly founded upon this position⁹. It had no chance to be listened to by Einstein, who could not accept its principle. Any progress in the debate on this question supposes from then on to try to leave aside the philosophical *diktat* of Bohr's reply, and keep considering only physical theory and the content of its concepts.

The question, as contemplated by Einstein, was to know whether the theory is complete in the indicated sense (it was to him the minimal sense for a

⁸ Einstein, Podolsky & Rosen [1935], Einstein [1948, 1949]. Cf. Paty [1988a, 1986 and in press, a].

⁹ Bohr [1935].

theory to be considered fundamental)¹⁰. If one can characterize an individual quantum system (be its theoretical representation probabilist or not), its state function has to represent it as such. If that is impossible because of some feature of the theory, then this theory can be only a statistical description (such was, indeed, the conclusion of the «EPR» argument). We will see further that this is also the point at stake with the *interference* of distinctive *individual* quantum systems when put actually in evidence. The EPR argument suppressed in principle the possibility to elude such a question in the name of an operationalist philosophical interpretation, and indeed the construction of rarefied beams later suppressed it effectively. Individualization for a system was indeed usually prevented by the alleged necessity to detect and measure it, if one wanted to know something of it (with a particle counter to know whether there came only one), and this act would destroy it immediately as a quantum system (it would project it on a classical particle state), forbidding all further knowledge of its quantum state (through the manifestation of wave properties).

In the EPR case, the system under study (U) was conceived in correlation to another one (V) while it did not maintain any dynamical interaction with it¹¹. The correlation, expressed by the conservation of a magnitude (A) used in the description of these systems, and known for the initial state formed by the two subsystems¹², allowed to determine the state of the first without perturbing it by a measurement, by deducing it from the state of the second, measured (supposedly) independently of it¹³. Measuring magnitude (A) of the second destroyed its state at the very moment of its determination, forbidding any meaning to the consideration of an alternative measurement for another magnitude (B) incompatible with the first : the initial system being no more available, no effective comparison can be made. But it would have nevertheless been logically possible, as a matter of principle, to perform the second measurement instead of the first, and it would have provided another state function for the second system ; from it, the first system would have been deduced, a priori different from the preceding result¹⁴. One could therefore have two different state functions to describe one and the same physical system : it would obviously be a theoretical

¹⁰ In particular in Einstein, Podolski & Rosen [1935], Einstein [1948, 1949]. See Paty [1986, 1988b, 1995a, and in press, a].

¹¹ The two systems U and V form at initial time one only system $U \oplus V$ and are allowed thereafter to move away from each other at arbitrary distances (for instance, two photons emitted in correlation by an atom).

¹² For instance, the overall momentum, $\vec{P} = \vec{P}_u + \vec{P}_v$, or the overall spin, $\vec{J} = \vec{J}_u + \vec{J}_v$ (its modulus J and one of its components, $J_1 = J_{u1} + J_{v1}$).

¹³ Measurement of magnitude A for the system V determines its state function, ψ_V^A , and the correlation gives the corresponding magnitude for the system U , and therefore the state function of the latter, ψ_U^A .

¹⁴ Let ψ_V^B be determined by the measurement of B , from which we deduce, without measuring, ψ_U^B . A priori, ψ_U^A and ψ_U^B are different, although they were not, *in principle*, perturbed, as they have not been directly measured.

weakness.

But this reasoning was dependent on a statement that did not belong to quantum formalism and that was at this time considered as optional : the separability of two far distant systems, that is to say their mutual independence in their respective locations. Einstein gave a precise definition of this *principle of separability*¹⁵, although recognizing that he added it to quantum theory. Without this principle, he believed, however, one could not characterize separately localized individual systems, unless one admitted a non-physical interaction (instantaneous action at a distance) between them. He concluded from this that quantum mechanics does not describe individual physical systems, but only statistical ensembles of systems, for which the objection does not hold¹⁶.

Further progress, both theoretical with John Bell's theorem (1964), and experimental, with experiments of correlation from a distance, has essentially consisted in analyzing local separability, a concept identified by Einstein, and in testing it for quantum systems. Bell's theorem on non locality demonstrated the existence of a contradiction between *local separability* and some predictions of *quantum mechanics* for systems of two correlated particles (*strong correlation* relationships for quantum systems expressed by *equalities* between averages for magnitudes were opposed to *weaker correlations* in the form of *inequalities* for the local separation case), and provided the sensitive relationships able to discriminate the local separability hypothesis and quantum theory¹⁷. From then on, experiments have decided in favor of quantum mechanics, in a hardly disputable manner, especially that of Freedman and Clauser, realized in 1972, and that of higher precision performed in 1981 by Alain Aspect¹⁸. *Local non separability* was henceforth established as a physical *fact*, a *general property* of quantum systems having been put in correlation, well identified from the phenomenal point of view¹⁹.

This property corresponds to a characteristic feature of state function in quantum mechanics : the state functions of subsystems that have been once correlated are not factorizable (i.e. independent of each other, i.e. separable). Having been linked together to form, even momentarily, one single system, two (sub-)quantum systems can not be dissociated : this «*entanglement*»²⁰ is a

¹⁵ In particular in Einstein [1948] and [1949]. See Paty [1995 and in press, a].

¹⁶ Ensembles of systems can admit a non-biunivocity of their state function, if the latter is only about mean values.

¹⁷ They are called «Bell's inequalities» : see Bell [1964, 1966, 1987]. They are relevant for the property of locality generally speaking, independently of they being or not related to determinist hidden variables, to which they had been linked in a first period. More general relationships have been obtained since then : Bell's theorem for locality without hidden-variables (Bell [1971], Eberhard [1977], Peres [1978], Stapp [1980]), and for more than two quantum correlated particles (Greenberger, Horne & Zeilinger [1989, 1990], Mermin [1990]).

¹⁸ Freedman & Clauser [1972], Aspect, Grangier & Roger [1981, 1982], Aspect, Dalibar & Roger [1982, Aspect [1983]. See the following reviews and analyses of the experimental results : Bell [1976a], Paty [1977, 1986], Clauser & Shimony [1978].

¹⁹ See Bohm & Hiley [1975], in Lopes & Paty [1977], p. 222 ; Paty [1988a], chap. 6, and Paty [1986].

²⁰ The use of this word, coined by Schrödinger in 1935 (Schrödinger [1935, 1984]), has been reactivated recently (Shimony [1993], d'Espagnat [1994], Cohen, Horne & Stachel [1997a & b], etc.).

fundamental property of quantum formalism, and possesses therefore a direct counterpart in phenomena, *local non separability*. This last can equally be considered as an aspect of *non locality* of quantum systems. The general and fundamental character of this property, and its inscription in the formulations that define in the theory the state of a system, incline us to see in it at the same time a *fact of experience* and a *principle* for quantum physics.

An important conceptual aspect of local non-separability is its place in the economy of quantum theory. One can analyze it, from the point of view of the conceptual and theoretical consistency adopted here (which may also be called of «critical realism»), in the following manner. As it is directly linked to the definition of quantum systems and of the magnitudes by which we represent their states, local non separability relies only on these, and does not have to refer to other magnitudes that would be defined outside of this theory. It bears on systems that, spatially speaking, are «extended systems», and for which the space variables have not, as an effect, a direct part in their definition ; in this sense, it is not concerned by special relativity, that it does not contradict, and has nothing to do with (instantaneous) actions at a distance²¹.

It is fair to say, however, that many physicists and philosophers of science would still disagree with this conclusion, that seems compelling from the point of view adopted here. It is, indeed, difficult to think physically without the help of spatial intuition, and this is probably the main reason of their dissatisfaction with «pure quantum reasoning». But who can say what kind of intuition is adequate for the quantum domain ? It seems to me that quantum physicists have developed over the years an adequate intuition in this respect, that is basically founded on quantum formalism as a practiced intellectual tool to explore and understand quantum phenomena (the epistemological implications of which we are exploring here, taking a point of view of general consistency). John Bell, who was reluctant to accept the above argument, which he viewed as too formal and even as a «verbal» solution²², admitted nevertheless non separability as a fundamental fact and eventually as a physical principle²³. But he would have preferred to have a dynamical interpretation of it. It seems to me, on the contrary, that as a principle it definitely does not need an explanation, but stands as a primary conceptual reference toward which the other quantum concepts must be consistently obliged (in a way similar to the principle of special relativity ruling the transformation laws of the concepts related with the motion of bodies).

As a fundamental quantum fact, one should perhaps consider that local non separability is to quantum physics as, for example, the principle of equivalence (of inertial and gravitational masses) is to the general relativity theory of gravitation. One can see it as a true *principle*, both a synthetic proposition based on experimental facts and a theoretical statement of a central, and perhaps

²¹ This aspect has been emphasized in the article quoted (Paty [1986], and already in 1980 (Paty [1981, 1982])). Bernard d'Espagnat seems to come also to the same conclusion in one of his recent books (d'Espagnat [1994], p. 430).

²² I remember my discussions with him : we disagreed on this point. This question represented to him an intellectual challenge whose difficulty remained untouched.

²³ Bell [1987]. I have quoted elsewhere (Paty [1988a], p. 245), a letter in this sense of John Bell to Alain Aspect.

foundational, importance ; it could serve to formulate quantum theory in a less formal manner than the usual presentation, which would make it come closer to the other physical theories, from which it parted until now in this respect.

Local non separability can be seen as an even wider theme of reflection, rejoining a cosmological perspective. One can, indeed, make rapprochments with other features of «disindividualization»²⁴ or of «desingularization» or, better, of *indifferentiation* such as *indistinguishability* (that partakes as well of the superposition principle), and perhaps as symmetries of matter, that are important features of primordial cosmology²⁵. One can also see in it, with David Bohm, the mark of a more general indivisible wholeness of material reality²⁶. On this one must nevertheless observe that to grasp an underlying order thought separations in such a wholeness are needed as a necessary approximation, without which the concept of wholeness would lose all utilizable physical content. Extended in an absolute manner to the whole Universe, the principle of non-separability would present the same kind of disadvantage as the one pointed out by Poincaré²⁷ regarding the principle of relativity of space if we were to formulate it with respect to all bodies of the Universe : being tautological it would not give us a hold on phenomena. But, at any rate, it might give us some hint on cosmological conditions, of the kind Einstein got for a closed and unlimited Universe²⁸ (for example, in quantum cosmology, some coherence condition for having finite time inside Planck's limit in the primordial Universe).

3

INDIVIDUAL SYSTEMS AND TRANSFORMATION OF PHYSICAL PROBABILITY

The phenomenon that is the simplest in its principle to characterize quantum properties is that of *interference*, which confirmed the wave-particle duality of matter and inspired Max Born's idea of the probabilist interpretation of the state function. This archetype phenomenon illustrates some fundamental aspects of the description of quantum systems and helps making explicit from the physical point of view the interpretation problems that had been raised.

The «orthodox» interpretation of complementarity and observationalism sees in it the necessity of wave-particle duality and the impossibility to go beyond it. The interference pattern (concentric rings, alternately obscure and bright), similar to those of classical waves, is due to the wave property of quantum systems ; whilst, on the other hand, the materialization

²⁴ This word is inadequate if by individuality one means a unity. Undifferentiated quantum systems can be counted : they keep *cardinality*.

²⁵ Paty [1999b].

²⁶ Bohm [1980].

²⁷ Poincaré [1912]. See Paty [1996].

²⁸ See Paty [1993], chap. 5 and 7.

of these varied intensity rings on the screen covered with a sensitized film comes from the corpuscular property of these systems (through their interaction with the grains of photographic emulsion, producing an image). The dual properties, contradictory if they are considered for individual «particles» or systems, can be reconciled as soon as one gives up a concern for causality of individual events, and minds the statistical aspect of the experiment. If one wanted to examine, in this experiment, the behavior of an individual quantum system, any meaning of it would be denied, according to the complementarity interpretation, in the name of the very definition of the systems. As a matter of fact, if one wanted to characterize a quantum system as individual, it would be necessary to submit it to a counting experiment, that would inform about which of the slits the quantum system has gone through ; by being localized in that way, the system would suffer a perturbation, and therefore lose its quantum aspect and its capacity to produce interferences.

Yet, in 1930 already, Paul Dirac, in his book *The principles of quantum mechanics*²⁹, indicated that, according to this theory, one photon interferes with itself and that this is the reason of the interference phenomenon. This is also the case for any quantum systems (particle, atom, etc.). The meaning of it would be that interference is a property of any individual quantum system, and that quantum physics is the theoretical description of such individual systems. The probabilistic turn of this description would not a priori be a hindrance for this purpose (after all, statistical mechanics does the same). However, the «complementarity explanation», to which we just referred, blunts and dissolves the force of this statement, by making of it a mere feature of the formalism, from its claiming of the impossibility by principle to observe it in experiments.

As for the ensemble interpretation of quantum mechanics, according to which the theory is only a statistical one (incomplete for Einstein, complete for others), it only knows averages, that have no physical meaning except for an ensemble of systems, and can not pronounce on the significance of individual quantum events.

However, since approximately two decades, experiments have been realized, and continuously improved thanks to technical advances, with *individual quantum systems* (photons, electrons, neutrons, atoms) that are known to be such without needing to be counted by detection on their path, and therefore without destruction of their quantum state. It has actually been possible to produce beams of such «particles» or quantum systems, extremely rarefied and with a high time definition (better than 0.1 ns), in such a manner that particles get to the interferometer one by one, spread in time, each having got across the experimental arrangement within an interval of time sufficiently small to ensure that the following one has not yet entered³⁰. One is then pretty confident that only one particle at a time has crossed the interference apparatus (and interfered with itself). The detection of impacts on the screen seems in the beginning to be at

²⁹ Dirac [1930].

³⁰ See, in particular, Pflugor & Mandel [1967], Grangier [1986]. The concepts of quantum theory of field, that permit the definition of states with a given number of particles, underly these experiments. It is necessary, for exemple, to prepare one-photon states of the electromagnetic field (Grangier [1986]).

random, but as time goes and many single «particles» having gone through the interferometer accumulate, the distribution of impacts is seen to obey a law : one obtains, in the end, the same interference pattern as in the traditional experiment with a beam of N identical particles crossing simultaneously the interferometer.

These results require that a physical meaning be attributed to an individual event in an interference experiment. Clearly, the final interference pattern with individual particles can only be obtained statistically, by the realization of a great number of successive one-particle or individual quantum system experiments. The result of N such experiments with single quantum systems gives the same result as a single experiment performed, in the same interferometer, with a beam of N identical systems. But the theoretical inference that one is allowed to make in the two cases is very different. The second case, of the traditional experiment with a great number of simultaneous systems, satisfies a frequentist and purely statistical interpretation of the probability given by the state function³¹.

But the effective occurrence of the first case, N experiments with an individual system identical each time, and represented by the same state function, assures us that each individual phenomenon, independently from the others, contributes to the final interference pattern. One is therefore led to conclude that *it is the individual quantum systems that make the phenomenon* and therefore that, in a way that remains to be specified, each individual phenomenon occurring with each (independent) system constitutes potentially the overall interference phenomenon revealed by the final pattern, obtained statistically. In other words, each phenomenon relative to an individual system is a *quantum phenomenon*, collected on the screen through a *classical measurement* process (the «photon» or quantum particle impact on a grain of silver bromide of the photographic emulsion). One is then inclined to consider that, just before interception on the screen, each of the individual systems having interfered with themselves is in a quantum superposition state. And that, as nothing distinguishes them from each other, all these individual systems in interference are strictly identical. From then on, the only remaining problem would be that of the measurement process : identical quantum systems provide, after detection, different results, but endowed with probabilities corresponding to the amplitude of probability of their state of superposition³².

As a result of what precedes, the ψ state function must be considered as the theoretical representation of an individual particle, which entails the

³¹ $P = |\psi|^2$.

³² Consider an initial individual system crossing a diaphragm with two slit a and b , and whose state is represented by $\varphi(x) = \frac{1}{\sqrt{2}} [\psi_a(x) + \psi_b(x)]$. Let z be the variable corresponding to various localisations on the screen, placed at a distance x from the diaphragm. The state $\varphi(x)$ of the individual interfering system can be considered as a linear superposition of states prepared along the values z_i of the variable (or magnitude) z : $\varphi(x) = \sum_{z_i} \alpha_i g(z_i)$. The probability of an

impact on the screen in z_i is $|\alpha_i|^2$.

following important consequence of its physical meaning : the *physical probability*, given by the ψ state function³³ (the latter being often named «probability amplitude», in a sense that can only be physical, since nothing of the kind exists in mathematical probabilities), is not liable to be reduced to statistics for ensembles of systems. It has a *theoretical function* from the physical point of view, as it is deduced from a magnitude having a direct physical meaning, the probability amplitude (i.e. the state function itself). One can therefore consider this probability as a *physical magnitude*, which makes it differ from probability in a merely mathematical sense, as well as from probability conceived physically as expressing a frequency³⁴.

³³ By the square of its modulus.

³⁴ Cf. Paty [1990].

4

INDISTINGUISHABILITY AND STATE FUNCTION

In quantum physics, the state function that represents a quantum system allows the complete description of all the properties attributed to this system, in such a way that systems represented by the same state function are effectively in the same state and are absolutely indistinguishable. That means that there does not exist any other possibility, external to the theory, to distinguish them. In other words, *a quantum «particle» has no other characteristics than those of its state*, differently from physical systems as described by other theories such as classical mechanics, thermodynamics or relativity theory. These theories describe *what happens* to physical objects that are in other respects defined outside of them. For example, the three body problem of classical astronomy is about the mechanical processes occurring to celestial objects that are supposed given. The theory bears not on these objects, but on their interaction properties. The Moon, the Sun and the Earth, for example, possess an identity - and an opacity - defined previously to the laws and equations under study of mechanics and astronomy.

The only theory, except quantum physics, for which the eventuality has been considered that it could be able by itself to describe its object, instead of getting it from outside, is the general theory of relativity, at all events in a further more elaborated formulation foreseen as a distant purpose (by A. Einstein and J. A. Wheeler notably), where it would be possible to describe in the same system of equations both a field and its source. Such was the «strong» meaning Einstein attached to the notion of theoretical *completeness*³⁵ and, to him, quantum mechanics was not a «complete theory», in this sense obviously, as its status of a framework theory rather than a dynamics suffices to show. But there was another, weaker meaning of the same notion, which he considered as crucial for the fundamental nature of quantum physics, as we have seen earlier. A theory would be «complete» in a minimal sense if it were able to describe fully its *object*, that is all the properties that can be physically considered about it. It was not the case, for Einstein, with quantum mechanics, because of EPR type correlations that, invalidating the principle of separability, excluded the description of individual systems³⁶.

We do not any more consider this argument in this form, such correlations having proved to be factual and to concern individual correlated systems. On the contrary, actually, completeness at least in the weaker sense would characterize, in principle, the description of *quantum systems* with the physical interpretation envisaged here. The main obstacle to this requirement seems today to remain the «quantum measurement problem». If one sets aside the latter for a moment, one can rightfully be struck by the purpose of quantum mechanics to get an *exact covering* of the *described system* by its *state function*, going even therefore, in a way, beyond the restricted completeness requirement.

³⁵ See Paty [1988b], [1993], chap 10, and [in press].

³⁶ Cf. Paty [1995a ; in press]. See above.

The most remarkable expression of this covering appears, finally, to be the property of *indistinguishability of identical quantum systems*. But is it a mere feature of the formalism, or a property of physical systems ? Both aspects, as always with quantum mechanics interpretation problems, seemed closely embedded and not easily disentangled. This property was identified on the eve of the constitution of wave and quantum mechanics, by Satyendra N. Bose and Albert Einstein for quantum systems of null or integer spin (photons and atoms named afterwards «bosons», obeying «Bose-Einstein statistics»), and by Enrico Fermi and Paul Dirac for quantum systems of half-integer spin (electrons, protons, and other «fermions », obeying «Fermi-Dirac statistics»). Indistinguishability of identical bosons (in the case, photons) appeared to be the real underlying reason of the quantification procedure for radiation energy exchanges in black body as performed by Planck in 1900³⁷ ; and indistinguishability of fermions (here, electrons) gave the explanation of the Pauli exclusion principle, that accounted for the constitution of atom levels in terms of state occupations by electrons.

This property, corresponding to two types of statistical (or probabilist) processing of quantum systems (the admission of several particles in a same state in the first case or, on the contrary, their mutual exclusion in the second one), opposed to the classical statistical processing *à la* Boltzmann of particles always distinguishable even when occupying a same state (for they possess a proper identity). Indistinguishability therefore limits drastically the possible state occupations. It indicates, actually, that quantum systems *do not occupy* states, but that *they are themselves* states, and are identified with their states³⁸. Indistinguishable quantum systems have no other element of identity than those participating the theoretical description of their state. The notion of state is identified with that of «particle» : a quantum «particle» (or system) *is its state* : it is not «in its state», as a classical system. This situation corresponds to a closer determination of the physical system by the theory, contrary to the idea that prevailed for quantum physics of a looser determination and a limitation of knowledge because of «indeterminacy» relations.

This formal property, indirectly dictated by factual reasons, and that finds also its expression with the principle of superposition³⁹, has proven to

³⁷ Already in 1911-1912, Ladislas Natanson and Paul Ehrenfest had diagnosed the non-classical character of the statistics corresponding to Planck's radiation law. See, for instance, Kastler [1981], Darrigol [1988, 1991], Petic [1991].

³⁸ This includes the invariant characteristics shared by the various possible states of a system, that contribute to define the system and its particular states corresponding to given magnitudes.

³⁹ Consider, in effect, a system of two identical quantum particles 1 and 2, each in its state, represented by the state functions ψ_1 and ψ_2 . The state function of their coupled system is symmetrical for the permutation of the particles in the case of Bose-Einstein statistics, hence : = $\psi_{12} = \frac{1}{\sqrt{2}}(\psi_1 \otimes \psi_2 + \psi_2 \otimes \psi_1) = \psi_{21}$. Nothing forbids identical (indistinguishable) particles 1 and 2 from being in the same state inside the system (identical bosons can accumulate in the same state inside a system). For the case of Fermi-Dirac statistics, the coupled state function is antisymmetric : $\psi_{12} = \frac{1}{\sqrt{2}}(\psi_1 \otimes \psi_2 - \psi_2 \otimes \psi_1) = -\psi_{21}$. If the identical fermions 1 and 2 were totally indistinguishable, occupying the same state in the system, then one would have :

correspond to fundamental physical properties of quantum systems that could be directly tested and that have implications to the macroscopic level itself.

Supraconductivity and *superfluidity* are such properties directly connected to indistinguishability. Bose-Einstein condensation, predicted since 1925 by Einstein from the indistinguishability of the identical for some kinds of atoms (it was, actually, the first theoretical description of a phase transition), was for a long time considered as being very far from possibilities of verification. Yet it has recently been put experimentally in evidence, thanks to the high technical realization of extreme colds and atoms trapping by laser rays⁴⁰. Tens of thousands of atoms are thus condensed in the lowest energy state (called «of the zero point»), nothing distinguishing them from each other : the superatom they then form corresponds to a fluid in absolute superfluidity state, without any viscosity, that can show itself at the macroscopic level (by an effect of *visible non locality*, the fluid occupying quasi-instantaneously all the space offered to it, rising on the container' walls). At this stage, restrictions claimed by the orthodox complementarity interpretation about the directly physical character of the state function appear rather ridiculous, and as an exercise of twisted rhetoric serving only to hide evidence.

One may invoke, for the exclusion principle equally - and therefore, for indistinguishability of identical fermions - direct consequences at a highly macroscopic level, concerning cosmic objects corresponding to definite phases of the evolution of stars. «White dwarfs» are compact stars in a state of equilibrium between the gravitational tendency to collapse and the pressure of degeneracy of electrons that cannot fall in the same fundamental state because of Pauli exclusion principle⁴¹. «Neutron stars» resist in the same way the collapse on themselves due to gravitation by the degeneracy pressure of the neutrons into which all atomic nuclear constituents have been transformed.

By its directly physical consequences, indistinguishability of identical quantum systems is indeed a physical property of these systems, and not only a feature of the theoretical formalism. It is described with exactness by quantum theory in terms of *state function* (submitted to the principle of superposition), and there is therefore, as we earlier suggested, a liaison of the property pointed at by *indistinguishability* (equivalence of particles of similar characteristics, occupying the same state within a system, that one can count but that nothing distinguishes)⁴² and the *theoretical description* by the state function of quantum mechanics (or, at a further stage, of quantum field theory). All that confirms us in the inclination to see indistinguishability not as a «lack», as would suggest the common intuition of the notion of «particle», taken from the immediate

$\psi_{12} = -\psi_{21} = 0$: two identical fermions cannot occupy the same state inside a quantum system (exclusion principle).

⁴⁰ Cornell & Wiemann [1998]. Cf. Griffin *et al.* [1995].

⁴¹ The mechanism was proposed by R. H. Fowler as soon as he knew the statistics studied by Paul Dirac, who was his student (cf. Doncel, Hermann, Michel & Pais [1987], p. 274).

⁴² From the point of view of arithmetics, concerning how to count or to identify by a number, such objects are characterized by cardinality, but not by ordinality. It has been proposed from a logic point of view to describe them with a set theory whose elements would possess this property, different from that of Zermelo-Frenkel (cf., for instance., French & Krause [1996]).

experience of bodies in our environment as well as from the habit of classical physics, but rather as a characteristic and determining physical *property*. For nothing authorizes us to think, about such objects, of properties that are not pointed out by the theory.

5

REAL PHYSICAL STATE AND SUPERPOSITION, MEASURED STATE AND PROJECTION.

The state of a quantum system, as we have tried to characterize it physically, is not identified with that obtained directly by one measurement alone. This last, indeed, is a reduction or at least a projection of the state physically defined on one of its components, according to the choice of the preparation of the system (by a complete set of compatible magnitudes)⁴³. A measurement device in the usual sense can only measure a classical magnitude ; with respect to the state of superposition that represents a system before the operation of measurement, it can only provide one of the components (one of the «eigenstates» of the measured set of magnitude). One should not be surprised by this as such is its function and its only ability.

The measurement apparatus is, as a matter of fact, by definition, a projection device (in the geometrical sense) on the various components of the state of the system. One has claimed that quantum measurement is a non causal interaction, but this is to pronounce a priori on the nature of the interaction between the quantum system and the macroscopic device. If one speaks rightly of a *rule of projection*, or eventually of *reduction*, this rule does not, up to now, mean any directly physical process and nothing allows it to be raised to the status of a physical principle. In the absence of a theory, in the proper sense, of quantum measurement, that would be a general theory of the interaction between quantum system and macroscopic measurement apparatus, one must hold it merely as a practical rule.

Each measurement provides a numerical value for the measured magnitude, one of its possible (classical) values (among the eigenvalues) with some frequency, given by the corresponding probability amplitude (eigenfunction). An experiment with a great number of identical systems, or a great number of independent experiments on such systems taken individually, provide all the spectrum of values of the magnitude with probabilities for each one (corresponding to the amplitudes in the superposition). From these results in terms of classical magnitudes, one infers the quantum superposition state that has been submitted to measurement, and of which one can reasonably think that it represents *the quantum system before measurement*, in one of the possible bases, the one chosen by preparation. The state function reconstituted in that way is not a simple catalogue of data, since the system that it represents has the capacity, capacity, a clearly physical one, to propagate, to evolve in the course of time, to

⁴³ Or, according to the usual terminology, «a complete set of observables that commute».

make interferences or to possibly get oscillating between different physical states (on which we shall give more details below). Measurement to determine the state will intervene only after these transformations, that owe nothing to man's hand or thought but everything to nature.

In summary, we propose to consider that the physical quantum states are the states expressed as superpositions themselves, that one can determine from the determination of their components, which reduces to magnitudes endowed with numerical values by classical measurement devices. Actually, this is nothing more than to take von Neumann and Dirac geometrical vector representations as meaning it : state vectors in Hilbert space are the physical one, represented on their various possible bases (determined from the preparations according to their possible sets of commuting magnitudes). As a vector, the system state is a basis-free geometrical representation of a physical state, and is more fundamental, because of its invariance, than its «contextual» components.

As an effect, physicists, familiarized by their practice with the thought of quantum systems, consider them in this manner : what is important to them is the representation of these systems' quantum states, i.e. the overall final reconstitution, and not the contingent and particular (classical) values obtained by measurement. These values are intermediate entities given by experiment, whose deep physical meaning is attained at only from their immediate translation in quantum terms, needed to come back to the description of the physical quantum system under study.

6

PHYSICAL PHENOMENA LINKED TO PROPAGATION OF SUPERPOSITION STATES

A physical state, as physicists consider it in their representation of quantum phenomena, and as they *think* it in their *theoretical work*, is given in an invariant form with respect to its «vector projections», while being generally presented at the same time as a state of superposition on one basis or the other. This is more general than keeping restricted to the consideration of measurement alone, which after all is nothing else that one of the moments of verification or of experimental test, and it is not a purely formal property : *this form rules the physical properties* of quantum systems. We have seen it for the phenomena evoked above, but one can also evoke a number of other ones, of a different nature, that show to what extent this is indeed the universal form of the description of all quantum systems. Two examples, both borrowed from elementary particle physics, will show it in a clear and striking manner, all the more as they have no classical analogous : these are the «*mixtures*» of particles states and the «*oscillations*» from one state to another, these mixings and oscillations being expressed directly in terms of state superpositions that

propagate⁴⁴.

The neutral «strange pseudoscalar» meson K^0 and its antiparticle, $\overline{K^0}$, are eigenstates of their «mass matrix» (M) and of the *strong interaction* hamiltonian (H_s), process through which they are produced (they are physical states in *associated production* conserving the «strangeness» magnitude, $S = +1$ for K^0 , $S = -1$ for $\overline{K^0}$ ⁴⁵ or for any other associated strange particle in the production interaction, for instance the «strange baryon» Λ^0). They have a different behaviour in their *decay through weak interaction*, with the strangeness non conserving hamiltonian H_F ⁴⁶. The eigenstates for such processes are the mesons as observed from their decays, characterized by proper lifetimes (τ) and decay modes, the short-lived K_S^0 ($\tau = 10^{-10}$ s) and the long-lived K_L^0 ($\tau = 10^{-8}$ s). The initial states K^0 (resp. $\overline{K^0}$) are expressed as linear superpositions of K_S^0 and K_L^0 states, which progressively transform according to the law of exponential decrease in time. If one considers a K^0 meson initially produced (actually, a beam of such mesons, appropriately selected), and worries about its state at a time t , the superposition containing initially the states K_S^0 and K_L^0 in equal parts impoverishes in K_S^0 , whose time decay is faster, and enriches in K_L^0 , that will in the end completely dominate. The then nearly pure beam of K_L^0 states can be written as a superposition of the states K^0 and $\overline{K^0}$ in equal proportions. One gets therefore, in the beam of K^0 mesons, a «regeneration» of $\overline{K^0}$ mesons that were absent in the initial beam. These can be detected through a strong interaction process with respect to which they are well defined, i.e. of which they are eigenstates.

Let us note, incidentally, that the qualification of eigenstate concerns definite states of a hamiltonian and other physical magnitudes that are not, here, of a classical nature. At this level, the identification of quantum systems in given states does not call for measurement in the classical sense. The latter is needed only at the end of the chain of experimental processes of the detection of «particles» typical of the considered interactions. In a general fashion, an eigenstate given for a set of compatible magnitudes can be projected (in the vector sense) on another (preparation) basis relative to another set of magnitudes incompatible (non commuting operators) with the first. This eigenstate of the first set of magnitudes will therefore be written as a superposition of eigenstates of the second set. In others words, the «preparation» of a quantum system concerns proper quantum magnitudes as well as magnitudes submitted to a classical determination by measurement. «Preparation» for measurement is only a particular case of «preparation» in general, that means the choice of a set of

⁴⁴ Strictly speaking, the representation of these «particles» makes use of the quantum theory of fields. However, the features of their properties that we discuss here are only those of the basic formalism of quantum mechanics (the definition of a state from physical magnitudes and the principle of superposition for the state functions).

⁴⁵ The magnitudes (the «observables», in the quantum jargon) H_s , M and S commute between themselves ($[H_s, S] = 0$, etc.) and have the same eigenstates.

⁴⁶ H_F and S do not commute ($[H_s, S] \neq 0$).

physical magnitudes corresponding to a set of eigenstates taken as referential (or as vector basis in the Hilbert space of their eigenfunctions).

One can also consider the behavior of these neutral K particles under the transformation by the CP operator⁴⁷, product of charge conjugation (C , that changes a particle into its antiparticle) and parity (P , or space symmetry) or, equivalently, by time reversal (T) operator, the equivalence ($CP = T$) being due to the conservation of the product CPT , following a theorem of the quantum theory of fields⁴⁸. If one represents the eigenstates of the CP magnitude by K_1^0 (with a corresponding eigenvalue $CP = +1$), and K_2^0 ($CP = -1$), and if the operator CP does not commute with the weak interaction hamiltonian (H_F)⁴⁹, the K_L^0 and K_S^0 states are different from the K_1^0 and K_2^0 states and they can be considered as linear superpositions of these states. The coefficients in the superpositions are functions of the parameters of CP violation in these weak interaction processes.

Such physical systems propagate with time between the moment of their production and that of their detection, and the state that is attributed to them during this course is that given by the state vector (invariant with respect to the basis), that is, for the chosen basis, the linear superposition, whose coefficients vary with time (let the function $\psi_K(t)$ be the representation of this state). That is to say that the *superposition* here is the *physical state*, without any circumlocution that would bring physical existence only to the state detected after observation or measurement. The quantum system under study (represented by the $\psi_K(t)$ state function) is analyzed by a detector placed on its line of flight, that projects it (in the geometrical sense of vector projection) at time t onto one of its components chosen by fixing the detection conditions («preparation»). From the frequencies for each detected state, that are a measure of their probabilities, one obtains the coefficients of the superposition or probability amplitudes (probabilities are the absolute squares of the coefficients), as in the usual case. One observes statistically, for K_L^0 , a given number of states in the $CP = +1$ mode (for example, $K_L^0 \rightarrow 2\pi$) and another one in the $CP = -1$ mode ($K_L^0 \rightarrow 3\pi$).

What is interesting for physicists, from a physical point of view, is not so much the final state observed at the detection, which choice is, as a matter of fact, purely contingent, as the indication it provides about the physical state of the K^0 meson *at a time t before its detection*, given by the basis-free or invariant state vector. This state vector is given, for each group of (compatible) magnitudes corresponding to a physical content (either M and H_S , or H_F , or CP), as a superposition of their eigenstates. Conversely, the knowledge of this state permits the characterization of the properties of these magnitudes (for instance, the degree of CP violation in the weak interaction process with a hamiltonian H_F)⁵⁰.

⁴⁷ Let us recall that in quantum theory the mathematical form of *physical magnitude* is a linear *operator* acting on the state function.

⁴⁸ Due to Gehrart Lüders, Wolfgang Pauli and Julian Schwinger, who established it around 1952-1955 (see Lüders [1952] and especially [1954], Pauli [1955], Schwinger [1951-1953]). See comments in Enz [1973], Doncel, Hermann, Michel & Pais [1987], Yang [1982].

⁴⁹ In fact, weak interaction does not conserve CP in these processes.

⁵⁰ The whole thought of «elementary particles» physics is, as quantum physics in general, ruled by the superposition principle. We could have taken other examples of state mixtures as

The so-called «oscillation» phenomena between quantum particles states are described and thought in a similar way⁵¹. Consider *neutrinos*, electrically neutral (fermion) «leptonic» particles existing under the form of three different species, ν_e , ν_μ , ν_τ , each one endowed with a distinct conservative magnitude, the leptonic, *electronic*, *muonic*, *tauc*, charges or quantum numbers, shared with the electrically charged corresponding particles, electron, muon, tauon⁵² (respectively e^- , μ^- , τ^-), together with which they constitute the three families of *leptons* (the most elementary «particles» of matter with the *quarks*). Their mass is very small, possibly null.

If the mass of neutrinos is not exactly zero, one can distinguish three states of mass, ν_1 , ν_2 and ν_3 , distinct from the states that represent the («leptonic») neutrinos observed through their «weak interactions» (ν_e , ν_μ and ν_τ and the corresponding antineutrinos). The latter can be described as linear superpositions of the mass states⁵³. Neutrinos emitted in nuclear reactions (in β decays of nuclei) are of the type ν_e (or $\bar{\nu}_e$). The evolution with time, during their course, of their state function, ψ_{ν_e} , is given by that of the amplitudes (or coefficients) associated with the states of the superposition. As a consequence, the proportion of the three mass states varies during the propagation ; as these mass states can themselves be put in the form of superpositions of the leptonic states, it entails that the initial neutrino (ν_e) transforms partly in neutrinos of the other species (ν_μ and ν_τ), with a given «oscillation length» (as one says wave

superpositions : the neutral states of «vector mesons» (ω^0 , ϕ^0 , with spin-parity $J^P = 1^-$) under the conservation of a given magnitude (for example under SU_2 «isospin» symmetry or SU_3 «unitary spin» symmetry), or the state superpositions of the neutral «intermediate bosons» (γ and B) of the gauge symmetry electroweak theory of A. Salam, S. Weinberg and S. Glashow (cf., p. ex. Paty [1970, 1985]). These bosons, and also the charged «intermediate bosons» (W^\pm), are initially supposed to have a vanishing mass as the photon, and their mixture, or superposition, is characterized by a coefficient (θ_{W-S}) called «Salam-Weinberg mixing angle», that is the parameter of the theory. The symmetry breaking generates the finite masses of the «physical» «intermediate bosons» (W^\pm , W^0), related to the mixing parameter (see, f. ex., H. Pietschmann & D. Haidt, in Gaillard & Nikolic [1977] ; Paty [1985]). All this however is happening inside the limits of the range of weak interaction, that is extremely small. The examples that we have presented in the text are more striking for our purpose, insofar as they correspond to phenomena that are manifested on large spatial distances, covered during the propagation, and for which one hardly could refrain to speak of *physical states*, beyond the mere mathematical formalism of the theory.

⁵¹ One example, hypothetical but theoretically founded, would be eventual oscillations of neutrons into antineutrons ($n \rightarrow \bar{n}$), through an interaction field violating baryonic number (such as required by the «Grand Unification» theories).

⁵² Or «heavy lepton» (with mass 1777 MeV, the muon mass being 106 MeV, and the electron mass 0,5 MeV ; the mass unit is MeV, million of electron-volts, in the appropriated unit system commonly used in subatomic physics).

⁵³ See, f. ex., Paty [1995b], Alexei Smirnov in Nguyen Khac, Ung & Lutz, Anne-Marie (eds.) [1994]. Leptonic numbers are no more completely conserved, and the heavier neutrinos can decay into a lighter neutrino together with other particles (a different process than «oscillations» considered here).

length)⁵⁴. Such effects (such *phenomena*) are actively searched for by physicists for the three types of neutrinos⁵⁵.

It is generally considered that physical neutrinos are those characterized by their properties in the (weak) interaction⁵⁶, through which their are produced or destroyed (interactions with other particles or eventual decays), that is to say that they are the «leptonic» neutrinos ν_e , ν_μ and ν_τ . Nevertheless, in the propagation of one or the other of these neutrinos, the effective *physical state* would be, at any instant of time, in the considered hypothesis (of non zero masses, and of some degree of leptonic numbers violation), due to the mentioned transformations, a *linear superposition of these states*, evolving in time in a determined way. The detection by (weak) interaction of one of the states allows, by comparison with the initial state (given by the choice of one of the three types of neutrinos), one to know the physical state at a chosen place on the covered distance (i.e. at a given time of flight). This detection is based on reactions of interaction where a neutrino transforms into the corresponding charged lepton ($\nu_e + n \rightarrow e^- + p$ and, similarly, $\nu_\mu + \dots \rightarrow \mu^- + \dots$, $\nu_\tau + \dots \rightarrow \tau^- + \dots$). These reactions require for production enough energy to create the mass of the charged leptons.

In the case of neutrinos originated from nuclear reactions, the energies are insufficient to create masses larger than that of the electron. The neutrinos ν_e transformed during their travel into ν_μ or in ν_τ will not give therefore reactions that would detect them and remain sterile. If one finds less ν_e than there were at the beginning, it might well be that the pure initial state has been transformed into a superposition of different neutrinos, of which only the projection on the ν_e state is detected. This is, for example, what is supposed to happen with solar neutrinos, whose proportion received on Earth is far less than what is expected if neutrinos continued on their way remaining identical to themselves⁵⁷. We would have there again (actually, the oscillation phenomenon has recently been definitely proven experimentally), an indubitable direct effect of the *physical character of a linear superposition state*.

The example (be it a real phenomenon or a simple possibility) gives indeed also evidence that the thought of such *states of superposition* is hereafter familiar to physicists. A superposition of states has to be understood as a simple change of basis relative to another set of mutually compatible physical magnitudes, corresponding to one of the possible «preparations». The physical state that physicists consider is not restrained *to that after the measurement* (otherwise it would only be the incident deficient neutrino) ; it is the state *that is*

⁵⁴ «Oscillations» are a function of neutrino mass differences, energies and covered distances.

⁵⁵ These experiments concern, besides nuclear reactor or solar neutrinos (essentially $\bar{\nu}_e$), atmospheric neutrinos (and antineutrinos) (mainly ν_μ) and those produced at particle accelerators (ν_μ and ν_τ).

⁵⁶ Neutrinos interact only through «weak interactions».

⁵⁷ ν_e neutrinos are detected by their capture by a nucleus with emission of an electron (or of a positon in the case of $\bar{\nu}_e$ antineutrinos). Neutrinos of other kinds resulting from oscillation are sterile for this type of reaction, and escape detection. But they are indeed part of the incident flux.

revealed to them by this measurement, and that contained besides another component that they do not detect but that can immediately be reconstituted. Recent observations (in 2002) on neutral currents induced by solar neutrinos, which are not dependent on mass threshold effects (as the neutrino is simply scattered by the nucleon target), have yielded the expected rate, confirming that the neutrino beam arriving on Earth is in a supersposition state of all the neutrino leptonic states. Of course, all these phenomena are studied with great numbers of «particles», but their description and their explanation must be understood in terms of properties of individual «particles», for the same reasons as those considered previously.

7

BEFORE DECOHERENCE,
SUPERPOSITION

It remains us to evoke another type of phenomenon of recent production and observation, «*decoherence*». We will not undertake here a thorough discussion of its implications and its interpretation. In particular, we will not pronounce (reserving it to another opportunity), whether this phenomenon gives or not a solution to the problem of measurement of quantum systems, and even neither whether it brings new views on the relationships between the «classical» and the «quantum». At least does it illustrate, to my eyes, by «visualizing» it, an important aspect : it makes us *see* a state of superposition propagating and, by this, it allows us to better *conceive* the possibility and the *physical reality* of such states⁵⁸.

The metastable state of superposition that has been observed recently for «mesoscopic systems»⁵⁹ is an «*entangled*» state made by coupling a Rydberg atom in a two-energy states superposition with an electromagnetic field (of few photons) in a two components superposition state. The field is a physical system that plays the role of the Schrödinger's cat of the famous thought experiment⁶⁰. The overall system is entangled (not factorizable in its various components), and this entanglement (that constitutes the «coherent state») is further multiplied through successive interactions with the various (quantum) elements of the environment (such as those that constitute the observation apparatus), so that in the end the initial coherence does not show anymore, the effect being absorbed rapidly («*decoherence*»)⁶¹. In such a production experiment of a coherent entangled state, one can vary the parameters which determine the degree of coherence of the entangled system : these parameters are the number of photons

⁵⁸ On the theoretical interpretations of the phenomena and of the experiments, see notably Zurek [1982, 1991], D'Espagnat [1994], Omnès [1994a & b].

⁵⁹ In the experiment performed at the Laboratoire de physique de l'Ecole Normale Supérieure, Paris : Haroche, Brune & Raimond [1997].

⁶⁰ Schrödinger [1935].

⁶¹ For a reflexion on this state of things, see Paty [2000a].

that make the electric field, and the time of propagation of the entangled system (which is the time elapsed between its production and its analysis to determine whether it is still in a coherent state). The coherent state itself manifests as such by some interference which can be observed through a correlation between pairs of the atom-analyzers at detection. Coherence can then be controlled, and the condition and time when coherence ceases marks the shift from quantum to « classical-type » behavior of the system. This shift is attributed to the many interactions occurring between the system and the quantum components of the environment. The simple original entangled system combines itself with the states of the latter (each one being itself in a linear superposition), giving rise to a further entanglement : as the process is going further, it leads in an irreversible way to a many component entangled system. Quantum non separability forbids to go back to the original components simply entangled, and that original entanglement is lost in the end, as it gets diluted in the multiple entangled overall system, and has become definitely unappreciable. In the end, the quantum character of the state under study has been lost, although the whole process has been considered from a purely quantum point of view. So to speak, a « classical » behavior (a non quantum one) has been generated from quantum states merged inside entangled multiplicities.

One sees that the process of decoherence is not to be identified with that of measurement, for it happens softly through the quantum interactions themselves, whereas measurement is a process which chooses at once one of the final states by suppressing the others : the continuous soft (natural) process is (artificially) interrupted by the apparatus arrangement itself, which favours at random one only of the components of the final state and destroys the superposition. So to speak, measurement is decoherence plus projection (reduction) on one only of the components of the initial state of the considered physical system. Nevertheless, decoherence helps understanding the initial stage of such a transition, which seems, in the final stage, be purely of a statistical mechanics and thermodynamics nature. But I don't want to comment further on this, letting it for another opportunity, and I content myself in observing that evidence for the process of coherence to decoherence is *per se* an evidence for the physical character of the coherent, entangled, i. e. quantum linear superposition, state, shown as propagating in space and time.

8

CONCLUSION.

All the physical phenomena examined so far persuade us that the state function ψ represents (or describes) completely the state of the physical system. We mean by « complete representation » adequacy and covering : there is nothing more in the physical system than what is comprised in its theoretical representation by the state function.

If we restrict the question of the theoretical representation of quantum systems to the mere quantum level where these systems exhibit properties and

interact with others systems of a similar nature, the concepts of state function, quantum system, quanton, quantized field, with the magnitudes that qualify them, are self-sufficient : they do not ask, for conception and handling in theoretical work, any physical or conceptual underlying classical basis such as that of a undulatory or corpuscular substance, distinguishable and localized. For the quantum physics of atomic and subatomic phenomena and quantized fields the «quantum level» where these concepts operate is the fundamental level, and, in particular, physical systems are effectively represented by their «state functions», and physical magnitudes by their «operators». At this level of representation, it is not necessary to go back, for each magnitude and each state, to the practical circumstances of their determination that refer ultimately to observations with the help of classical apparatuses.

For theoretical thought at the quantum level, the classical systems constituting these apparatuses are only intermediary in the process of constitution of data, that are in the end translated in quantum terms. The data being acquired, the quantum domain let itself be conceived and explored in full conceptual and theoretical independence with respect to the classical domain.

This consideration does not diminish the problem of the quantum-to-classical relationship : it simply puts it aside, provisionally, as a fundamental problem. It is an epistemological and philosophical decision, taken in order to give the quantum domain and its theoretical representation the largest autonomy with respect to particular philosophical perspectives on knowledge. It has often been considered that (physical) knowledge is to be referred to observation, in the name of a primacy of perception in characterizing phenomena. However, contemporary reflection on science, and particularly on the various areas of physics, has led to conceive the relations of concepts and theories to *perception* as most indirect. The demand for *intelligibility* requires, as we suggested in the beginning, a direct and close connection with the *understanding*, that undertakes its theoretical elaborations by following a process of *rational construction* that is linked only in a mediate manner with the forms of *perception*. As it were, the phenomena under consideration, regarding the conceptualization and the theoretical insight obtained of them, are first brought to the understanding and secondly to the perception. If we refer these *phenomena* to (quantum) *objects*, that means that the latter are rationally constructed before being secondarily and indirectly perceived.

The question of the *physical meaning of magnitudes*, among which the representative state function of a system stands in the forefront, is henceforth more directly illuminated than by the current («orthodox») interpretation, conceiving this meaning *through reference to measurement*. The reference, according to the view here proposed, is to quantum phenomena, whose access is an indirect one but that is knowable by a rational construction, consistent and supported by data coming in the last instance from the perceptual (observation and experiment). Consequently, nothing opposes considering the state function in the form of a superposition (but basis-invariant) *describing effectively the state* of a physical system, evolving in the course of time.

The notion of *quantum physical state* differs from the current idea of a *physical state*, referred generally to magnitudes that are directly observable

through instruments ruled by the laws of classical physics. The difference between a physical phenomenon (or system) at the quantum level and a phenomenon (or a system) at the classical level is that *the second is closer (if not homogeneous) to its conditions of observation referred to perception*, whilst the first remains radically *distant from them* and is definitely *heterogeneous* to them⁶². This formulation of the difference between the classical and the quantum domains is free of philosophical bias about knowledge : it has the advantage of not limiting arbitrarily the capacity of the quantum to be intelligible. If they are dissimilar in their relationship with *perception*, their links to *understanding* are not of a different nature : all concepts of physics, classical as well as quantum ones, are expressed by magnitudes that are constructed (by man's mind) and abstract⁶³.

That a *quantum state* be accessible to experiment only indirectly does not affect the possibility *to get knowledge of it*. *Magnitudes* that characterize it are not either directly accessible, since they are not endowed with numerical values. To take into account all the elements considered in what precedes, we must therefore conceive an *extension of meaning*, to the quantum domain, of the notions of *physical magnitude* and of *physical state*, beyond the meaning usually accepted for them with classical physics (including the theory of relativity). This extension, legitimated by the *phenomena* (with a sense of this term that does not reduce them to mere objects of perception but that conceives them according to their capacity to be brought to *knowledge*), is actually already realized in practice by the main properties of the very formalism of quantum theory⁶⁴.

Such extensions of meaning have been a common procedure in mathematics as well as in physics, if we look back into their history : an example among many others in mathematics is the extension of the concept of number from integer to fractional, to irrational and then to imaginary and complex numbers ; as for physics, consider only motion, force, energy and also the extension of finite magnitudes to differential ones.... In all cases, such extensions were not the least obvious, and led to hard scientific and philosophical debates and controversies.

By proposing this extension of meaning for the concept of physical magnitude to forms that are not endowed with numerical values, to states that are linear superpositions of eigenstates, in order to insure epistemological *aseity* (self-contentness) for the quantum domain and its theoretical representation, we give primacy to understanding over perception, which is driven to an ancillary status. This is a pragmatic decision, that avoids deciding on the fundamental problem that still remains open of the relationship between the classical and the quantum, but that allows us at the same time to think with full legitimacy a wide range of phenomena, that might well be the base of all others. But, about this, we can not

⁶² There still remains, anyhow, between a *physical system* qualified as such, be it a classical or a quantum one, and its *conditions of observation*, a difference of nature. I want only to underline here that the working modes of measurement devices are referred to classical phenomena.

⁶³ Cf. Paty [1988a, and 2000a].

⁶⁴ Intuitively perceived by such theoreticians as Dirac, who extended the notion of commutative magnitudes expressed by ordinary numbers (*c*-numbers), to non-commutative ones (*q*-numbers) (Dirac [1926a & b, 1928]. Cf. Mehra & Rechenberg [1982], vol. 4, p. 162 sq, Darrigol [1992]), it has not, however, been explicitly legitimated as such, which ensured the permanence of the dominant philosophical interpretation (cf. Paty [2000a]).

swear, considering the present state of our knowledge. We can only relate it to the more fundamental and general question, still standing and in evolution, of the unity of physical phenomena and of a unified approach to them. But such an approach might precisely still be doomed to remain out of reach of present theories, until a deeper penetration of the unity of physical phenomena is obtained, through a sound unification of the fundamental interaction fields of matter.

To solve outside of this perspective, if it would prove possible, the «quantum problem of measurement», that is to say the nature of the relationship of the quantum and the classical, would be finally only of a limited interest. With the practical rule connecting, through probabilities, quantum magnitudes and their state functions with the corresponding classical entities determined from measurement devices, we have the minimal algorithm needed to place on a pragmatism basis the quasi autonomous existence of two coherent, intelligible, domains of physical reality, referred to their proper and specific phenomena and objects : the classical and the quantum.

ACKNOWLEDGEMENT

To John Stachel, who contributed so much (and still does) to pave the way to a deeper understanding of the rational exigencies of physics, through his pioneer einsteinian studies, particularly on the lifelong “Einstein's struggle” with quanta, and his exemplary harmonious practice of physical intuition and elaborated mathematical formalization in considering as well the physics of today as that of the past, always combined with sound epistemological insights. The reflections proposed above are a tribute to his enlightening and stimulating endeavours as well as - I hope - a testimony of intellectual kinship.

REFERENCES

- AGAZZI, Evandro (ed.) [1997]. *Realism and quantum physics*, Rodopi, Amsterdam, 1997.
- ASPECT, Alain, GRANGIER, Philippe & ROGER, Gérard [1981]. Experimental tests of realistic local theories via Bell's theorem, *Physical Review Letters* 47, 1981, 460-463.
- ASPECT, Alain, GRANGIER, Philippe & ROGER, Gérard [1982]. Experimental realization of Einstein-Podolsky-Rosen-Bohm *Gedankenexperiment* : a new violation of Bell's inequalities, *Physical Review Letters* 49, 1982, 91-94.
- ASPECT, Alain, DALIBAR, Jean & Roger, Gérard [1982]. Experimental tests of Bell's inequalities using time-varying analyzers, *Physical Review Letters* 49, 1982, 1804-1807.
- ASPECT, Alain [1983]. *Trois tests expérimentaux des inégalités de Bell par mesure de polarisation de photons*, Thèse de doctorat ès-sciences physiques, Université Paris-Sud, Orsay, 1983.
- BELL, John S. [1964]. On the Einstein-Podolsky-Rosen paradox, *Physics* 1,

1964,195-200. Repris in Bell 1987, pp. 14-21.

- [1966]. On the problem of hidden variables in quantum mechanics, *Review of Modern Physics* 38, 1966, 447-452. Also in Bell 1987, pp. 1-13.

- [1971]. Introduction to the hidden-variable question, in d'Espagnat [1971], p. 171-181 ; reprinted in Bell [1987], p. -.

- [1973]. Subject and object, in Mehra [1973], p. 687-690.

- [1976a]. Einstein-Podolsky-Rosen experiments in *Proceedings of the Symposium on Frontier problems in high energy physics, in honour of Gilberto Bernardini on his 70 th birthday, Pisa, 1976* , Scuola Normale Superiore, Pisa, pp. 33-45. Also in Bell 1987, pp. 81-92.

- [1976b]. The measurement theory of Everett and de Broglie's pilot wave, in Flato, M. et al. (eds.), *Quantum mechanics, determinism, causality, and particles*, Reidel, Dordrecht, 1976, p. 11-17. Also in Bell [1987], pp. 93-99.

- [1982]. On the impossible pilot wave, *Foundations of physics* 12, 1982 (n°10), 989-999. Also in Bell [1987], pp. 159-168.

- [1984]. Beables for Quantum Field Theory, preprint, CERN-TH 4035/84 (1984, Aug. 2th), in Bell [1987], p. 173-180.

- [1987]. *Speakable and unspeakable in quantum mechanics*, Cambridge University Press, Cambridge, 1987.

- [1989]. Against measurement, in Miller [1989], pp. 17-32.

BEN-DOV, Yoav [1987]. Bell's version of the «pilot-wave» theory, *Fundamenta Scientiae* 8, 1987, 331-343.

- [1988]. *Versions de la mécanique quantique sans réduction de la fonction d'onde*, Thèse de doctorat, Université Paris-13, 1988.

- [1989]. De Broglie's causal interpretation of quantum mechanics, *Annales de la Fondation Louis de Broglie* 14, 1989, 349-360.

BIMBOT, René & PATY, Michel [1996]. Vingt cinq années d'évolution de la physique nucléaire et des particules, in Yoccoz [1996], p. 12-99.

BOHM, David [1980]. *Wholeness and the implicate order*, Routledge and Kegan Paul, London, 1980.

BOHM, David & HILEY, David [1975]. On the Intuitive Understanding of Non-Locality as Implied by Quantum Theory, *Foundations of Physics* 5, 1975, 93-109 ; also in Leite Lopes & Paty [1977], p. 206-225.

BOHR, Niels [1935]. Can quantum-mechanical description of physical reality be considered complete ?, *Physical Review* 48, 1935, 696-702.

- [1958]. *Atomic physics and human knowledge*, New York, Wiley, 1958. French transl. by Edmond Bauer and Roland Omnès, *Physique atomique et connaissance humaine*, Paris, Gauthier-Villars ; new ed. established by Catherine Chevalley, Paris, Gallimard, 1991.

BROGLIE, Louis de [1953]. *Louis de Broglie, physicien et penseur*, Albin Michel, Paris, 1953.

BROWN, Laurie M., PAIS, Abraham & PIPPARD, Brian (eds.) [1995]. *Twentieth century physics*, 3 vols., Philadelphia Institute of Physics, New York, 1995.

BUNGE, Mario [1973]. *Philosophy of physics*, Reidel, 1973.

CLAUSER, John F. & SHIMONY, Abner [1978]. Bell's theorem : experimental tests and implications, *Reports on Progress in Physics* 41, 1978, 1881-1927.

COHEN, Robert S., HORNE, Michael & STACHEL, John (eds.) [1997a]. *Experimental Metaphysics. Quantum Mechanical Studies for Abner Shimony, Vol. One*, Kluwer, Dordrecht, 1997. .

COHEN, Robert S., HORNE, Michael & STACHEL, John (eds.) [1997b]. *Potentiality, Entanglement and passion-at-a-distance. Quantum Mechanical Studies for Abner Shimony, Vol. Two*, Kluwer, Dordrecht, 1997.

CORNELL, Eric & WIEMANN, Carl [1998]. Bose-Einstein condensation, *Scientific American* n° , 1997?, -. French tr., La condensation de Bose-Einstein, *Pour la science*, n° 247, mai 1998, 92-97.

COSTA, Newton da [1997]. *Logiques classiques et non classiques*, french transl. from portuguese (Brazil) and completed by Jean-Yves Béziau, Masson, Paris, 1997.

DARRIGOL, Olivier [1988]. Statistics and combinatorics in early quantum theory, *Historical Studies in the Physical Science* 19, 1988, 17-80.

- [1991]. Statistics and combinatorics in early quantum theory, II : Early symptoma of indistinguishability and holism, *Historical studies in the physical sciences* 21 (2), 1991, 237-298.

- [1992]. *From c-Numbers to q-Numbers. The classical Analogy in the History of Quantum Theory*, University of California Press, Berkeley, 1992.

DIRAC, Paul A. M. [1926a]. On the theory of quantum mechanics, *Proceedings of the Royal society of London A* 112, 1926, 661-677.

- [1926b]. The physical interpretation of the quantum dynamics, *Proceedings of the Royal society of London A* 113, 1926, 621-641.

- [1930]. *The principles of quantum mechanics*, Clarendon Press, Oxford, 1930. 4th ed., 1958.

DONCEL, Manuel ; HERMANN, Armin ; MICHEL, Louis & PAIS, Abraham [1987]. *Symmetries in physics (1600-1980)*, Universitat Autònoma de Barcelona, Bellaterra (Barcelona), 1987.

EBERHARD, P. H. [1977]. Bell's theorem without hidden variables, *Nuovo Cimento* 38B, 1977, 75-80.

EINSTEIN, Albert [1948]. Quantenmechanik und Wirklichkeit, *Dialectica* 2, 1948, 35-39.

- [1949]. Reply to criticism. Remarks concerning the essays brought together in this cooperative volume, in Schilpp 1949, p. 663-693. (Engl. transl. by P. A. Schilpp on the original in german, Bemerkungen zu den in diesen bande Vereinigten Arbeiten, in Schilpp [1949] german edition, p. 493-511).

- [1953]. Einleitende Bemerkungen über Grundbegriffe. Remarques préliminaires sur les principes fondamentaux (French transl. by Marie-Antoinette Tonnelat), in de Broglie [1953], p. 4-15.

- [1989-1993]. *Oeuvres choisies*, French transl. by the transl. group of ENS

Fontenay-St-Cloud *et al.*, éd. by Françoise Balibar *et al.*, Seuil/éd. du CNRS, Paris, 1989-1993, 6 vols.

EINSTEIN, Albert, PODOLSKY, Boris & ROSEN, Nathan [1935]. Can quantum-mechanical description of physical reality be considered complete ?, *Physical Review*, ser. 2, XLVII, 1935, 777-780.

EINSTEIN, Albert & BORN, Max [1969]. *Briefwechsel 1916-1955*, Nymphenburger Verlagshandlung, München, 1969.

ENZ, Charles P. (ed.) [1973]. W. Pauli's Scientific Work, in Mehra, J. (ed.), *The Physicist's Conception of Nature*, Reidel, Dordrecht, p. 766-799.

D'ESPAGNAT, Bernard (ed.) [1971]. *Foundations of Quantum Mechanics. Proceedings of the International School of Physics Enrico Fermi, Course XLIX*, Academic Press, New York, 1971.

D'ESPAGNAT, Bernard [1984]. Nonseparability and the Tentative Descriptions of Reality, *Physics Report*, 110, 1984, 201-264.

D'ESPAGNAT, Bernard [1994]. *Le réel voilé. Analyse des concepts quantiques*, Fayard, Paris, 1994.

FLATO, Moshe *et al.* (eds.), *Quantum mechanics, determinism, causality, and particles*, Reidel, Dordrecht, 1976.

FREEDMAN, Stuart J. & CLAUSER, John F. [1972]. Experimental test of local-hidden variable theories, *Physical Review Letters* 28, 1972, 938-941.

FREIRE Jr., Olival [1995]. *A emergencia da totalidade. David Bohm e a controversia dos quanta*, Doctoral Thesis, Dpto de História, Universidade de São Paulo, 1995.

FREIRE Jr., Olival, PATY, Michel & ROCHA BARROS, Alberto [2000]. Physique quantique et causalité selon Bohm. Un cas d'accueil défavorable, in Kragh, H. Marage, P. & Vanpaemel, G. (eds), *On the History of Modern Physics*, Archives de l'Académie internationale d'histoire des sciences, XXè Congrès International d'Histoire des Sciences, Liège, 1997, Actes, Collection de Travaux de l'Académie Internationale d'Histoire des Sciences, Brepols, Liège, 1999, in press.

FRENCH, Steven & KRAUSE, Decio [1999]. The Logic of Quanta, in Cao, Tian Yu (ed.), *Conceptual Foundations of Quantum Field Theory*, Cambridge University Press, 1999, 324-342.

GHIRARDI, Gian Carlo & WEBER, T. [1997]. An interpretation which is appropriate for dynamical reduction theories, in Cohen, Horne & Stachel [1997b], p. 89-104.

GRANGIER, Philippe [1986]. *Etude expérimentale de propriétés non-classiques de la lumière ; interférences à un seul photon*, Thèse de doctorat ès-sciences physiques, Université Paris-Sud, Orsay, 1986.

GREENBERGER, D. M., HORNE, M.A., ZEILINGER, A., [1989]. Going beyond Bell's theorem, in Kafatos, M. (ed.), *Bell's Theorem, Quantum Theory and Conceptions of the Universe*, Kluwer, Dordrecht, 1989, p. 73 sq.

- [1990]. Bell's theorem without inequalities, *American Journal of Physics*, 58, 1990, p. 1131 sq.

GRIFFIN, A, SNOKE, D.W. & STRINGARI, S. (eds.) [1995]. *Bose-Einstein*

condensation, Cambridge University Press, 1995.

HAROCHE, Serge, BRUNE, M. & RAIMOND, Jean-Michel [1997]. Experiments with single atoms in a cavity : entanglement, Schrödinger's cats and decoherence, *Philosophical Transactions of the Royal Society of London A*, 1997, 355, 2367-2380.

JAMMER, Max [1974]. *The Philosophy of Quantum Mechanics. The Interpretations of Quantum Mechanics in Historical Perspective*, Wiley, New York, 1974.

JAUCH, José-Maria [1968]. *Foundations of Quantum Mechanics*, Addison-Wesley, Reading (Mass.), 1968.

KASTLER, Alfred [1981]. On the historical development of the indistinguishability concept for microparticles, in Merwe [1981], p. 607-623.

LEITE LOPES, José & PATY, Michel (eds.) [1977]. *Quantum mechanics, a half century later*, Reidel, Dordrecht, 1977.

LÜDERS, Gerhart [1952]. Zum Bewegungsumkehr in quantisierten Feldtheorien, *Zeitschrift für Physik* 133, 1952, 325-339.

- [1954]., *Kgl .Danske Videnskab. Selskab, Mat.-Fys.Medd.* 28, 1954, n°5.

MEHRA, Jagdish (ed.) [1973]. *The Physicist's Conception of Nature*, Reidel, Dordrecht, 1973.

MEHRA, Jagdish and RECHENBERG, Helmut [1982]. *The historical development of quantum theory*, 7 vols., Springer, New York, 1982.

MERMIN, N. D. [1990]. Quantum Mystery Revisited, *American Journal of Physics*, 58, 1990, p. 731 sq.

MERWE, Alwyn van der (ed.) [1981] *Old and new questions in physics, cosmology, philosophy and theoreticall biology. Essays in honour of Wolfgang Yourgrau*, Plenum Press, New-York, 1981.

MILLER, Arthur (ed.) [1989]. *Sixty-two years of uncertainty. Historical, philosophical and physical inquiries into the Foundations of Quantum mechanics* (Proceedings of the International School of History of Science, Majorana Centre for Scientific Culture, Erice-Sicilia, 5-15 aug. 1989), Plenum Press, New York, 1989.

NEUMANN, John von [1932]. *Mathematische Grundlagen der Quantenmechanik*, Springer, Berlin, 1932 ; 1969 ; Dover, New-York, 1943. Engl. transl., *The Mathematical Foundations of Quantum Mechanics*, Princeton University Press, Princeton, 1955.

OMNES, Roland [1994a]. *Philosophie de la science contemporaine*, Gallimard, Paris, 1994.

- [1994b]. *The interpretation of Qiantum Mechanics*, Princeton University Press, Princeton, 1994.

PATY, Michel [1970]. Désintégrations électromagnétiques des bosons, *Ecole internationale de physique des particules élémentaires, Herceg-Novi*, 1970, 80 p.

- [1977]. The recent attempts to verify quantum mechanics, in Leite Lopes & Paty [1977], p. 261-289.

- [1981]. Interventions sur les *Implications conceptuelles de la physique quantique*, *Colloque de la Fondation Hugot du Collège de France, juin 1980*, *Journal de Physique, Supplément, fasc. 3, Colloque n° 2*, 1981, pp. C2 37-40, 78-80, 96-98, 109-111, 116-118.
 - [1982]. L'inséparabilité quantique en perspective, ou: Popper, Einstein et le débat quantique aujourd'hui, *Fundamenta Scientiae* 3, 1982, 79-92.
 - [1985]. Symétrie et groupes de transformation dans les théories contemporaines de la matière : jalons épistémologiques, *Colloque Abel-Galois, Lille, 21-25 février 1983, Première partie*, Institut de Recherches de Mathématiques Avancées (IRMA), Lille, 1985, fasc 5, 54 p.
 - [1986]. La non-séparabilité locale et l'objet de la théorie physique, *Fundamenta Scientiae* 7, 1986, 47-87.
 - [1988a]. *La matière dérobée. L'appropriation critique de l'objet de la physique contemporaine*, Archives contemporaines, Paris, 1988.
 - [1988b]. Sur la notion de complétude d'une théorie physique, in Fleury, Norbert ; Joffily, Sergio ; Martins Simões, José A. & Troper, A. (eds), *Leite Lopes Festschrift. A pioneer physicist in the third world* (Dedicated to J. Leite Lopes on the occasion of his seventieth birthday), World Scientific Publishers, Singapore, 1988, p. 143-164.
 - [1990]. Reality and Probability in Mario Bunge's *Treatise*, in Dorn, Georg and Weingartner, Paul (eds.), *Studies on Mario Bunge's Treatise*, Poznan studies in the philosophy of the sciences and humanities, Rodopi, Amsterdam-Atlanta, 1990, p. 301-322.
 - [1993]. *Einstein philosophe. La physique comme pratique philosophique*, Presses Universitaires de France, Paris, 1993.
 - [1995a]. The nature of Einstein's objections to the Copenhagen interpretation of quantum mechanics, *Foundations of physics* 25, 1995, n°1 (january), 183-204.
 - [1995b]. Les neutrinos, *Encyclopædia Universalis*, vol.12, 1995, p. 294-300.
 - [1997]. Predicate of existence and predictability for a theoretical object in physics, in Agazzi [1997], p. 97-130.
 - [2000a]. Interprétations et significations en physique quantique, *Revue Internationale de philosophie* (Brussels), n° 212, 2 (juin) 2000, 199-242. (Interpretations and significations in quantum physics).
 - [2000b]. The quantum and the classical domains as provisional parallel coexistents, *Synthese* (Dordrecht, Boston), 125, n° 1-2, oct.-nov. 2000, 179-200 (in French, Steven ; Krause, Décio & Doria, Francisco, (eds), *In honour of Newton da Costa*).
 - [2001]. Physical quantum states and the meaning of probability, as Chapter 14, in Galavotti, Maria Carla ; Suppes, Patrick and Costantini, Domenico, (eds.), *Stochastic Causality*, CSLI Publications (Center for Studies on Language and Information), Stanford (Ca, USA), 2001, p. 235-255.
 - [in press]. *Einstein, les quanta et le réel (critique et construction théorique)*.
- PAULI, Wolfgang [1955]. Exclusion Principle, Lorentz Group and Reflection of Space-Time and Charge, in Pauli, W., Rosenfeld, Léon & Weisskopf, Victor

(eds.), *Niels Bohr and the Development of Physics*, Pergamon, London, 1955, p. 30-51.

PERES, Asher [1978]. Unperformed experiments have no results, *American Journal of Physics* 46, 1978, 745-747.

PESIC, Peter D. [1991]. The principle of identity and the foundation of quantum theory. 1. The Gibbs paradox ; 2. The role of identity in the formation of quantum theory, *American Journal of Physics* 59 (11), 1991, 971-974, 975-979.

PFLEGOR, R. L. & MANDEL, L. [1967]. Interference of independent photon beam, *Physical Review* 159, 1967, 1084-1088.

POPPER, Karl R. [1959]. *The Logic of Scientific Discovery*, Hutchinson, London, 1959 ; 1968. (Engl. transl. modif. and augm. from the original in German, *Logik der Forschung*, Springer, 1935).

SCHILPP, Paul-Arthur (ed.) [1949]. *Albert Einstein : philosopher-scientist*, The library of living philosophers, Open Court, LaSalle (Ill.), 1949. Ré-ed. 1970. Transl. in German, *Albert Einstein als Philosoph und Naturforscher*, Kohlhammer Verlag, Stuttgart, 1955 (original texts of Einstein and Pauli).

SCHRÖDINGER, Erwin [1935]. Die gegenwärtige Situation in der Quantenmechanik, *Die Naturwissenschaften* 23, 1935, 807-812 ; 824-828 ; 844-849. Also in Schrödinger [1984], vol. 3, p. 484-501.

- [1984]. *Gesammelte Abhandlungen. Collected papers*, 4 vols., Verlag der Oesterreichischen Akademie der Wissenschaften/ Vieweg und Sohn, Braunschweig/Wien, 1984.

SCHWEBER, Sam S. [1994]. *QED and the men who made it*, Princeton University Press, Princeton, 1994.

SCHWINGER, Julian [1951-1953]. The theory of quantized fields, I, *Physical Review* 82, 1951, 914-927 ; II, *Ibid.* 91, 1953, 713-728 ; reprinted in Schwinger, J. ed., *Selected papers on electrodynamics*, Dover, New York, 1958, p. 342-355, 356-371. *Ibid.*, III, *Ibid.*, 91, 1953, 728-740 ; IV, *Ibid.*, 92, 1953, 1283-1299 ; reprinted in Schwinger, J., *Selected papers (1937-1976)*, ed. by Flato, M, Fronsdal, C. & Milton, K.A., Reidel, Dordrecht, 1979, p. 43-55 ; 56-72.

SEIDENGART, Jean et SCZECINIARZ, Jean-Jacques (éds.) [2000]. *Cosmologie et philosophie. En hommage à Jacques Merleau-Ponty*, numéro spécial de *Epistémologiques. Philosophie, sciences, histoire. Philosophy, Science, History* (Paris, São Paulo) 1, n°1-2, 2000.

SHIMONY, Abner [1993]. *Search for a naturalistic world view*, Cambridge University Press, Cambridge, 1993, 2 vols.

STAPP, Henry [1980]. Locality and reality, *Foundations of Physics* 10, 1980, 767-795.

WEYL, Hermann [1931]. *Gruppentheorie und Quantenmechanik*, Engl. transl. (on the 2nd rev. German ed.) by H. P. Robertson, *The Theory of Groups and Quantum Mechanics*, Dover, New York, 1950.

WHEELER, John A. & ZUREK, Wojciech H. (eds.) [1983]. *Quantum theory of measurement*, Princeton University Press, Princeton, 1983.

WIGNER, Eugen P. [1967]. *Symmetries and reflections*, Indiana University Press, Bloomington, 1967.

YANG, C. N. [1982]. The discrete symmetries P, T and C, in *Colloque International sur l'histoire de la physique des particules (Paris, 21-23 july 1982)*, *Journal de Physique*, Colloque C8, Supplé au n° 12, Tome 43, déc. 1982, 439-449.

YOCCOZ, Jean (éd.), *Physique subatomique : 25 ans de recherche à l'IN2P3, la science, les structures, les hommes*, Editions Frontières, Gif-sur-Yvette, 1996.

ZUREK, Wojcieh H. [1982]. Environment-induced superselection rules, *Physical Review D*, 24, 1981, 1516-1525.

ZUREK, Wojcieh H. [1991]. Decoherence and the Transition from Quantum to Classical, *Physics Today*, 44, oct. 1991, 36-44.