ANALYTICAL NON-RELATIVISTIC SCF-HARTREE-FOCK WAVE FUNCTIONS FOR THE GROUND STATE AND FOR EXCITED STATES OF In AND Tl IONS

R. Bauer, K. Differt, L. Schwan

► To cite this version:

R. Bauer, K. Differt, L. Schwan. ANALYTICAL NON-RELATIVISTIC SCF-HARTREE-FOCK WAVE FUNCTIONS FOR THE GROUND STATE AND FOR EXCITED STATES OF In AND Tl IONS. Journal de Physique Colloques, 1976, 37 (C7), pp.C7-177-C7-180. 10.1051/jphyscol:1976740 . jpa-00216897

HAL Id: jpa-00216897

<https://hal.science/jpa-00216897>

Submitted on 1 Jan 1976

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ANALYTICAL NON-RELATIVISTIC SCF-HARTREE-FOCK WAVE FUNCTIONS FOR THE GROUND STATE AND FOR EXCITED STATES OF In AND Tl IONS

R. BAUER, K. DIFFERT

Institut für Physik am Max-Planck-Institut für Metallforschung, Stuttgart, und Institut für Theoretische und Angewandte Physik der Universität Stuttgart, Pfaffenwaldring 57/VI, D-7000 Stuttgart 80, F. R. G.

and

L. SCHWAN

Physikalisches Institut, Teilinstitut 2, der Universität Stuttgart, F. R. G.

Résumé. — Des solutions Hartree-Fock non relativistes en forme de développements analytiques utilisant des fonctions orbitales du type Slater ont été obtenues pour l'état fondamental de In^+ , In^{++} , Tl^+ , Tl^{++} et pour des états excités de In^+ . Les fonctions d'onde et les intégrales de recouvrement des fonctions orbitales extérieures de ces métaux avec l'ion Cl^- sont présentées. Dans l'approximation non relativiste les ions de Tl possèdent essentiellement la même taille et les mêmes intégrales de recouvrement que les ions de In. Des corrections relativistes ont été estimées et sont discutées ; pour Tl^+ ces corrections sont à peu près trois fois plus grandes que pour In^+ . Dans le cas des centres de défauts de substitution d'ions métalliques monovalents dans KCl les déplacements des plus proches voisins sont estimés pour l'état fondamental et pour l'état excité. Tous ces déplacements se révèlent positifs, indiquant un déplacement des voisins radialement vers l'extérieur.

Abstract. — Non-relativistic Hartree-Fock solutions for the ground state of In^+ , In^{++} , Tl^+ , Tl^{++} and for excited states of In^+ have been obtained in the form of analytical expansions in Slater-type orbitals. The wave functions and overlap integrals of the outer metal orbitals with the chlorine ion are presented. In the non-relativistic approximation the Tl ions have essentially the same size and the same overlap integrals as the In ions. Relativistic corrections are estimated and discussed ; for Tl^+ they are almost three times larger than for In^+ . The nearest neighbour displacements for substitutional defect centres of the monovalent ions in KCl are derived for the ground state and for an excited state. All displacements come out as positive, indicating an outward radial displacement of the neighbours.

1. Introduction. — In an attempt to improve the understanding of heavy metal ion centers in ionic solids, we have calculated non-relativistic Hartree-Fock (HF) wave functions for the ground states of In^+ , In^{++} , Tl^+ , Tl^{++} , and for several excited states of In^+ and In^0 , using Roothaan's method [1, 2] of analytical expansion of the orbital wave functions in Slater-type orbitals (STOs). In this representation, the orbitals are described by a limited number of parameters and may hence be easily communicated and are readily applicable in molecular or solid state problems.

In this note we present atomic Hartree-Fock data and some overlap integrals of the substitutional metal ions with neighbouring ions of the crystal environment in KCl. We then discuss corrections due to relativistic effects. Finally the data are used to estimate the sign and magnitude of the radial displacements of the nearest-neighbours (nn) of the defect centers.

2. Results of the Hartree-Fock calculations. — The STO basis, being similar to that used by Synek *et al.* [3] for Nd^{3+} , may be characterized as an extended double zeta basis set yielding the accuracy of the near HF limit. The basis set comprises 30 STOs for the cases of Tl^+ and Tl^{++} . The program handles the full nonlocal exchange interaction. Further details on the numerical calculations and the accuracy of the results will be given elsewhere.

Figure 1 presents the probability amplitude of the Tl^+ 6s orbital in the (x, y, z) plane in a vertical section through the nucleus. It shows the five radial nodes of a 6s function and the rapid oscillations near the nucleus.

In Table I, the mean orbital radii $\langle r \rangle$ for the outer shell of the ionic ground states are given, and the overlap integrals between the substitutional metal ions and a n.n. chlorine ion in KCl are listed.

The following figures show the radial part of the

FIG. 1. — Probability amplitude of the Tl^+ 6s HF orbital in the (x, y, z) plane in a vertical section through the nucleus. The sharp negative central peak describes the finite amplitude of the s state at the nucleus.

FIG. 2. — Radial part of HF orbitals of Cl^- and of the ground states of K^+ $1S_0$ (solid lines) and In^+ $1S_0$ (dashed lines). The centres of the ions are located at the crossing points of the coordinate axes and are separated by the interatomic distance in KCl , $r_0 = 3.12 \text{ \AA} = 5.896 \text{ a.u.}$

metal orbitals and their overlap with the nn chlorine 3s and 3p orbitals. The nuclei are located in the two crossing points of the coordinate axes. The radial functions are multiplied by the radial distance r from their respective nucleus. The even and odd parities are indicated for the Cl^- 3s and 3p orbitals, respectively.

Figure 2 compares the 5s and 4d orbitals of In^+ (dashed lines) with the outer 3s and 3p shells of K^+ representing the situation in the perfect lattice. It is seen that the In^+ ion is considerably larger than K^+ , forming much larger overlap integrals (cf. Table I), though as the In^+ overlap does not depend so strongly on the interatomic distance.

Figure 3 shows the 6s, 5d and 4f orbitals of Tl^+ (full lines) and Tl^{++} (dashed lines). The HF radial extensions of the Tl ions are nearly the same as those of the In ions, their mean radii differing but by 3 %.

The reduction of the occupation number of the outer s shell from two to one in going from Tl^+ $6s^2$ to Tl^{++} $6s^1$ reduces its mean radius $\langle r \rangle$ by only 7 % and has a quite small effect on the shape of the 5d shell. The 6s mean overlap integrals differ by 17 per cent.

In Figure 4 the outer orbitals of several excited states of In^+ are collected. The 5p state lying only 5 eV above the ground state is somewhat more extended than the

FIG. 3. — Radial part of HF orbitals of Cl^- and of the ground states of Tl^+ $1S_0$ (solid lines) and Tl^{++} $2S_{1/2}$ (dashed lines). The ions are separated by the interatomic distance in KCl .

5s orbital, forming quite large overlap integrals of the order of 0.2 to 0.3. The 5d and 6s states lying 10 eV above the ground state are widely extended having their maxima near the nn nucleus. Hence the $|3p_\sigma\rangle$ overlap is very small due to the odd parity of the 3p function, while the $|3p_\pi\rangle$ and $|3s\rangle$ overlaps are of the order of 0.4. The excited state orbitals of neutral

TABLE I

Mean orbital radius $\langle r \rangle$ and overlap integrals for the outer orbital of ionic ground states of K, In, and Tl. The overlap integrals are calculated for the nearest-neighbour distance in KCl , $r_0 = 3.12 \text{ \AA}$

Atomic State	$\text{K}^+ \text{Ar}$	$\text{In}^+ 5s^2$	$\text{In}^{++} 5s^1$	$\text{Tl}^+ 6s^2$	$\text{Tl}^{++} 6s^1$
Orbital nl	$1S_0$ 3p	$1S_0$ 5s	$2S_{1/2}$ 5s	$1S_0$ 6s	$2S_{1/2}$ 6s
$\langle r \rangle [\text{\AA}]$	0.757 4	1.429	1.325	1.486	1.382
$\langle nl_\sigma \text{Cl}_{3p\sigma}^- \rangle$	0.070 5	0.182 5	0.162 8	0.187 5	0.167 3
$\langle nl_\sigma \text{Cl}_{3s}^- \rangle$	0.025 7	0.085 5	0.062 4	0.092 0	0.064 9

FIG. 4. — Radial part of HF orbitals of excited states of In^+ and their overlap with the Cl^- 3p orbital in KCl. The shown 5s function belongs to the states $5s^1$ $5d^1$ and $5s^1$ $6s^1$.

In^0 not shown here are even more extended, their mean radii being 20 per cent larger than those of the corresponding states of In^+ .

3. Relativistic corrections. — The relativistic effects due to the high electron velocities in the inner $s_{1/2}$ and $p_{1/2}$ shells (jj coupling) of large Z atoms cause a spatial tightening and an increase of the binding of these orbitals. This effect is transferred via the orthogonality constraints of the radial parts of the orbitals to the outer $s_{1/2}$ and $p_{1/2}$ electrons, whereas in a compensating reaction the $p_{3/2}$, d, and f shell electrons loosen slightly and extend radially [4, 5].

The relativistic corrections for several atomic observables of the HF ground states of all neutral atoms are given in [5]; from these data and from a comparison between the calculated HF ionization energies and the optical ionization potentials the corrections for the ions may be found.

TABLE II

Relativistic corrections for the orbital energy ε , the mean radius $\langle r \rangle$, and the mean nearest-neighbour overlap $\bar{S}(\text{Me}^+ | \text{Cl}^-)$ in KCl, in per cent of the Hartree-Fock values

Per cent	$\Delta\varepsilon$	$\Delta\langle r \rangle$	$\Delta\bar{S}(\text{Me}^+ \text{Cl}^-)$
$\text{Ga}^+ 4s^2$	2.4	— 1.5	— 3.5
$\text{In}^+ 5s^2$	7.0	— 4	— 9
$\text{Tl}^+ 6s^2$	17.0	— 11	— 24

Table II shows the relativistic corrections obtained for the orbital eigenvalue ε , the mean orbital radius $\langle r \rangle$, and an average of the overlap integrals with the nn Cl^- ion in KCl. The corrections for the overlap integrals are calculated from those for $\langle r \rangle$

using the relation between the change in $\langle r \rangle$ and the change in the interatomic overlap found by comparing the data for the monovalent and divalent ions given in Table I.

It is seen that the corrections for Tl ($Z = 81$) are almost three times larger than those for In ($Z = 49$) and seven times larger than those for Ga ($Z = 31$). In the non-relativistic approximation, In and Tl are almost identical in all their properties, cf. table I. The relativistic effects yield a much stronger additional binding for the Tl 6s shell than for the In 5s shell; they turn Tl into a monovalent metal forming CsCl-type ionic solids, while In stays trivalent like Ga and Al and forms preferentially covalent bonds. The Tl ions eventually have a smaller size than the In ions, the ionic radius and the mean nn overlap of Tl^+ being smaller by 6 and 15 per cent, respectively, than those of In^+ .

4. The equilibrium configuration of In and Tl centres in KCl. — The equilibrium displacements in the lattice about a substitutional isovalent metal ion centre may be obtained to a first approximation by contrasting the short range forces of the substituted cation of the perfect lattice (K^+) with those of the substituent cation (In^+ or Tl^+). In the framework of the Löwdin. Approximation [6, 7] of the HF scheme for ionic solids, the short range interaction is roughly proportional to the square of the interatomic overlap, and the forces in addition depend on the steepness of its variation with the interatomic distance. From our overlap data and from an estimate of the Van der Waals strengths being considerably larger for In^+ and especially for Tl^+ than for K^+ representing the perfect lattice, we obtain the nn displacements of substitutional centres given in table III. The displacements being related to the perfect lattice positions are all positive, indicating a radial outward shift both for the ground states and for the excited states of the centres. The vibronic interaction with the lattice has not yet been taken into account on this stage of approximation.

TABLE III

Equilibrium displacements of the nearest neighbours of substitutional In^+ and Tl^+ centres in KCl. The displacements refer to the perfect lattice positions and are given in per cent of the nn distance, $r_0 = 3.12 \text{ \AA}$

Radial displ.	K ⁺	In ⁺	Tl ⁺	In ⁺	5s ¹	5p ¹	Tl ⁺ 6s ¹	6p ¹
	ground states			excited states				
	σ	π	σ	π	σ	π	σ	π
$u_{s,r} + v_{d,w}$	0	6.5	4.0	9.5	2.5	6.0	1.7	

The relaxation of the excited states $ns^1 np^1$ consists of an additional outward movement of the two σ neighbours and an inward relaxation of the four π

neighbours, leading to a strong tetragonal distortion of the lattice. Due to the stronger relativistic effects for Tl causing a radial contraction of the $s_{1/2}$ and $p_{1/2}$ orbitals, the displacements associated with the Tl centres are smaller than those of the In centres.

Further details on the derivation of the lattice distortions will be given elsewhere. A more thorough treatment of the short range forces and the inclusion of other interactions not yet considered, such as the

polarization energy originating from the fields of deformation dipoles, will probably not alter the relative sizes and the sign of the reported displacements.

Note added in proof. — A detailed account of the subject of this note is given by the same authors in *Semiconductors and Insulators* 1 (1977).

Acknowledgments. — This work is part of the program of the « *Sonderforschungsbereich 67 : Defekturen* ».

References

- [1] Roothaan, C. C. J., *Rev. Mod. Phys.* **23** (1951) 69.
- [2] Roothaan, C. C. J., Bagus, P. S., in *Methods of Computational Physics*, B. Alder *et al.* (Pergamon) 1963, Vol. 2.
- [3] Synek, M., Grossgut, P., *Phys. Rev. A* **1** (1970) 1.
- [4] Desclaux, J. P., *J. J. Quant. Chem.* **6** (1972) 25.
- [5] Desclaux, J. P., *Atomic Data and Nuclear Data Tables* **12** (1973) 312.
- [6] Abarenkov, I. V., Antonova, I. M., *Phys. Status Solidi* **38** (1970) 783.
- [7] Bauer, R., *Phys. Status Solidi (b)* **50** (1972) 225 ; Bauer, R., *J. Nonmetals* **1** (1973) 257.

DISCUSSION

R. J. Friauf. — The effect of the Madelung potential and the neighboring ions on the Cl^- wave functions is by no means negligible in KCl. The overlap integrals for K^+Cl^- , and even more so for Cl^-Cl^- , are reduced by factors of 2 to 5.

R. Bauer. — The large $\text{nn Me}^+\text{-Cl}^-$ overlaps are not affected much by a reordering of the anion p orbital ; the electrons in the anion orbital have no great chance to escape from overlapping with the extended cation s orbitals. Moreover, if the magnitude of the overlap is decreased by reordering, the steepness of its dependence on the interatomic dis-

tance is increased, thus compensating for a change in the force and essentially in the force constants.

A. Laforgue. — How do you calculate the relativistic term ?

R. Bauer. — We correct our self-consistent Hartree-Fock data for the relativistic effects. The corrections are obtained from the ratios between the Dirac-Fock data and the Hartree-Fock data for the neutral atom ground state given in ref. [5]. These corrections are scaled to the ionic ones using the corrections of optical data. Details of this procedure will be published in *Semiconductors and Insulators* 1 (1977).