INSTABILITIES INDUCED BY RESONANT ABSORPTION OF AN ELECTROMAGNETIC WAVE IN AN INHOMOGENEOUS PLASMA

J.C. Adam, A. Gourdin Servenire and G. Laval

Centre de Physique Thorique de l'Ecole Polytechnique⁺, Plateau de Palaiseau, 91128 Palaiseau Cedex, France

Rsum - Dans le cas d'une incidence oblique, une onde lectromagntique cohrente gnre dans un plasma inhomogne des ondes longitudinales. Ces ondes longitudinales se propagent non-linairement dans le plasma pour des flux d'intensit modre. Ces non-linarits sont responsables d'instabilits paramtriques couplant les ondes de Langmuir aux ondes ioniques. On observe alors une transition entre des comportements priodiques et chaotiques correspondant des missions de solitons. La consquence de ce comportement sur les ondes rflchies et le plasma sont discuts.

Abstract - For oblique incidence and appropriate polarization, a coherent electromagnetic wave generates longitudinal Langmuir waves in an inhomogeneous plasma. These longitudinal waves propagate non-linearly through the plasma even for rather low incident energy flux. This non-linearity induces instabilities by coupling with two other Langmuir waves and an ion sound waves. For weak non-linearity or strong damping, periodic time dependent solutions are found. They correspond to periodic emissions of soliton-like perturbations. For stronger non-linearity, the solitons are emitted in a chaotic way. The consequence of these behaviors on the reflected electromagnetic wave and the plasma will be discussed.

I - INTRODUCTION

In an inhomogeneous plasma, when the electric field of an electromagnetic wave has a component along the density gradient, large amplitude longitudinal waves are excited at points where the plasma frequency coincides with the wave frequency. This mechanism is called resonant absorption. It takes place for obliquely incident p-polarized waves. The large amplitude longitudinal wave can be modulationally unstable. The absorption mechanism becomes non-stationary. We describe here the coupling between these unstable longitudinal waves and the electromagnetic waves.

II - THE ABSORPTION MECHANISM

A plane electromagnetic wave is supposed to be obliquely incident on an inhomogeneous plasma. The plasma frequency $\omega_p(x)$ coincides with the wave frequency ω_o for $x=0$ and is an increasing function of x such that $\omega_p(x) \rightarrow 0$ as $x \rightarrow -\infty$.

As in Ref. [1], the wave vector $\vec{k}_o$ of the incident wave is lying in the x,y plane and the electric field is contained into the same plane. We assume

$$\left| \frac{1}{k_o} \frac{d}{dx} L_n \omega_p \right| \ll 1 \quad k_o \lambda_D \ll 1 \quad (1)$$

where λ_D is the Debye length.

When collisional and thermal effects are negligible, the cold plasma approximation can be used so that we may write

⁺ "Groupe de Recherche du C.N.R.S. n 48"

$$\vec{j} = i\epsilon_0 \frac{\omega_p^2}{\omega} \vec{E} \quad \text{with} \quad \vec{E} = \vec{E}_0 e^{-i\omega_0 t}$$

and the equation for the B_z magnetic field is

$$\partial_x \epsilon^{-1} \partial_x B_z + \epsilon^{-1} \partial_y^2 B_z + k_0^2 B_z = 0 \quad (2)$$

with $\epsilon = 1 - \omega_p^2/\omega^2$.

By assuming $B_z = B(x) e^{-ik_0 \sin \theta y - i\omega t}$, Equation (2) becomes

$$\partial_x \epsilon^{-1} \partial_x B + k_0^2 [1 - \epsilon^{-1} \sin^2 \theta] B = 0 \quad (3)$$

Such an equation is singular for $x=0$. In the vicinity of this point, Eq. 3 has a regular solution y_1 and a singular one y_2 . In order to find their behavior in this region, we set $\epsilon = -x/L$ in Eq. 3 where L is the local density gradient length. We find

$$y_1 = k_0^2 x^2 \left[1 + \frac{k_0^2 \sin^2 \theta}{8} x^2 + \frac{k_0^2 L^{-1}}{15} x^3 + \dots \right] \quad (4)$$

$$y_2 = \sin^2 \theta y_1 \ln(k_0 x) + 2 \left[1 + \frac{k_0^2 L^{-1}}{3} x^3 + \dots \right]$$

The appropriate boundary conditions cannot be fulfilled by y_1 since by multiplying Eq. 3 by B^* and integrating on x we obtain

$$\int_{x_0}^{x_1} dx \left\{ \epsilon^{-1} \left| \frac{\partial B}{\partial x} \right|^2 - k_0^2 [1 - \epsilon^{-1} \sin^2 \theta] |B|^2 \right\} = \left[B^* \epsilon^{-1} \frac{\partial B}{\partial x} \right]_{x_0}^{x_1} \quad (5)$$

For $x_1 \rightarrow +\infty$ and $x_0 \rightarrow 0$, the right hand side vanishes whenever $B = y_1$. Thus we have generally $B = \lambda y_1 + \mu y_2$ with $\mu \neq 0$. But we also have

$$E_x = -C \sin \theta \epsilon^{-1} B \quad (6)$$

and $\mu \neq 0$ implies $B(0) \neq 0$. Thus E_x is diverging.

The physical meaning of this divergence results from the resonant excitation of plasma waves at the critical density where $\omega_0 = \omega_p$. The model must be improved in order to broaden this resonance. The excitation of these plasma waves and their subsequent absorption by the plasma is then the dominant absorption mechanism of the incident electromagnetic wave.

III - EQUATIONS FOR THE PLASMA WAVES

Since the typical scale length of plasma waves is λ_D , conditions (1) insures that we can neglect the x dependence of the electromagnetic field in the plasma wave propagation. For all high frequency quantities a_h we set

$$a_h(x, t) = \text{Re} \{ a(x, t) \exp(-i\omega_0 t) \} \quad (7)$$

where the time dependence of the envelope a is slow enough to be neglected in the electromagnetic field equations but at the resonance for the longitudinal waves. Thus we shall be able to take into account the time dependence induced by the instabilities.

We take into account the finite group velocity of longitudinal plasma waves and the low frequency modification of the plasma density n induced by the ponderomotive effect of the high frequency electric field. From

$$\partial_y B_z = \mu_0 j_x + \frac{1}{c^2} \partial_t E_x \quad (8)$$

and

$$\partial_t j_x = (n_0 + n_\ell) \frac{e^2}{m_e} E_x - \frac{3\epsilon_0 T}{m_e} \partial_x^2 E_x \quad (9)$$

we obtain for the fields envelopes

$$k_y \omega_0 c^2 B(0) = \omega_{pe}^2 \left[\frac{x}{L} + \frac{n_\ell}{n_0} \right] E - 2i\gamma \omega_p E - 3V_e^2 \partial_x^2 E - 2i\omega_p \partial_t E \quad (10)$$

where E is the envelope of E_x , $\gamma(x)$ a damping rate, V_e the electron thermal velocity and where we have used a linear approximation for the x dependence of ω_p^2 . The low frequency density perturbation n_ℓ are given by

$$\partial_t^2 n_\ell - C_s^2 \partial_x^2 n_\ell = n(0) C_s^2 \partial_x^2 \frac{\epsilon_0 |E|^2}{n_c T} \quad (11)$$

Knowing $B(0)$, we can solve the non-linear equations (10) and (11). However, $B(0)$ is connected to E as we show in the next section.

IV - CONNECTING FORMULAS

Inside the resonant layer at critical density we have

$$i\omega_0 B = \frac{\partial E_y}{\partial x} - ik_0 \sin \theta E.$$

B is almost constant across the resonant layer and E is very large over a small distance η across $x = 0$. Thus we may write

$$[E_y]_{-\eta}^{+\eta} = -ik_0 \sin \theta \int_{-\eta}^{+\eta} E(x) dx$$

For from the resonance, we have again

$$E_y = \frac{c^2}{i\omega_0 \epsilon} \partial_x B$$

and with $\epsilon \approx -x/L$, we obtain

$$\left[\frac{1}{x} \partial_x B \right]_{-\eta}^{+\eta} = -\frac{k_o^2 \sin \theta}{LC} \int_{-\eta}^{+\eta} E(x) dx \quad (12)$$

For $x > 0$, a solution of Eq. 3 such that $B \rightarrow 0$ for $x \rightarrow \infty$ is of the form

$$B = \ell y_1 + y_2$$

where ℓ is real.

For $x < 0$, a solution of Eq. 3 such that it corresponds for $x \rightarrow -\infty$ to a pure in going wave of amplitude unity and given phase is

$$B = \alpha y_1 + \beta y_2$$

and for a pure outgoing wave,

$$B = \alpha^* y_1 + \beta^* y_2.$$

Using Eq. 5, we find that

$$\text{Im}[4L \beta^* \alpha k_o] = -1 \quad (13)$$

which is the condition for conservation of the energy flux. If B_o and B'_o are the magnetic component of the incident and reflected wave, the appropriate solution of Eq. 3 for $x < 0$ is

$$B_- = (B_o \alpha + B'_o \alpha^*) y_1 + (B_o \beta + B'_o \beta^*) y_2$$

and for $x > 0$

$$B_+ = \rho(\ell y_1 + y_2)$$

We have $B_-(0) = B_+(0)$ which gives with Eq. 12 :

$$B'_o = -B_o \frac{\beta \ell - \alpha}{\beta \ell - \alpha^*} - \frac{\sin^2 \theta}{2Lc[\beta \ell - \alpha^*]} \int_{-\eta}^{+\eta} E dx \quad (14)$$

$$\rho = + \frac{B(0)}{2} = \frac{(\beta \ell - \alpha^*)^{-1}}{2L} \left[\frac{B_o}{2k_o} - \frac{\sin^2 \theta \beta^*}{c} \int E dx \right] \quad (15)$$

For a given B_o , Eq. 15 provides the value of $B(0)$ to be inserted in Eq. 10. Solving Eq. 10 determines $E(x)$ from which we obtain the reflected field B'_o by using Eq. 14.

V - THE WEAK ABSORPTION CASE

It is well known^[2] that the absorption coefficient remains much smaller than unity unless $(k_o L)^{1/3} \sin \theta \sim 1$. In such cases ρ depends only weakly on E . Thus we may assume that $B(0)$ is given, does not depend on time and we solve Eqs. 10, 11. This

approximation has been used^{1,3} to obtain stationary solutions. It has been shown that non-linear profile modifications by ponderomotive effects lead to a reduction of the absorption coefficient a . We find

$$\frac{a}{a_0} = 0.8 p^{-0.90} \quad \text{for } 2 < p < 100 \quad (16)$$

where a_0 is the linear absorption and

$$p = \left(\frac{L}{\lambda_D}\right)^2 \frac{\epsilon_0 |E_d|^2}{n_0 T}$$

with $E_d = \sin \theta C B(0)$.

If Eq. 10 is solved numerically, keeping the time dependence of E , it is found that this stationary solution is unstable for $p > 1$ for all physically relevant values of the parameters. It can be stabilized only by introducing large values of γ such that the non-linear effects become negligible and we recover $a \approx a_0$.

For $p \approx 1$, the time dependent solution looks like periodic emission of solitons from the resonant region where $x \approx 0$ (Fig. 1). These solitons convect towards the low density plasma where they are absorbed by Landau damping. For high values of p , the soliton emission becomes chaotic (Fig. 2)

Figures 1 (a) - Space time representation of $E(t)$ for $p = 1.33$. Each curve shows the x -dependence of E at given time t . This time can be found at the intersection of the curve with a vertical axis.

(b) - Spatial average value of E for $p = 1.33$ as a function of time.

Figures 2 (a) - Space time representation of $E(t)$ for $p = 38$. Each curves shows the x -dependence of E at given time t . This time can be found at the intersection of the curve with a vertical axis.

(b) - Spatial average value of E for $p = 38$ as a function of time.

VI - COUPLING

In order to couple the solutions of Eqs. 10, 11 to the radiation field, we may use Eq. 14 to determine the amplitude of the reflected field. It is easier to determine the absorption coefficient by noticing that it corresponds to the work of the electric field on the plasma particles. We find that the absorbed energy flux is given by

$$\pi = \frac{ky}{2\mu_0} \operatorname{Im} \int dx E^* B_z(0) \quad (17)$$

On Figs. 1 and 2, we have plotted the ratio $a/a_0 = \pi/\pi_0$ where a is the absorption coefficient, a_0 and π_0 being the values of a and π in the usual linear case.

It is seen that a oscillates with a frequency much larger than the amplitude time scale. It must be noticed also that for large p , the average value of a is larger than the stationary state estimation given by Eq. 16.

As regards the high frequency oscillation of a , we can understand its origin in a particular case. For weak amplitudes of the incident waves and large L , we can have

$$\partial_t^2 n_\ell \ll c_s^2 \partial_x^2 n_\ell$$

so that Eq. 11 reduces to

$$n_\ell = -n_0 \partial_x^2 \left(\frac{\epsilon_0 |E|^2}{n_c T} \right) \quad (18)$$

and Eq. 10 becomes

$$E_d = \left[\frac{x}{L} - \frac{\epsilon_0 |E|^2}{n_c T} \right] E - 3\lambda_D^2 \partial_x^2 E - \frac{2i}{\omega_p} [\partial_t E + \gamma E] \quad (19)$$

For $E_d = \gamma = 0$, Eq. 10 can be transformed into a non-linear Schrödinger equation⁽⁴⁾ which can be solved by the inverse scattering method. Numerically, we find that solitons are emitted in the vicinity of the origin. These solitons reach a large amplitude before convecting towards $x \rightarrow -\infty$. Thus a solution of Eq. 19 with $E_d = 0$ has some relevance with the real problem, since, part of the time, E_d and γE are small perturbations. Such a solution can be written

$$E = a e^{-i[\delta\omega t + \alpha(xt - \lambda^2 t^3)]} \operatorname{sech}[b(x + \gamma t^2)]$$

with $\frac{\delta\omega}{\omega} = -\frac{\epsilon a^2}{2n_c T}$, $\alpha = \frac{\omega p}{2L}$, $\lambda^2 = \frac{v^2}{L}$, $b = \lambda_D^{-1} \sqrt{\frac{\epsilon_0 a^2}{n_c T}}$, $\gamma = \frac{\lambda_D^2}{L} \omega_p^2$

which describes a soliton centered in $x = 0$ at $t = 0$ and uniformly accelerated towards the low density. For such a solution we find

$$\int_{-\infty}^{+\infty} E dx = i e^{-i\delta\omega t} \frac{a}{b} \pi [\operatorname{sh}(\pi a t / b)]^{-1}$$

Thus, by inserting this result in Eq. 14 we find that such a soliton radiates at a frequency $\omega_0 + \delta\omega$. The intensity of the radiation do not depend on the soliton amplitude but the duration of the emission is $(\epsilon_0 a^2 / n_c T)^{1/2} (2L/\lambda_D)$. These results

explain the fast oscillation of a and shows that the reflected light should have a red-shifted component.

CONCLUSION

When the ponderomotive effect is the dominant non-linear mechanism, resonant absorption of electromagnetic waves in an inhomogeneous medium is a non stationary process for $p > 1$ and weak damping rates. The non-linear evolution shows a transition from a periodic to a disordered regime. Soliton like perturbation plays a dominant role in the absorption mechanism as well as in the frequency spectrum of reflected light.

REFERENCES

1. ADAM, J.C., GOURDIN-SERVENIERE, A., LAVAL, G., Phys. Fluids 25 (1982) 376.
2. GINZBURG, V.L., The Propagation of Electromagnetic Waves in Plasmas (Pergamon, N.Y. 1970), 2nd Ed.
3. GILDENBURG, V.B., FRAINAN, G.M., Zh. Eksp. Teor. Fiz. 69 (1975) 1601
Sov. Phys. J.E.T.P. 42 (1975) 816.
4. CHEN, H.C., LIU, C.S., Phys. Rev. Lett. 37 (1976) 693.