

A CHARACTERIZATION OF DIMENSION FREE CONCENTRATION IN TERMS OF TRANSPORTATION INEQUALITIES

NATHAEL GOZLAN

UNIVERSITÉ PARIS EST,
Laboratoire Analyse et Mathématiques Appliquées,
UMR CNRS 8050,
5 bd Descartes, 77454 Marne la Vallée Cedex 2, France

ABSTRACT. The aim of this paper is to show that a probability measure μ on \mathbf{R}^d concentrates independently of the dimension like a gaussian measure if and only if it verifies Talagrand's \mathbf{T}_2 transportation-cost inequality. This theorem permits us to give a new and very short proof of a result of Otto and Villani. Generalizations to other types of concentration are also considered. In particular, one shows that the Poincaré inequality is equivalent to a certain form of dimension free exponential concentration. The proofs of these results rely on simple Large Deviations techniques.

1. INTRODUCTION

One says that a probability measure μ on \mathbf{R}^d has the gaussian dimension free concentration property if there are three non-negative constants a , b and r_o such that for every integer n , the product measure μ^n verifies the following inequality:

$$(1.1) \quad \forall r \geq r_o, \quad \mu^n(A + rB_2) \geq 1 - be^{-a(r-r_o)^2},$$

for all measurable subset A of $(\mathbf{R}^d)^n$ with $\mu^n(A) \geq 1/2$ denoting by B_2 the Euclidean unit ball of $(\mathbf{R}^d)^n$.

The first example is of course the standard Gaussian measure on \mathbf{R} for which the inequality (1.1) holds true with the sharp constants $r_o = 0$, $a = 1/2$ and $b = 1/2$. Gaussian concentration is not the only possible behavior ; for example, if $p \in [1, 2]$ the probability measure $d\mu_p(x) = Z_p^{-1}e^{-|x|^p} dx$ verifies a concentration inequality similar to (1.1) with r^2 replaced by $\min(r^p, r^2)$. In recent years many authors developed various functional approaches to the concentration of measure phenomenon. For example, the Logarithmic-Sobolev inequality is well known to imply (1.1) ; this is the renowned Herbst argument (which is explained, for example, in Chapter 5 of Ledoux's book [Led01]). Among the many functional inequalities yielding concentration estimates let us mention: Poincaré inequalities [GM83, BL97], Logarithmic-Sobolev inequalities([Led96, BG99]), modified Logarithmic-Sobolev inequalities

Date: April 18, 2008.

1991 Mathematics Subject Classification. 60E15, 60F10 and 26D10.

Key words and phrases. Concentration of measure, Transportation-cost inequalities, Sanov's Theorem, Logarithmic-Sobolev inequalities.

[BL97, BZ05, GGM05, BR06], Transportation-cost inequalities [Mar86, Tal96, BG99, Sam00, BL00, OV00, BGL01, Goz07], inf-convolution inequalities [Mau91, LW08], Latała-Oleskiewicz inequalities [Bec89, LO00, BR03, BCR06]... Several surveys and monographs are now available on this topic (see for instance [Led01], [ABC⁺00] or [Vil03, Vil08]). This large variety of tools and points of view raises the following natural question: is one of these functional inequalities equivalent to say (1.1) ?

In this paper, one shows with a certain generality that Talagrand's transportation-cost inequalities are equivalent to dimension free concentration of measure. Let us give a flavor of our results in the Gaussian case. Let us first define the optimal quadratic transportation-cost on $P(\mathbf{R}^d)$ (the set of probability measures on \mathbf{R}^d). For all ν and μ in $P(\mathbf{R}^d)$, one defines

$$(1.2) \quad \mathcal{T}_2(\nu, \mu) = \inf_{\pi} \int |x - y|_2^2 d\pi(x, y),$$

where π describes the set $P(\nu, \mu)$ of probability measures on $\mathbf{R}^d \times \mathbf{R}^d$ having ν and μ for marginal distributions. One says that μ verifies the inequality $\mathbf{T}_2(C)$, if

$$(1.3) \quad \forall \nu \in P(\mathbf{R}^d), \quad \mathcal{T}_2(\nu, \mu) \leq C H(\nu | \mu),$$

where $H(\nu | \mu)$ is the relative entropy of ν with respect to μ defined by $H(\nu | \mu) = \int \log \left(\frac{d\nu}{d\mu} \right) d\nu$ if ν is absolutely continuous with respect to μ and $+\infty$ otherwise. The idea of controlling an optimal transportation-cost by the relative entropy to obtain concentration first appeared in Marton's works [Mar86, Mar96]. The inequality \mathbf{T}_2 was then introduced by Talagrand in [Tal96], where it was proved to be fulfilled by Gaussian probability measures. In particular, if $\mu = \gamma$ is the standard Gaussian measure on \mathbf{R} , then the inequality (1.3) holds true with the sharp constant $C = 2$.

The following theorem is the main result of this work.

Theorem 1.4. *Let μ be a probability measure on \mathbf{R}^d and $a > 0$; the following propositions are equivalent:*

- (1) *There are $r_o, b \geq 0$ such that for all n the probability μ^n verifies (1.1),*
- (2) *The probability measure μ verifies $\mathbf{T}_2(1/a)$.*

The example of the standard Gaussian measure γ proves that the relation between the constants is sharp. The fact that (2) implies (1) is well known and follows from a nice and general argument of Marton. The proof of the converse is surprisingly easy and relies on a very simple Large Deviations argument. We think that this new result confirms the relevance of the Large Deviations point of view for functional inequalities initiated by Léonard and the author in [GL07] and pursued in [GLWY07] by Guillin, Léonard, Wu and Yiao. Moreover Theorem 1.4 turns out to be a quite powerful tool. For example, the famous result by Otto and Villani stating that the Logarithmic-Sobolev inequality (**LSI**) implies the \mathbf{T}_2 inequality (see [OV00, Theorem 1]) is a direct consequence of Theorem 1.4 (see Theorem 3.6 and its proof).

The paper is organized as follows. In section 2, we give a brief account on the Large Deviations phenomenon entering the game. In section 3, we focus on the case of Gaussian concentration and prove Theorem 1.4 in an abstract Polish setting. In section 4, one considers non-Gaussian concentrations and relates them to other transportation-cost inequalities.

In section 5, we prove the equivalence between Poincaré inequality and dimension free concentration of the exponential type. The section 6, is devoted to remarks concerning known criteria for transportation-cost inequalities.

Acknowledgements: I want to warmly acknowledge Patrick Cattiaux, Arnaud Guillin, Michel Ledoux and Paul-Marie Samson for their valuable comments about this work.

2. SOME PRELIMINARIES ON LARGE DEVIATIONS

In this section, we consider the following abstract framework: (\mathcal{X}, ρ) is a Polish space and the set of probability measures on \mathcal{X} is denoted by $\mathbb{P}(\mathcal{X})$. Let μ be a probability measure on \mathcal{X} and $(X_i)_i$ an i.i.d sequence of random variables with law μ defined on some probability space (Ω, \mathbb{P}) . The empirical measure L_n is defined for all integer n by

$$L_n = \frac{1}{n} \sum_{i=1}^n \delta_{X_i},$$

where δ_x stands for the Dirac mass at point x .

According to Varadarajan's Theorem (see for instance [Dud89, Theorem 11.4.1]), with probability 1 the sequence $(L_n)_n$ converges to μ in $\mathbb{P}(\mathcal{X})$ for the topology of weak convergence, this means that there is a measurable subset \mathcal{N} of Ω with $\mathbb{P}(\mathcal{N}) = 0$ such that for all $\omega \notin \mathcal{N}$,

$$\int f dL_n(\omega) \xrightarrow{n \rightarrow +\infty} \int f d\mu,$$

for all bounded continuous f on \mathcal{X} .

The topology of weak convergence can be metrized by various metrics. Here, one will consider the Wasserstein metrics. Let $p \geq 1$ and define

$$\mathbb{P}_p(\mathcal{X}) = \left\{ \nu \in \mathbb{P}(\mathcal{X}) \text{ s.t. } \int \rho(x_o, x)^p d\nu(x) < +\infty, \text{ for some } x_o \in \mathcal{X} \right\}.$$

For all probability measures $\nu_1, \nu_2 \in \mathbb{P}_p(\mathcal{X})$, define

$$\mathcal{I}_p(\nu_1, \nu_2) = \inf_{\pi} \int \rho(x, y)^p d\pi(x, y) \quad \text{and} \quad W_p(\nu_1, \nu_2) = (\mathcal{I}_p(\nu_1, \nu_2))^{1/p}$$

where π describes the set $P(\nu_1, \nu_2)$ of couplings of ν_1 and ν_2 .

According to e.g [Vil03, Theorems 7.3 and 7.12], W_p is a metric on $\mathbb{P}_p(\mathcal{X})$ and for every sequence μ_n in $\mathbb{P}_p(\mathcal{X})$, $W_p(\mu_n, \mu) \rightarrow 0$ if and only if μ_n converges to μ for the weak topology and $\int \rho(x_o, x)^p d\mu_n \rightarrow \int \rho(x_o, x)^p d\mu$, for some (and thus any) $x_o \in \mathcal{X}$.

From these considerations, one can conclude that if $\mu \in \mathbb{P}_p(\mathcal{X})$, then $W_p(L_n, \mu) \rightarrow 0$ with probability one, and in particular, $\mathbb{P}(W_p(L_n, \mu) \geq t) \rightarrow 0$ when $n \rightarrow +\infty$, for all $t > 0$. Moreover, supposing that $\mu \in \mathbb{P}_p(\mathcal{X})$, with $p > 1$, it is easy to check that the sequence $W_p(L_n, \mu)$ is bounded in $\mathbb{L}_p(\Omega, \mathbb{P})$, thus it is uniformly integrable and consequently $\mathbb{E}[W_p(L_n, \mu)] \rightarrow 0$. This is summarized in the following proposition:

Proposition 2.1. *If $\mu \in \mathbb{P}_p(\mathcal{X})$, then the sequence $W_p(L_n, \mu) \rightarrow 0$ almost surely (and thus in probability) and if $p > 1$, then the convergence is in \mathbb{L}_1 : $\mathbb{E}[W_p(L_n, \mu)] \rightarrow 0$.*

On the other hand, Sanov's Theorem (see e.g [DZ98, Theorem 6.2.10]) says that for all good sets A , $\mathbb{P}(L_n \in A)$ behaves like $e^{-nH(A|\mu)}$ when n is large, where $H(A|\mu)$ stands for the infimum of $H(\cdot|\mu)$ on A . So, when A does not contain μ , $H(A|\mu) > 0$ and this probability tends to 0 exponentially fast. With this in mind, one can expect that $\mathbb{P}(W_p(L_n, \mu) > t)$ behaves like $e^{-nH(t)}$, where $H(t) = \inf \{H(\nu|\mu) : \nu \text{ s.t. } W_p(\nu, \mu) > t\}$. The following result validates partially this heuristic, stating that $\mathbb{P}(W_p(L_n, \mu) > t)$ tends to 0 not faster than $e^{-nH(t)}$.

Theorem 2.2. *If $\mu \in \mathbb{P}_p(\mathcal{X})$, then for all $t > 0$,*

$$\liminf_{n \rightarrow +\infty} \frac{1}{n} \log \mathbb{P}(W_p(L_n, \mu) > t) \geq -\inf \{H(\nu|\mu) : \nu \in \mathbb{P}_p(\mathcal{X}) \text{ s.t. } W_p(\nu, \mu) > t\}.$$

For the sake of completeness, an elementary proof of this result will be displayed in the appendix. As in [GL07], the use of this Large Deviations technique will be the key step in the proof of Theorem 1.4.

3. THE GAUSSIAN CASE

3.1. An abstract version of Theorem 1.4. As in the preceding section, (\mathcal{X}, ρ) will be a Polish space. The product space \mathcal{X}^n will be equipped with the following metric:

$$\rho_2^n(x, y) = \left[\sum_{i=1}^n \rho(x^i, y^i)^2 \right]^{1/2}$$

(here $x = (x^1, x^2, \dots, x^n)$ with $x^i \in \mathcal{X}$ for all i).

In the general case, one says that a probability measure μ on (\mathcal{X}, ρ) verifies the dimension free Gaussian concentration property, if there are $r_o, a, b \geq 0$ such that for all n the probability μ^n verifies

$$(3.1) \quad \forall r \geq r_o, \quad \mu^n(A^r) \geq 1 - be^{-a(r-r_o)^2},$$

for all measurable $A \subset \mathcal{X}^n$ such that $\mu_n(A) \geq 1/2$, where A^r denotes the r -enlargement of A defined by

$$A^r = \{x \in \mathcal{X}^n \text{ such that there is } \bar{x} \in A \text{ with } \rho_2^n(x, \bar{x}) \leq r\}$$

Of course, when $\mathcal{X} = \mathbf{R}^d$ is equipped with its Euclidean metric one has $A^r = A + rB_2$ and one recovers the definition (1.1).

Theorem 3.2. *Let $\mu \in \mathbb{P}_2(\mathcal{X})$ and $a > 0$; the following propositions are equivalent:*

- (1) *There are $r_o, b \geq 0$ such that for all n the probability μ^n verifies (3.1),*
- (2) *The probability μ verifies $\mathbf{T}_2(1/a)$.*

Let us recall the definition of the \mathbf{T}_1 transportation-cost inequality. One says that a probability measure μ on \mathcal{X} verifies $\mathbf{T}_1(C)$, if

$$\forall \nu \in \mathbb{P}(\mathcal{X}), \quad W_1(\nu, \mu) \leq \sqrt{CH(\nu|\mu)}.$$

According to Jensen's inequality, the inequality $\mathbf{T}_1(C)$ is weaker than $\mathbf{T}_2(C)$; it was completely characterized in terms of square exponential integrability in [DGW04].

The proof of the following well known result makes use of the so called Marton's argument.

Proposition 3.3 (Marton). *If μ verifies $\mathbf{T}_1(C)$, then for all measurable subset A of \mathcal{X} , such that $\mu(A) \geq 1/2$*

$$\forall r \geq r_o, \quad \mu(A^r) \geq 1 - e^{-C^{-1}(r-r_o)^2},$$

where $r_o = \sqrt{C \log(2)}$.

Proof. Consider a subset A of \mathcal{X} and define $d\mu_A = \mathbb{1}_A d\mu(x)/\mu(A)$. Let $B = \mathcal{X} \setminus A^r$ and define μ_B accordingly. Since the distance between two points of A and B is always more than r , one has $W_1(\mu_A, \mu_B) \geq r$. The triangle inequality and the transportation-cost inequality $\mathbf{T}_1(C)$ yield

$$\begin{aligned} r \leq W_1(\mu_A, \mu_B) &\leq W_1(\mu_A, \mu) + W_1(\mu_B, \mu) \\ &\leq \sqrt{C \mathbb{H}(\mu_A | \mu)} + \sqrt{C \mathbb{H}(\mu_B | \mu)} \\ &= \sqrt{C \log(1/\mu(A))} + \sqrt{C \log(1/\mu(B))}. \end{aligned}$$

Rearranging terms gives the result. \square

Proof of Theorem 3.2. Let us show that (2) implies (1). The main point is that \mathbf{T}_2 tensorizes ; this means that if μ verifies $\mathbf{T}_2(1/a)$ then μ^n verifies $\mathbf{T}_2(1/a)$ on the space \mathcal{X}^n equipped with ρ_2^n . The reader can find a general result concerning tensorization properties of transportation-cost inequalities in [GL07, Theorem 5]. Jensen's inequality implies that $W_1^2 \leq \mathcal{T}_2$ and consequently μ^n verifies $\mathbf{T}_1(1/a)$ (on \mathcal{X}^n equipped with ρ_2^n) for all n . Applying Proposition 3.3 to μ^n gives (1.1) with $r_o = \sqrt{\log(2)/a}$, $b = 1$ and a .

Let us show that (1) implies (2). For every integer n , and $x \in \mathcal{X}^n$, define $L_n^x = n^{-1} \sum_{i=1}^n \delta_{x^i}$. The map $x \mapsto W_2(L_n^x, \mu)$ is $1/\sqrt{n}$ -Lipschitz. Indeed, if $x = (x^1, \dots, x^n)$ and $y = (y^1, \dots, y^n)$ are in \mathcal{X}^n , then thanks to the triangle inequality,

$$|W_2(L_n^x, \mu) - W_2(L_n^y, \mu)| \leq W_2(L_n^x, L_n^y).$$

According to the convexity property of $\mathcal{T}_2(\cdot, \cdot)$ (see e.g [Vil08, Theorem 4.8]), one has

$$\mathcal{T}_2(L_n^x, L_n^y) \leq \frac{1}{n} \sum_{i=1}^n \mathcal{T}_2(\delta_{x^i}, \delta_{y^i}) = \frac{1}{n} \sum_{i=1}^n \rho(x^i, y^i)^2 = \frac{1}{n} \rho_2^n(x, y)^2,$$

which proves the claim.

Now, let $(X_i)_i$ be an i.i.d sequence of law μ and let L_n be its empirical measure. Let m_n be the median of $W_2(L_n, \mu)$ and define $A = \{x : W_2(L_n^x, \mu) \leq m_n\}$. Then $\mu^n(A) \geq 1/2$ and it is easy to show that $A^r \subset \{x : W_2(L_n^x, \mu) \leq m_n + r/\sqrt{n}\}$. Applying (3.1) to A gives

$$\forall r \geq r_o, \quad \mathbb{P}(W_2(L_n, \mu) > m_n + r/\sqrt{n}) \leq b \exp(-a(r - r_o)^2).$$

Equivalently, as soon as $\sqrt{n}(u - m_n) \geq r_o$, one has

$$\mathbb{P}(W_2(L_n, \mu) > u) \leq b \exp(-a(\sqrt{n}(u - m_n) - r_o)^2).$$

Now, since $W_2(L_n, \mu)$ converges to 0 in probability (see Proposition 2.1), the sequence m_n goes to 0 when n goes to $+\infty$. Consequently,

$$\forall u > 0, \quad \limsup_{n \rightarrow +\infty} \frac{1}{n} \log \mathbb{P}(W_2(L_n, \mu) > u) \leq -au^2.$$

The final step is given by Large Deviations. According to Theorem 2.2,

$$\liminf_{n \rightarrow +\infty} \frac{1}{n} \log \mathbb{P}(W_2(L_n, \mu) > u) \geq -\inf \{H(\nu | \mu) : \nu \in \mathbf{P}_2(\mathcal{X}) \text{ s.t. } W_2(\nu, \mu) > u\}.$$

This together with the preceding inequality yields

$$\inf \{H(\nu | \mu) : \nu \in \mathbf{P}_2(\mathcal{X}) \text{ s.t. } W_2(\nu, \mu) > u\} \geq au^2$$

or in other words,

$$aW_2(\nu, \mu)^2 \leq H(\nu | \mu),$$

and this achieves the proof. \square

Let us make a remark on the proof. The careful reader will notice that the second part of the proof applies if one replaces $W_2(\cdot, \mu)$ by any application $\Phi : \mathbf{P}(\mathcal{X}) \rightarrow \mathbf{R}^+$ which is continuous with respect to the weak topology, verifies $\Phi(\mu) = 0$, and is such that for all integer n , the map $\mathcal{X}^n \rightarrow \mathbf{R}^+ : x \mapsto \Phi(L_n^x)$ is $1/\sqrt{n}$ -Lipschitz for the metric ρ_2^n on \mathcal{X}^n . For such an application Φ , one can show, with exactly the same proof, that the dimension free Gaussian concentration property (3.1) implies that $a\Phi^2(\nu) \leq H(\nu | \mu)$, for all ν and it could be that this new inequality is stronger than \mathbf{T}_2 . Actually, it is not the case. Namely, it is an easy exercise to show that if Φ verifies the above listed properties, then $\Phi(\nu) \leq W_2(\nu, \mu)$, for all ν , and so the choice $\Phi = W_2$ is optimal.

3.2. Otto and Villani's Theorem. Our aim is now to recover and extend a theorem by Otto and Villani stating that the Logarithmic-Sobolev inequality is stronger than Talagrand's \mathbf{T}_2 inequality.

Let us recall that a probability measure μ on \mathcal{X} verifies the Logarithmic-Sobolev inequality with constant $C > 0$ (**LSI**(C) for short) if

$$\text{Ent}_\mu(f^2) \leq C \int |\nabla f|^2 d\mu,$$

for all locally Lipschitz f , where the entropy functional is defined by

$$\text{Ent}_\mu(f) = \int f \log f d\mu - \int f d\mu \log \left(\int f d\mu \right), \quad f \geq 0,$$

and the length of the gradient is defined by

$$(3.4) \quad |\nabla f|(x) = \limsup_{y \rightarrow x} \frac{|f(x) - f(y)|}{\rho(x, y)}$$

(when x is an isolated point, we put $|\nabla f|(x) = 0$).

In [OV00, Theorem 1], Otto and Villani proved that if a probability measure μ on a Riemannian manifold M , satisfies the inequality **LSI**(C) then it also satisfies the inequality $\mathbf{T}_2(C)$. Their proof was rather involved and uses partial differential equations, optimal transportation results, and fine observations relating relative entropy and Fisher information. A simpler proof, as well as a generalization, was proposed by Bobkov, Gentil and Ledoux in [BGL01]. It makes use of the dual formulation of transportation-cost inequalities discovered by Bobkov and Götze in [BG99] and relies on hypercontractivity properties of the Hamilton-Jacobi semi group put in light in the same paper [BGL01]. Otto and Villani's result was successfully generalized by Wang on paths spaces in [Wan04]. More recently, Lott and Villani showed

that implication $\mathbf{LSI} \Rightarrow \mathbf{T}_2$ remains true on a length space provided the measure μ satisfies a doubling condition and a local Poincaré inequality (see [LV07, Theorem 1.8]).

The converse implication $\mathbf{T}_2 \Rightarrow \mathbf{LSI}$ is sometimes true. For example, it is the case when μ is a Log-concave probability measure (see [OV00, Corollary 3.1]). However, in the general case, \mathbf{T}_2 and \mathbf{LSI} are not equivalent. In [CG06], Cattiaux and Guillin give an example of a probability measure verifying \mathbf{T}_2 and not \mathbf{LSI} .

With Theorem 3.2 in hand, one could think that the implication $\mathbf{LSI} \Rightarrow \mathbf{T}_2$ is now completely straightforward. Namely, it is well known that the Logarithmic-Sobolev inequality implies dimension free Gaussian concentration ; since this latter is equivalent to Talagrand's \mathbf{T}_2 inequality it should be clear that the Logarithmic-Sobolev inequality implies \mathbf{T}_2 . It is effectively the case on reasonable spaces such as \mathbf{R}^d but in the general case, a subtle technical question was not taken into account in the preceding line of reasoning. Namely, if μ verifies the $\mathbf{LSI}(C)$ inequality, then according to the additive property of the Logarithmic-Sobolev inequality, one can conclude that the product measure μ^n verifies

$$(3.5) \quad \text{Ent}_{\mu^n}(f^2) \leq C \int \sum_{i=1}^n |\nabla_i f|^2(x) d\mu^n(x),$$

where the length of the 'partial derivative' $|\nabla_i f|$ is defined according to (3.4). The problem is that, in this very abstract setting, $\sum_i |\nabla_i f|^2(x)$ and $|\nabla f|^2(x)$ (computed with respect to ρ_2^n) may be different. The tensorized Logarithmic-Sobolev inequality will yield concentration inequalities for functions such that $\sum_i |\nabla_i f|^2(x) \leq 1$ μ^n -almost everywhere and this class of functions may not contain 1-Lipschitz functions for the ρ_2^n metric. Nevertheless, this difficulty can be circumvented as shown in the following theorems.

Theorem 3.6. *Let μ be a probability measure on \mathcal{X} and suppose that for all integer n the function F_n defined on \mathcal{X}^n by $F_n(x) = W_2(L_n^x, \mu)$ verifies*

$$(3.7) \quad \sum_{i=1}^n |\nabla_i F_n|^2(x) \leq 1/n, \text{ for } \mu^n \text{ almost every } x \in \mathcal{X}^n.$$

If μ verifies the inequality $\mathbf{LSI}(C)$, then μ verifies the inequality $\mathbf{T}_2(C)$.

We have seen during the proof of Theorem 3.2 that the functions F_n are $1/\sqrt{n}$ -Lipschitz for the metric ρ_2^n . Suppose that $\mathcal{X} = \mathbf{R}^d$ or a Riemannian manifold M , then according to Rademacher's Theorem, F_n is almost everywhere differentiable on $(\mathbf{R}^d)^n$ (resp. M^n) with respect to the Lebesgue measure. It is thus easy to show that condition (3.7) is fulfilled when μ is absolutely continuous with respect to Lebesgue measure. This permits us to recover Otto and Villani's result as stated in [OV00].

Proof. As we said above the product measure μ^n verifies the inequality (3.5). Apply this inequality to $f = e^{\frac{s}{2}F_n}$, with $s \in \mathbf{R}^+$. It is easy to show that $|\nabla_i e^{\frac{s}{2}F_n}| = \frac{s}{2} e^{\frac{s}{2}F_n} |\nabla_i F_n|$, thus, using condition (3.7), one sees that the right hand side of (3.5) is less than $C \frac{s^2}{4n} \int e^{sF_n} d\mu^n$. Letting $Z(s) = \int e^{sF_n} d\mu^n$, one gets the differential inequality:

$$\frac{Z'(s)}{sZ(s)} - \frac{\log Z(s)}{s^2} \leq \frac{C}{4n}.$$

Integrating this yields:

$$\forall s \in \mathbf{R}^+, \quad Z(s) = \int e^{sF_n} d\mu^n \leq e^{s \int F_n d\mu^n + \frac{Cs^2}{4n}}.$$

This implies that

$$\mathbb{P}(W_2(L_n, \mu) \geq t + \mathbb{E}[W_2(L_n, \mu)]) \leq e^{-nt^2/C}.$$

According to Proposition 2.1, $\mathbb{E}[W_2(L_n, \mu)] \rightarrow 0$. Arguing exactly as in proof of Theorem 3.2, one concludes that the inequality $\mathbf{T}_2(C)$ holds. \square

With an extra assumption on the support of μ , one shows in the following theorem that the implication $\mathbf{LSI} \Rightarrow \mathbf{T}_2$ is true with a relaxed constant:

Theorem 3.8. *Let μ be a probability measure on \mathcal{X} such that*

$$(3.9) \quad \forall k \in \mathbf{R}, \quad \forall u \neq v \in \mathcal{X}, \quad \mu \{x \in \mathcal{X} \text{ s.t. } \rho^2(x, u) - \rho^2(x, v) = k\} = 0.$$

If μ verifies the inequality $\mathbf{LSI}(C)$ then μ satisfies $\mathbf{T}(2C)$.

The condition (3.9) first appeared in a paper by Cuesta-Albertos and Tuero-Díaz on optimal transportation. Roughly speaking, this assumption guaranties the uniqueness of the Monge-Kantorovich Problem of transporting μ on a probability measure ν with finite support (see [CATD93, Theorem 3]). For μ on \mathbf{R}^d , the condition (3.9) amounts to say that μ does not charge hyperplanes. We think that working better it would be possible to obtain the right constant C instead of $2C$.

Proof. We will use a sort of symmetrization argument. First observe that the probability measure $\mu^n \times \mu^n$ verifies the following Logarithmic-Sobolev inequality:

$$\text{Ent}_{\mu^n \times \mu^n}(f^2) \leq C \sum_{i=1}^n |\nabla_{i,1} f|^2(x, y) + |\nabla_{i,2} f|^2(x, y) d\mu^n(x) d\mu^n(y)$$

for all $f : \mathcal{X}^n \times \mathcal{X}^n \rightarrow \mathbf{R} : (x, y) \mapsto f(x, y)$, where $|\nabla_{i,1} f|$ (resp. $|\nabla_{i,2} f|$) denotes the length of the gradient with respect to the x^i -coordinate (resp. the y^i -coordinate).

Define $G_n(x, y) = W_2(L_n^x, L_n^y)$ for all $x, y \in \mathcal{X}^n$. One wants to apply the tensorized Logarithmic-Sobolev inequality to the function G_n . To do so one needs to compute the length of its partial derivatives. Let us explain how to compute $L = |\nabla_{1,1} G_n|(a, b)$, for instance. For every $z \in \mathcal{X}$, let $za = (z, a^2, \dots, a^n)$; obviously,

$$L = \limsup_{z \rightarrow a^1} \frac{|W_2(L_n^{za}, L_n^b) - W_2(L_n^a, L_n^b)|}{\rho(z, a^1)} = \frac{1}{2W_2(L_n^a, L_n^b)} \limsup_{z \rightarrow a^1} \frac{|\mathcal{T}_2(L_n^{za}, L_n^b) - \mathcal{T}_2(L_n^a, L_n^b)|}{\rho(z, a^1)}.$$

According to the condition (3.9), the probability measure μ is diffuse; so the probability of points $x \in \mathcal{X}^n$ having distinct coordinates is one. So, one can suppose without restriction that the coordinates of a (resp. b) are all different. If z is sufficiently close to a^1 , the coordinates of za are all distinct too. According to e.g [Vil03, Example p. 5], the optimal transport of L_n^a on L_n^b is given by a permutation, this means that there is at least one permutation σ of $\{1, \dots, n\}$ such that

$$\mathcal{T}_2(L_n^a, L_n^b) = n^{-1} \sum_{i=1}^n \rho(a^i, b^{\sigma(i)})^2.$$

Let us denote by S the set of these permutations and define accordingly the set S_z of permutations realizing the optimal transport of L_n^{za} on L_n^b .

Without loss of generality, one can suppose that S is a singleton. Indeed, let σ and $\tilde{\sigma}$ be two distinct permutations and consider

$$H_{\sigma, \tilde{\sigma}} = \left\{ x \in \mathcal{X}^n : \sum_{i=1}^n \rho(x^i, b^{\sigma(i)})^2 = \sum_{i=1}^n \rho(x^i, b^{\tilde{\sigma}(i)})^2 \right\}.$$

Applying Fubini's Theorem together with the condition (3.9), one gets easily that $\mu^n(H_{\sigma, \tilde{\sigma}}) = 0$. This readily proves the claim. In the sequel we will set $S = \{\sigma^*\}$.

Now we claim that if z is sufficiently close to a^1 , then $S_z = \{\sigma^*\}$. Indeed, let

$$\varepsilon_o = \min_{\sigma \neq \sigma^*} \left\{ n^{-1} \sum_{i=1}^n \rho(a^i, b^{\sigma(i)})^2 - \mathcal{T}_2(L_n^a, L_n^b) \right\} > 0;$$

then there is a neighborhood V of a^1 such that for all $z \in V$, one has

$$\left| \mathcal{T}_2(L_n^{za}, L_n^b) - \mathcal{T}_2(L_n^a, L_n^b) \right| \leq \varepsilon_o/3$$

and for all permutation σ ,

$$\left| n^{-1} \sum_{i=1}^n \rho((za)^i, b^{\sigma(i)})^2 - n^{-1} \sum_{i=1}^n \rho(a^i, b^{\sigma(i)})^2 \right| \leq \varepsilon_o/3.$$

Now, if $z \in V$ and $\sigma \in S_z$, one has

$$n^{-1} \sum_{i=1}^n \rho(a^i, b^{\sigma(i)})^2 \leq n^{-1} \sum_{i=1}^n \rho((za)^i, b^{\sigma(i)})^2 + \varepsilon_o/3 = \mathcal{T}_2(L_n^{za}, L_n^b) + \varepsilon_o/3 \leq \mathcal{T}_2(L_n^a, L_n^b) + 2\varepsilon_o/3.$$

By the definition of the number ε_o , one concludes that $\sigma = \sigma^*$, which proves the claim.

Now, if $z \in V$, then

$$\frac{|\mathcal{T}_2(L_n^{za}, L_n^b) - \mathcal{T}_2(L_n^a, L_n^b)|}{\rho(z, a^1)} = \frac{|\rho(z, b^{\sigma^*(1)})^2 - \rho(a^1, b^{\sigma^*(1)})^2|}{n\rho(z, a^1)} \leq \frac{1}{n} \left(\rho(z, b^{\sigma^*(1)}) + \rho(a^1, b^{\sigma^*(1)}) \right).$$

So letting $z \rightarrow a^1$, yields $L \leq \frac{\rho(a^1, b^{\sigma^*(1)})}{nW_2(L_n^a, L_n^b)}$.

Doing the same for the other partial derivatives yields:

$$\sum_{i=1}^n |\nabla_{i,1} G_n|^2(a, b) \leq \frac{\sum_{i=1}^n \rho(a^i, b^{\sigma^*(i)})^2}{n^2 \mathcal{T}_2(L_n^a, L_n^b)} = \frac{1}{n}.$$

Finally,

$$\sum_{i=1}^n |\nabla_{i,1} G_n|^2(a, b) + |\nabla_{i,2} G_n|^2(a, b) \leq \frac{2}{n},$$

for $\mu^n \times \mu^n$ almost every $a, b \in \mathcal{X}^n \times \mathcal{X}^n$.

Now reasoning as in the proof of Theorem 3.6, one concludes that

$$\mathbb{P}(W_2(L_n^X, L_n^Y) > t + \mathbb{E}[W_2(L_n^X, L_n^Y)]) \leq e^{-nt^2/(2C)}.$$

On the other hand, an easy adaptation of Proposition 2.2 yields

$$\liminf_{n \rightarrow +\infty} \frac{1}{n} \log \mathbb{P} (W_2(L_n^X, L_n^Y) > t + \mathbb{E} [W_2(L_n^X, L_n^Y)]) \geq \\ - \inf \{ \mathbf{H}(\nu_1 | \mu) + \mathbf{H}(\nu_2 | \mu) : \nu_1, \nu_2 \in \mathbf{P}_2(X) \text{ s.t. } W_2(\nu_1, \nu_2) > t \}.$$

From this follows as before that

$$\mathcal{T}_2(\nu_1, \nu_2) \leq 2C (\mathbf{H}(\nu_1 | \mu) + \mathbf{H}(\nu_2 | \mu))$$

holds for all probability measures ν_1, ν_2 belonging to $\mathbf{P}_2(X)$. Taking $\nu_2 = \mu$ gives the inequality $\mathbf{T}_2(2C)$. \square

Our next goal is to recover and extend a result of Lott and Villani. Following [LV07], one says that a probability measure μ on \mathcal{X} verifies the inequality $\mathbf{LSI}^+(C)$ if

$$\text{Ent}_\mu(f^2) \leq C \int |\nabla^- f|^2 d\mu,$$

holds true for all locally Lipschitz f , where the subgradient norm $|\nabla^- f|$ is defined by

$$|\nabla^- f|(x) = \limsup_{y \rightarrow x} \frac{[f(y) - f(x)]_+}{\rho(x, y)},$$

with $[a]_+ = \max(a, 0)$. Since $|\nabla^- f| \leq |\nabla f|$, the inequality \mathbf{LSI}^+ is stronger than \mathbf{LSI} ; more precisely, $\mathbf{LSI}^+(C) \Rightarrow \mathbf{LSI}(C)$.

Theorem 3.10. *If μ verifies the inequality $\mathbf{LSI}^+(C)$, then μ verifies $\mathbf{T}_2(C)$.*

This result was first obtained by Lott and Villani using the Hamilton-Jacobi method. This approach forced them to make many assumptions on \mathcal{X} and μ . In particular, in [LV07, Theorem 1.8] \mathcal{X} was supposed to be a compact length space and a doubling condition was imposed on μ . The result above shows that the implication $\mathbf{LSI}^+ \Rightarrow \mathbf{T}_2$ is in fact always true. The following proof uses an argument which I learned from Paul-Marie Samson.

Proof. The inequality \mathbf{LSI}^+ tensorizes, so μ^n verifies

$$\text{Ent}_{\mu^n}(f^2) \leq C \int \sum_{i=1}^n |\nabla_i^- f|^2 d\mu^n.$$

Take $f = e^{\frac{s}{2} F_n}$, $s \in \mathbf{R}^+$ with $F_n(x) = W_2(L_n^x, \mu)$. Once again, it is easy to check that $|\nabla_i^- e^{\frac{s}{2} F_n}| = \frac{s}{2} e^{\frac{s}{2} F_n} |\nabla_i^- F_n|$ (note that the function $x \mapsto e^{sx}$ is non decreasing). Reasoning as in the proof of Theorem 3.6, it is enough to show that $\sum_i |\nabla_i^- F_n|^2(x) \leq 1/n$ for μ^n -almost all $x \in \mathcal{X}^n$. Let us show how to compute $|\nabla_1^- F_n|$. Let $z \in X$, $a = (a^1, \dots, a^n) \in \mathcal{X}^n$ and set $za = (z, a^2, \dots, a^n)$, then

$$|\nabla_1^- F_n|(a) = \frac{1}{2F_n(a)} \limsup_{z \rightarrow a^1} \frac{[\mathcal{T}_2(L_n^{za}, \mu) - \mathcal{T}_2(L_n^a, \mu)]_+}{\rho(z, a^1)}.$$

Let $\pi \in P(L_n^a, \mu)$ be an optimal coupling; it is not difficult to see that one can write $\pi(dx, dy) = p(x, dy) L_n^a(dx)$, where $p(a^i, dy) = \nu_i(dy)$ with ν_1, \dots, ν_n probability measures

on \mathcal{X} such that $n^{-1}(\nu_1 + \dots + \nu_n) = \mu$. Let \tilde{p} be defined as p with z in place of a^1 ; then $\tilde{\pi} = \tilde{p}(x, dy)L_n^{za}(dy)$ belongs to $P(L_n^{za}, \mu)$ (but is not necessary optimal). One has

$$\begin{aligned} \mathcal{T}_2(L_n^{za}, \mu) - \mathcal{T}_2(L_n^a, \mu) &\leq \int \rho(x, y)^2 d\tilde{\pi}(x, y) - \int \rho(x, y)^2 d\pi(x, y) \\ &= \frac{1}{n} \sum_{i=1}^n \int \rho((za)^i, y)^2 d\nu_i(y) - \frac{1}{n} \sum_{i=1}^n \int \rho(a^i, y)^2 d\nu_i(y) \\ &= \frac{1}{n} \int \rho(z, y)^2 - \rho(a^1, y)^2 d\nu_1(y) \\ &\leq \frac{1}{n} \rho(z, a^1) \int \rho(z, y) + \rho(a^1, y) d\nu_1(y). \end{aligned}$$

Since the function $x \mapsto [x]_+$ is non decreasing, one has

$$\frac{[\mathcal{T}_2(L_n^{za}, \mu) - \mathcal{T}_2(L_n^a, \mu)]_+}{\rho(z, a^1)} \leq \frac{1}{n} \int \rho(z, y) + \rho(a^1, y) d\nu_1(y).$$

Letting $z \rightarrow a^1$ yields $|\nabla_1^- F_n(a)|^2 \leq \frac{\int \rho(a^1, y)^2 d\nu_1(y)}{n^2 \mathcal{T}_2(L_n^a, \mu)}$. Doing the same computations for the

other derivatives (with the same optimal coupling π), one gets $|\nabla_i^- F_n(a)|^2 \leq \frac{\int \rho(a^i, y)^2 d\nu_i(y)}{n^2 \mathcal{T}_2(L_n^a, \mu)}$.

Summing these inequalities gives $\sum_i |\nabla_i^- F_n|^2(a) \leq 1/n$ for all $a \in \mathcal{X}^n$, which achieves the proof. \square

4. GENERALIZATIONS TO NON GAUSSIAN CONCENTRATION

4.1. A first generalization for super-Gaussian concentration. The following theorem can be established with exactly the same proof as Theorem 1.4. We leave the proof to the reader.

Theorem 4.1. *Let μ be a probability measure on \mathcal{X} , $p \geq 2$ and $a > 0$. The following propositions are equivalent:*

- (1) *There are $r_o, b \geq 0$ such that for every n the probability measure μ^n verifies for all A subset of \mathcal{X}^n with $\mu^n(A) \geq 1/2$,*

$$(4.2) \quad \forall r \geq r_o, \quad \mu^n(A^r) \geq 1 - be^{-a(r-r_o)^p},$$

where the enlargement A^r is performed with respect to the metric ρ_p^n on \mathcal{X}^n defined by

$$\forall x, y \in \mathcal{X}^n, \quad \rho_p^n(x, y) = \left[\sum_{i=1}^n \rho(x^i, y^i)^p \right]^{1/p}.$$

- (2) *The probability measure μ verifies the following transportation cost inequality:*

$$\forall \nu \in \mathbb{P}_p(\mathcal{X}), \quad \mathcal{T}_p(\nu, \mu) \leq a^{-1} \mathbb{H}(\nu | \mu).$$

4.2. Talagrand's two level concentration inequalities. Our approach is sufficiently flexible to be adapted to various forms of concentration. We do not want to enter in too general (and maybe useless) generalizations. We will content to give one more example not covered by the preceding theorems. We want to find the transportation-cost inequality equivalent to Talagrand's two level concentration inequalities which are well adapted to concentration rates between exponential and Gaussian.

Let us say that a probability measure μ on \mathbf{R}^d satisfies a two level dimension free concentration inequality of order $p \in [1, 2]$ if there are two non-negative constants a and b such that for every n the inequality

$$(4.3) \quad \forall r \geq 0, \quad \mu^n \left(A + \sqrt{r}B_2 + \sqrt[p]{r}B_p \right) \geq 1 - be^{-ar},$$

holds for all measurable subset A of $(\mathbf{R}^d)^n$ such that $\mu^n(A) \geq 1/2$, where B_2 and B_p are the standard unit balls of $(\mathbf{R}^d)^n$. Inequalities of this form appear in [Tal94], where it is proved that the measure $d\mu_p(x) = Z_p^{-1}e^{-|x|^p}$, $p \geq 1$ verifies such a bound.

The transportation-cost adapted to this kind of concentration is defined for all probability measures ν_1, ν_2 on $(\mathbf{R}^d)^n$ by

$$\mathcal{T}_{2,p}(\nu, \mu) = \inf_{\pi \in P(\nu_1, \nu_2)} \int \sum_{i=1}^n \sum_{j=1}^d \alpha_p(x_j^i - y_j^i) d\pi(x, y)$$

where $\alpha_p(u) = \min(|u|^2, |u|^p)$ (here $x = (x^1, \dots, x^n)$ with $x^i \in \mathbf{R}^d$ for all i).

Theorem 4.4. *Let μ be a probability measure on \mathbf{R}^d and $p \in [1, 2]$. The following propositions are equivalent:*

- (1) *The two level concentration (4.3) holds for some non-negative a, b independent of n .*
- (2) *The probability measure μ verifies the transportation-cost inequality*

$$\forall \nu \in P(\mathbf{R}^d), \quad \mathcal{T}_{2,p}(\nu, \mu) \leq C H(\nu | \mu),$$

for some constant C .

More precisely, if (4.3) holds for some constants a, b , then the transportation-cost inequality holds with the constant $C = 288/a$. Conversely, if the transportation-cost inequality holds for some constant C , then (4.3) is true for $b = 2$ and $a = 1/(2C)$.

The following lemma collects different facts that are needed in the proof.

Lemma 4.5.

- (1) *For all $x, y \geq 0$, $\alpha_p(x + y) \leq 2\alpha_p(x) + 2\alpha_p(y)$.*
- (2) *For all integer $n \geq 1$ and all probability measures ν_1, ν_2 and ν_3 on $(\mathbf{R}^d)^n$,*

$$\mathcal{T}_{2,p}(\nu_1, \nu_3) \leq 2\mathcal{T}_{2,p}(\nu_1, \nu_2) + 2\mathcal{T}_{2,p}(\nu_2, \nu_3).$$

- (3) *For all integer $n \geq 1$ and all $r \geq 0$, define*

$$B_{2,p}(r) = \left\{ x \in (\mathbf{R}^d)^n : \sum_{i=1}^n \sum_{j=1}^d \alpha_p(x_j^i) \leq r \right\}.$$

Then for all $p \in [1, 2]$,

$$\frac{1}{12} (\sqrt{r}B_2 + \sqrt[p]{r}B_p) \subset B_{2,p}(r) \subset \sqrt{r}B_2 + \sqrt[p]{r}B_p$$

Proof. The first point is easy to check. The second point follows from the first one by integration ; the detailed argument can be found in the proof of [Goz07, Proposition 4]. The third point is Lemma 2.3 of [Tal94]. \square

Proof of Theorem 4.4. Let us recall the proof of (2) implies (1). According to the tensorization property, for all n and all probability measure ν on $(\mathbf{R}^d)^n$,

$$\mathcal{T}_{2,p}(\nu, \mu^n) \leq C \mathbf{H}(\nu | \mu^n)$$

holds. Take A and B in $(\mathbf{R}^d)^n$ and define $d\mu_A^n = \mathbf{1}_A d\mu/\mu^n(A)$ and $d\mu_B^n = \mathbf{1}_B d\mu/\mu^n(B)$. According to point (2) of Lemma 4.5, and the transportation-cost inequality satisfied by μ^n , one has

$$\begin{aligned} \mathcal{T}_{2,p}(\mu_A^n, \mu_B^n) &\leq 2\mathcal{T}_{2,p}(\mu_A^n, \mu^n) + 2\mathcal{T}_{2,p}(\mu_B^n, \mu^n) \leq 2C \mathbf{H}(\mu_A^n | \mu^n) + 2C \mathbf{H}(\mu_B^n | \mu^n) \\ &= -2C \log(\mu^n(A)\mu^n(B)). \end{aligned}$$

Define

$$c_{2,p}(A, B) = \inf \{r \geq 0 \text{ s.t. } (A + B_{2,p}(r)) \cap B \neq \emptyset\}$$

then $\mathcal{T}_{2,p}(\mu_A^n, \mu_B^n) \geq c_{2,p}(A, B)$ and so

$$\mu^n(A)\mu^n(B) \leq e^{-c_{2,p}(A,B)/2C}.$$

Now, if $\mu^n(A) \geq 1/2$ and $B = (\mathbf{R}^d)^n \setminus (A + B_{2,p}(r))$, one has $c_{2,p}(A, B) = r$ and so $\mu^n(A + B_{2,p}(r)) \geq 1 - 2e^{-r/2C}$. Using point (3) of Lemma 4.5 gives $\mu^n(A + \sqrt{r}B_2 + \sqrt[p]{r}B_p) \geq 1 - 2e^{-r/2C}$.

Now let us prove the converse. Let $(X_i)_i$ be an i.i.d sequence of law μ and let L_n be its empirical measure. Consider $A = \{x \in (\mathbf{R}^d)^n \text{ s.t. } \mathcal{T}_{2,p}(L_n^x, \mu) \leq m_n\}$ where m_n denotes the median of $\mathcal{T}_{2,p}(L_n, \mu)$. According to point (3) of Lemma 4.5, $A + \sqrt{r}B_2 + \sqrt[p]{r}B_p \subset A + 12B_{2,p}(r)$. Let $x \in A + 12B_{2,p}(r)$; there is some $\bar{x} \in A$ such that

$$\sum_{i=1}^n \sum_{j=1}^d \alpha_p \left(\frac{x_j^i - \bar{x}_j^i}{12} \right) \leq r$$

(here $x = (x^1, x^2, \dots, x^n)$ with $x^i \in \mathbf{R}^d$). Since $\alpha_p(x/12) \geq \alpha_p(x)/144$, one gets $\mathcal{T}_{2,p}(L_n^x, L_n^{\bar{x}}) \leq 144r/n$. According to point (2) of Lemma 4.5, $\mathcal{T}_{2,p}(L_n^x, \mu) \leq 2\mathcal{T}_{2,p}(L_n^x, L_n^{\bar{x}}) + 2\mathcal{T}_{2,p}(L_n^{\bar{x}}, \mu) \leq 2m_n + 288r/n$. Consequently, the following holds for all n :

$$\forall r \geq 0, \quad \mathbb{P}(\mathcal{T}_{2,p}(L_n, \mu) \geq 2m_n + 288r/n) \leq be^{-ar}.$$

Reasoning as in the proof of Theorem 1.4, one concludes that

$$\forall \nu \in \mathbf{P}(\mathbf{R}^d), \quad \mathcal{T}_{2,p}(\nu, \mu) \leq \frac{288}{a} \mathbf{H}(\nu | \mu).$$

\square

5. POINCARÉ INEQUALITY AND EXPONENTIAL CONCENTRATION

In this section, one considers more carefully the case $p = 1$ of the preceding one. Let us recall that a probability measure μ on \mathbf{R}^d satisfies the Poincaré inequality with constant $C > 0$ if

$$(5.1) \quad \text{Var}_\mu(f) \leq C \int |\nabla f|_2^2 d\mu$$

for all smooth f .

The following theorem proves the equivalence between Poincaré inequality, dimension free exponential concentration and the corresponding transportation-cost inequality.

Theorem 5.2. *Let μ be a probability measure on \mathbf{R}^d . The following propositions are equivalent:*

- (1) *The probability measure μ verifies Poincaré inequality with a constant C_1 .*
- (2) *The probability measure μ verifies for some constants $a, b > 0$*

$$\forall r \geq 0, \quad \mu^n(A + D_{2,1}(r)) \geq 1 - be^{-ar},$$

for all subset A of $(\mathbf{R}^d)^n$ such that $\mu^n(A) \geq 1/2$, where the set $D_{2,1}(r)$ is defined by

$$D_{2,1}(r) = \left\{ x \in (\mathbf{R}^d)^n \text{ s.t. } \sum_{i=1}^n \alpha_1(|x^i|_2) \leq r \right\}.$$

- (3) *The probability measure μ verifies the following transportation-cost inequality for some constant $C_2 > 0$*

$$\forall \nu \in \mathcal{P}(\mathbf{R}^d), \quad \mathcal{T}_{\text{SG}}(\nu, \mu) = \inf_{\pi} \int \alpha_1(|x - y|_2) d\pi(x, y) \leq C_2 \mathcal{H}(\nu | \mu).$$

More precisely:

- (1) implies (2) with $a = \kappa \max(C_1, \sqrt{C_1})^{-1}$, κ being a universal constant.
- (2) implies (3) with $C_2 = 2/a$.
- (3) implies (1) with $C_1 = C_2/2$.

The equivalence between (1) and (3) was first obtained by Bobkov, Gentil and Ledoux in [BGL01, Corollary 5.1] with the Hamilton-Jacobi approach. The equivalence of (1) and (2) (or (2) and (3)) seems to be new.

Proof. According to (a careful reading of) [BL97, Corollary 3.2], (1) implies (2) with $b = 1$ and a depending only on C_1 ; one can take $a = \kappa \max(C_1, \sqrt{C_1})^{-1}$, where κ is a universal constant. According to (a slightly different version of) Theorem 4.4 with $p = 1$, (2) implies (3) (with $C_2 = 2/a$). It remains to prove that (3) implies (1). This last point is classical; let us simply sketch the proof. The transportation-cost inequality is equivalent to the following property: for all bounded f on \mathbf{R}^d ,

$$\int e^{Qf} d\mu \leq e^{\int f d\mu},$$

where $Qf(x) = \inf_{y \in \mathbf{R}^d} \{f(y) + C_2^{-1} \alpha_1(|x - y|_2)\}$ (for a proof of this fact see e.g the proof of (3.15) in [BG99] or [GL07, Corollary 1]). Let f be a smooth function and apply the preceding

inequality to tf . When t goes to 0, it can be shown that

$$Q(tf)(x) - tf(x) = -\frac{C_2 t^2}{4} |\nabla f|_2^2(x) + o(t^2),$$

so $\int e^{Q(tf)} d\mu = 1 + t \int f d\mu + \frac{t^2}{2} \int f^2 d\mu - \frac{C_2 t^2}{4} \int |\nabla f|_2^2 d\mu + o(t^2)$. On the other hand, $e^{t \int f d\mu} = 1 + t \int f d\mu + \frac{t^2}{2} (\int f d\mu)^2$. One concludes, that

$$\text{Var}_\mu(f) \leq \frac{C_2}{2} \int |\nabla f|^2 d\mu,$$

which achieves the proof. □

6. REMARKS

6.1. The (τ) property. Transportation-cost inequalities are closely related to the so called (τ) property introduced by Maurey in [Mau91]. If $c(x, y)$ is a non negative function defined on some product space $\mathcal{X} \times \mathcal{X}$ and μ is a probability measure on \mathcal{X} , one says that (μ, c) has the (τ) property if for all non-negative f on \mathcal{X} ,

$$\int e^{Q_c f} d\mu \cdot \int e^{-f} d\mu \leq 1,$$

where $Q_c f(x) = \inf_{y \in \mathcal{X}} \{f(y) + c(x, y)\}$. The recent paper by Latała and Wojtaszczyk [LW08] provides an excellent introduction together with a lot of new results concerning this class of inequalities.

The (τ) property is in fact a sort of dual version of the transportation-cost inequality. This was first observed by Bobkov and Götze in [BG99]. In the case of \mathbf{T}_2 , one can show that if μ verifies $\mathbf{T}_2(C)$ then $(\mu, (2C)^{-1}|x-y|_2^2)$ has the (τ) property and conversely, if $(\mu, C^{-1}|x-y|_2^2)$ has the (τ) property, then μ verifies $\mathbf{T}_2(C)$. A general statement can be found in [Goz08, Proposition 4.17].

6.2. Sufficient conditions for transportation-cost inequalities. Several sufficient conditions for transportation-cost inequalities are known. Let us recall some of them. In [Goz07, Theorem 5], the author proved the following result:

Theorem 6.1. *Let μ be a symmetric probability measure on \mathbf{R} of the form $d\mu(x) = e^{-V(x)} dx$, with V a smooth function such that $\lim_{x \rightarrow +\infty} \frac{V''(x)}{V'(x)^2} = 0$. Let $p \geq 1$; if V is such that*

$$\limsup_{x \rightarrow +\infty} \frac{x^{p-1}}{V'(x)} < +\infty, \text{ then } \mu \text{ verifies the transportation-cost inequality}$$

$$\forall \nu \in \mathbf{P}(\mathbf{R}), \quad \inf_{\pi \in P(\nu, \mu)} \int \alpha_p(x-y) d\pi(x, y) \leq C H(\nu | \mu),$$

where $\alpha_p(u) = u^2$ if $|u| \leq 1$ and $\alpha_p(u) = |u|^p$ if $|u| \geq 1$.

The case $p = 2$ was first established by Cattiaux and Guillin in [CG06] with a completely different proof. Other cost functions α can be considered in place of the α_p . Furthermore, if μ satisfies Cheeger's inequality on \mathbf{R} , then a necessary and sufficient condition is known for the transportation-cost inequality associated to α (see [Goz07, Theorem 2]).

On \mathbf{R}^d , a relatively weak sufficient condition for \mathbf{T}_2 (and other transportation-cost inequalities) was established by the author in [Goz08] (Theorem 4.8 and Corollary 4.13). Define $\omega^{(d)} : \mathbf{R}^d \rightarrow \mathbf{R}^d : (x_1, \dots, x_d) \mapsto (\omega(x_1), \dots, \omega(x_d))$, where $\omega(u) = \varepsilon(u) \max(|u|, u^2)$ with $\varepsilon(u) = 1$ when u is non-negative and -1 otherwise. If the image of μ under the map $\omega^{(d)}$ verifies the Poincaré inequality, then μ satisfies \mathbf{T}_2 . It can be shown that this condition is strictly weaker than the condition μ verifies **LSI** (see [Goz08, Theorem 5.9]).

Other sufficient conditions were obtained by Bobkov and Ledoux in [BL00] with an approach based on the Prekopa-Leindler inequality, or in [CEGH04] by Cordero-Erausquin, Gangbo and Houdré with an optimal transportation method.

APPENDIX

The following proposition is quite classical in Large Deviations theory. It can be found in Deuschell and Stroock's book [DS89, Exercise 3.3.23, p. 76].

Proposition 6.2. *Let $A \subset \mathcal{P}(\mathcal{X})$ be such that $\{x \in \mathcal{X}^n : L_n^x \in A\}$ is measurable. Then for every probability measure ν on \mathcal{X} absolutely continuous with respect to μ and such that $\nu^n(x : L_n^x \in A) > 0$, one has*

$$(6.3) \quad \frac{1}{n} \log \left(\mu^n(L_n \in A) e^{nH(\nu|\mu)} \right) \geq -H(\nu|\mu) \frac{\nu^n(L_n \in A^c)}{\nu^n(L_n \in A)} + \frac{1}{n} \log \nu^n(L_n \in A) - \frac{1}{n e \nu^n(L_n \in A)}$$

Proof. Let $h = \frac{d\nu^n}{d\mu^n}$ and $B = \{x \in \mathcal{X}^n : L_n^x \in A \text{ and } h(x) > 0\}$. Then,

$$\mu^n(L_n \in A) \geq \mu^n(B) = \int_B h(x) d\nu^n(x) = \nu^n(B) \frac{\int_B e^{-\log h(x)} d\nu^n(x)}{\nu^n(B)}.$$

Applying Jensen's inequality gives

$$\log \mu^n(L_n \in A) \geq \log \nu^n(B) - \frac{\int_B \log h(x) d\nu^n}{\nu^n(B)}.$$

Since $H(\nu^n|\mu^n) = \int \log h(x) d\nu^n$, one concludes that

$$(6.4) \quad \log \mu^n(L_n \in A) \geq \log \nu^n(B) - \frac{H(\nu^n|\mu^n)}{\nu^n(B)} + \frac{\int_{B^c} \log h(x) h(x) d\mu^n}{\nu^n(B)}$$

But for all $x > 0$, $x \log x \geq -1/e$, so

$$(6.5) \quad \frac{\int_{B^c} \log h(x) h(x) d\mu^n}{\nu^n(B)} \geq -\frac{\mu^n(B)}{e \nu^n(B)} \geq -\frac{1}{e \nu^n(B)}.$$

Putting (6.5) into (6.4) and using

$$H(\nu^n|\mu^n) = n H(\nu|\mu) \quad \text{and} \quad \nu^n(B) = \nu^n(L_n \in A),$$

gives the desired inequality. \square

Proof of Theorem 2.2. Let $t \geq 0$ and define $A = \{\nu \in \mathcal{P}_p(\mathcal{X}) \text{ s.t. } W_p(\nu, \mu) > t\}$. Take $\nu \in A$ such that $H(\nu|\mu) < +\infty$. If $(Y_i)_i$ is an i.i.d sequence of law ν , and $L_n^Y = n^{-1} \sum_{i=1}^n \delta_{Y_i}$, then L_n^Y converges to ν almost surely for the W_p distance and so $\nu^n(L_n \in A) = \mathbb{P}(W_p(L_n^Y, \mu) > t) \rightarrow$

$\mathbb{P}(W_p(\nu, \mu) > t) = 1$, when n tends to $+\infty$. Applying Proposition 6.2 to A and ν and taking the limit when n goes to $+\infty$, gives

$$\liminf_{n \rightarrow +\infty} \frac{1}{n} \log \mathbb{P}(W_p(L_n, \mu) > t) \geq -H(\nu | \mu).$$

Optimizing over ν gives the result. \square

REFERENCES

- [ABC⁺00] C. Ané, S. Blachère, D. Chafaï, P. Fougères, I. Gentil, F. Malrieu, C. Roberto, and G. Scheffer. *Sur les inégalités de Sobolev logarithmiques*, volume 10 of *Panoramas et Synthèses [Panoramas and Syntheses]*. Société Mathématique de France, Paris, 2000.
- [BCR06] F. Barthe, P. Cattiaux, and C. Roberto. Interpolated inequalities between exponential and Gaussian, Orlicz hypercontractivity and isoperimetry. *Rev. Mat. Iberoamericana*, 22(3):993–1067, 2006.
- [Bec89] W. Beckner. A generalized Poincaré inequality for Gaussian measures. *Proc. Amer. Math. Soc.*, 105:397–400, 1989.
- [BG99] S.G. Bobkov and F. Götze. Exponential integrability and transportation cost related to logarithmic Sobolev inequalities. *J. Funct. Anal.*, 163(1):1–28, 1999.
- [BGL01] S.G. Bobkov, I. Gentil, and M. Ledoux. Hypercontractivity of Hamilton-Jacobi equations. *J. Math. Pures Appl. (9)*, 80(7):669–696, 2001.
- [BL97] S.G. Bobkov and M. Ledoux. Poincaré’s inequalities and Talagrand’s concentration phenomenon for the exponential distribution. *Probab. Theory Related Fields*, 107(3):383–400, 1997.
- [BL00] S.G. Bobkov and M. Ledoux. From Brunn-Minkowski to Brascamp-Lieb and to logarithmic Sobolev inequalities. *Geom. Funct. Anal.*, 10(5):1028–1052, 2000.
- [BR03] F. Barthe and C. Roberto. Sobolev inequalities for probability measures on the real line. *Studia Math.*, 159(3):481–497, 2003.
- [BR06] F. Barthe and C. Roberto. Modified Logarithmic-Sobolev inequalities. Preprint, 2006.
- [BZ05] S.G. Bobkov and B. Zegarlinski. Entropy bounds and isoperimetry. *Mem. Amer. Math. Soc.*, 176(829):x+69, 2005.
- [CATD93] J. A. Cuesta-Albertos and A. Tueró-Díaz. A characterization for the solution of the Monge-Kantorovich mass transference problem. *Statist. Probab. Lett.*, 16(2):147–152, 1993.
- [CEGH04] D. Cordero-Erausquin, W. Gangbo, and C. Houdré. Inequalities for generalized entropy and optimal transportation. In *Recent advances in the theory and applications of mass transport*, volume 353 of *Contemp. Math.*, pages 73–94. Amer. Math. Soc., Providence, RI, 2004.
- [CG06] P. Cattiaux and A. Guillin. Talagrand’s like quadratic transportation cost inequalities. *J. Math. Pures Appl.*, 86(9):341–361, 2006.
- [DGW04] H. Djellout, A. Guillin, and L. Wu. Transportation cost-information inequalities and applications to random dynamical systems and diffusions. *Ann. Probab.*, 32(3B):2702–2732, 2004.
- [DS89] J.D. Deuschel and D.W. Stroock. *Large deviations*, volume 137 of *Pure and Applied Mathematics*. Academic Press Inc., Boston, MA, 1989.
- [Dud89] R.M. Dudley. *Real analysis and probability*. The Wadsworth & Brooks/Cole Mathematics Series. Wadsworth & Brooks/Cole Advanced Books & Software, Pacific Grove, CA, 1989.
- [DZ98] A. Dembo and O. Zeitouni. *Large deviations techniques and applications. Second edition*. Applications of Mathematics 38. Springer Verlag, 1998.
- [GGM05] I. Gentil, A. Guillin, and L. Miclo. Modified logarithmic Sobolev inequalities and transportation inequalities. *Probab. Theory Related Fields*, 133(3):409–436, 2005.
- [GL07] N. Gozlan and C. Léonard. A large deviation approach to some transportation cost inequalities. *Probab. Theory Related Fields*, 139(1-2):235–283, 2007.
- [GLWY07] A. Guillin, C. Léonard, L.M. Wu, and N. Yiao. Transportation-information inequalities for Markov processes. Preprint. Available on the ArXiv, 2007.
- [GM83] M. Gromov and V.D. Milman. A topological application of the isoperimetric inequality. *Amer. J. Math.*, 105:843–854, 1983.
- [Goz07] N. Gozlan. Characterization of Talagrand’s like transportation-cost inequality on the real line. *J. Funct. Anal.*, 250(2):400–425, 2007.

- [Goz08] N. Gozlan. Poincaré inequalities and dimension free concentration of measure. Preprint, 2008.
- [Led96] M. Ledoux. On Talagrand's deviation inequalities for product measures. *ESAIM Probab. Statist.*, 1:63–87, 1996.
- [Led01] M. Ledoux. *The concentration of measure phenomenon*, volume 89 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2001.
- [LO00] R. Latała and K. Oleszkiewicz. Between Sobolev and Poincaré. In *geometric aspects of Functional Analysis. Lect. Notes Math.*, 1745:147–168, 2000.
- [LV07] J. Lott and C. Villani. Hamilton-Jacobi semigroup on length spaces and applications. *J. Math. Pures Appl. (9)*, 88(3):219–229, 2007.
- [LW08] R. Latała and J. O. Wojtaszczyk. On the infimum convolution inequality. Preprint. Available on the ArXiv, 2008.
- [Mar86] K. Marton. A simple proof of the blowing-up lemma. *IEEE Trans. Inform. Theory*, 32(3):445–446, 1986.
- [Mar96] K. Marton. Bounding \bar{d} -distance by informational divergence: a method to prove measure concentration. *Ann. Probab.*, 24(2):857–866, 1996.
- [Mau91] B. Maurey. Some deviation inequalities. *Geom. Funct. Anal.*, 1(2):188–197, 1991.
- [OV00] F. Otto and C. Villani. Generalization of an inequality by Talagrand and links with the logarithmic Sobolev inequality. *J. Funct. Anal.*, 173(2):361–400, 2000.
- [Sam00] P.M. Samson. Concentration of measure inequalities for Markov chains and Φ -mixing processes. *Ann. Probab.*, 28(1):416–461, 2000.
- [Tal94] M. Talagrand. The supremum of some canonical processes. *Amer. J. Math.*, 116(2):283–325, 1994.
- [Tal96] M. Talagrand. Transportation cost for Gaussian and other product measures. *Geom. Funct. Anal.*, 6(3):587–600, 1996.
- [Vil03] C. Villani. *Topics in Optimal Transportation*. Graduate Studies in Mathematics 58. American Mathematical Society, Providence RI, 2003.
- [Vil08] C. Villani. Optimal transport, old and new. In preparation, 2008.
- [Wan04] F-Y Wang. Probability distance inequalities on Riemannian manifolds and path spaces. *J. Funct. Anal.*, 206(1):167–190, 2004.

UNIVERSITÉ PARIS EST - LABORATOIRE D'ANALYSE ET DE MATHÉMATIQUES APPLIQUÉES (UMR CNRS 8050), 5 BD DESCARTES, 77454 MARNE LA VALLÉE CEDEX 2, FRANCE

E-mail address: nathael.gozlan@univ-mlv.fr