

## Obscured High Mass X-Ray Binaries and Supergiant Fast X-ray Transients: Infrared Observations of *INTEGRAL* Sources

S. Chaty

*Laboratoire AIM, CEA/DSM - CNRS - Université Paris Diderot, IRFU/Service d'Astrophysique, Bât. 709, CEA-Saclay, FR-91191 Gif-sur-Yvette Cedex, France*

**Abstract.** A new type of high-energy binary system has been revealed by the *INTEGRAL* satellite. These sources are being unveiled by means of multi-wavelength optical, near- and mid-infrared observations. Among these sources, two distinct classes are appearing: the first one is constituted of intrinsically obscured high-energy sources, of which IGR J16318-4848 seems to be the most extreme example. The second one is populated by the so-called supergiant fast X-ray transients, with IGR J17544-2619 being the archetype. We report here on multi-wavelength optical to mid-infrared observations of a sample constituted of 21 *INTEGRAL* sources. We show that in the case of the obscured sources our observations suggest the presence of absorbing material (dust and/or cold gas) enshrouding the whole binary system. We finally discuss the nature of these two different types of sources, in the context of high energy binary systems.

### 1. Introduction

The *INTEGRAL* observatory has performed a detailed survey of the Galactic plane. The ISGRI detector on the IBIS imager has discovered many new high energy sources, most of which have been reported in Bird et al. (2007)<sup>1</sup>. The most important result of *INTEGRAL* to date is the discovery of many new high energy sources – concentrated in the Galactic plane, and in the Norma arm (see e.g. Chaty & Filliatre 2005) – exhibiting common characteristics which previously had rarely been seen. Many of them are high mass X-ray binaries (HMXBs) hosting a neutron star orbiting around an O/B companion, in most cases a supergiant star. They divide into two classes: some of the new sources are very obscured, exhibiting a huge intrinsic and local extinction, and the others are HMXBs hosting a supergiant star and exhibiting fast and transient outbursts – an unusual characteristic among HMXBs. These are therefore called Supergiant Fast X-ray Transients (SFXTs, Negueruela et al. 2006; Sguera et al. 2005). High-energy observations are not sufficient to reveal the nature of the newly discovered sources, since the *INTEGRAL* localisation ( $\sim 2'$ ) is not accurate enough to unambiguously pinpoint the source at other wavelengths. Once X-ray satellites such as *XMM-Newton*, *Chandra*, or *Swift* provide an arcsecond position, the hunt for the optical counterpart of the source is open. However, the high level of absorption towards the galactic plane makes the near-infrared (NIR) domain more efficient for identifying these sources. We first report on multi-wavelength observations of two sources, one belonging to each class described above, and we then give general results on *INTEGRAL* sources, before discussing them and concluding.

---

<sup>1</sup>See also <http://isdc.unige.ch/~rodrigue/html/igrsources.html>




Figure 1. Optical to MIR SEDs of IGR J16318-4848 (left) and IGR J17544-2619 (right), including data from ESO/NTT, VISIR on VLT/UT3 and *Spitzer* (Rahoui et al. 2008). IGR J16318-4848 exhibits a MIR excess, interpreted by Rahoui et al. (2008) as the signature of a strong stellar outflow coming from the sgB[e] companion star (Filliatre & Chaty 2004). On the other hand, IGR J17544-2619 is well fitted with only a stellar component corresponding to the O9Ib companion star spectral type (Pellizza et al. 2006).

## 2. Observations and Results

The multiwavelength observations were performed at the European Southern Observatory (ESO), using Target of Opportunity (ToO) and Visitor modes, in 3 domains: optical (400 – 800 nm) with EMMI, NIR (1 – 2.5  $\mu\text{m}$ ) with SOFI, both instruments at the focus of the 3.5m New Technology Telescope (NTT) at La Silla, and mid-infrared (MIR, 5 – 20  $\mu\text{m}$ ) with the VISIR instrument on Melipal, the 8m Unit Telescope 3 (UT3) of the Very Large Telescope (VLT) at Paranal (Chile). With these observations we performed accurate astrometry, photometry and spectroscopy on a sample constituted of 21 *INTEGRAL* sources, aiming at identifying their counterparts and the nature of the companion star, deriving their distance, and finally characterising the presence and temperature of their circumstellar medium.

### 2.1. IGR J16318-4848: Extreme among the Obscured High-Energy Sources

IGR J16318-4848 was the first source discovered by IBIS/ISGRI on *INTEGRAL* on 29 January 2003 (Courvoisier et al. 2003). *XMM-Newton* observations revealed an unusually high level of absorption:  $N_{\text{H}} \sim 2 \times 10^{24} \text{ cm}^{-2}$  (Matt & Guainazzi 2003). The accurate localisation by *XMM-Newton* allowed Filliatre & Chaty (2004) to rapidly trigger ToO photometric and spectroscopic observations in optical/NIR, leading to the discovery of the optical counterpart and to the confirmation of the NIR one (Walter et al. 2003). The extremely bright NIR source ( $K_s = 7.20$  magnitudes) exhibits an unusually strong intrinsic absorption in the optical ( $A_v = 17.4$  magnitudes), much stronger than the absorption along the line of sight ( $A_v = 11.4$  magnitudes), but still 100 times lower than the absorption in X-rays. This led Filliatre & Chaty (2004) to suggest that the material absorbing in X-rays was concentrated around the compact object, while the material absorbing in optical/NIR was enshrouding the whole system. The NIR spectroscopy revealed an unusual spectrum, with many strong emission lines, originating from a highly complex and stratified circumstellar environment of various densities and temperatures, suggesting the presence of an envelope and strong stellar outflow responsible for the absorption. Only luminous early-type stars such as su-




Figure 2. Scenarios illustrating two possible configurations of *INTEGRAL* sources: a neutron star orbiting around a supergiant star on a circular orbit (left image); and on an eccentric orbit (right image), accreting from the clumpy stellar wind of the supergiant. The accretion of matter is persistent in the case of the obscured sources, as in the left image, where the compact object orbits inside the cocoon of dust enshrouding the whole system. On the other hand, the accretion is intermittent in the case of SFXTs, which might correspond to a compact object on an eccentric orbit, as in the right image. A 3D animation of these sources is available on the website:

<http://www.aim.univ-paris7.fr/CHATY/Research/hidden.html>

pergiant sgB[e] show such extreme environments, and Filliatre & Chaty (2004) concluded that IGR J16318-4848 was an unusual HMXB. By combining these optical and NIR data with MIR observations, and fitting these observations with a model of a sgB[e] companion star, Rahoui et al. (2008) showed that IGR J16318-4848 exhibits a MIR excess (see Figure 1, left panel), that they interpreted as being due to the strong stellar outflow emanating from the sgB[e] companion star. They found that the companion star had a temperature of  $T = 22200$  K and radius  $R_{\star} = 20.4R_{\odot}$ , and an extra component of temperature  $T = 1100$  K and radius  $R = 10R_{\star}$ , with  $A_v = 17.6$  magnitudes. By taking a typical orbital period of 10 days and a mass of the companion star of  $20 M_{\odot}$ , we obtain an orbital separation of  $50 R_{\odot}$ , smaller than the extension of the extra component, suggesting that this component enshrouds the whole binary system, as would do a cocoon of gas/dust (see Figure 2, left panel). In summary, IGR J16318-4848 is an HMXB system, located at a distance between 1 and 6 kpc, hosting a compact object (probably a neutron star) and a sgB[e] star (it is therefore the second HMXB with a sgB[e] star, after CI Cam; Clark et al. 1999). The most striking facts are (i) the compact object seems to be surrounded by absorbing material and (ii) the whole binary system seems to be surrounded by a dense and absorbing circumstellar

material envelope or cocoon, made of cold gas and/or dust. This source exhibits such extreme characteristics that it might not be fully representative of the other obscured sources.

## 2.2. IGR J17544-2619: Archetype of the Supergiant Fast X-ray Transients

SFXTs constitute a new class of sources identified among the recently discovered *INTEGRAL* sources, exhibiting these common characteristics: rapid outbursts lasting only hours, a faint quiescent emission, high energy spectra requiring a BH or NS accretor, and O/B supergiant companion stars. IGR J17544-2619, a bright recurrent transient X-ray source discovered by *INTEGRAL* on 17 September 2003 (Sunyaev et al. 2003), seems to be their archetype. Observations with *XMM-Newton* have shown that it exhibits a very hard X-ray spectrum, and a relatively low intrinsic absorption ( $10^{22}$  cm $^{-2}$ , González-Riestra et al. 2004). Its bursts last for hours, and in between bursts it exhibits long quiescent periods, which can reach more than 70 days (Zurita Heras & Chaty, in prep.). The compact object is probably a neutron star (in't Zand 2005). Pellizza et al. (2006) managed to get optical/NIR ToO observations only one day after the discovery of this source. They identified a likely counterpart inside the *XMM-Newton* error circle, confirmed by an accurate localization from *Chandra*. Spectroscopy showed that the companion star was a blue supergiant of spectral type O9Ib, with a mass of  $25 - 28M_{\odot}$  and temperature of  $T \sim 31000$  K; the system is therefore an HMXB (Pellizza et al. 2006). Rahoui et al. (2008) combined optical, NIR and MIR observations and showed that they could accurately fit the observations with a model of an O9Ib star, with temperature  $T = 31000$  K and radius  $R_{\star} = 21.9R_{\odot}$ . They derived an absorption  $A_v = 6.1$  magnitudes and a distance  $D = 3.6$  kpc. The source does not exhibit any MIR excess (see Figure 1, right panel, Rahoui et al. 2008). In summary, IGR J17544-2619 is an HMXB at a distance of  $\sim 3.6$  kpc, constituted of an O9Ib supergiant, with a mild stellar wind and a compact object which is probably a neutron star, without any MIR excess.

## 3. General Results on *INTEGRAL* Sources and Discussion

To better characterize this population, Chaty et al. (2008) and Rahoui et al. (2008) studied a sample of 21 *INTEGRAL* sources belonging to both classes described above. Some results are reported in Table 1. The optical/NIR study, through accurate astrometry, photometry, and spectroscopy, allowed Chaty et al. (2008) to identify the counterparts, and to show that most of these systems are HMXBs containing massive and luminous early-type companion stars. By combining MIR photometry, and fitting their optical–MIR spectral energy distributions, Rahoui et al. (2008) showed that (i) most of these sources exhibit an intrinsic absorption and (ii) three of them exhibit a MIR excess, which they suggest is due to the presence of a cocoon of dust and/or cold gas enshrouding the whole binary system (see also Chaty & Rahoui 2006). Nearly all the *INTEGRAL* HMXBs for which both spin and orbital periods have been measured are located in the upper part of the Corbet diagram (Corbet 1986). They are wind accretors, typical of supergiant HMXBs, and X-ray pulsars exhibiting longer pulsation periods and higher absorption (by a factor  $\sim 4$ ) as compared to the average of previously known HMXBs (Bodaghee et al. 2007). This extra absorption might be due to the presence of a cocoon of dust/cold gas enshrouding the whole binary system in the case of the obscured sources. The intrinsic properties of the supergiant companion star could therefore explain some properties of these sources. However, differences exist between obscured sources and SFXTs, which might be explained by the geometry of the binary systems, and/or the extension of the wind/cocoon enshrouding either the companion star or the whole system. Indeed, obscured sources are naturally explained by a compact object orbiting inside a cocoon of dust and/or cold gas, while the fast X-ray behaviour of SFXTs needs a clumpy

Table 1. Results on the sample of *INTEGRAL* sources; more details are given in Chaty et al. (2008). We indicate respectively the name of the sources, the region of the Galaxy in the direction which they are located, their spin and orbital period, the interstellar, optical-IR, and X-ray derived column density respectively (in units of  $10^{22} \text{ cm}^{-2}$ ), their spectral type, nature and reference. Type abbreviations: AGN = Active Galactic Nucleus, B = Burster, BHC = Black Hole Candidate, CV = Cataclysmic Variable, D = Dipping source, H = High Mass X-ray Binary system, IP = Intermediate polar, L = Low Mass X-ray Binary, O = Obscured source, P = Persistent source, S = Supergiant Fast X-ray Transient, T: Transient source, XP: X-ray Pulsar. Reference are: c: Chaty et al. (2008), co: Combi et al. (2006), f: Filliatre & Chaty (2004), h: Hannikainen et al. (2007), m1: Masetti et al. (2004) m2: Masetti et al. (2006) n1: Negueruela et al. (2005), n2: Negueruela et al. (2006), n3: Nespoli et al. (2007), p: Pellizza et al. (2006), t: Tomsick et al. (2006).

Source	Reg	$P_s$ (s)	$P_o$ (d)	$Nh_{is}$	$Nh_{IR}$	$Nh_X$	SpT	Type	Ref
IGR J16167-4957	No			2.2	0.23	0.5	A0	CV/IP	t,m2
IGR J16195-4945	No			2.18	2.9	7	OB	H?/S?/O	t
IGR J16207-5129	No			1.73	2.0	3.7	BOI	H/O	t,m2
IGR J16318-4848	No			2.06	3.3	200	sgB[e]	H/O/P	f
IGR J16320-4751	No	1250	8.96(1)	2.14	6.6	21	sgOB	H/XP/T/O	c
IGR J16358-4726	No	5880		2.20	3.3	33	sgB[e]?	H/XP/T/O	c
IGR J16393-4643	No	912	3.6875(6)	2.19	2.19	24.98	BIV-V?	H/XP/T	c
IGR J16418-4532	No	1246	3.753(4)	1.88	2.7	10	sgOB?	H/XP/S	c
IGR J16465-4507	No	228		2.12	1.1	60	B0.5I	H/S	n1
IGR J16479-4514	No			2.14	3.4	7.7	sgOB	H/S?	c
IGR J16558-5203	-	-	-	-	-	-	Sey1.2	AGN	m2
IGR J17091-3624	GC			0.77	1.03	1.0	L	L/BHC	c
IGR J17195-4100	GC			0.77		0.08		CV/IP	t,m2
IGR J17252-3616	GC	413	9.74(4)	1.56	3.8	15	sgOB	H/XP/O	c
IGR J17391-3021	GC			1.37	1.7	29.98	O8Iab(f)	H/S/O	n2
IGR J17544-2619	GC		$\geq 70?$	1.44	1.1	1.4	O9Ib	S	p
IGR J17597-2201	GC			1.17	2.84	4.50	L	L/B/D/P	c
IGR J18027-1455	-	-	-	-	-	-	Sey1	AGN	m1,co
IGR J18027-2016	GC	139	4.5696(9)	1.04	1.53	9.05	sgOB	H/XP/T	c
IGR J18483-0311	GC	21.05	18.55	1.62	2.45	27.69	sgOB?	H/XP	c
IGR J19140+0951			13.558(4)	1.68	2.9	6	sgB0.5I	H/O	n3,h

stellar wind environment, to account for fast and transient accretion phenomena (see Figure 2, left and right panels respectively, and Chaty & Rahoui 2006). These results show the existence in our Galaxy of a dominant population of a previously rare class of high-energy binary systems: supergiant HMXBs, some exhibiting a high intrinsic absorption (Chaty et al. 2008; Rahoui et al. 2008). A careful study of this population, recently revealed by *INTEGRAL*, will provide a better understanding of the formation and evolution of short-living HMXBs. Furthermore, stellar population models will henceforth have to take these objects into account, to assess a realistic number of high-energy binary systems in our Galaxy. Our final word is that only a multiwavelength study reveal the nature of these obscured high-energy sources.

**Acknowledgments.** SC would like to thank the organisers for their invitation to report on these exciting results about newly discovered *INTEGRAL* sources, and for organising an interesting workshop, in a nice place, fruitful for scientific discussions and new ideas to arise.

## References

- Bird, A. J., Malizia, A., Bazzano, A., Barlow, E. J., Bassani, L., Hill, A. B., Bélanger, G., Capitanio, F., Clark, D. J., Dean, A. J., Fiocchi, M., Götz, D., Lebrun, F., Molina, M., Produit, N., Renaud, M., Sguera, V., Stephen, J. B., Terrier, R., Ubertini, P., Walter, R., Winkler, C., and Zurita, J. (2007). The Third IBIS/ISGRI Soft Gamma-Ray Survey Catalog. *ApJS*, 170:175–186.
- Bodaghee, A., Courvoisier, T. J.-L., Rodriguez, J., Beckmann, V., Produit, N., Hannikainen, D., Kuulkers, E., Willis, D. R., and Wendt, G. (2007). A description of sources detected by *INTEGRAL* during the first 4 years of observations. *A&A*, 467:585–596.
- Chaty, S. & Filliatre, P. (2005). Revealing the Nature of the Obscured High Mass X-ray Binary IGR J16318-4848. *Ap&SS*, 297:235–244.
- Chaty, S. & Rahoui, F. (2006). Optical to Mid-infrared observations revealing the most obscured high-energy sources of the Galaxy. In *Procs. of 6th INTEGRAL workshop, Moscow, Russia*. in press (astro-ph/0609474).
- Chaty, S., Rahoui, F., Foellmi, C., Rodriguez, J., Tomsick, J. A., and Walter, R. (2008). The most obscured high-energy sources of the galaxy brought to light by optical/infrared observations. *A&A*. in press.
- Clark, J. S., Steele, I. A., Fender, R. P., and Coe, M. J. (1999). Near IR spectroscopy of candidate B[e]/X-ray binaries. *A&A*, 348:888–896.
- Combi, J. A., Ribó, M., Martí, J., and Chaty, S. (2006). Multi-wavelength properties of the high-energy bright Seyfert 1 galaxy IGR J18027-1455. *A&A*, 458:761–766.
- Corbet, R. H. D. (1986). The three types of high-mass X-ray pulsator. *MNRAS*, 220:1047–1056.
- Courvoisier, T. J.-L., Walter, R., Rodriguez, J., Bouchet, L., and Lutovinov, A. A. (2003). Igr J16318-4848. *IAU Circ.*, 8063:3–+.
- Filliatre, P. & Chaty, S. (2004). The Optical/Near-Infrared Counterpart of the *INTEGRAL* Obscured Source IGR J16318-4848: An sgB[e] in a High-Mass X-Ray Binary? *ApJ*, 616:469–484.
- González-Riestra, R., Oosterbroek, T., Kuulkers, E., Orr, A., and Parmar, A. N. (2004). XMM-Newton observations of the *INTEGRAL* X-ray transient IGR J17544-2619. *A&A*, 420:589–594.
- Hannikainen, D. C., Rawlings, M. G., Muhli, P., Vilhu, O., Schultz, J., and Rodriguez, J. (2007). The nature of the infrared counterpart of IGR J19140+0951. *MNRAS*, 380:665–668.
- in’t Zand, J. J. M. (2005). Chandra observation of the fast X-ray transient IGR J17544-2619: evidence for a neutron star? *A&A*, 441:L1–L4.
- Masetti, N., Morelli, L., Palazzi, E., Galaz, G., Bassani, L., Bazzano, A., Bird, A. J., Dean, A. J., Israel, G. L., Landi, R., Malizia, A., Minniti, D., Schiavone, F., Stephen, J. B., Ubertini,

- P., and Walter, R. (2006). Unveiling the nature of INTEGRAL objects through optical spectroscopy. V. Identification and properties of 21 southern hard X-ray sources. *A&A*, 459:21–30.
- Masetti, N., Palazzi, E., Bassani, L., Malizia, A., and Stephen, J. B. (2004). Unveiling the nature of three INTEGRAL sources through optical spectroscopy. *A&A*, 426:L41.
- Matt, G. & Guainazzi, M. (2003). The properties of the absorbing and line-emitting material in IGR J16318 - 4848. *MNRAS*, 341:L13–L17.
- Negueruela, I., Smith, D. M., and Chaty, S. (2005). Optical counterpart to IGR J16465-4507. *The Astronomer's Telegram*, 429:1–+.
- Negueruela, I., Smith, D. M., Reig, P., Chaty, S., and Torrejón, J. M. (2006). Supergiant Fast X-ray Transients: a new class of high mass X-ray binaries unveiled by INTEGRAL. In Wilson, A., editor, *ESA Special Publication*, volume 604 of *ESA Special Publication*, pages 165–170.
- Nespoli, E., Fabregat, J., and Mennickent, R. (2007). K-band spectroscopy of AX J1841.0-0536 and IGR J19140+0951. *The Astronomer's Telegram*, 982:1–+.
- Pellizza, L. J., Chaty, S., and Negueruela, I. (2006). IGR J17544-2619: a new supergiant fast X-ray transient revealed by optical/infrared observations. *A&A*, 455:653–658.
- Rahoui, F., Chaty, S., Lagage, P.-O., and Pantin, E. (2008). The most obscured high-energy sources of the galaxy brought to light by optical/infrared observations. *A&A*. in press.
- Sguera, V., Barlow, E. J., Bird, A. J., Clark, D. J., Dean, A. J., Hill, A. B., Moran, L., Shaw, S. E., Willis, D. R., Bazzano, A., Ubertini, P., and Malizia, A. (2005). INTEGRAL observations of recurrent fast X-ray transient sources. *A&A*, 444:221–231.
- Sunyaev, R. A., Grebenev, S. A., Lutovinov, A. A., Rodriguez, J., Mereghetti, S., Gotz, D., and Courvoisier, T. (2003). New source IGR J17544-2619 discovered with INTEGRAL. *The Astronomer's Telegram*, 190:1–+.
- Tomsick, J. A., Chaty, S., Rodriguez, J., Foschini, L., Walter, R., and Kaaret, P. (2006). Identifications of Four INTEGRAL Sources in the Galactic Plane via Chandra Localizations. *ApJ*, 647:1309–1322.
- Walter, R., Rodriguez, J., Foschini, L., de Plaa, J., Corbel, S., Courvoisier, T. J.-L., den Hartog, P. R., Lebrun, F., Parmar, A. N., Tomsick, J. A., and Ubertini, P. (2003). INTEGRAL discovery of a bright highly obscured galactic X-ray binary source IGR J16318-4848. *A&A*, 411:L427–L432.