

Diffusion versus absorption in semilinear parabolic problems¹

Andrey Shishkov

Institute of Applied Mathematics and Mechanics of NAS of Ukraine, R. Luxemburg str. 74, 83114 Donetsk, Ukraine

Email: *shishkov@iamm.ac.donetsk.ua* .

Laurent Véron

Laboratoire de Mathématiques et Physique Théorique, CNRS UMR 6083, Faculté des Sciences, 37200 Tours, France.

Email: *veronl@univ-tours.fr*

Abstract. We study the limit, when $k \rightarrow \infty$, of the solutions $u = u_k$ of (E) $\partial_t u - \Delta u + h(t)u^q = 0$ in $\mathbb{R}^N \times (0, \infty)$, $u_k(\cdot, 0) = k\delta_0$, with $q > 1$, $h(t) > 0$. If $h(t) = e^{-\omega(t)/t}$ where $\omega > 0$ satisfies to $\int_0^1 \sqrt{\omega(t)}t^{-1}dt < \infty$, the limit function u_∞ is a solution of (E) with a single singularity at $(0, 0)$, while if $\omega(t) \equiv 1$, u_∞ is the maximal solution of (E). We examine similar questions for equations such as $\partial_t u - \Delta u^m + h(t)u^q = 0$ with $m > 1$ and $\partial_t u - \Delta u + h(t)e^u = 0$.

Diffusion versus absorption dans des problèmes paraboliques semi-linéaires

Résumé. Nous étudions la limite, quand $k \rightarrow \infty$, des solutions $u = u_k$ de (E) $\partial_t u - \Delta u + h(t)u^q = 0$ dans $\mathbb{R}^N \times (0, \infty)$, $u_k(\cdot, 0) = k\delta_0$ avec $q > 1$, $h(t) > 0$. Nous montrons que si $h(t) = e^{-\omega(t)/t}$ où $\omega > 0$ vérifie $\int_0^1 \sqrt{\omega(t)}t^{-1}dt < \infty$, la fonction limite u_∞ est une solution of (E) avec une singularité isolée en $(0, 0)$, alors que si $\omega(t) \equiv 1$, u_∞ est la solution maximale de (E). Nous examinons des questions semblables pour des équations des type suivants $\partial_t u - \Delta u^m + h(t)u^q = 0$ avec $m > 1$ et $\partial_t u - \Delta u + h(t)e^u = 0$.

Version française abrégée

Soit $q > 1$ et $h : \mathbb{R}_+ \mapsto \mathbb{R}_+$ une fonction continue, croissante telle que $h(t) > 0$ pour $t > 0$. Il est facile de vérifier que toute solution positive u de

$$(1) \quad \partial_t u - \Delta u + h(t)u^q = 0 \quad \text{dans } \mathbb{R}^N \times]0, +\infty[$$

satisfait à

$$(2) \quad u(x, t) \leq U(t) := \left((q-1) \int_0^t h(s) ds \right)^{-1/(q-1)} \quad \forall (x, t) \in \mathbb{R}^N \times]0, +\infty[.$$

Si $h \in L^1(0, 1, t^{Nq/2} dt)$, il est classique que pour tout $k > 0$ il existe une unique solution (dite fondamentale) $u = u_k$ de (1) sur $\mathbb{R}^N \times]0, +\infty[$ vérifiant $u_k(\cdot, 0) = k\delta_0$.

Par le principe du maximum $k \mapsto u_k$ est croissant et deux cas peuvent se produire:

- (i) ou bien $u_\infty = \lim_{k \rightarrow \infty} u_k = U$. *Explosion initiale complète.*
- (ii) ou bien u_∞ est une solution de (1) singulière en $(0, 0)$ vérifiant $\lim_{t \rightarrow 0} u_\infty(x, t)$

¹To appear in *C. R. Acad. Sci. Paris, Ser. I.*

= 0 pour tout $x \neq 0$. *Explosion initiale ponctuelle.*

Theorem 1. (I) Si $h(t) = e^{-\sigma/t}$ pour un $\sigma > 0$, alors $u_\infty = U$.

(II) Si $h(t) = e^{-\omega(t)/t}$ où ω est monotone croissante sur $]0, +\infty[$ et vérifie, pour un $\alpha \in [0, 1[$, $\inf\{\omega(t)/t^\alpha : 0 < t \leq 1\} > 0$ et $\int_0^1 \sqrt{\omega(t)} t^{-1} dt < \infty$, alors u_∞ a une explosion initiale ponctuelle.

Dans le cas de l'équation

$$(3) \quad \partial_t u - \Delta u + h(t)e^u = 0 \quad \text{dans } \mathbb{R}^N \times]0, +\infty[,$$

toute solution u satisfait à

$$(4) \quad u(x, t) \leq \tilde{U}(t) := -\ln \left(\int_0^t h(s) ds \right) \quad \forall (x, t) \in \mathbb{R}^N \times]0, +\infty[,$$

et l'existence d'une solution fondamentale $u = u_k$ est assurée si $h(t) = e^{-b(t)}$ avec $\lim_{t \rightarrow +\infty} t^{N/2} b(t) = +\infty$.

Theorem 2. (I) Si $h(t) = e^{-e^\sigma/t}$ pour un $\sigma > 0$, alors $u_\infty = \tilde{U}$.

(II) Si $h(t) = e^{-e^{\omega(t)}/t}$ où ω vérifie les conditions du Théorème 1, alors u_∞ a une explosion initiale ponctuelle.

Nos méthodes nous permettent aussi de traiter l'équation des milieux poreux avec absorption.

Main results

Let $q > 1$ and $h : (0, \infty) \mapsto (0, \infty)$ be a continuous nondecreasing function. It is easy to prove that any positive solution u of

$$(1) \quad \partial_t u - \Delta u + h(t)u^q = 0 \quad \text{dans } \mathbb{R}^N \times (0, +\infty)$$

verifies

$$(2) \quad u(x, t) \leq U(t) := \left((q-1) \int_0^t h(s) ds \right)^{-1/(q-1)} \quad \forall (x, t) \in \mathbb{R}^N \times (0, \infty).$$

If $h \in L^1(0, 1, t^{Nq/2} dt)$, it is classical that, for any $k > 0$, there exists a unique solution (called fundamental) $u = u_k$ of (1) sur $\mathbb{R}^N \times (0, \infty)$ such that $u_k(\cdot, 0) = k\delta_0$. By the maximum principle $k \mapsto u_k$ is increasing and the following alternative occurs:

(i) either $u_\infty = \lim_{k \rightarrow \infty} u_k = U$. *Complete initial blow-up.*

(ii) or u_∞ is a solution of (1) singular at $(0, 0)$ such that

$$\lim_{t \rightarrow 0} u_\infty(x, t) = 0 \quad \text{for all } x \neq 0. \quad \textit{Single-point initial blow-up.}$$

Theorem 1. (I) If $h(t) = e^{-\sigma/t}$ for some $\sigma > 0$, then $u_\infty = U$.

(II) If $h(t) = e^{-\omega(t)/t}$ where ω is nondecreasing on $(0, +\infty)$ and verifies, for some $\alpha \in [0, 1)$, $\inf\{\omega(t)/t^\alpha : 0 < t \leq 1\} > 0$ and

$$(3) \quad \int_0^1 \frac{\sqrt{\omega(t)} dt}{t} < +\infty,$$

then u_∞ has single-point initial blow-up.

Concerning equation

$$(4) \quad \partial_t u - \Delta u + h(t)e^u = 0 \quad \text{dans } \mathbb{R}^N \times (0, +\infty),$$

any solution u verifies

$$(5) \quad u(x, t) \leq \tilde{U}(t) := -\ln \left(\int_0^t h(s) ds \right) \quad \forall (x, t) \in \mathbb{R}^N \times (0, +\infty).$$

and the existence of a fundamental solution $u = u_k$ is ensured if $h(t) = e^{-b(t)}$ where $\lim_{t \rightarrow +\infty} t^{N/2} b(t) = +\infty$.

Theorem 2. (I) If $h(t) = O(e^{-e^{\sigma/t}})$ for some $\sigma > 0$, then $u_\infty = \tilde{U}$.

(II) If $h(t) = e^{-e^{\omega(t)/t}}$ where ω satisfies the conditions of Theorem 1, then u_∞ has single-point initial blow-up.

Our methods apply to equations of porous media type

$$(6) \quad \partial_t u - \Delta u^m + h(t)u^q = 0 \quad \text{in } \mathbb{R}^N \times (0, \infty),$$

with $m > 1$, $q > 1$ and $h : (0, \infty) \mapsto (0, \infty)$ is nondecreasing. As above, any positive solution satisfies (2). If $h \in L^1((0, 1; t^{-(q-1)/(m-1+2N^{-1})} dt))$, for any $k > 0$ there exists a solution $u = u_k$ of (6) such that $u_k(\cdot, 0) = k\delta_0$. Since $k \mapsto u_k$ is increasing, the same alternative as in case of (1) occurs concerning u_∞ .

Theorem 3. Assume $q > m > 1$. (I) If $h(t) = O(t^{(q-m)/(m-1)})$, then $u_\infty = U$.

(II) If $h(t) = t^{(q-m)/(m-1)}\omega^{-1}(t)$ where ω is nondecreasing and positive on $(0, +\infty)$ and verifies

$$(7) \quad \int_0^1 \frac{\omega^\theta(t) dt}{t} < +\infty,$$

where

$$\theta = \frac{m^2 - 1}{(N(m-1) + 2(m+1))(q-1)},$$

then u_∞ has single-point initial blow-up.

Sketch of the proofs. The complete initial blow-up results are proved by constructing local subsolutions by modifying the very singular solutions of some related equations. Since for equation (1), the proof is already given in [3] we shall outline the (more complicated) construction for equation (4).

Lemma 4. If $h(t) = \sigma t^{-2} e^{\sigma t^{-1} - e^{-\sigma/t}}$ for some $\sigma > 0$, complete initial blow-up occurs for equation (4).

Proof. Writing $h(t) = e^{-a(t)}$ is first observed that fundamental solutions u_k of (4) exist for all $k > 0$ if $\lim_{t \rightarrow 0} t^{N/2} a(t) = \infty$. For $\ell > 1$, let $v = v_{\infty, \ell}$ be the very singular solution of

$$(8) \quad \partial_t v - \Delta v + ct^{\alpha_\ell} v^\ell = 0$$

in $\mathbb{R}^N \times (0, \infty)$, where α_ℓ and c are positive constants. The choice of $\alpha_\ell = (N+2)/(\ell-1)/2 - 1$ ensures the existence of $v_{\infty, \ell}$. Furthermore, if we write

$$v_{\infty, \ell}(x, t) = \left(\frac{2c}{N+2} \right)^{1/(\ell-1)} t^{-(1+N/2)} f_\ell(x/\sqrt{t}),$$

then $f_\ell(\eta) \leq 1$ for $\eta \in \mathbb{R}^N$ and

$$(9) \quad \Delta f_\ell + \frac{1}{2} Df_\ell \cdot \eta + \frac{N+2}{2} f_\ell - f_\ell^\ell = 0.$$

By the maximum principle $0 < f_\ell < f_{\ell'} \leq 1$ for $\ell' > \ell > 1$. For the particular choice $\ell^* = (N+4)/(N2)$, we can use the expression of the asymptotic expansion of the very singular solution given in [1],

$$f_{\ell^*}(\eta) = C|\eta|^2 e^{-|\eta|^2/4} (1 + o(1)) \text{ as } |\eta| \rightarrow \infty,$$

from which follows $f_\ell(\eta) \geq f_{\ell^*}(\eta) \geq \delta^*(|\eta|^2 + 1)e^{-|\eta|^2/4}$ for some $\delta^* > 0$, any $\eta \in \mathbb{R}^N$ and $\ell \geq \ell^*$. Thus there exists $\delta > 0$ depending only on N such that

$$(10) \quad v_{\infty, \ell}(x, t) \geq \delta c^{1/(\ell-1)} t^{-1-N/2} (|x|^2 + t) e^{-|x|^2/4t} \quad \forall (x, t) \in \mathbb{R}^N \times (0, \infty).$$

Because any positive solution u of (4) satisfies (5), we have to prove that we can fix c and $\tau > 0$ such that

$$(11) \quad ct^{\alpha_\ell}(\rho^\ell + 1) \geq h(t)e^\rho \quad \forall (t, \rho) \in (0, \tau] \times [0, \tilde{V}(t)].$$

Writing h under the form $h(t) = -\omega'(t)e^{\omega(t)}$ where $\omega(t) = e^{\gamma(t)}$ and γ is a positive decreasing C^1 function, infinite at $t = 0$, we first notice that it is sufficient to prove this inequality for $\rho = \tilde{U}(t)$, and in that case

$$(12) \quad ct^{\alpha_\ell}(e^{\ell\gamma(t)} + 1) \geq -\gamma'(t)e^{\gamma(t)} \quad \forall t \in (0, \tau].$$

We take now $\gamma(t) = \sigma/t$, and prove that there exists $\beta > 0$, depending only on N such that, for any $0 < \tau \leq \beta\sigma$, estimate (11) holds with

$$c = e^{(1-\ell)\sigma/\tau - 2^{-1}(\ell(N+2)-N)\ln \tau}.$$

The maximum principle and (11) imply that for any $\ell > 1$ and $k > 0$ the solutions $u = u_k$ of (4) and $v = \tilde{v}_k$ of

$$\partial_t v - \Delta v + ct^{\alpha_\ell}(v^\ell + 1) = 0$$

with initial data $k\delta_0$ verifies $0 \leq \tilde{v}_{k, \ell} \leq u_k$, on $(0, \tau]$. Therefore $v_{\infty, \ell} \leq u_\infty + ct^{\alpha_\ell+1}/(\alpha_\ell + 1)$ on $(0, \tau]$ leads to

$$u_\infty(x, \tau) \geq \delta\tau^{-1-N/2} (|x|^2 + \tau) e^{\frac{4\sigma-|x|^2}{4\tau} - \frac{\ell(N+2)-N}{2(1-\ell)} \ln \tau}$$

Thus $\lim_{\tau \rightarrow 0} u_\infty(x, \tau) = \infty$, locally uniformly in $B_{2\sqrt{\sigma}}$, which implies the result.

The proof of Theorem 2 follows from the fact that for any $\sigma > \sigma' > 0$ there exists an interval $(0, \theta]$ where $\sigma't^{-2}e^{\sigma't^{-1}-e^{-\sigma'/t}} \geq e^{-\sigma'/t}$.

The single-point initial blow-up is proved by local energy methods. Because of their high degree of technicality we shall just give a short sketch of them in the simplest case of Theorem 1. For $k > 0$, let $u_k = u$ be the solution of the next result.

$$(13) \quad \begin{cases} \partial_t u - \Delta u + h(t)|u|^{q-1}u = 0 & \text{in } \mathbb{R}^N \times (0, \infty) \\ u(x, 0) = u_{0, k}(x) = M_k^{1/2} k^{-N/2} \eta_k(x) & \forall x \in \mathbb{R}^N, \end{cases}$$

where $\eta_k \in C(\mathbb{R}^N)$ is nonnegative, has compact support in $B_{k^{-1}}$, converges weakly to δ_0 as $k \rightarrow \infty$, and $\{M_k\}$ satisfies $\lim_{k \rightarrow \infty} k^{-N/2} M_k = \infty$. Furthermore it can be assumed that $\|\eta_k\|_{L^2} \leq c_0 k^{N/2}$. The single-point initial blow-up will be a consequence of

Lemma 5. For any $\delta > 0$ there exists $C = C(\delta)$ such that:

$$(14) \quad \sup_{t \in [0,1]} \int_{|x| \geq \delta} u_k^2(x,t) dx + \int_0^1 \int_{|x| \geq \delta} (|\nabla u_k|^2 + u_k^2) dx dt \leq C \quad \forall k \in \mathbb{N}.$$

Proof. For $r \in (0, 1)$, $\tau \geq 0$ we set $\Omega(\tau) = \{x \in \mathbb{R}^N : |x| > \tau\}$, $Q^r(\tau) = \Omega(\tau) \times (0, r]$, $Q_r(\tau) = \Omega(\tau) \times (r, 1)$ and $Q_r = \mathbb{R}^N \times (r, 1)$, and denote

$$I_1(r) = \iint_{Q_r} |\nabla u|^2 dx dt, \quad I_2(r) = \iint_{Q_r} u^2 dx dt, \quad I_3(r) = \iint_{Q_r} h(t) |u|^{q+1} dx dt.$$

If we multiply the equation by $u(x,t)e^{(r-t)/(2-r)}$, integrate on Q_r and use Hölder's inequality, we get, since h is nondecreasing,

$$(15) \quad \int_{\mathbb{R}^N} u^2(x, 1) dx + I_1(r) + I_2(r) + I_3(r) \leq c \int_{\mathbb{R}^N} u^2(x, r) dx \\ \leq c\tau^{\frac{N(q-1)}{q+1}} h(r)^{\frac{-2}{q+1}} (-I_3'(r))^{\frac{2}{q+1}} + c \int_{\Omega(\tau)} u^2(x, r) dx.$$

Let $\tau \mapsto \mu(\tau)$ be a smooth decreasing function, we define

$$E_1^\mu(r, \tau) = \iint_{Q^r(\tau)} (|\nabla u|^2 + \mu^2 u^2(x, t)) e^{-\mu^2 t} dx dt, \\ E_2(r, \tau) = \iint_{Q^r(\tau)} u^2 dx dt \quad \text{and} \quad f_\mu(r, \tau) = \sup\{e^{-\mu^2 t} \int_{\Omega(\tau)} u^2(x, t) dx : 0 \leq t \leq r\}$$

and $f(r) = f_0(r, 0)$. Then we introduce a parameter in the equation as in [4] by multiplying it by $u(x, t) \exp(-\mu^2(\tau)t)$ and integrating in the domain $Q^r(\tau)$ with $\tau > k^{-1}$ $Q^r(\tau)$ and $\tau > k^{-1}$. After some simple computations we deduce:

$$f_\mu(r, \tau) + 2E_1^\mu(r, \tau) \leq \frac{2}{\mu} \int_0^\tau \int_{\partial\Omega(\tau)} (|\nabla u|^2 + \mu^2 u^2(x, t)) e^{-\mu^2 t} dS dt \quad \forall \tau > k^{-1}.$$

Assuming $1 - 2\mu'/\mu^2 > 1/2$, we deduce from last inequality:

$$f_\mu(r, \tau) + E_1^\mu(r, \tau) \leq -\frac{2}{\mu(\tau)} \frac{dE_1^\mu(r, \tau)}{d\tau} \quad \forall \tau > k^{-1},$$

and by integration

$$f_\mu(r, \tau_2) \left(e^{\int_{\tau_1}^{\tau_2} \frac{\mu(\tau) d\tau}{2}} - 1 \right) + E_1^\mu(r, \tau_2) e^{\int_{\tau_1}^{\tau_2} \frac{\mu(\tau) d\tau}{2}} \leq E_1^\mu(r, \tau_1) \quad \forall \tau_2 > \tau_1 > k^{-1}.$$

The choice $\mu(\tau) = r^{-1}(\tau - k^{-1})/8$ ($\tau > k^{-1}$) yields to

$$(16) \quad \int_{\Omega(\tau)} u^2(x, r) dx + \iint_{Q^r(\tau)} \left(|\nabla_x u|^2 + \frac{(\tau - k^{-1})^2}{64r^2} u^2 \right) dx dt \leq c_1 e^{-\frac{(\tau - k^{-1})^2}{64r}} \\ \times \iint_{Q^r(\tau_0^k)} \left(|\nabla_x u|^2 + \frac{u^2}{2r} \right) dx dt \quad \forall \tau \geq \tilde{\tau}_0^k := k^{-1} + 8\sqrt{r} > \tau_0^k := k^{-1} + 4\sqrt{2r}.$$

We will need standard global energy estimate of solution of problem (13) too:

$$(17) \quad \int_{\mathbb{R}^N} |u(x, r)|^2 dx + \int_{Q^r} (|\nabla_x u|^2 + |u|^2 + h(t)|u|^{q+1}) dx dt \\ \leq c \|u_{0,k}\|_{L_2(\mathbb{R}^N)}^2 \leq \bar{c} M_k \quad \forall r > 0.$$

Estimating the right-hand side terms in (15) and (16) by (17), we derive:

$$(18) \quad \begin{aligned} (i) \quad & \sum_{i=1}^3 I_i(r) \leq c_1 \tau^{\frac{N(q-1)}{q+1}} h(r)^{\frac{-2}{q+1}} (-I_3'(r))^{\frac{2}{q+1}} + \frac{c_2 M_k}{r} e^{-(\tau-k^{-1})^2/64r} \quad \forall \tau \geq \tilde{\tau}_0^k(r) \\ (ii) \quad & f_0(r, \tau) + E_1^0(r, \tau) + \frac{\tau - k^{-1}}{64r^2} E_2(r, \tau) \leq \frac{c_2 M_k}{r} e^{-(\tau-k^{-1})^2/64r} \quad \forall \tau \geq \tilde{\tau}_0^k(r). \end{aligned}$$

Next we choose $M_k = e^{e^k}$, fix $\epsilon_0 \in (0, e^{-1})$ and define a pair (r_k, τ_k) by the following relations: $r_k = \sup\{r : I_1(r) + I_2(r) + I_3(r) > 2M_k^{\epsilon_0}\}$; $c_2 r_k^{-1} \exp(-\frac{\tau_k^2}{64r_k}) M_k = M_k^{\epsilon_0} \Leftrightarrow \tau_k = 8\sqrt{r_k(1-\epsilon_0)e^k + \ln(c_2/r_k)}$. Taking $\tau = \tau_k + k^{-1}$ in (18)-i and solving the corresponding O.D.E. yields the estimate:

$$(19) \quad \sum_{i=1}^3 I_i(r) \leq c_3(\tau_k + k^{-1})(H(r))^{-2/(q-1)} \quad \forall r \leq r_k, \quad H(r) = \int_0^r h(s)ds.$$

If we write $h(t) = e^{-\omega(t)/t}$, the assumption $\inf\{\omega(t)/t^\alpha : 0 < t \leq 1\} > 0$ implies that $H(r) \geq c_0 e^{-\omega(r)/r} r^2 / \omega(r)$ and, replacing τ_k by its expression, (19) turns into

$$\sum_{i=1}^3 I_i(r) \leq c_4 \left(\sqrt{r_k(1-\epsilon_0)e^k + \ln(c_2/r_k)} + k^{-1} \right)^N \left(\frac{\omega(r)e^{-\omega(r)/r}}{r^2} \right)^{2/(q-1)} \quad \forall r \leq r_k.$$

Thus $r_k \leq b_k$, where b_k is solution of equation:

$$c_4 \left(\sqrt{r_k(1-\epsilon_0)e^k + \ln(c_2/b_k)} + k^{-1} \right)^N \left(\frac{\omega(b_k)e^{-\omega(b_k)/b_k}}{b_k^2} \right)^{2/(q-1)} = 2M_k^{\epsilon_0} = 2e^{\epsilon_0 e^k}.$$

From this inequality using additionally assumption on $\omega(t)$, we obtain inequalities: $c_5 e^k \geq \omega(b_k)/b_k \geq c_6 e^k$, $c_6 > 0$; $b_k \geq e^{-c_7 k}$, $c_7 > 0$. These inequalities yield:

$$(20) \quad \tau_k \leq c_8 \sqrt{\omega(c_9 e^{-k})}.$$

Using the definition of r_k , inequality (18)-ii, the fact that $3M_k^{\epsilon_0} \leq \bar{c}M_{k-1} \quad \forall k \geq k_0(\bar{c})$ (\bar{c} is from (17), $0 < \epsilon_0 < e^{-1}$), we deduce the main result of first round of computations:

$$(21) \quad \sum_{i=1}^3 I_i(r_k) + f_0(r_k, \tau_k + k^{-1}) + \sum_{i=1}^2 E_i(r_k, \tau_k + k^{-1}) \leq 3M_k^{\epsilon_0} \leq \bar{c}M_{k-1}.$$

Next we organize the second round of estimates with $\mu(\tau) = (\tau - \tau_k - k^{-1})/8$, r_{k-1} and τ_{k-1} be defined similarly as r_k and τ_k , up to the change of indices, using obtained estimate (21) instead of (17). As result we derive:

$$(22) \quad \sum_{i=1}^3 I_i(r_{k-1}) + f_0(r_{k-1}, \tau_k + \tau_{k-1} + k^{-1}) + \sum_{i=1}^2 E_i(r_{k-1}, \tau_k + \tau_{k-1} + k^{-1}) \leq \bar{c}M_{k-2}.$$

Fixing arbitrary $n > k_0(\bar{c})$ and repeating the above described round of computations $k - n$ times, we obtain:

$$(23) \quad \sum_{i=1}^3 I_i(r_n) + f_0 \left(r_n, \sum_{j=0}^{k-n} \tau_{k-j} + k^{-1} \right) + \sum_{i=1}^2 E_i \left(r_n, \sum_{j=0}^{k-n} \tau_{k-j} + k^{-1} \right) \leq \bar{c}M_{n-1},$$

and, since by induction τ_{k-j} satisfies (20) with k replaced by $k-j$, we obtain

$$(24) \quad \sum_{j=0}^{k-n} \tau_{k-j} \leq c_8 \sum_{j=0}^{k-n} \sqrt{\omega(c_9 e^{-(k-j)})} \leq c_{10} \int_{c_9 e^{-k}}^{c_9 e^{-n}} \frac{\sqrt{\omega(s)} ds}{s}.$$

We denote $\tau^*(n) = \lim_{k \rightarrow \infty} c_8 \sum_{j=0}^{k-n} \sqrt{\omega(c_9 e^{-(k-j)})}$. We derive from (23) by letting $k \rightarrow \infty$,

$$(25) \quad \sup_{0 < t \leq r_n} \int_{|x| \geq \tau^*(n)} u^2(x, t) dx + \int_0^{r_n} \int_{|x| \geq \tau^*(n)} (|Du|^2 + u^2) dx dt \leq \bar{c} M_{n-1}.$$

Due to assumption (4) $\tau^*(n) \rightarrow 0$ as $n \rightarrow \infty$, therefore inequality (25) implies the result.

REFERENCES

- [1] Brezis H, Peletier L. A. and Terman D. A very singular solution of the heat equation with absorption, Arch. Rat. Mech. Anal. **96** (1985), 185-209.
- [2] Galaktionov V. A. and Shishkov A.E., *Saint-Venant's principle in blow-up for higher-order quasilinear parabolic equations*, Proc. Roy. Soc. Edinburgh Sect. A **133** (2003), 1075-1119.
- [3] Marcus M. and Véron L., *Initial trace of positive solutions to semilinear parabolic inequalities*, Adv. Nonlinear Studies **2** (2002), 395-436.
- [4] Oleinik O. A. and Radkevich E.V., *Method of introducing of a parameter in evolution equation*, Russian Math. Survey **33** (1978), 7-74.
- [5] Peletier L. A. and Terman D., *A very singular solution of the porous media equation with absorption*, J. Diff. Equ. **65** (1986), 396-410.
- [6] Shishkov A. E., *Propagation of perturbation in a singular Cauchy problem for degenerate quasilinear parabolic equations*, Sbornik: Mathematics **187:9** (1996), 1391-1440.
- [7] Shishkov A. E., *Dead cores and instantaneous compactification of the supports of energy solutions of quasilinear parabolic equations of arbitrary order*, Sbornik: Mathematics **190:12** (1999), 1843-1869.