



HAL
open science

Modeling secondary organic aerosol formation through cloud processing of organic compounds

J. Chen, R. J. Griffin, A. Grini, Pierre Tulet

► **To cite this version:**

J. Chen, R. J. Griffin, A. Grini, Pierre Tulet. Modeling secondary organic aerosol formation through cloud processing of organic compounds. *Atmospheric Chemistry and Physics Discussions*, 2007, 7 (3), pp.8951-8982. hal-00328209

HAL Id: hal-00328209

<https://hal.science/hal-00328209>

Submitted on 18 Jun 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Modeling in-cloud
SOA formation

J. Chen et al.

Modeling secondary organic aerosol formation through cloud processing of organic compounds

J. Chen¹, R. J. Griffin^{1,2}, A. Grini³, and P. Tulet⁴

¹Institute for the Study of Earth, Oceans, and Space, University of New Hampshire, Durham, NH, USA

²Department of Earth Sciences, University of New Hampshire, Durham, NH, USA

³Department of Geoscience, University of Oslo, Pb 1022 Blindern, 0315, Oslo, Norway

⁴CNRM/GMEI Météo France, Toulouse, France

Received: 30 May 2007 – Accepted: 11 June 2007 – Published: 26 June 2007

Correspondence to: J. Chen (jianjunc@cisunix.unh.edu)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

Abstract

Interest in the potential formation of secondary organic aerosol (SOA) through reactions of organic compounds in condensed aqueous phases is growing. In this study, the potential formation of SOA from irreversible aqueous-phase reactions of organic species in clouds was investigated. A new proposed aqueous-phase chemistry mechanism (AqChem) is coupled with the existing gas-phase Caltech Atmospheric Chemistry Mechanism (CACM) and the Model to Predict the Multiphase Partitioning of Organics (MPMPO) that simulate SOA formation. AqChem treats irreversible organic reactions that lead mainly to the formation of carboxylic acids, which are usually less volatile than the corresponding aldehydic compounds. Zero-dimensional model simulations were performed for tropospheric conditions with clouds present for three consecutive hours per day. Zero-dimensional model simulations show that 48-h averaged SOA formation are increased by 27% for a rural scenario with strong monoterpene emissions and 7% for an urban scenario with strong emissions of aromatic compounds, respectively, when irreversible organic reactions in clouds are considered. AqChem was also incorporated into the Community Multiscale Air Quality Model (CMAQ) version 4.4 with CACM/MPMPO and applied to a previously studied photochemical episode (3–4 August 2004) focusing on the eastern United States. The CMAQ study indicates that the maximum contribution of SOA formation from irreversible reactions of organics in clouds is $0.28 \mu\text{g m}^{-3}$ for 24-h average concentrations and $0.60 \mu\text{g m}^{-3}$ for one-hour average concentrations at certain locations. On average, domain-wide surface SOA predictions for the episode are increased by 8.6% when irreversible, in-cloud processing of organics is considered.

1 Introduction

Atmospheric particulate matter (PM) is associated with adverse human health effects (Pope and Dockery, 2006), decreases in visibility (Malm, 1989), and global climate

Modeling in-cloud SOA formation

J. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

forcing (Charlson et al., 1992, Jones et al., 1994). Organic aerosol (OA) is a ubiquitous and important constituent of atmospheric PM (Murphy et al., 1998; Turpin et al., 2000). OA consists of primary OA (POA), which is directly emitted from various sources, and secondary OA (SOA), which is formed in the atmosphere from the oxidation of volatile organic compounds (VOCs).

The processes leading to SOA formation typically are viewed as gas-phase oxidation of VOCs followed by nucleation/gas-particle partitioning of low-volatility products (Pankow, 1994; Odum et al., 1996; Griffin et al., 1999). Oligomer and/or polymer formation in the aerosol phase has also been shown to be important with regard to SOA formation (Jang et al., 2002; Kalberer et al., 2004). Besides these processes, it has also been hypothesized that fog/cloud processing, which leads to substantial sulfate formation, could also be a source for SOA (Blando and Turpin, 2000). The process of SOA formation from aqueous-phase processing of organic compounds in clouds involves formation of potential low-volatility products through aqueous-phase reactions of organics, followed by subsequent gas-particle partitioning of the low-volatility products post evaporation of the hydrometeor (Blando and Turpin, 2000; Kanakidou et al., 2005). Recent experimental and modeling studies indeed demonstrated that SOA potentially can be formed from aqueous-phase processing of organic compounds in clouds (Carlton et al., 2006; Loeffler et al., 2006). Ambient particle size distribution measurements also showed the occurrence of droplet-mode organics (Blando et al., 1998; Yao et al., 2002), which, similarly to droplet-mode sulfate, are most likely formed from cloud processing of organic materials (Blando and Turpin, 2000). Several modeling studies (Warneck, 2003; Ervens et al., 2004; Lim et al., 2005) have demonstrated the formation of low-molecular weight dicarboxylic acid from cloud processing of organics. Such dicarboxylic acids have been found in atmospheric aerosols in various regions (Kawamura and Ikushima, 1993; Decesari et al., 2000). In addition, Claeys et al. (2004) showed that multiphase acid-catalyzed organic reactions with hydrogen peroxide provided a new route for SOA formation from isoprene and hypothesized that such a mechanism could also provide a pathway for SOA formation from monoterpenes

**Modeling in-cloud
SOA formation**

J. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

and their oxidation products. Most recently, Heald et al. (2006) analyzed the covariance of water soluble particulate organics with other species in the free troposphere over the eastern United States, with the results suggesting aqueous-phase SOA generation involving biogenic precursors.

5 Given the increasing interest in potential SOA formation through the aqueous-phase processing of organic compounds, this paper evaluates the significance of SOA formation from aqueous-phase reactions of organics in cloud droplets through a modeling study. The study involves the development of an aqueous-phase chemistry mechanism (AqChem) that specifically includes treatment of potential organic reactions that
10 lead to production of semi-volatile organic compounds. Because of our incomplete knowledge on the aqueous-phase chemistry for organics (Kanakidou et al., 2005; Gencsler and Varga, 2005), AqChem treats organic reactions based on previous work by Ervens et al. (2004) and Lim et al. (2005) for organics of carbon number up to four and the simple protocol used by Aumont et al. (2000) for organics of carbon number
15 greater than four. Potential reversible oligomerization reactions (Hastings et al., 2005; Loeffler et al., 2006) are not included because of a current lack of a quantitative description of these phenomena. AqChem utilizes an existing gas-phase chemistry mechanism, the Caltech Atmospheric Chemistry Mechanism (CACM) (Griffin et al., 2002) that provides the prediction of gas-phase production of organic compounds. It
20 is also linked with a SOA module, the Model to Predict the Multi-phase Partitioning of Organics (MPMPO) (Griffin et al., 2003), such that the partitioning between gas and aerosol phases of semi-volatile organic products from aqueous-phase reactions is determined thermodynamically. AqChem was used in a zero-dimensional model as well as a three-dimensional model to study the magnitude of potential SOA formation in
25 clouds.

**Modeling in-cloud
SOA formation**J. Chen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2 Methods

2.1 Development of the AqChem mechanism

AqChem was developed based on previous work (Aumont et al., 2000; Warneck et al., 2003; Ervens et al., 2004; Lim et al., 2005) and to work specifically with the CACM mechanism and the MPMPO module. The main purpose of AqChem is to treat the irreversible organic reactions that could potentially lead to the production of semi-volatile organic compounds in clouds.

In AqChem, the reactions for organics of carbon number up to four that are treated in the CACM mechanism were adopted from previous work by Ervens et al. (2004) and Lim et al. (2005) and are shown schematically in Fig. 1a. The main purpose of the reaction scheme for organics of carbon number up to four is to provide formation routes for oxalic and pyruvic acids. Briefly, organics in the cloud phase are assumed to be oxidized only by hydroxyl radical (OH). Glycoladehyde and glyoxal are converted to oxalic acid via glyoxylic acid (Ervens et al., 2004; Lim et al., 2005). Methylglyoxal is converted to pyruvic acid, which decomposes to form acetic acid. Ethanol and acetaldehyde are converted to acetic acid as well. A portion of acetic acid is converted to oxalic acid, while the rest decomposes to formaldehyde (Lim et al., 2005). No reactions are treated for methyl vinyl ketone and methacrolein in the aqueous phase because of their low solubility in water (Ervens et al., 2004).

Within CACM, there are approximately 140 semi-volatile organic compounds of carbon number greater than four (Griffin et al., 2002; Chen and Griffin, 2005). Instead of treating potential aqueous-phase chemistry for these organic compounds individually, an organic surrogate approach similar to that used in MPMPO was adopted here. In the MPMPO module, each semi-volatile organic compound is lumped into one of 11 organic surrogate species (Griffin et al., 2003; Griffin et al., 2005; Chen et al., 2006). MPMPO calculates the partitioning between the gas and aerosol phases for these 11 surrogate species. Similarly, AqChem treats the aqueous-phase chemistry for four of the 11 surrogates because of their potential to lead to less volatile organic

Modeling in-cloud SOA formation

J. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Modeling in-cloud
SOA formation**

J. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

products (e.g., carboxylic acids). These four surrogate species are surrogate 3 (S3, 3-hydroxy-2, 4-dimethyl-2, 4-hexadienal), surrogate 5 (S5, 2-hydroxyl-3-isopropyl-6-keto-heptanal), surrogate 7 (S7, 2,4-dimethyl-3-formyl-benzoic acid), and surrogate 10 (S10, 1-methyl-1-hydroxy-2-nitrato-4-isopropyl-cyclohexane) in the MPMPO module (Griffin et al., 2003; Griffin et al., 2005). In this way, although only four aqueous-phase chemical reactions were added for organic compounds of carbon number greater than four, 58 organic compounds that are lumped into these four surrogates in the MPMPO undergo further reactions in the aqueous phase. The products of aqueous-phase reactions for these four MPMPO surrogates were based on the protocol used by Aumont et al. (2000). Aumont et al. (2000) assumed that (1) OH is the only oxidant; (2) an aldehyde moiety is converted to a carboxylic acid moiety; and (3) the carbon skeleton is not changed during oxidation. Following the protocol of Aumont et al. (2000), the oxidation of S3 by OH leads to the formation of 3-hydroxy-2, 4-dimethyl-2, 4-hexadienalic acid (S3PD). The oxidation product of S5 is 2-hydroxyl-3-isopropyl-6-keto-heptanoic acid (S5PD). S7 is converted to 3, 5-dimethyl-1, 4-dibenzoic acid (S7PD). Oxidation of S10 by OH in the aqueous phase leads to a product (S10PD, 1-methyl-1-hydroxy-2-nitrato-3-oxo-4-isopropyl-cyclohexane) that has one more ketone group than surrogate 10. The aqueous-phase oxidation rates by OH for these surrogate species were estimated based on the method of Dutot et al. (2003). The reaction scheme for these surrogate species is shown in Fig. 1b.

AqChem also includes a compact treatment of aqueous-phase hydrogen-oxygen and sulfur chemistry mainly based on Ervens et al. (2003) and Ervens et al. (2004). Such a compact treatment includes the important reactions for determining aqueous OH levels and aqueous sulfate formation but is computationally less demanding than detailed treatments, which is especially important for three-dimensional air quality simulations. Tables 1 and 2 list the irreversible and equilibrium reactions, respectively, that are included in AqChem. Gas and aqueous-phase mass transfer was treated according to the resistance model of Schwartz (1986) by considering mass accommodation coefficient (α), gas-phase diffusion constants (D_g), and Henry's law constants (K_H) for each

soluble species, values of which are shown in Table 3.

After the evaporation of cloud droplets, the partitioning between the gas and aerosol phases for the oxidation products formed from the aqueous-phase reactions is computed by the MPMPO module. This is shown schematically in Fig. 2. If clouds are not present, the amounts of semi-volatile organic compounds computed from the CACM mechanism are used directly as input to the MPMPO module to compute SOA formation, as in previous applications (Chen et al., 2006). If clouds are present, AqChem is first called to simulate the cloud processing of organics, followed by a call to MPMPO to calculate SOA formation. The organic oxidation products in clouds were lumped into existing surrogates in the MPMPO module. Oxalic and pyruvic acids were lumped into surrogate 1 (oxalic acid) of MPMPO. S3PD, S5PD, S7PD, and S10PD were lumped into surrogate 2 (S2, 2-methyl-5-formyl-2, 4-hexadiendioic acid), surrogate 4 (S4, 2-hydroxy-3-isopropyl-5-keto-3-hexenoic acid), surrogate 6 (S6, 3, 5-dimethyl-2-nitro-4-hydroxy-benzoic acid), and surrogate 5 (S5, 2-hydroxy-3-isopropyl-6-keto-heptanal) of MPMPO, respectively. Overall, the aqueous products S3PD, S5PD, S7PD, and S10PD are less volatile than their precursors, as in the treatment used by Gelencser and Varga (2005). Vapor pressure and K_H estimates for the surrogate species in MPMPO can be found in Pun et al. (2002) and Griffin et al. (2005).

The kinetic processor KPP2.1 (Damian et al., 2002; Daescu et al., 2003; Sandu et al., 2003) was used to process AqChem and generate the associated numerical codes. A Rosenbrock solver in KPP2.1 was used to solve the differential equations associated with the mechanism (Sandu et al., 1997).

2.2 Zero-dimensional model simulations

A zero-dimensional model was constructed to investigate potential SOA formation due to irreversible aqueous-phase organic chemistry under scenarios for which cloud is present for three consecutive hours per day (Lim et al., 2005). The zero-dimensional model considers emissions, dry deposition, and gas- and aqueous-phase chemistry for chemical species. The mass balance of the gas- and aqueous-phase species is

Modeling in-cloud SOA formation

J. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

described by (Pandis and Seinfeld, 1989; Lim et al., 2005)

$$\frac{dC_g}{dt} = Q_g - S_g + \frac{E}{Z} - \frac{\nu C_g}{Z} - LkC_g + \frac{LkC_a}{K_H RT} \quad (1)$$

$$\frac{dC_a}{dt} = Q_a - S_a + kC_g - \frac{kC_a}{K_H RT} \quad (2)$$

where C_g is the gas-phase concentration (mole L^{-1} of air), C_a is the aqueous-phase concentration (M), Q_g (mole L^{-1} of air sec^{-1}) and Q_a (M sec^{-1}) are the gas-phase and aqueous-phase production rate, respectively, S_g (mole L^{-1} of air sec^{-1}) and S_a (M sec^{-1}) are the gas-phase and aqueous-phase loss rate, respectively, E (mole dm^{-2} sec^{-1}) is the emission rate of the gas-phase species, Z (dm) is the boundary layer height, ν (dm sec^{-1}) is the dry deposition velocity, L is the cloud liquid water volume fraction, K_H (M atm^{-1}) is the effective Henry's law constant, R (L atm $mole^{-1}$ K^{-1}) is the ideal gas constant, T (K) is the temperature, and k (sec^{-1}) is the mass transfer coefficient,

$$k = \left(\frac{a^2}{3D_g} + \frac{4a}{3v\alpha} \right)^{-1} \quad (3)$$

where α is the mass accommodation coefficient, D_g (cm^2 s^{-1}) is the gas-phase diffusion constant, a is the cloud droplet radius (cm), and v is the mean molecular speed (cm s^{-1}).

The zero-dimensional model couples CACM, AqChem, and MPMPO. Emissions and dry deposition of gas-phase species were implemented within the gas-phase chemistry mechanism. As with AqChem, the latest version of CACM with emissions and dry deposition of species was processed by KPP2.1, and a Rosenbrock solver was used for solving the differential equations. If clouds are present, the simulations of gas-phase and aqueous-phase chemistry processes are split every 12 min, following the treatment

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

in the three-dimensional model that will be discussed subsequently. If clouds are not present, only gas-phase chemistry is simulated every 12 min. At the end of every 12 min, MPMPO is called to simulate SOA formation.

For the zero-dimensional model simulation, conditions are based on the study of Lim et al. (2005). The boundary layer height was assumed to be fixed at 1000 m. Temperature and relative humidity (RH) are 298 K and 75%, respectively, if clouds are not present. Clouds are assumed to be present through the entire boundary layer for hours 13–16 of each day. Cloud water content was assumed to be 0.5 g m^{-3} , with a uniform cloud droplet diameter of $10 \mu\text{m}$. Temperature and RH are assumed to be 288 K and 100%, respectively, when clouds are present. In addition, a constant cloud pH of 4.5 was assumed. Photolysis rates were assumed to vary semisinusoidally between hour 6 and hour 18 of each day, with a peak at hour 12.

Two emissions scenarios were studied. Scenario 1 features strong biogenic monoterpene emissions for a rural area, and Scenario 2 features strong anthropogenic emissions of aromatic compounds for an urban area. Emission rates and deposition velocities of gas-phase hydrocarbons were extracted from previous three-dimensional model simulations (Chen et al., 2006; Mao et al., 2006). These are shown in Tables 4 and 5, respectively. In order to simulate SOA formation in MPMPO, total POA concentrations (fractions of individual POA species are consistent between the two scenarios and are based on those used in Chen et al., 2006), aerosol water content, and aerosol pH for both scenarios were assumed to be constant at $1.0 \mu\text{g m}^{-3}$, $30 \mu\text{g m}^{-3}$, and 3.0, respectively. This simplified treatment omits considerations of emission, deposition, and size distribution of aerosols. In addition, the zero-dimensional model simulations were performed for four days for each scenario. The first two days were used as initialization days, while the last two days were used for analysis.

2.3 Three-dimensional model simulation

AqChem was incorporated into the Community Multiscale Air Quality (CMAQ) model version 4.4 with CACM and MPMPO (Chen et al., 2006). In addition, the computa-

Modeling in-cloud SOA formation

J. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

tionally efficient version of the MPMPO module (Tulet et al., 2006) was used. CMAQ originally employed the RADM aqueous-phase chemistry mechanism, mainly for the purpose of predicting aqueous-phase sulfate formation in grid resolved clouds and/or sub-grid convective clouds (Byun and Ching, 1999). In the CMAQ model, the aqueous-phase chemistry mechanism is called when the liquid water content of clouds exceeds $10^{-5} \text{ kg m}^{-3}$. Cloud droplet diameter was assumed to be $10 \mu\text{m}$ (Lim et al., 2005). In addition, CMAQ assumes that accumulation mode particles serve as cloud condensation nuclei and Aitken mode particles form interstitial aerosol that can be scavenged by clouds. Therefore, all accumulation mode particles and some fraction of Aitken mode particles are included in the resulting aqueous phase of the hydrometeors. Chemical species (e.g., nitrate, oxalic acid, S3, S5, S7, and S10) of cloud-incorporated particles are also subject to aqueous-phase chemistry. SOA formation due to both the gas-phase oxidation and the aqueous-phase oxidation is distributed between accumulation and Aitken mode particles using the proportion of preexisting OA in these two modes.

CMAQ with CACM and MPMPO was applied previously to an episode (3–4 August 2004 with spin-up days of 1–2 August 2004) over the eastern United States (Chen et al., 2006). In this study, CMAQ with CACM and MPMPO and with the newly developed AqChem was applied to the same episode to study the significance of SOA formation from the consideration of irreversible aqueous-phase organic chemistry in clouds. Model inputs (e.g., emissions and meteorological fields) and model configurations (e.g., model domain and vertical layers) for this episode are described in Mao et al. (2006) and Chen et al. (2006).

3 Results

3.1 Zero-dimensional model simulations

Figure 3a shows the SOA predictions from the zero-dimensional model simulation for Scenario 1, which features strong monoterpene emissions for a rural area. Averaged

Modeling in-cloud SOA formation

J. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

over 48 h, total SOA prediction considering SOA formation from cloud-phase organic reactions is $5.7 \mu\text{g m}^{-3}$, which is approximately 27% higher than the prediction without considering aqueous-phase chemistry in clouds. At the end of simulation, total SOA prediction with aqueous-phase chemistry in clouds is $8.7 \mu\text{g m}^{-3}$, which is 32% more than that from the simulation without clouds. Compared to the SOA prediction without consideration of the aqueous-phase chemistry, total SOA prediction with aqueous-phase chemistry increases substantially during the hours 13–16 and hours 37–40, during which clouds are present. This suggests substantial rapid processing of organic compounds in cloud droplets and that such processing leads to formation of SOA.

As shown in Fig. 3a, the increase of SOA prediction from aqueous-phase chemistry is due primarily to the increase of SOA formation from S4 and S5 of MPMPPO, which is attributed to the oxidation of S5 to form S5PD (lumped to S4) and the oxidation of S10 to form S10PD (lumped to S5) in AqChem, respectively. Because S4 and S5 are less volatile and more soluble than S5 and S10, respectively, more SOA is formed. Both S5 and S10 are surrogates for oxidation products of monoterpenes.

Figure 3b shows the SOA production from Scenario 2 using the same zero-dimensional model. Scenario 2 represents an urban environment with strong anthropogenic emissions (e.g., nitrogen oxides (NO_x) and aromatic compounds). If aqueous-phase chemistry in clouds is considered, the SOA concentrations predicted at the end of the simulation period and averaged over 48 hours are $9.23 \mu\text{g m}^{-3}$ and $6.33 \mu\text{g m}^{-3}$, respectively, which are 11% and 7% higher than that from simulation without consideration of cloud chemistry. Individually, the increase of SOA prediction by considering aqueous-phase organic chemistry is due to the increase of SOA contributions from S2, S6, and, to a smaller extent, S1, which is then attributed to the aqueous-phase conversion of S3, S7, and organic compounds of carbon number less than four. S2, S6, and the majority of S1 are surrogates for oxidation products from aromatic compounds. Based on the aqueous-phase organic chemistry proposed in this study, the effect of aqueous-phase chemistry in clouds on SOA formation is less important for aromatic compounds than it is for monoterpene species.

**Modeling in-cloud
SOA formation**

J. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3.2 Three-dimensional model simulation

Figure 4 shows the difference between 24-hour average SOA predictions at the surface for 3–4 August 2004 (UTC) when the SOA formation from aqueous-phase organic chemistry in cloud droplets is considered and not. In addition, the absolute values of SOA predictions at the surface layer for these two days are also shown if the SOA formation from aqueous-phase organic chemistry in clouds is considered. Consideration of SOA formation from aqueous-phase organic chemistry increases 24-h average SOA predictions. The maximum difference in 24-h average SOA predictions within the domain is $0.28 \mu\text{g}/\text{m}^3$ for both days. Averaged over the entire domain, the relative increases of 24-h average SOA predictions with consideration of SOA formation from aqueous-phase organic chemistry in clouds are 8.7% and 8.5% for 3 August and 4 August 2004, respectively.

The increase of SOA predictions if aqueous-phase organic chemistry is considered is due mainly to the cloud processing of S5 and S10, which leads to the formation of less volatile products. Both S5 and S10 are lumped species for oxidation products of monoterpenes. This is consistent with the fact that the majority of SOA predicted within the domain is from biogenic monoterpenes (Chen et al., 2006).

The average relative increase across the domain is smaller than that from the zero-dimensional model simulation for Scenario 1 in which strong monoterpene emissions are considered. This is expected because the zero-dimensional model simulation represents an ideal situation in which a site is influenced by biogenic monoterpene emissions and is exposed to clouds for three hours in the afternoon (12.5% of the time of each day). Examination of the meteorological inputs for the chosen episode indicates that resolved clouds occupy approximately 10–20% of surface area during the episode. In addition, CMAQ internally calculates sub-grid convective clouds. Therefore, on average, each grid has the probability of 10–20% of being exposed to cloud processing, which is close to the cloud exposure time used in the zero-dimensional model. However, locations influenced strongly by monoterpene emissions may not be

Modeling in-cloud SOA formation

J. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

necessarily exposed to cloud processing. This is believed to be the major reason for a smaller relative SOA increase in the three-dimensional model simulation compared to the zero-dimensional model simulations if SOA formation from aqueous-phase organic chemistry in clouds is considered.

5 Figure 5 indicates the time series of the SOA predictions with or without consideration of SOA formation from the aqueous-phase organic chemistry in clouds at the two sites which have the maximum difference in 24-h average SOA predictions for 3 August and 4 August 2004, respectively. At these two sites, the relative difference in SOA prediction with or without consideration of SOA formation from aqueous-phase organic chemistry are as high as 36% and 43%, respectively, indicating that accounting for SOA formation from aqueous-phase organic chemistry in clouds could be important at these locations. As shown in Fig. 5, the increase of SOA formation by considering aqueous-phase organic chemistry is due mainly to the increase of SOA formation from S4 and S5, which is again attributed to the aqueous-phase oxidation of S5 and S10, which are surrogates for oxidation products of monoterpenes. This implies that the increase of SOA formation with consideration of aqueous-phase organic chemistry in clouds is due to the cloud processing of monoterpene oxidation products. These two sites are influenced by transport of monoterpene or monoterpene oxidation products from the New England region. In addition, these two sites are also exposed to resolved clouds almost throughout the entire episode. Therefore, SOA formation from aqueous-phase organics can be important when a site is influenced by biogenic monoterpene emissions or products and also exposed to cloud processing for significant time periods.

25 An interesting feature of Fig. 5 is that the increase in the difference in the SOA predictions usually occurs during afternoon hours. This is a direct result of OH being the only oxidant considered in the aqueous phase. Typically, OH concentrations in clouds reach a peak in the early afternoon (Ervens et al., 2003), which facilitates the conversion of organic compounds in cloud droplets and further formation of SOA. Differences in SOA prediction with or without consideration of SOA formation through cloud pro-

**Modeling in-cloud
SOA formation**

J. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

cessing was also examined for three specific sites: Thompson Farm, New Hampshire (semi-rural), Boston, Massachusetts (urban), and Appledore Island, Maine (marine area influenced by continental outflow) and was found to be approximately 5%. These sites were chosen because of different emissions patterns and relative influences of primary emissions and secondary chemistry.

Figure 6 indicates additional SOA formation due to the in-cloud organic chemistry at layer 14 of the modeling domain. In addition, for comparative purposes, the total SOA predictions from the simulations with consideration of SOA formation from aqueous-phase organic chemistry are also shown for layer 14. The CMAQ modeling domain includes 21 vertical layers using a σ -pressure system extending from the surface to 10 000 Pa. Although the actual altitude for layer 14 varies, according to the U.S. standard atmosphere, layer 14 corresponds to the altitude between 1800–2300 meters. Layer 14 thus roughly represents the lower free troposphere. The maximum additional SOA formation due to aqueous-phase organic chemistry in clouds is $0.25 \mu\text{g m}^{-3}$ for both days, only slightly smaller than the maximum difference at the surface. Similar to the surface layer, the increase of SOA predictions is due primarily to cloud processing of gas-phase monoterpene oxidation products, indicating the potential of SOA formation from an aqueous-phase mechanism involving biogenic precursors as suggested by Heald et al. (2006). Averaged across the modeling domain for layer 14, the relative difference in SOA predictions is 18.8% and 15.3%, respectively for 3 August and 4 August, which is much larger than the relative difference at the surface layer. This is due to the fact that less SOA is predicted to form from gas-phase oxidation of organic compounds in the free troposphere (Heald et al., 2005).

4 Discussion

An aqueous-phase chemistry mechanism was developed to study the potential of SOA formation in cloud droplets. The AqChem mechanism incorporates a compact treatment of hydrogen-oxygen and sulfur chemistry but treats in more detail organic chem-

Modeling in-cloud SOA formation

J. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

istry based on recent work of Aumont et al. (2000), Ervens et al. (2004), and Lim et al. (2005) in an effort to represent the potential irreversible organic reactions that may lead to low-volatility products. AqChem was coupled to the existing CACM mechanism and MPMPO module. Zero-dimensional model simulations indicated that consideration of SOA formation from organic chemistry in cloud droplets could lead to increases of SOA prediction by approximately 27% and 7% for a rural scenario with strong monoterpene emissions and an urban scenario with strong aromatic compound emissions, respectively. Three-dimensional simulations showed an average of 8–9% increase in the SOA predictions at the surface when the aqueous-phase organic chemistry in clouds is considered. However, the relative increase of SOA prediction at certain locations could be much higher than the domain-wide, averaged surface increase. Overall, based on the proposed AqChem, the modeling work in this study suggests that SOA formation due to aqueous-phase organic chemistry in clouds could be important at locations that are strongly influenced by monoterpene emissions and where clouds are present for significantly long time, especially in the afternoon when photochemistry is strongest.

A key limitation of this study is the simple treatment of organic reactions in the aqueous phase for organic compounds of carbon number greater than four. Herrmann et al. (2005) presented a much more detailed treatment of organic chemistry in their aqueous-phase chemistry mechanism, although such a mechanism is probably too complex for incorporation into three-dimensional models. More laboratory studies are also needed to characterize the mechanism of aqueous-phase reactions, especially for intermediate gas-phase oxidation products, which are soluble in water and have the potential to lead to low-volatility products.

Another limitation is the lack of treatment of potentially reversible oligomerization processes, especially for small molecules like glyoxal and methyl glyoxal (Hastings et al., 2005; Loeffler et al., 2006). These processes are not included because of a current lack of a quantitative description.

In addition, future studies need to consider the aqueous-phase processing of organic compounds in deliquescent particles. Unlike short-lived clouds, deliquescent particles

**Modeling in-cloud
SOA formation**

J. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

provide additional time for processing organic compounds and could be more likely to lead to SOA, despite having significantly smaller water contents (Claeys et al., 2004). Such treatment would need to consider the interactions of different ions and molecules in the particles, which poses significantly more difficulty than processes in the cloud droplets.

Despite these limitations, the method used in this study provides a framework to study SOA formation due to aqueous-phase processing of organic compounds and underscores the potential of SOA formation via cloud processing of organic compounds. New findings of aqueous-phase organic chemistry should be incorporated into this framework when such quantitative information becomes available.

Acknowledgements. Financial support for this work was provided by the Electric Power Research Institute and by the Office of Oceanic and Atmospheric Research of the National Oceanic and Atmospheric Administration under AIRMAP grants NA03OAR4600122 and NA04OAR4600154. We thank H. Mao for provision of model input files for the eastern United States. The development and availability of KPP2.1 by A. Sandu and his coworkers is greatly appreciated.

References

- Aumont, B., Madronich, S., Bey, I., and Tyndall, G. S.: Contribution of secondary VOC to the composition of aqueous atmospheric particles; a modeling approach, *J. Atmos. Chem.*, 35, 59–75, 2000.
- Blando, J. D., Porcja, R. J., Li, T. H., Bowman, D., Lioy, P. J., and Turpin, B. J.: Secondary formation and the Smoky Mountain organic aerosol: An examination of aerosol polarity and functional group composition during SEAVS, *Environ. Sci. Technol.*, 32, 604–613, 1998.
- Blando, J. D. and Turpin, B. J.: Secondary organic aerosol formation in cloud and fog droplets: a literature evaluation of plausibility, *Atmos. Environ.*, 34, 1623–1632, 2000.
- Byun, D. W. and Ching, J. K. S.: Science algorithms of the EPA Models-3 Community Multiscale Air Quality (CMAQ) modeling system, Report EPA/600/R-99/030, Office of Research and Development, United States Environmental Protection Agency, Washington, DC, 1999.

Modeling in-cloud SOA formation

J. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Modeling in-cloud
SOA formation**

J. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Carlton, A. G., Turpin, B. J., Lim, H. J., Altieri, K. E., and Seitzinger, S.: Link between isoprene and secondary organic aerosol (SOA): Pyruvic acid oxidation yields low volatility organic acids in clouds, *Geophys. Res. Lett.*, 33, L06822, doi:10.1029/2005GL025374, 2006.
- Charlson, R. J., Schwartz, S. E., Hales, J. M., Cess, R. D., Coakley Jr., J. A., Hansen, J. E., and Hofmann D. J.: Climate forcing by anthropogenic aerosols, *Science*, 255, 423–430, 1992.
- Chen, J. and Griffin, R. J.: Modeling secondary organic aerosol formation from oxidation of α -pinene, β -pinene, and d-limonene, *Atmos. Environ.*, 39, 7731–7744, 2005.
- Chen, J., Mao, H., Talbot, R. W., and Griffin, R. J.: Application of the CACM and MPMPO modules using the CMAQ model for the eastern United States, *J. Geophys. Res.*, 111, D23S25, doi:10.1029/2006JD007603, 2006.
- Claeys, M., Wang, W., Ion, A. C., Kourtchev, I., Gelencser, A., and Maenhaut, W.: Formation of secondary organic aerosols from isoprene and its gas-phase oxidation products through reaction with hydrogen peroxide, *Atmos. Environ.*, 38, 4093–4098, 2004.
- Daescu, D., Sandu, A., and Carmichael, G. R.: Direct and adjoint sensitivity analysis of chemical kinetic systems with KPP: II Numerical validations and applications, *Atmos. Environ.*, 37, 5097–5114, 2003.
- Damian, V., Sandu, A., Damian, M., Potra, F., and Carmichael, G. R.: The Kinetic preprocessor KPP – A software environment for solving chemical kinetics, *Comput. Chem. Eng.*, 26, 1567–1579, 2002.
- Decesari, S., Facchini, M. C., Fuzzi, S., and Tagliavini, E.: Characterization of water-soluble organic compounds in atmospheric aerosol: A new approach, *J. Geophys. Res.*, 105, 1481–1489, 2000.
- Dutot, A., Rude, J., and Aumont, B.: Neural network method to estimate the aqueous rate constants for the OH reactions with organic compounds, *Atmos. Environ.*, 37, 269–276, 2003.
- Ervens, B., George, C., Williams, J. E., et al.: CAPRAM 2.4 (MODAC mechanism): An extended and condensed tropospheric aqueous phase mechanism and its application, *J. Geophys. Res.*, 108, 4426, doi:10.1029/2002JD002202, 2003.
- Ervens, B., Feingold, G., Frost, G. J., and Kreidenweis, S. M.: A modeling study of aqueous production of dicarboxylic acids: 1. Chemical pathways and speciated organic mass production, *J. Geophys. Res.*, 109, D15205, doi:10.1029/2003JD004387, 2004.
- Gelencser, A. and Varga, Z.: Evaluation of the atmospheric significance of multiphase reactions in atmospheric secondary organic aerosol formation, *Atmos. Chem. Phys.*, 5, 2823–2831,

2005,

<http://www.atmos-chem-phys.net/5/2823/2005/>.

Griffin, R. J., Cocker III, D. R., Flagan, R. C., and Seinfeld, J. H.: Organic aerosol formation from the oxidation of biogenic hydrocarbons, *J. Geophys. Res.*, 104, 3555–3567, 1999.

5 Griffin, R. J., Dabdub, D., and Seinfeld, J. H.: Secondary organic aerosol, 1. Atmospheric chemical mechanism for production of molecular constituents, *J. Geophys. Res.*, 107, 4332, doi: 10.1029/2001JD000541, 2002

Griffin, R. J., Dabdub, D., and Seinfeld, J. H.: Development and initial evaluation of a dynamic species-resolved model for gas-phase chemistry and size-resolved gas/particle partitioning associated with secondary organic aerosol formation, *J. Geophys. Res.*, D05304, doi:10.1029/2004JD005219, 2005.

Griffin, R. J., Nguyen, K., Dabdub, D., and Seinfeld, J. H.: A coupled hydrophobic-hydrophilic model for predicting secondary organic aerosol formation, *J. Atmos. Chem.*, 44, 171–190, 2003.

15 Hastings, W. P., Koehler, C. A., Bailey, E. L., and DeHaan, D. O.: Secondary organic aerosol formation by glyoxal hydration and oligomer formation: Humidity effects and equilibrium shifts during analysis, *Environ. Sci. Technol.*, 39, 8728–8735, 2005.

Heald, C. L., Jacob, D. J., Park, R. J., Russell, L. M., Huebert, B. J., Seinfeld, J. H., Liao, H., and Weber, R. J.: A large organic aerosol source in the free troposphere missing from current models, *Geophys. Res. Lett.*, L18809, doi:10.1029/2005GL023831, 2005.

20 Heald, C. L., Jacob, D. J., Turquety, S., et al.: Concentrations and sources of organic carbon aerosols in the free troposphere over North America, *J. Geophys. Res.*, 111, D23S47, doi:1029/2006JD007705, 2006.

Herrmann, H., Tilgner, A., Barzaghi, P., Majdik, Z., Gligorovski, S., Poulain, L., and Monod, A.: Towards a more detailed description of tropospheric aqueous phase organic chemistry: CAPRAM 3.0, *Atmos. Environ.*, 39, 4351–4363, 2005.

Jang, M., Czoschke, N. M., Lee, S., and Kamens, R. M.: Heterogeneous atmospheric aerosol production by acid-catalyzed particle-phase reactions, *Science*, 298, 814–817, 2002.

25 Jones, A., Roberts, L., and Slingo, A.: A climate model study of indirect radiative forcing by anthropogenic sulphate aerosols, *Nature*, 370, 450–453, 1994.

30 Kalberer, M., Paulsen, D., Sax, M., et al.: Identification of polymers as major components of atmospheric organic aerosols, *Science*, 303, 1659–1662, 2004.

Kanakidou, M., Seinfeld, J. H., Pandis, S. N., et al.: Organic aerosol and global climate model-

ACPD

7, 8951–8982, 2007

Modeling in-cloud SOA formation

J. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

ing: a review, *Atmos. Chem. Phys.*, 5, 1053–1123, 2005,
<http://www.atmos-chem-phys.net/5/1053/2005/>.

Kawamura, K. and Ikushima, K.: Seasonal changes in the distribution of dicarboxylic acids in the urban atmosphere, *Environ. Sci. Technol.*, 27, 2227–2235, 1993.

5 Lim, H. J., Carlton, A. G., and Turpin, B. J.: Isoprene forms secondary organic aerosol through cloud processing: model simulations, *Environ. Sci. Technol.*, 39, 4441–4446, 2005.

Loeffler, K. W., Koehler, C. A., Paul, N. M., and DeHaan, D. O.: Oligomer formation in evaporating aqueous glyoxal and methyl glyoxal solutions, *Environ. Sci. Technol.*, 40, 6318–6323, 2006.

10 Malm, W. C.: Atmospheric haze: Its sources and effects on visibility in rural areas of the continental United States, *Environ. Monit. Assess.*, 12, 203–225, 1989.

Mao, H., Talbot, R., Troop, D., Johnson, R., Businger, S., and Thompson, A. M.: Smart balloon observations over the North Atlantic: Part II ? O₃ data analysis and modeling, *J. Geophys. Res.*, 111, D23S56, doi:10.1029/2005JD006507, 2006.

15 Martin, L. R. and Good, T. W.: Catalyzed oxidation of sulfur dioxide in solution: the iron-manganese synergism, *Atmos. Environ.*, 25A, 2395–2399, 1991.

Murphy, D. M., Thomson, D. S., and Mahoney, M. J.: In situ measurements of organics, meteoritic material, mercury, and other elements in aerosols at 5 to 19 kilometers, *Science*, 282, 1664–1669, 1998.

20 Odum, J. R., Hoffmann, T., Bowman, F., Collins, D., Flagan, R. C., and Seinfeld, J. H.: Gas/particle partitioning and secondary organic aerosol yields, *Environ. Sci. Technol.*, 30, 2580–2585, 1996.

Pandis, S. N. and Seinfeld, J. H.: Sensitivity analysis of a chemical mechanism for aqueous-phase atmospheric chemistry, *J. Geophys. Res.*, 94, 1105–1126, 1989.

25 Pankow, J. F.: An absorption model of the gas/aerosol partitioning involved in the formation of secondary organic aerosol, *Atmos. Environ.*, 28, 189–193, 1994.

Pope, C. A. and Dockery, D. W.: Health effects of fine particulate air pollution: Lines that connect, *J. Air. Waste Manage. Assoc.*, 56, 709–742, 2006.

30 Pun, B. K., Griffin, R. J., Seigneur, C., and Seinfeld, J. H.: Secondary organic aerosol: II: Thermodynamic model for gas/particle partitioning of molecular constituents, *J. Geophys. Res.*, 107, 4333, doi:10.1029/2001JD000542, 2002.

Sandu, A., Daescu, D., and Carmichael, G. R.: Direct and adjoint sensitivity analysis of chemical kinetic systems with KPP: I – Theory and software tools, *Atmos. Environ.*, 37, 5083–5096,

**Modeling in-cloud
SOA formation**

J. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2003.

Sandu, A., Verwer, J. G., Blom, J. G., Spee, E. J., Carmichael, G. R., and Potra, F. A.: Benchmarking stiff ODE solvers for atmospheric chemistry problems II: Rosenbrock methods, *Atmos. Environ.*, 31, 3459–3472, 1997.

5 Schwartz, S. E.: Mass-transport considerations pertinent to aqueous-phase reactions of gases in liquid-water clouds, in *Chemistry of Multiphase Atmospheric Systems*, edited by: Jaeschke, W., Springer, New York, 1986.

Tulet, P., Grini, A., Griffin, R. J., and Petitcol, S.: ORILAM-SOA: A computationally efficient model for predicting secondary organic aerosols in 3D atmospheric models, *J. Geophys. Res.*, D23208, doi:10.1029/2006JD007152, 2006.

10 Turpin, B. J., Saxena, P., and Andrews, E.: Measuring and simulating particulate organics in the atmosphere: Problems and prospects, *Atmos. Environ.*, 34, 2983–3013, 2000.

Warneck, P.: The relative importance of various pathways for the oxidation of sulfur dioxide and nitrogen dioxide in sunlit continental fair weather clouds, *Phys. Chem. Chem. Phys.*, 1, 5471–5483, 1999.

15 Warneck, P.: In-cloud chemistry opens pathway to the formation of oxalic acid in the marine atmosphere, *Atmos. Environ.*, 37, 2423–2427, 2003.

Yao, X., Fang, M., and Chan, C. K.: Size distributions and formation of dicarboxylic acids in atmospheric particles, *Atmos. Environ.*, 36, 2099–2107, 2002.

20 Zaveri, R.: Development and evaluation of a comprehensive tropospheric chemistry model for regional and global application, dissertation, Virginia Polytechnic Institute and State University, Blacksburg, Virginia, 1997.

ACPD

7, 8951–8982, 2007

Modeling in-cloud SOA formation

J. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

Table 1. Aqueous-phase irreversible reactions included in AqChem.

Num.	Reactions	k ^a	E/R (K)	Ref. ^b
1	$O_3 + O_2^- \rightarrow 2O_2 + OH + OH^-$	1.5e9	2200	2
2	$H_2O_2 + OH \rightarrow HO_2 + H_2O$	3.0e7	1680	1
3	$H_2O_2 + hv \rightarrow 2OH$			1
4	$HO_2 + O_2^- \rightarrow H_2O_2 + O_2 + OH^-$	9.7e7	1060	1
5	$HO_2 + HO_2 \rightarrow H_2O_2 + O_2$	8.3e5	2720	1
6	$OH + HO_2 \rightarrow H_2O + O_2$	1.0e10		1
7	$OH + O_2^- \rightarrow OH^- + O_2$	1.1e10	2120	1
8	$O_3 + OH \rightarrow HO_2 + O_2$	1.0e8		1
9	$NO_3^- + hv \rightarrow NO_2 + OH + OH^-$			1
10	$S(IV) + O_2 \xrightarrow{Fe^{3+}, Mn^{2+}} S(VI)$	$2.6e3[Fe^{3+}] + 7.5e2[Mn^{2+}]$ $+ 1.0e10[Fe^{3+}][Mn^{2+}]$ (if PH \leq 5.0) $7.5e2[Mn^{2+}]$ $2.0e10[Fe^{3+}][Mn^{2+}]$ (if PH $>$ 5.0)		5,6
11	$HSO_3^- + H_2O_2 + H^+ \rightarrow SO_4^{2-} + H_2O + 2 H^+$	6.9e7	4000	1,4
12	$HSO_3^- + O_3 \rightarrow HSO_4^- + O_2$	3.7e5	5530	1,4
13	$SO_2 \cdot H_2O + O_3 \rightarrow HSO_4^- + O_2 + H^+$	2.4e4		1,4
14	$SO_3^{2-} + O_3 \rightarrow SO_4^{2-} + O_2$	1.5e9	5280	1,4
15	$HCHO + HSO_3^- \rightarrow HOCH_2SO_3^-$	4.5e2	2660	4
16	$HCHO + SO_3^{2-} \rightarrow HOCH_2SO_3^- + OH^-$	5.4e6	2530	4
17	$HOCH_2SO_3^- + OH^- \rightarrow CH_2(OH)_2 + SO_3^{2-}$	4.6e3	4880	4
18	$HOCH_2SO_3^- + OH^- \rightarrow HOCHSO_3^- + H_2O$	3.0e8		4
19	$HOCHSO_3^- + O_2 \rightarrow HCOOH + HSO_3^- + HO_2$	2.5e9		4
20	$CH_3OH + OH^- \rightarrow HCHO + HO_2$	1.0e9	580	1
21	$CH_2(OH)_2 + OH^- \rightarrow HCOOH + HO_2 + H_2O$	1.1e9	1020	1
22	$CH_3OO + HO_2 \rightarrow CH_3OOH + O_2$	4.3e5		4
23	$CH_3OO + O_2^- \rightarrow CH_3OOH + OH^- + O_2$	5.0e7		4
24	$CH_3OOH + OH^- \rightarrow CH_3OO + H_2O$	3.0e7	1680	1
25	$CH_3OOH + OH^- \rightarrow HO_2 + HCOOH$	6.0e6	1680	1
26	$HCOOH + OH^- \rightarrow CO_2 + HO_2 + H_2O$	1.3e8	1000	1
27	$HCOO^- + OH^- \rightarrow CO_2 + H_2O$	1.0e9	1000	1
28	$CH_3CHO + OH^- \rightarrow CH_3COOH + HO_2$	3.6e9		1
29	$CH_3CH(OH)_2 + OH^- \rightarrow CH_3COOH + HO_2$	1.2e9		1
30	$CH_3CH_2OH + OH^- \rightarrow CH_3CHO + HO_2$	1.9e9		1
31	$CH_2OHCHO + OH^- \rightarrow (CH_2(OH)_2)_2 + HO_2$	1.2e9		2
32	$(CH_2(OH)_2)_2 + OH^- \rightarrow CHOCOOH + HO_2$	1.1e9	1516	1
33	$CHOCOOH + OH^- \rightarrow (COOH)_2 + HO_2$	3.6e8	1000	2
34	$CHOCOO^- + OH^- \rightarrow (COOH)_2 + HO_2$	2.9e9	4300	2
35	$CH_3COOH + OH^- \rightarrow 0.85CHOCOOH + 0.15 CH_2(OH)_2$	1.6e7		3

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modeling in-cloud
SOA formation

J. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

Table 1. Continued.

Num.	Reactions	k^a	E/R (K)	Ref. ^b
36	$\text{CH}_3\text{COO}^- + \text{OH} \rightarrow 0.85\text{CHOCOOH} + 0.15 \text{CH}_2(\text{OH})_2$	8.5e7		3
37	$(\text{COOH})_2 + \text{OH} \rightarrow \text{HO}_2 + 2\text{CO}_2 + \text{H}_2\text{O}$	1.4e6		2
38	$\text{HC}_2\text{O}_4^- + \text{OH} \rightarrow \text{HO}_2 + 2\text{CO}_2 + \text{H}_2\text{O}$	1.9e8	2800	2
39	$\text{C}_2\text{O}_4^{2-} + \text{OH} \rightarrow \text{HO}_2 + 2\text{CO}_2 + \text{H}_2\text{O}$	1.6e8	4300	2
40	$\text{CH}_3\text{COCHO} + \text{OH} \rightarrow \text{CH}_3\text{COCO}^\bullet\text{OH} + \text{HO}_2$	1.1e9	1600	2
41	$\text{CH}_3\text{COCO}^\bullet\text{OH} + \text{OH} \rightarrow \text{CH}_3\text{COOH} + \text{HO}_2 + \text{CO}_2$	6.0e7		3
42	$\text{CH}_3\text{COCO}^\bullet\text{O}^- + \text{OH} \rightarrow \text{CH}_3\text{COO}^- + \text{HO}_2 + \text{CO}_2$	6.0e7		3
43 ^c	$\text{S3} + \text{OH} \rightarrow \text{S3PD} + \text{HO}_2$	4.98e9		7
44 ^c	$\text{S5} + \text{OH} \rightarrow \text{S5PD} + \text{HO}_2$	5.76e9		7
45 ^c	$\text{S7} + \text{OH} \rightarrow \text{S7PD} + \text{HO}_2$	4.26e9		7
46 ^c	$\text{S7}^- + \text{OH} \rightarrow \text{S7PD} + \text{HO}_2$	4.08e9		7
47 ^c	$\text{S10} + \text{OH} \rightarrow \text{S10PD}$	1.23e10		7

^a: in appropriate units of M and s⁻¹^b: References: 1: Ervens et al. (2003); 2: Ervens et al. (2004); 3: Lim et al. (2005); 4: Warneck (1999); 5: Martin and Good (1991); 6: Zaveri (1999); 7: Dutot et al. (2003).^c: S3: 3-hydroxy-2, 4-dimethyl-2, 4-hexadienal; S3PD: 3-hydroxy-2, 4-dimethyl-2, 4-hexadienalic acid; S5: 2-hydroxyl-3-isopropyl-6-keto-heptanal; S5PD: 2-hydroxyl-3-isopropyl-6-keto-heptanoic acid; S7: 2,4-dimethyl-3-formyl-benzoic acid; S7PD: 3, 5-dimethyl-1, 4-dibenzoic acid; S10: 1-methyl-1-hydroxy-2-nitrato-4-isopropyl- cyclohexane; S10PD: 1-methyl-1, 3-dihydroxy-2-nitrato-4-isopropyl-cyclohexane

**Modeling in-cloud
SOA formation**

J. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 2. Aqueous-phase equilibrium reactions included in AqChem.

Num.	Equilibrium reactions	K (M)	E/R (K)	Ref. ^a
1	$\text{H}_2\text{O} \leftrightarrow \text{H}^+ + \text{OH}^-$	1.8e-16	6800	1
2	$\text{CO}_2\text{-H}_2\text{O} \leftrightarrow \text{HCO}_3^+ + \text{H}^-$	7.7e-7	-1000	2
3	$\text{HCO}_3^+ \leftrightarrow \text{CO}_3^{2-} + \text{H}^-$	4.84e-11	-1760	2
4	$\text{HO}_2 \leftrightarrow \text{H}^+ + \text{O}_2^-$	1.6e-5		1
5	$\text{HNO}_4 \leftrightarrow \text{H}^+ + \text{NO}_4^-$	1.0e-5		1
6	$\text{HNO}_3 \leftrightarrow \text{H}^+ + \text{NO}_3^-$	22.0	-1800	1
7	$\text{HONO} \leftrightarrow \text{H}^+ + \text{NO}_2^-$	5.3e-4	1760	1
8	$\text{NH}_3\text{-H}_2\text{O} \leftrightarrow \text{NH}_4^+ + \text{OH}^-$	1.75e-5	560	1
9	$\text{SO}_2\text{-H}_2\text{O} \leftrightarrow \text{HSO}_3^- + \text{H}^+$	1.73e-2	-1940	1
10	$\text{HSO}_3^- \leftrightarrow \text{SO}_3^{2-} + \text{H}^+$	6.22e-8	-1960	1
11	$\text{H}_2\text{SO}_4 \leftrightarrow \text{HSO}_4^- + \text{H}^+$	1000		1
12	$\text{HSO}_4^- \leftrightarrow \text{SO}_4^{2-} + \text{H}^+$	1.02e-2	-2700	1
13	$\text{HCHO} + \text{H}_2\text{O} \leftrightarrow \text{CH}_2(\text{OH})_2$	36.0	-4030	1
14	$\text{HCOOH} \leftrightarrow \text{HCOO}^- + \text{H}^+$	1.77e-4	-12	1
15	$\text{CH}_3\text{CHO} + \text{H}_2\text{O} \leftrightarrow \text{CH}_3\text{CH}(\text{OH})_2$	2.46e-2	-2500	1
16	$\text{CH}_3\text{COOH} \leftrightarrow \text{H}^+ + \text{CH}_3\text{COO}^-$	1.75e-5	-46	1
17	$\text{CHOCOOH} \leftrightarrow \text{CHOCOO}^- + \text{H}^+$	6.6e-4		2
18	$\text{COOHCOOH} \leftrightarrow \text{HC}_2\text{O}_4^- + \text{H}^+$	6.4e-2		1
19	$\text{HC}_2\text{O}_4^- \leftrightarrow \text{C}_2\text{O}_4^{2-} + \text{H}^+$	5.25e-5		1
20	$\text{CH}_3\text{COCOOH} \leftrightarrow \text{CH}_3\text{COCOO}^- + \text{H}^+$	4.07e-3		2
21	$\text{S7} \leftrightarrow \text{S7}^- + \text{H}^+$	7.335e-5		3
22 ^b	$\text{S3PD} \leftrightarrow \text{S3PD}^- + \text{H}^+$	3.7e-5		3
23 ^b	$\text{S5PD} \leftrightarrow \text{S5PD}^- + \text{H}^+$	6.52e-4		3
24 ^b	$\text{S7PD} \leftrightarrow \text{S7PD}^- + \text{H}^+$	1.7e-3		3

^a: References: 1: Ervens et al. (2003); 2: Ervens et al. (2004); 3: Pun et al. (2002).

^b: See Table 1 for molecular definition of S7, S3PD, S5PD, and S7PD.

Table 3. Uptake parameters for gas-phase species in AqChem.

Species ^a	$K_{H,298K}$ (M atm ⁻¹)	$\Delta H/R$ (K)	α	D_g (m ² s ⁻¹)	Ref. ^b
CO ₂	3.11e-2	-2423	2.0e-4	1.55e-5	1
NH ₃	60.7	-3920	0.04	2.3e-5	1
O ₃	1.14e-2	-2300	0.05	1.48e-5	1
HO ₂	9.0e3		0.01	1.04-5	1
OH	25.0	-5280	0.05	1.53e-5	1
H ₂ O ₂	1.02e5	-6340	0.11	1.46e-5	1
HNO ₃	2.1e5	-8700	0.054	1.32e-5	1
HONO	49.0	-4880	0.5	1.30e-5	1
HNO ₄	3.0e4		0.1	1.30e-5	1
SO ₂	1.24	-3247	3.5e-2	1.28e-5	1
H ₂ SO ₄	2.1e5		0.07	1.30e-5	1
HCHO	2.5		0.02	1.64e-5	1
CH ₃ OH	220	-5390	0.015	1.16e-5	1
CH ₃ OO	310		3.8e-3	1.35e-5	1
CH ₃ OOH	310		3.8e-3	1.31e-5	1
HCOOH	5530	5630	0.10	1.0e-5	2
CH ₃ CHO	11.4	-6254	0.03	1.22e-5	1
CH ₃ CH ₂ OH	190	-6290	8.2e-3	0.95e-5	1
CH ₂ OHCHO	4.14e4		0.03	1.95e-5	2
CH ₃ COOH	5500	-5890	0.019	1.24e-5	2
COOHCOOH	3.26e6		0.019	1.24e-5	3
CHOCHO	3.0e5		3.8e-3	1.31e-5	2
CHOCOOH	9000		0.1	1.0e-5	2
CH ₃ COCHO	3710		0.023	1.15e-5	2
CH ₃ COCOOH	3.11e5	5100	0.1	1.0e-5	2
S3 ^c	6.01e6		0.05	1.0e-5	4,5
S5 ^c	3.90e7		0.05	1.0e-5	4,5
S7 ^c	4.91e6		0.05	1.0e-5	4,5
S10 ^c	1.18e6		0.05	1.0e-5	4,5

^a: N₂O₅ is assumed to be completely dissolved in water and to form two moles of HNO₃.

^b: References: 1: Ervens et al. (2003); 2: Ervens et al. (2004); 3: Lim et al. (2005); 4: Pun et al. (2002); 5: Aumont et al. (2000).

^c: See Table 1 for molecular definition of S3, S5, S7, and S10.

Modeling in-cloud SOA formation

J. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modeling in-cloud SOA formation

J. Chen et al.

Table 4. Twenty-four-hour average emission rates (moles $\text{m}^{-2} \text{s}^{-1}$) for two scenarios used in zero-dimensional model simulations.

Species	Emissions for Scenario 1	Emissions for Scenario 2
NO	1.36e-9	3.27e-8
NO ₂	1.42e-10	3.63e-9
CO	1.33e-8	3.40e-7
HCHO	1.33e-11	6.73e-10
ALD1 ^a	2.63e-10	1.06e-9
OLEL ^a	1.38e-9	1.49e-9
AROL ^a	5.56e-11	7.18e-10
AROH ^a	6.60e-11	1.33e-9
ISOP ^a	2.64e-9	1.83e-9
TERP ^a	8.30e-10	1.87e-11
ALKL ^a	3.04e-9	1.11e-8
SO ₂	5.06e-11	6.38e-9

^a: ALD1: lumped small-carbon-number aldehydes; OLEL: lumped alkenes C₃-C₆; AROL: lumped low-SOA-yield aromatic species; AROH: lumped high-SOA-yield aromatic species; ISOP: isoprene; TERP: monoterpenes; ALKL: lumped alkanes C₂-C₆.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Modeling in-cloud SOA formation

J. Chen et al.

Table 5. Twenty-four-hour average dry deposition velocities (dm sec^{-1}) for two scenarios used in zero-dimensional model simulations.


Species	Dry Deposition (urban)	Dry deposition (rural)
NO	9.08e-4	2.52e-3
NO ₂	7.45e-4	2.03e-3
O ₃	1.46e-3	2.71e-3
H ₂ O ₂	4.77e-3	7.41e-3
NO ₃	3.26e-2	1.48e-2
HNO ₃	3.41e-2	1.75e-2
HONO	3.94e-3	6.88e-3
N ₂ O ₅	3.00e-2	1.56e-2
CO	1.26e-3	2.83e-3
SO ₂	3.82e-3	6.28e-3
HCHO	3.20e-3	6.67e-3
ALD ^a	2.03e-3	4.03e-3
ORA ^a	3.63e-3	6.06e-3
PAN ^a	8.89e-4	1.95-e3

^a: ALD for general aldehydes, ORA for general organic acids, and PAN for general peroxy acetyl nitrates.


[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Modeling in-cloud
SOA formation

J. Chen et al.



a.)



b.)

Fig. 1. Schematic of organic reactions in clouds: **(a)** organics of carbon number less than four (based on Ervens et al., 2004 and Lim et al., 2005); **(b)** organics of carbon number greater than four (based on Aumont et al., 2000) (S2–S7 and S10 are surrogates in the MPMPO module).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Modeling in-cloud
SOA formation**

J. Chen et al.




Fig. 2. Flow diagram for SOA modeling with/without consideration of the aqueous-phase chemistry in clouds.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

EGU

Modeling in-cloud
SOA formation

J. Chen et al.




Fig. 3. SOA predictions from box-model simulations: **(a)** Scenario 1 (S4: SOA of surrogate species 4 of MPMPO; S5: SOA of surrogate species 5 of MPMPO) **(b)** Scenario 2 (S2: SOA of surrogate species 2 of MPMPO; S6: SOA of surrogate species 6 of MPMPO). “w aq” indicates simulation with consideration of the aqueous-phase organic chemistry in clouds; “w/o aq” indicates simulation without consideration of the aqueous-phase organic chemistry in clouds.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Modeling in-cloud
SOA formation

J. Chen et al.




Fig. 4. Spatial distribution of surface SOA predictions: **(a)** difference between 24-h average SOA prediction on 3 August 2004, with/without aqueous-phase organic chemistry simulation in clouds; **(b)** 24-h average SOA prediction on 3 August 2004, with aqueous-phase organic chemistry simulation in clouds; **(c)** difference between 24-h average SOA prediction on 4 August 2004, with/without aqueous-phase organic chemistry simulation in clouds; **(d)** 24-h average SOA prediction on 4 August 2004, with aqueous-phase organic chemistry simulation in clouds.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Modeling in-cloud
SOA formation

J. Chen et al.




Fig. 5. Time profiles of SOA predictions at sites with maximum difference in 24-h averaged SOA predictions with or without consideration of aqueous-phase organic chemistry in clouds on 3 August 2004 in the northern Gulf of Maine (a) and 4 August 2004 in the northern Gulf of Maine (b).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Modeling in-cloud
SOA formation

J. Chen et al.




Fig. 6. Spatial distribution of SOA predictions of layer 14: **(a)** difference in 24-h average SOA prediction on 3 August 2004, with/without aqueous-phase organic chemistry simulation in clouds; **(b)** 24-h average SOA prediction on 3 August 2004, with aqueous-phase organic chemistry simulation in clouds; **(c)** difference in 24-h average SOA prediction on 4 August 2004, with/without aqueous-phase organic chemistry simulation in clouds; **(d)** 24-h average SOA prediction on 4 August 2004, with aqueous-phase organic chemistry simulation in clouds.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)