

Equipartition of energy for the wave equation associated to the Dunkl-Cherednik Laplacian

Fatma Ayadi*

Abstract. This paper is concerned with energy properties of the wave equation associated to the Dunkl-Cherednik Laplacian. We establish the conservation of the total energy, the strict equipartition of energy under suitable assumptions and the asymptotic equipartition in the general case.

Mathematics Subject Index 2000: Primary 35L05; Secondary 22E30, 33C67, 35L65, 58J45

Keywords and phrases: Dunkl-Cherednik operator, wave equation, energy, equipartition

1. Introduction

We use [10] as a reference for the Dunkl-Cherednik theory. Let \mathfrak{a} be a Euclidean vector space of dimension d equipped with an inner product $\langle \cdot, \cdot \rangle$. Let \mathcal{R} be a crystallographic root system in \mathfrak{a} , \mathcal{R}^+ a positive subsystem and W the Weyl group generated by the reflections $r_\alpha(x) = x - 2\frac{\langle \alpha, x \rangle}{\|\alpha\|^2}\alpha$ along the roots $\alpha \in \mathcal{R}$. We let $k : \mathcal{R} \rightarrow [0, +\infty)$ denote a multiplicity function on the root system \mathcal{R} , and $\rho = \frac{1}{2} \sum_{\alpha \in \mathcal{R}^+} k_\alpha \alpha$. We note that k is W -invariant. The Dunkl-Cherednik operators are the following differential-difference operators, which are deformations of partial derivatives and still commute pairwise :

$$T_\xi f(x) = \partial_\xi f(x) - \langle \rho, \xi \rangle f(x) + \sum_{\alpha \in \mathcal{R}^+} k_\alpha \frac{\langle \alpha, \xi \rangle}{1 - e^{-\langle \alpha, x \rangle}} \{f(x) - f(r_\alpha x)\}.$$

Given an orthonormal basis $\{\xi_1, \dots, \xi_d\}$ of \mathfrak{a} , the Dunkl-Cherednik Laplacian is defined by

$$L f(x) = \sum_{j=1}^d T_{\xi_j}^2 f(x).$$

More explicit formulas for L exist but they will not be used in this paper. The Laplacian L is selfadjoint with respect to the measure $\mu(x)dx$ where

$$\mu(x) = \prod_{\alpha \in \mathcal{R}^+} \left| 2 \sinh \frac{\langle \alpha, x \rangle}{2} \right|^{2k_\alpha}.$$

*This work is part of my master thesis, done under the supervision of M. Sifi (Université de Tunis El Manar) and J.-Ph. Anker (Université d'Orléans), within the program CMCU 07G 1501 *Aspects analytique et probabiliste de la théorie de Dunkl*

Consider the wave equation

$$\begin{cases} \partial_t^2 u(t, x) = L_x u(t, x), \\ u(0, x) = f(x), \partial_t|_{t=0} u(t, x) = g(x), \end{cases} \quad (1)$$

with smooth and compactly supported initial data (f, g) . Let us introduce:

- the *kinetic energy* $\mathcal{K}[u](t) = \frac{1}{2} \int_{\mathfrak{a}} |\partial_t u(x, t)|^2 \mu(x) dx$,
- the *potential energy* $\mathcal{P}[u](t) = -\frac{1}{2} \int_{\mathfrak{a}} Lu(x, t) \overline{u(x, t)} \mu(x) dx$,
- the *total energy* $\mathcal{E}[u](t) = \mathcal{K}[u](t) + \mathcal{P}[u](t)$.

In this paper we prove

- the conservation of the total energy:

$$\mathcal{E}[u](t) = \text{constant}, \quad (2)$$

- the strict equipartition of energy, under the assumptions that the dimension d is odd and that all the multiplicities k_α are integers:

$$\mathcal{K}[u](t) = \mathcal{P}[u](t) = \frac{1}{2} \mathcal{E}[u] \text{ for } |t| \text{ large}, \quad (3)$$

- the asymptotic equipartition of energy, for arbitrary d and \mathbb{R}^+ -valued multiplicity function k :

$$\mathcal{K}[u](t) \rightarrow \frac{1}{2} \mathcal{E}[u] \text{ and } \mathcal{P}[u](t) \rightarrow \frac{1}{2} \mathcal{E}[u] \text{ as } |t| \text{ goes to } \infty. \quad (4)$$

The proofs follow [6] and use the Fourier transform in the Dunkl-Cherednik setting, which we will recall in next section. We mention that during the past twenty years, several works were devoted to Huygens' principle and equipartition of energy for wave equations on symmetric spaces and related settings. See for instance [3], [8, ch. V, § 5], [6], [7], [2], [4], [5], [1].

2. Generalized hypergeometric functions and Dunkl-Cherednik transform

Opdam [9] introduced the following special functions, which are deformations of exponential functions $e^{\langle \lambda, x \rangle}$, and the associated Fourier transform.

Theorem 2.1. *There exist a neighborhood U of 0 in \mathfrak{a} and a unique holomorphic function $(\lambda, x) \mapsto G_\lambda(x)$ on $a_{\mathbb{C}} \times (\mathfrak{a} + iU)$ such that*

$$\begin{cases} T_\xi G_\lambda(x) = \langle \lambda, \xi \rangle G_\lambda(x) \quad \forall \xi \in \mathfrak{a}, \\ G_\lambda(0) = 1. \end{cases}$$

Moreover, the following estimate holds on $a_{\mathbb{C}} \times \mathfrak{a}$:

$$|G_\lambda(x)| \leq |W|^{\frac{1}{2}} e^{\|\operatorname{Re} \lambda\| \|x\|}.$$

Definition 2.2. Let f be a nice function on \mathfrak{a} , say f belongs to the space $\mathcal{C}_c^\infty(\mathfrak{a})$ of smooth functions on \mathfrak{a} with compact support. Its Dunkl-Cherednik transform is defined by

$$\mathcal{F}f(\lambda) = \int_{\mathfrak{a}} f(x) G_{-i w_0 \lambda}(w_0 x) \mu(x) dx.$$

Here w_0 denotes the longest element in the Weyl group W .

The involvement of w_0 in the definition of \mathcal{F} is related to the below skew-adjointness property of the Dunkl-Cherednik operators with respect to the inner product

$$\langle f, g \rangle = \int_{\mathfrak{a}} f(x) \overline{g(x)} \mu(x) dx, \quad f, g \in \mathcal{C}_c^\infty(\mathfrak{a}).$$

Lemma 2.3. The adjoint of T_ξ is $-w_0 T_{w_0 \xi} w_0$:

$$\langle T_\xi f, g \rangle = \langle f, -w_0 T_{w_0 \xi} w_0 g \rangle.$$

As an immediate consequence, we obtain:

Corollary 2.4. For every $\xi, \lambda \in \mathfrak{a}$ and $f \in \mathcal{C}_c^\infty(\mathfrak{a})$, we have

$$\mathcal{F}(T_\xi f)(\lambda) = i \langle \lambda, \xi \rangle \mathcal{F}f(\lambda),$$

and therefore

$$\mathcal{F}(Lf)(\lambda) = -\|\lambda\|^2 \mathcal{F}f(\lambda).$$

Next we will recall from [9] three main results about the Dunkl-Cherednik transform (see also [10]). For $R > 0$, let $\mathcal{C}_R^\infty(\mathfrak{a})$ be the space of smooth functions on \mathfrak{a} vanishing outside the ball $B_R = \{x \in \mathfrak{a} \mid \|x\| \leq R\}$. We let $\mathcal{H}_R(\mathfrak{a}_\mathbb{C})$ denote the space of holomorphic functions h on the complexification $\mathfrak{a}_\mathbb{C}$ of \mathfrak{a} such that, for every integer $N > 0$,

$$\sup_{\lambda \in \mathfrak{a}_\mathbb{C}} (1 + \|\lambda\|)^N e^{-R \|\operatorname{Im} \lambda\|} |h(\lambda)| < +\infty.$$

Theorem 2.5. (Paley-Wiener) The transformation \mathcal{F} is an isomorphism of $\mathcal{C}_R^\infty(\mathfrak{a})$ onto $\mathcal{H}_R(\mathfrak{a}_\mathbb{C})$, for every $R > 0$.

The Plancherel formula and the inversion formula of \mathcal{F} involve the complex measure $\nu(\lambda) d\lambda$ with density

$$\nu(\lambda) = \prod_{\alpha \in \mathcal{R}_0^+} \frac{\Gamma(i \langle \lambda, \check{\alpha} \rangle + k_\alpha)}{\Gamma(i \langle \lambda, \check{\alpha} \rangle)} \frac{\Gamma\left(\frac{i \langle \lambda, \check{\alpha} \rangle + k_\alpha}{2} + k_{2\alpha}\right)}{\Gamma\left(\frac{i \langle \lambda, \check{\alpha} \rangle + k_\alpha}{2}\right)} \frac{\Gamma(-i \langle \lambda, \check{\alpha} \rangle + k_\alpha)}{\Gamma(-i \langle \lambda, \check{\alpha} \rangle + 1)} \frac{\Gamma\left(\frac{-i \langle \lambda, \check{\alpha} \rangle + k_\alpha}{2} + k_{2\alpha} + 1\right)}{\Gamma\left(\frac{-i \langle \lambda, \check{\alpha} \rangle + k_\alpha}{2}\right)},$$

where $\mathcal{R}_0^+ = \{\alpha \in \mathcal{R}^+ \mid \frac{\alpha}{2} \notin \mathcal{R}\}$ is the set of positive indivisible roots, $\check{\alpha} = 2\|\alpha\|^{-2}\alpha$ the coroot corresponding to α , and $k_{2\alpha} = 0$ if $2\alpha \notin \mathcal{R}$. Notice that ν is an analytic function on \mathfrak{a} , with a polynomial growth and which extends meromorphically to $\mathfrak{a}_\mathbb{C}$. It is actually a polynomial if the multiplicity function k is integer-valued and it has poles otherwise.

Theorem 2.6. (Inversion formula) *There is a constant $c_0 > 0$ such that, for every $f \in \mathcal{C}_c^\infty(\mathfrak{a})$,*

$$f(x) = c_0 \int_{\mathfrak{a}} \mathcal{F}f(\lambda) G_{i\lambda}(x) \nu(\lambda) d\lambda.$$

Theorem 2.7. (Plancherel formula) *For every $f, g \in \mathcal{C}_c^\infty(\mathfrak{a})$,*

$$\int_{\mathfrak{a}} f(x) \overline{g(x)} \mu(x) dx = c_0 \int_{\mathfrak{a}} \mathcal{F}f(\lambda) \tilde{\mathcal{F}}g(\lambda) \nu(\lambda) d\lambda,$$

where $\tilde{\mathcal{F}}g(\lambda) := \overline{\mathcal{F}(w_0g)(w_0\lambda)} = \int_{\mathfrak{a}} \overline{g(x)} G_{i\lambda}(x) \mu(x) dx$.

3. Conservation of energy

This section is devoted to the proof of (2). Via the Dunkl-Cherednik transform, the wave equation (1) becomes

$$\begin{cases} \partial_t^2 \mathcal{F}u(t, \lambda) = -\|\lambda\|^2 \mathcal{F}u(t, \lambda), \\ \mathcal{F}u(0, \lambda) = \mathcal{F}f(\lambda), \partial_t|_{t=0} \mathcal{F}u(t, \lambda) = \mathcal{F}g(\lambda), \end{cases}$$

and its solution satisfies

$$\mathcal{F}u(t, \lambda) = (\cos t\|\lambda\|) \mathcal{F}f(\lambda) + \frac{\sin t\|\lambda\|}{\|\lambda\|} \mathcal{F}g(\lambda). \quad (5)$$

By means of the Paley-Wiener Theorem 2.5, in [1, p. 52-53] the author proves the following finite speed propagation property:

Assume that the initial data f and g belong to $\mathcal{C}_R^\infty(\mathfrak{a})$. Then the solution $u(t, x)$ belongs to $\mathcal{C}_{R+|t|}^\infty(\mathfrak{a})$ as a function of x .

Let us express the potential and kinetic energies defined in the introduction via the Dunkl-Cherednik transform. Using the Plancherel formula and Corollary 2.4, we have

$$\mathcal{P}[u](t) = \frac{c_0}{2} \int_{\mathfrak{a}} \|\lambda\|^2 \mathcal{F}u(t, \lambda) \tilde{\mathcal{F}}u(t, \lambda) \nu(\lambda) d\lambda. \quad (6)$$

Moreover, since the Dunkl-Cherednik Laplacian is W -invariant, it follows that $(w_0u)(t, x) = u(t, w_0x)$ is the solution to the wave equation (1) with the initial data w_0f and w_0g . Thus

$$\tilde{\mathcal{F}}u(t, \lambda) = (\cos t\|\lambda\|) \tilde{\mathcal{F}}f(\lambda) + \frac{\sin t\|\lambda\|}{\|\lambda\|} \tilde{\mathcal{F}}g(\lambda). \quad (7)$$

Now, by substituting (5) and (7) in (6), we get

$$\begin{aligned} \mathcal{P}[u](t) &= \frac{c_0}{2} \int_{\mathfrak{a}} \|\lambda\|^2 (\cos t\|\lambda\|)^2 \mathcal{F}f(\lambda) \tilde{\mathcal{F}}f(\lambda) \nu(\lambda) d\lambda \\ &+ \frac{c_0}{2} \int_{\mathfrak{a}} (\sin t\|\lambda\|)^2 \mathcal{F}g(\lambda) \tilde{\mathcal{F}}g(\lambda) \nu(\lambda) d\lambda \\ &+ \frac{c_0}{4} \int_{\mathfrak{a}} \|\lambda\| (\sin 2t\|\lambda\|) \{ \mathcal{F}f(\lambda) \tilde{\mathcal{F}}g(\lambda) + \mathcal{F}g(\lambda) \tilde{\mathcal{F}}f(\lambda) \} \nu(\lambda) d\lambda. \end{aligned} \quad (8)$$

Similarly to $\mathcal{P}[u]$, we can rewrite the kinetic energy as

$$\mathcal{K}[u](t) = \frac{c_0}{2} \int_a \partial_t \mathcal{F}u(t, \lambda) \partial_t \tilde{\mathcal{F}}u(t, \lambda) \nu(\lambda) d\lambda.$$

Using the following facts

$$\begin{cases} \partial_t \mathcal{F}u(t, \lambda) = -\|\lambda\| (\sin t\|\lambda\|) \mathcal{F}f(\lambda) + (\cos t\|\lambda\|) \mathcal{F}g(\lambda), \\ \partial_t \tilde{\mathcal{F}}u(t, \lambda) = -\|\lambda\| (\sin t\|\lambda\|) \tilde{\mathcal{F}}f(\lambda) + (\cos t\|\lambda\|) \tilde{\mathcal{F}}g(\lambda), \end{cases}$$

we deduce that

$$\begin{aligned} \mathcal{K}[u](t) &= \frac{c_0}{2} \int_a \|\lambda\|^2 (\sin t\|\lambda\|)^2 \mathcal{F}f(\lambda) \tilde{\mathcal{F}}f(\lambda) \nu(\lambda) d\lambda \\ &\quad + \frac{c_0}{2} \int_a (\cos t\|\lambda\|)^2 \mathcal{F}g(\lambda) \tilde{\mathcal{F}}g(\lambda) \nu(\lambda) d\lambda \\ &\quad - \frac{c_0}{4} \int_a \|\lambda\| (\sin 2t\|\lambda\|) \{ \mathcal{F}f(\lambda) \tilde{\mathcal{F}}g(\lambda) + \mathcal{F}g(\lambda) \tilde{\mathcal{F}}f(\lambda) \} \nu(\lambda) d\lambda. \end{aligned} \quad (9)$$

By suming up (8) and (9), we obtain the conservation of the total energy:

$$\mathcal{E}[u](t) = \frac{c_0}{2} \int_a \{ \|\lambda\|^2 \mathcal{F}f(\lambda) \tilde{\mathcal{F}}f(\lambda) + \mathcal{F}g(\lambda) \tilde{\mathcal{F}}g(\lambda) \} \nu(\lambda) d\lambda = \mathcal{E}[u](0).$$

That is $\mathcal{E}[u](t)$ is independent of t .

4. Equipartition of energy

This section is devoted to the proof of (3) and (4). Using the classical trigonometric identities for double angles, we can rewrite the identities (8) and (9) respectively as

$$\begin{aligned} \mathcal{P}[u](t) &= \frac{c_0}{4} \int_a \{ \|\lambda\|^2 \mathcal{F}f(\lambda) \tilde{\mathcal{F}}f(\lambda) + \mathcal{F}g(\lambda) \tilde{\mathcal{F}}g(\lambda) \} \nu(\lambda) d\lambda \\ &\quad + \frac{c_0}{4} \int_a (\cos 2t\|\lambda\|) \{ \|\lambda\|^2 \mathcal{F}f(\lambda) \tilde{\mathcal{F}}f(\lambda) - \mathcal{F}g(\lambda) \tilde{\mathcal{F}}g(\lambda) \} \nu(\lambda) d\lambda \\ &\quad + \frac{c_0}{4} \int_a \|\lambda\| (\sin 2t\|\lambda\|) \{ \mathcal{F}f(\lambda) \tilde{\mathcal{F}}g(\lambda) + \mathcal{F}g(\lambda) \tilde{\mathcal{F}}f(\lambda) \} \nu(\lambda) d\lambda \end{aligned}$$

and

$$\begin{aligned} \mathcal{K}[u](t) &= \frac{c_0}{4} \int_a \{ \|\lambda\|^2 \mathcal{F}f(\lambda) \tilde{\mathcal{F}}f(\lambda) + \mathcal{F}g(\lambda) \tilde{\mathcal{F}}g(\lambda) \} \nu(\lambda) d\lambda \\ &\quad - \frac{c_0}{4} \int_a (\cos 2t\|\lambda\|) \{ \|\lambda\|^2 \mathcal{F}f(\lambda) \tilde{\mathcal{F}}f(\lambda) - \mathcal{F}g(\lambda) \tilde{\mathcal{F}}g(\lambda) \} \nu(\lambda) d\lambda \\ &\quad - \frac{c_0}{4} \int_a \|\lambda\| (\sin 2t\|\lambda\|) \{ \mathcal{F}f(\lambda) \tilde{\mathcal{F}}g(\lambda) + \mathcal{F}g(\lambda) \tilde{\mathcal{F}}f(\lambda) \} \nu(\lambda) d\lambda. \end{aligned}$$

Hence

$$\begin{aligned} \mathcal{P}[u](t) - \mathcal{K}[u](t) &= \\ &= \frac{c_0}{2} \int_a (\cos 2t\|\lambda\|) \{ \|\lambda\|^2 \mathcal{F}f(\lambda) \tilde{\mathcal{F}}f(\lambda) - \mathcal{F}g(\lambda) \tilde{\mathcal{F}}g(\lambda) \} \nu(\lambda) d\lambda \\ &\quad + \frac{c_0}{2} \int_a \|\lambda\| (\sin 2t\|\lambda\|) \{ \mathcal{F}f(\lambda) \tilde{\mathcal{F}}g(\lambda) + \mathcal{F}g(\lambda) \tilde{\mathcal{F}}f(\lambda) \} \nu(\lambda) d\lambda. \end{aligned} \quad (10)$$

Introducing polar coordinates in \mathfrak{a} , (10) becomes

$$\mathcal{P}[u](t) - \mathcal{K}[u](t) = \frac{c_0}{2} \int_0^{+\infty} \{ \cos(2tr) \Phi(r) + \sin(2tr) r \Psi(r) \} r^{d-1} dr, \quad (11)$$

where

$$\begin{aligned} \Phi(r) &= \int_{S(\mathfrak{a})} \{ r^2 \mathcal{F}f(r\sigma) \tilde{\mathcal{F}}f(r\sigma) - \mathcal{F}g(r\sigma) \tilde{\mathcal{F}}g(r\sigma) \} \nu(r\sigma) d\sigma, \\ \Psi(r) &= \int_{S(\mathfrak{a})} \{ \mathcal{F}f(r\sigma) \tilde{\mathcal{F}}g(r\sigma) + \mathcal{F}g(r\sigma) \tilde{\mathcal{F}}f(r\sigma) \} \nu(r\sigma) d\sigma, \end{aligned}$$

and $d\sigma$ denotes the surface measure on the unit sphere $S(\mathfrak{a})$ in \mathfrak{a} . Let $\gamma_0 \in (0, +\infty]$ be the width of the largest horizontal strip $|\operatorname{Im} z| < \gamma_0$ in which $z \mapsto \nu(z\sigma)$ is holomorphic for all directions $\sigma \in S(\mathfrak{a})$.

Lemma 4.1. (i) $\Phi(z)$ and $\Psi(z)$ extend to even holomorphic functions in the strip $|\operatorname{Im} z| < \gamma_0$.

(ii) If $\gamma_0 < +\infty$, the following estimate holds in every substrip $|\operatorname{Im} z| \leq \gamma$ with $\gamma < \gamma_0$: For every $N > 0$, there is a constant $C > 0$ (depending on $f, g \in \mathcal{C}_R^\infty(\mathfrak{a})$, N and γ) such that

$$|\Phi(z)| + |\Psi(z)| \leq C |z|^{|\mathcal{R}_0^+|} (1+|z|)^{-N} e^{2R|\operatorname{Im} z|}.$$

(iii) If $\gamma_0 = +\infty$, the previous estimate holds uniformly in \mathbb{C} .

Proof. (i) follows from the definition of Φ and Ψ . Let us turn to the estimates (ii) and (iii). On one hand, according to the Paley-Wiener Theorem (Theorem 2.5), all transforms $\mathcal{F}f(z\sigma)$, $\tilde{\mathcal{F}}f(z\sigma)$, $\mathcal{F}g(z\sigma)$, $\tilde{\mathcal{F}}g(z\sigma)$ are $O(\{1+|z|\}^{-N} e^{R|\operatorname{Im} z|})$. On the other hand, let us discuss the behavior of the Plancherel measure. Consider first the case where all multiplicities are integers. Without loss of generality, we may assume that $k_\alpha \in \mathbb{N}^*$ and $k_{2\alpha} \in \mathbb{N}$ for every indivisible root α . Then

$$\begin{aligned} \nu(\lambda) &= \text{const.} \prod_{\alpha \in \mathcal{R}_0^+} \langle \lambda, \check{\alpha} \rangle \{ \langle \lambda, \check{\alpha} \rangle + i(k_\alpha + 2k_{2\alpha}) \} \\ &\quad \times \prod_{0 < j < k_\alpha} \{ \langle \lambda, \check{\alpha} \rangle^2 + j^2 \} \prod_{0 \leq \tilde{j} < k_{2\alpha}} \{ \langle \lambda, \check{\alpha} \rangle^2 + (k_\alpha + 2\tilde{j})^2 \} \end{aligned}$$

is a polynomial of degree $2|k| = 2 \sum_{\alpha \in \mathcal{R}_+} k_\alpha$. In general,

$$\nu(\lambda) = \text{const.} \pi(\lambda) \tilde{\nu}(\lambda),$$

where

$$\pi(\lambda) = \prod_{\alpha \in \mathcal{R}_0^+} \langle \lambda, \check{\alpha} \rangle$$

is a homogeneous polynomial of degree $|\mathcal{R}_0^+|$ and

$$\tilde{\nu}(\lambda) = \prod_{\alpha \in \mathcal{R}_0^+} \frac{\Gamma(i\langle \lambda, \check{\alpha} \rangle + k_\alpha)}{\Gamma(i\langle \lambda, \check{\alpha} \rangle + 1)} \frac{\Gamma(\frac{i\langle \lambda, \check{\alpha} \rangle + k_\alpha}{2} + k_{2\alpha})}{\Gamma(\frac{i\langle \lambda, \check{\alpha} \rangle + k_\alpha}{2})} \frac{\Gamma(-i\langle \lambda, \check{\alpha} \rangle + k_\alpha)}{\Gamma(-i\langle \lambda, \check{\alpha} \rangle + 1)} \frac{\Gamma(\frac{-i\langle \lambda, \check{\alpha} \rangle + k_\alpha}{2} + k_{2\alpha} + 1)}{\Gamma(\frac{-i\langle \lambda, \check{\alpha} \rangle + k_\alpha}{2})}$$

is an analytic function which never vanishes on \mathfrak{a} . Notice that $z \mapsto \nu(z\sigma)$ or $\tilde{\nu}(z\sigma)$ has poles for generic directions $\sigma \in S(\mathfrak{a})$ as soon as some multiplicities are not integers. Using Stirling's formula

$$\Gamma(\xi) \sim \sqrt{2\pi} \xi^{\xi - \frac{1}{2}} e^{-\xi} \quad \text{as } |\xi| \rightarrow +\infty \text{ with } |\arg z| < \pi - \varepsilon,$$

we get the following estimate for the Plancherel density, in each strip $|\operatorname{Im} z| < \gamma$ with $0 < \gamma < \gamma_0$:

$$|\nu(z\sigma)| \leq C |z|^{|\mathcal{R}_0^+|} (1+|z|)^{2|k|-|\mathcal{R}_0^+|}.$$

The estimates (ii) and (iii) follow easily from these considerations. \blacksquare

Proposition 4.2. *Assume that the dimension d is odd and that all multiplicities are integers. Then there exists a constant $C > 0$ (depending on the initial data $f, g \in \mathcal{C}_R^\infty(\mathfrak{a})$) such that, for every $\gamma \geq 0$ and $t \in \mathbb{R}$,*

$$|\mathcal{P}[u](t) - \mathcal{K}[u](t)| \leq C e^{2\gamma(R-|t|)}.$$

Proof. Evenness allows us to rewrite (11) as follows :

$$\mathcal{P}[u](t) - \mathcal{K}[u](t) = \frac{c_0}{4} \int_{-\infty}^{+\infty} e^{i2tr} \{ \Phi(r) - ir\Psi(r) \} r^{d-1} dr.$$

Let us shift the contour of integration from \mathbb{R} to $\mathbb{R} \pm i\gamma$, according to the sign of t , and estimate the resulting integral, using Lemma 4.1.iii. As a result, the difference of energy

$$\mathcal{P}[u](t) - \mathcal{K}[u](t) = \frac{c_0}{4} e^{-2\gamma|t|} \int_{-\infty}^{+\infty} e^{i2tr} \{ \Phi(r \pm i\gamma) - i(r \pm i\gamma) \Psi(r \pm i\gamma) \} (r \pm i\gamma)^{d-1} dr$$

is $O((1+\gamma)^{-N} e^{2\gamma(R-|t|)})$. \blacksquare

As an immediate consequence of the above statement and in view of the fact that $\gamma_0 = \infty$ when k is integer valued, we deduce the strict equipartition of energy (3) for $|t| \geq R$, by letting $\gamma \rightarrow \infty$.

Henceforth, we will drop the above assumption on k . By resuming the proof of Proposition 4.2 and using Lemma 4.1.ii instead of Lemma 4.1.iii, we obtain the following result.

Proposition 4.3. *Assume that the dimension d is odd. Then, for every $0 < \gamma < \gamma_0$, there is a constant $C > 0$ (depending on the initial data $f, g \in \mathcal{C}_R^\infty(\mathfrak{a})$) such that*

$$|\mathcal{P}[u](t) - \mathcal{K}[u](t)| \leq C e^{-2\gamma|t|} \quad \forall t \in \mathbb{R}.$$

As a corollary, we obtain the asymptotic equipartition of energy (4) in the odd dimensional case, with an exponential rate of decay. In the even dimensional case, the expression (11) cannot be handled by complex analysis and we proceed differently.

Proposition 4.4. *Assume that the dimension d is even. Then there is a constant $C > 0$ (depending on the initial data $f, g \in \mathcal{C}_R^\infty(\mathfrak{a})$) such that*

$$|\mathcal{P}[u](t) - \mathcal{K}[u](t)| \leq C (1+|t|)^{-d-|\mathcal{R}_0^+|} \quad \forall t \in \mathbb{R}.$$

Proof. The problem lies in the decay at infinity. According to lemma 4.1, $\Phi(r)$ and $\Psi(r)$ are divisible by r^D , where $D = |\mathcal{R}_0^+|$. Let us integrate (11) $d+D$ times by parts. This way

$$\int_0^{+\infty} \cos(2tr) r^{d-1} \underbrace{\left\{ \int_{S(\mathfrak{a})} \mathcal{F}g(r\sigma) \tilde{\mathcal{F}}g(r\sigma) \nu(r\sigma) d\sigma \right\}}_{\tilde{\Phi}(r)} dr$$

becomes

$$\pm \frac{1 \text{ or } 0}{(2t)^{d+D}} \frac{(d+D)!}{(D+1)!} \left(\frac{\partial}{\partial r}\right)^{D+1} \tilde{\Phi}(r) \Big|_{r=0} \pm \int_0^{+\infty} \frac{\cos(2tr) \text{ or } \sin(2tr)}{(2t)^{d+D}} \left(\frac{\partial}{\partial r}\right)^{d+D} \{r^{d-1} \tilde{\Phi}(r)\} dr$$

which is $O(|t|^{-d-D})$. Similarly

$$\int_0^{+\infty} \cos(2tr) r^{d+1} \left\{ \int_{S(\mathfrak{a})} \mathcal{F}f(r\sigma) \tilde{\mathcal{F}}f(r\sigma) \nu(r\sigma) d\sigma \right\} dr = O(|t|^{-d-D-2})$$

and

$$\int_0^{+\infty} \cos(2tr) r^d \Psi(r) dr = O(|t|^{-d-D-1}).$$

This concludes the proof of Proposition 4.4. ■

As a corollary, we obtain the asymptotic equipartition of energy (4) in the even dimensional case, with a polynomial rate of decay.

Remark 4.5. *Our result may not be optimal. In the W -invariant case, one obtains indeed the rate of decay $O(\{1+|t|\}^{-d-2|\mathcal{R}_0^+|})$ as in [6].*

References

- [1] Ben Said, S., *Huygens' principle for the wave equation associated with the trigonometric Dunkl-Cherednik operators*, Math. Research Letters **13** (2006), no. 1, 43–58.
- [2] Ben Said, S. and Ørsted, B., *The wave equation for Dunkl operators*, Math. (N.S.) **16** (2005), no. 3–4, 351–391.
- [3] Branson, T. and Ólafsson, G., *Equipartition of energy for waves in symmetric space*, J. Funct. Anal. **97** (1991), no. 2, 403–416.
- [4] Branson, T. Ólafsson, G. Pasquale, A., *The Paley–Wiener theorem and the local Huygens' principle for compact symmetric spaces*, Math. (N.S.) **16** (2005), no. 3–4, 393–428.
- [5] Branson, T. and Ólafsson, G., *The Paley–Wiener theorem for the Jacobi transform and the local Huygens' principle for root systems with even multiplicities*, Math. (N.S.) **16** (2005), no. 3–4, 429–442.
- [6] Branson, T. Ólafsson, G. Schlichtkrull, H., *Huygens' principle in Riemannian symmetric spaces*, Math. Ann. **301** (1995), no. 3, 445–462.
- [7] El Kamel, J. Yacoub, C., *Huygens' principle and equipartition of energy for the modified wave equation associated to a generalized radial Laplacian*, Ann. Math Blaise Pascal **12** (2005), no. 1, 147–160.

- [8] Helgason, S., *Geometric analysis on symmetric spaces*, Math. Surveys Monographs 39, Amer. Math. Soc. (1994).
- [9] Opdam, E. M. , *Harmonic analysis for certain representations of graded Hecke algebras*, Acta. Math. 175 (1995), 75–121.
- [10] Opdam, E. M. , *Lecture notes on Dunkl operators for real and complex reflection groups*, Math. Soc. Japan Mem. 8 (2000).

F. Ayadi

Université de Tunis El Manar

Faculté des Sciences de Tunis

Département de Mathématiques

2092 Tunis El Manar, Tunisie

ayfatm@yahoo.fr

&

Université d'Orléans & CNRS

Fédération Denis Poisson (FR 2964)

Laboratoire MAPMO (UMR 6628)

Bâtiment de Mathématiques

B.P. 6759, 45067 Orléans cedex 2,

France

Fatma.Ayadi@etu.univ-orleans.fr

Received

and in final form