

journal of thrombosis and haemostasis

Mosaicism in men in haemophilia : is it exceptional? Impact on genetic counselling.



Journal:	<i>Journal of Thrombosis and Haemostasis</i>
Manuscript ID:	JTH-2008-00983.R1
Manuscript Type:	Letter - to the Editor
Date Submitted by the Author:	n/a
Complete List of Authors:	COSTA, Catherine; AP-HP, Groupe Henri Mondor-Albert Chenevier, Laboratoire de Génétique Moléculaire; INSERM U841, Département de Génétique, équipe 11, Université Paris 12 FRANCES, Anne-Marie; Centre Hospitalier Intercommunal Toulon La Seyne sur mer, Genetique Medicale LETOURNEAU, Sylvia; Centre Hospitalier Universitaire AP-HP Groupe Henri Mondor-Albert Chenevier, Laboratoire de genetique Moleculaire GIRODON-BOULANDET, Emmanuelle; Centre Hospitalier Intercommunal Toulon La Seyne sur mer, Genetique Medicale; INSERM U841, departement de genetique equipe 11 Université Paris 12 GOOSSENS, Michel; Centre Hospitalier Universitaire AP-HP Groupe Henri Mondor-Albert Chenevier, Laboratoire de genetique Moleculaire; INSERM U841, departement de genetique equipe 11 Université Paris 12
Key Words:	genetic counselling, haemophilia, mosaic

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Mosaicism in men in haemophilia : is it exceptional? Impact on genetic counselling.

C Costa^{*,†}, AM Frances[‡], S Letourneau^{*}, E Girodon-Boulandet^{*,†} and M Goossens^{*,†}.

* AP-HP, Groupe Henri Mondor-Albert Chenevier, Laboratoire de Génétique Moléculaire, † INSERM U841, Département de Génétique, équipe 11, Université Paris 12, Créteil F-94000, ‡ Centre Hospitalier Intercommunal Toulon la Seyne sur mer, Génétique Médicale, Toulon, France.

short title: mosaicism in men in haemophilia

key words : genetic counselling, haemophilia, mosaic

For Peer Review

1
2
3 Haemophilia A is an X-linked bleeding disorder caused by a wide range of mutations in the
4 factor VIII (*F8*) gene [1]. About one third of cases are due to a *de novo* mutation. The
5
6
7
8 majority are thought to occur in a single germ cell but some, occurring during early
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Haemophilia A is an X-linked bleeding disorder caused by a wide range of mutations in the factor VIII (*F8*) gene [1]. About one third of cases are due to a *de novo* mutation. The majority are thought to occur in a single germ cell but some, occurring during early embryogenesis, produce a germline and/or somatic mosaic. In haemophilia, somatic mosaicism has been generally observed in women and seems to represent a fairly common event [2]. We report here a case of exceptional mosaicism in the asymptomatic maternal grandfather of a haemophilia A patient.

The proband has severe haemophilia A with factor (F)VIIIc levels <1% and no previous family history of the disorder. Gene mutation studies were performed in order to identify the deleterious mutation and offer genetic counselling to the mother and the family. The mutation p.Arg336X in exon 8 was identified in the proband by direct sequencing and subsequently searched for in the mother and maternal grandmother. It was found only in the mother, suggesting a *de novo* germline mutation in one of the grandparents or a *de novo* somatic mutation early during embryogenesis in the mother. The maternal aunt, who had not been tested, was at first reassured as being probably not a carrier. Several years later, when undergoing medically assisted procreation because of the infertility of her partner, a genetic test was performed. Unexpectedly, the mutation p.Arg336X was identified, leading to a modification of her status as being a carrier of severe haemophilia A. The presence of the mutation in the two sisters thus first suggested the grandmother was a carrier with a somatic mosaicism. The absence of the mutation in her peripheral blood as well as in her buccal and uroepithelial cells, which have different embryological origins, then raised the question of the mechanism of occurrence of this mutation. Linkage analysis, using intragenic and extragenic markers linked to the *F8* gene, actually showed that the deleterious allele originated from the asymptomatic maternal grandfather whose FVIIIc was normal FVIIIc=96% (Fig1).

1
2
3 Somatic mosaicism in the grandfather was then hypothesized. However, it is well known that
4 somatic mosaicism may be difficult to detect with conventional methods such as direct
5 sequencing. Mutation-enrichment procedures, not used during routine test analyses, are often
6 required [2]. Nowadays, due to technology progress, methods presenting higher sensitivity are
7 available. One of them, denaturing-high-liquid-pressure-chromatography (DHPLC) was used
8 in this family. DHPLC is well known for its efficiency to detect heteroduplexes that are DNA
9 molecules containing mismatched base pairs and created during amplification reaction (PCR)
10 when a mutation is present in heterozygosity. Under partial denaturation, heteroduplexes are
11 eluted from the column by an acetonitrile gradient flow before homoduplexes [3]. Analysis of
12 the grandfather's leucocytes, buccal and uroepithelial cells showed the presence of the
13 mutated allele with a proportion estimated between 15-20% (fig 1). Karyotype analysis
14 showed a normal 46,XY karyotype, ruling out Klinefelter syndrome.

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
The presence of the mutation in all tested grandpaternal tissues and in his two daughters
suggested that the mutation had arisen very early during embryonic development.

The distinction between isolated or sporadic cases is of major importance in genetic
counselling. A case may appear to be isolated because family size is small; DNA testing may
help for carrier diagnosis but negative results will not rule out the possibility of an occult
mosaic. In a recent study only a small number (11%) of maternal grandmothers of isolated
cases had the mutation in their white blood cells, while 85% of mothers were carriers, which
favours the hypothesis that isolated cases may have originated as a *de novo* germline mutation
in one of the grandparents or a *de novo* somatic mutation early during embryogenesis in the
proband's mother [2, 4]. Somatic mosaicism has been found in around 10% of mothers of
isolated cases [2] and in 13% of patients' mothers and grandmothers in a study which used
mutation enrichment procedures [5]. These results indicate that mosaicism is a fairly
common event in haemophilia, but is still underestimated due to the limited sensitivities of the

1
2
3 methods for detection of mosaicism and probably also because the distinction between carrier
4 and 50% mosaicism is difficult. It is of note that most of the time, mosaicism has been
5
6 reported in families with point mutations while only once in an isolated case with intron 22
7
8 inversion. [2, 5, 6]. Somatic mosaicism in families with apparent *de novo* mutations is
9
10 however rarely explored in women, and grandfathers are usually not considered. In this
11
12 present case we have been questioned because the proband's mother and aunt were carriers
13
14 while the grandmother was not. In the literature only three cases of mosaicism in men have
15
16 been reported, all of them in grandfathers in families with point mutations [7-9]. Our case
17
18 underlies that somatic mosaicism in men is probably underestimated because of the
19
20 difficulties of obtaining blood sample from grandfathers, and points to the need for testing
21
22 men in such apparent isolated cases. In these situations and even if the mutation is
23
24 characterized, linkage analysis remains a precious help to identify the origin of the deleterious
25
26 allele.

27
28 Assessment of mosaicism in mothers of apparent isolated cases is now part of genetic
29
30 counselling. It also seems important now to take into account the risk of mosaicism in
31
32 grandfathers as well as grandmothers with a view to the genetic counselling of all their
33
34 daughters.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1-HAMSTeRS haemophilia A Mutation database: <http://europium.csc.mrc.ac.uk/>

2- Leuer M, Oldenburg J, Lavergne JM, Ludwig M, Fregin A, Eigel A, Ljung R, Goodeve A, Peake I, Olek K. Somatic mosaicism in hemophilia A: a fairly common event. *Am J Hum Genet* 2001; **69**: 75-87.

3- Xiao W, Oefner PJ. Denaturing high-performance liquid chromatography: A review. *Hum Mutat.* 2001;**17**:439-74.

4- Kasper CK, Lin JC. Prevalence of sporadic and familial haemophilia. *Haemophilia.* 2007; **13**: 90-2.

5- Peyvandi F, Jayandharan G, Chandy M, Srivastava A, Nakaya SM, Johnson MJ, Thompson AR, Goodeve A, Garagiola I, Lavoretano S, Menegatti M, Palla R, Spreafico M, Tagliabue L, Asselta R, Duga S, Mannucci PM. Genetic diagnosis of haemophilia and other inherited bleeding disorders. *Haemophilia* 2006; **12**: 82-89.

6- Oldenburg J, Ros S, El-Maarri O, Leuer M, Olek K, Müller CR, and Schwaab R. De novo *factor VIII* gene intron 22 inversion in a female carrier presents as a somatic mosaicism. *Blood* 2000; **96**:2905-2906.


7- Casey GJ, Rodgers SE, Hall JR, Rudzki Z, Lloyd JV. Grandpaternal mosaicism in a family with isolated haemophilia A. *Br J Haematol* 1999; **107**: 560-2.

1
2
3 8- Lebo RV, Koerper MA, Kim JH, Chueh J, Golbus MS. Prenatal diagnosis of hemophilia
4 involving grandpaternal mosaicism. *Am J Med Genet* 1993; **47**: 401-4.
5
6
7

8
9
10 9- Cutler JA, Mitchell MJ, Smith MP, Savidge GF. Germline mosaicism resulting in the
11 transmission of severe hemophilia B from a grandfather with a mild deficiency. *Am J Med*
12 *Genet A*. 2004; **129**:13-5.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Figure 1



review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Legend to Figure 1 : A-Family pedigree, haplotypes and mutation studies. B-
Electropherogram obtained by direct sequencing. C-Denaturing High Liquid Chromatography
(DHPLC) elution profiles of carrier, control and grandfather: in female carriers,
heteroduplexes, which contain mismatched base pairs, are eluted first (left peaks) followed by
the homoduplexes (right peaks); normal control have only one right peak. Analysis of the
grandfather's leucocytes and buccal cells shows a right peak corresponding to the normal
allele and a small left peak indicative of the presence of the mutated allele with a proportion
estimated between 15-20%.
The mutation is detected with higher sensitivity with DHPLC compared to direct sequencing.