



HAL
open science

A note on Renner monoids

Eddy Godelle

► **To cite this version:**

| Eddy Godelle. A note on Renner monoids. 2009. hal-00373601

HAL Id: hal-00373601

<https://hal.science/hal-00373601>

Preprint submitted on 6 Apr 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A NOTE ON RENNER MONOIDS

GODELLE EDDY

ABSTRACT. We extend the result obtained in [3] to every Renner monoid: we provide Renner monoids with a monoid presentation and we introduce a length function which extends the Coxeter length function and which behaves nicely.

INTRODUCTION

The notion of a *Weyl group* is crucial in Linear Algebraic Group Theory [4]. The seminal example occurs when one considers the *algebraic group* $GL_n(K)$. In that case, the associated Weyl group is isomorphic to the group of monomial matrices, that is to the permutation group S_n . Weyl groups are special examples of *finite Coxeter groups*. Hence, they possess a group presentation of a particular type, and an associated length function. It turns out that this presentation and this length function are deeply related with the geometry of the associated algebraic group. Linear Algebraic Monoid Theory, mainly developed by Putcha, Renner and Solomon, has deep connections with Algebraic Group Theory. In particular, the *Renner monoid* [14] plays the role that the Weyl group does in Linear Algebraic Group Theory. As far as I know, in the case of Renner monoids, there is no known theory that plays the role of Coxeter Group Theory. Therefore it is natural to look for such a theory, and, therefore, to address the question of monoid presentations for Renner monoids. In [3], we considered the particular case of the *rook monoid* defined by Solomon [18]. We obtained a presentation of this group and introduced a length function, that is nicely related to the Hecke Algebra of the rook monoid. Our objective here is to consider the general case. We obtain a presentation of every Renner monoid and introduce a length function. In the case of the rook monoid, we recover the results obtained in [3]. Our length function is not the classical length function on Renner monoids [14]. We remark that the former shares with the latter several nice geometrical and combinatorial properties. We believe that this new length function is nicely related to the Hecke algebra, as in the special case of the rook monoid.

Let us postpone to the next sections some definitions and notations, and state here our main results. Consider the Renner monoid $R(M)$ of a linear algebraic monoid M . Denote by W the unit group of $R(M)$ and consider its associated Coxeter system (W, S) . Denote by Λ a *cross section lattice* of the monoid $E(R(M))$ of idempotent elements of $R(M)$, and by Λ_\circ the set of elements of Λ that are distinct from the unity. Denote finally by λ the associated *type map* of $R(M)$; roughly speaking, this is a map that describes the action of W on $E(R(M))$.

Theorem 0.1. *The Renner monoid $R(M)$ admits the monoid presentation whose generating set is $S \cup \Lambda_\circ$ and whose defining relations are:*

$$\begin{aligned} (\text{COX1}) \quad & s^2 = 1, & s \in S; \\ (\text{COX2}) \quad & |s, t\rangle^m = |t, s\rangle^m, & (\{s, t\}, m) \in \mathcal{E}(\Gamma); \\ (\text{TYM1}) \quad & se = es, & e \in \Lambda_\circ, s \in \lambda^*(e); \\ (\text{TYM2}) \quad & se = es = e, & e \in \Lambda_\circ, s \in \lambda_*(e); \\ (\text{TYM}) \quad & \underline{ew}f = e \wedge_w f, & e, f \in \Lambda_\circ, w \in G^\dagger(e) \cap D^\dagger(f). \end{aligned}$$

We define the length ℓ on $R(M)$ in the following way: if s lies in S , we set $\ell(s) = 1$; if e lies in Λ we set $\ell(e) = 0$. Then, we extend ℓ by additivity to the free monoid of words on $S \cup \Lambda_\circ$. If w lies in $R(M)$, its length $\ell(w)$ is the minimal length of its word representatives on $S \cup \Lambda_\circ$. In Section 2, we investigate the properties of this length function. In particular we prove that it is nicely related to the classical *normal form* defined on $R(M)$, and we also prove

Proposition 0.2. *Let T be a maximal torus of the unit group of M . Fix a Borel subgroup B that contains T . Let w lie in $R(M)$ and s lie in S . Then,*

$$BsBwB = \begin{cases} BwB & \text{if } \ell(sw) = \ell(w); \\ BswB & \text{if } \ell(sw) = \ell(w) + 1; \\ BswB \cup BwB & \text{if } \ell(sw) = \ell(w) - 1. \end{cases}$$

This article is organized as it follows. In section 1, we first recall the backgrounds on Algebraic Monoid Theory and on Coxeter Theory. Then, we prove Theorem 0.1. In Section 2, we consider several examples of Renner monoids and deduce explicit presentations from Theorem 0.1. In Section 3, we focus on the length function.

1. PRESENTATION FOR RENNER MONOIDS

Our objective in the present section is to associate a monoid presentation to every Renner monoid. The statement of our result and its proof require some properties of algebraic monoid theory and of Coxeter group theory. In Section 1.1, we introduce Renner monoids and we state the results we need about algebraic monoids. In Section 1.2 we recall the definition of Coxeter groups and some of its well-known properties. Using the two preliminary sections, we can prove Theorem 0.1 in Section 1.3. This provides a monoid presentation to every Renner monoid.

We fix an algebraically closed field \mathbb{K} . We denote by M_n the set of all $n \times n$ matrices over \mathbb{K} , and by GL_n the set of all invertible matrices in M_n . We refer to [11, 14, 17] for the general theory and proofs involving linear algebraic monoids and Renner monoids; we refer to [4] for an introduction to linear algebraic groups. If X is a subset of M_n , we denote by \bar{X} its closure for the Zariski topology.

1.1. Algebraic Monoid Theory. We introduce here the basic definitions and notation on Algebraic Monoid Theory that we shall need in the sequel.

1.1.1. Regular monoids and reducible groups.

Definition 1.1 (Algebraic monoid). An *algebraic monoid* is a submonoid of M_n , for some positive integer n , that is closed for the Zariski topology. An algebraic monoid is *irreducible* if it is irreducible as a variety.

It is very easy to construct algebraic monoids. Indeed, the Zariski closure $M = \overline{G}$ of any submonoid G of M_n is an algebraic monoid. The main example occurs when for G one considers an algebraic subgroup of GL_n . It turns out that in this case,

the group G is the unit group of M . Conversely, if M is an algebraic monoid, then its unit group $G(M)$ is an algebraic group. The monoid M_n is the seminal example of an algebraic monoid, and its unit group GL_n is the seminal example of an algebraic group.

One of the main differences between an algebraic group and an algebraic monoid is that the latter have idempotent elements. In the sequel we denote by $E(M)$ the set of idempotent elements of a monoid M . We recall that M is *regular* if $M = E(M)G(M) = G(M)E(M)$, and that M *has a zero element* if there exists an element 0 such that $0 \times m = m \times 0 = 0$ for every m in M . The next result, which is the starting point of the theory, was obtained independently by Putcha and Renner in 1982.

Theorem 1.2. *Let M be an irreducible algebraic monoid with a zero element. Then M is regular if and only if $G(M)$ is reductive.*

The order \leq on $E(M)$ defined by $e \leq f$ if $ef = fe = e$, provides a natural connection between the Borel subgroups of $G(M)$ and the idempotent elements of M :

Theorem 1.3. *Let M be a regular irreducible algebraic monoid with a zero element. Let $\Gamma = (e_1, \dots, e_k)$ be a maximal increasing sequence of distinct elements of $E(M)$. (i) The centralizer $Z_{G(M)}(\Gamma)$ of Γ in $G(M)$ is a maximal torus of the reductive group $G(M)$. (ii) Set*

$$B^+(\Gamma) = \{b \in G(M) \mid \forall e \in \Gamma, be = ebe\},$$

$$B^-(\Gamma) = \{b \in G(M) \mid \forall e \in \Gamma, eb = ebe\}.$$

Then, $B^-(\Gamma)$ and $B^+(\Gamma)$ are two opposed Borel subgroups that contain $Z_{G(M)}(\Gamma)$.

1.1.2. Renner monoid.

Definition 1.4 (Renner monoid). Let M be a regular irreducible algebraic monoid with a zero element. If T is a Borel subgroup of $G(M)$, then we denote its normalizer by $N_{G(M)}(T)$. The *Renner monoid* $R(M)$ of M is the monoid $\overline{N_{G(M)}(T)}/T$.

It is clear that $R(M)$ does not depend on the choice of the maximal torus of $G(M)$.

Example 1.5. Consider $M = M_n(K)$, and choose the maximal torus \mathbb{T} of diagonal matrices. The Renner monoid is isomorphic to the monoid of matrices with at most one nonzero entry, that is equal to 1, in each row and each column. This monoid is called the rook monoid R_n [19]. Its unit group is the group of monomial matrices, which is isomorphic to the symmetric group S_n .

It is almost immediate from the definition that we have

Proposition 1.6. *Let M be a regular irreducible algebraic monoid with a zero element, and fix a maximal torus T of $G(M)$. The Renner monoid $R(M)$ is a finite factorisable inverse monoid. In particular, the set $E(R(M))$ is a commutative monoid and a lattice for the partial order \leq defined by $e \leq f$ when $ef = e$. Furthermore, there is a canonical order preserving isomorphism of monoids between $E(R(M))$ and $E(\overline{T})$.*

1.2. Coxeter Group Theory. Here we recall some well-known facts about Coxeter groups. We refer to [1] for general theory and proof.

Definition 1.7. Let Γ be a finite simple labelled graph whose labels are positive integers greatest or equal than 3. We denote by S the vertex set of Γ . We denote by $\mathcal{E}(\Gamma)$ the set of pairs $(\{s, t\}, m)$ such that either $\{s, t\}$ is an edge of Γ labelled by m or $\{s, t\}$ is not an edge of Γ and $m = 2$. When $(\{s, t\}, m)$ belongs to $\mathcal{E}(\Gamma)$, we denote by $|s, t|^m$ the word $sts \cdots$ of length m . The *Coxeter group* $W(\Gamma)$ associated with Γ is defined by the following group presentation

$$\left\langle S \mid \begin{array}{l} s^2 = 1 \\ |s, t|^m = |t, s|^m \end{array} \quad \begin{array}{l} s \in S \\ (\{s, t\}, m) \in \mathcal{E}(\Gamma) \end{array} \right\rangle$$

In this case, one says that (W, S) is a *Coxeter system*.

Proposition 1.8. *Let M be a regular irreducible algebraic monoid with a zero element, and denote by G its unit group. Fix a maximal torus T included in B . Then*

- (i) *the Weyl group $W = N_G(T)/T$ of G is a finite Coxeter group.*
- (ii) *The unit group of $R(M)$ is the Weyl group W .*

Remark 1.9. Gathering the results of Propositions 1.6 and 1.8 we get

$$R(M) = E(\overline{T}) \cdot W = W \cdot E(\overline{T}).$$

Definition 1.10. Let (W, S) be a *Coxeter system*. Let w belong to W . The *length* $\ell(w)$ of w is the minimal integer k such that w has a word representative of length of k on the alphabet S . Such a word is called a *reduced word representative* of w .

In the sequel, we use the following classical result [1].

Proposition 1.11. *Let (W, S) be a Coxeter system and I, J be subsets of S . Let W_I and W_J be the subgroups of W generated by I and J respectively.*

- (i) *The pairs (W_I, I) and (W_J, J) are Coxeter systems.*
- (ii) *For every element w which belongs to W there exists a unique element \hat{w} of minimal length in the double-class $W_J w W_I$. Furthermore there exists w_1 in W_I and w_2 in W_J such that $w = w_2 \hat{w} w_1$ with $\ell(w) = \ell(w_1) + \ell(\hat{w}) + \ell(w_2)$.*

Note that (ii) holds when I or J are empty.

1.3. Cross section. Our objective here is to prove Theorem 0.1. We first need to precise the notation used in this theorem. In all this section, we assume M is a regular irreducible algebraic monoid with a zero element. We denote by G the unit group of M . We fix a maximal torus T of G and we denote by W the Weyl group $N_G(T)/T$ of G . We denote by S the standard generating set associated with the canonical Coxeter structure of the Weyl group W .

1.3.1. The Cross Section Lattice. To describe the generating set of our presentation, we need to introduce the *cross section lattice*, which is related to Green's relations. The latter are classical tools in semigroup theory. Let us recall the definition of Relation \mathcal{J} . The \mathcal{J} -class of an element a in M is the double coset MaM . The set $\mathcal{U}(M)$ of \mathcal{J} -classes carries a natural partial order \leq defined by $MaM \leq MbM$ if $MaM \subseteq MbM$. It turns out that the map $e \mapsto MeM$ from $E(M)$ to $\mathcal{U}(M)$ induces a one-to-one correspondence between the set of W -orbits on $E(\overline{T})$ and the

set $\mathcal{U}(M)$. The existence of this one-to-one correspondence leads to the following definition:

Definition 1.12 (cross section lattice). A subset Λ of $E(\overline{T})$ is a *cross section lattice* if it is a transversal of $E(\overline{T})$ for the action of W such that the bijection $e \mapsto MeM$ from Λ onto $\mathcal{U}(M)$ is order preserving.

It is not immediatly clear that such a cross section lattice exists. Indeed it is, and

Theorem 1.13. [11, Theorem 9.10] *For every Borel subgroup B of G that contains T , we set*

$$\Lambda(B) = \{e \in E(\overline{T}) \mid \forall b \in B, be = ebe\}.$$

The map $B \mapsto \Lambda(B)$ is a bijection between the set of Borel subgroups of G that contain T and the set of cross section lattices of $E(T)$.

Example 1.14. Consider $M = M_n$. Consider the Borel subgroups \mathbb{B} of invertible upper triangular matrices and \mathbb{T} the maximal torus of invertible diagonal matrices. Denote by e_i the diagonal matrix $\begin{pmatrix} Id_i & 0 \\ 0 & 0 \end{pmatrix}$ of rank i . Then, the set $\Lambda(\mathbb{B})$ is $\{e_0, \dots, e_n\}$. One has $e_i \leq e_{i+1}$ for every index i .

Remark 1.15. (i) Let Γ be a maximal chain of idempotent elements of \overline{T} and consider the Borel subgroup $B^+(\Gamma)$ defined in Theorem 1.3. It follows from the definitions that we have $\Gamma \subseteq \Lambda(B^+(\Gamma))$.

(ii)[11, Def. 9.1] A cross section lattice is a sublattice of $E(\overline{T})$.

1.3.2. *Normal form and type map.* In order to state the defining relations of our presentation, we turn now to the notion of a type map. We fix a Borel subgroup of G that contains T . We write Λ for $\Lambda(B)$. We consider Λ as a sublattice of $E(R(M))$ (cf. Proposition 1.6).

Notation 1.16. [14] *Let e belong to Λ .(i) We set*

$$\lambda(e) = \{s \in S \mid se = es\}.$$

The map $\lambda : e \mapsto \lambda(e)$ is called the type map of the reductive monoid M .

(ii) *We set $\lambda_*(e) = \bigcap_{f \geq e} \lambda(f)$ and $\lambda^*(e) = \bigcap_{f \geq e} \lambda(f)$.*

(iii) *We set $W(e) = \{w \in W \mid we = ew\}$, $W_*(e) = \{w \in W(e) \mid we = e\}$ and $W^*(e) = \{w \in W(e) \mid we \neq e\}$.*

Proposition 1.17. [14, Lemma 7.15] . *Then*

(i) $\lambda_*(e) = \{s \in S \mid se = es = e\}$ and $\lambda^*(e) = \{s \in S \mid se = es \neq e\}$.

(ii) *The sets $W(e)$, $W_*(e)$ and $W^*(e)$ are the standard parabolic subgroups of W generated by the sets $\lambda(e)$ and $\lambda_*(e)$ and $\lambda^*(e)$, respectively. Furthermore, $W(e)$ is the direct product of $W_*(e)$ and $W^*(e)$.*

Notation 1.18. [14] *By Propositions 1.11 and 1.17, for every w in W and every e, f in Λ , each of the sets $wW(e)$, $W(e)w$, $wW_*(e)$, $W_*(e)w$ and $W(e)wW(f)$ has a unique element of minimal length. We denote by $D(e)$, $G(e)$, $D_*(e)$ and $G_*(e)$ the set of elements w of W that are of minimal length in their classes $wW(e)$, $W(e)w$, $wW_*(e)$ and $W_*(e)w$, respectively. Note that the set of elements w of W that are of minimal length in their double class $W(e)wW(f)$ is $G(e) \cap D(f)$.*

1.3.3. *Properties of the cross section lattice.* As in previous sections, we fix a Borel subgroup B of G that contains T , and denote by Λ the associated cross section lattice contained in $E(R(M))$. We use the notation $\mathcal{E}(\Gamma)$ of Section 1.2. We set $\Lambda_\circ = \Lambda - \{1\}$. To make the statement of Proposition 1.24 clear we need a preliminary result.

Lemma 1.19. *Let e_1, e_2 lie in $E(T)$ such that $e_1 \leq e_2$. There exists f_1, f_2 in Λ with $f_1 \leq f_2$ and w in W such that $wf_1w^{-1} = e_1$ and $wf_2w^{-1} = e_2$.*

Proof. Let Γ be a maximal chain of $E(R(M))$ that contains e_1 and e_2 . The Borel subgroup $B^+(\Gamma)$ contains the maximal torus T . Therefore, there exists w in W such that $w^{-1}B^+(\Gamma)w = B$. This implies that $w^{-1}\Lambda(B^+(\Gamma))w = \Lambda$. We conclude using Remark 1.15(i). \square

Lemma 1.20. *Let h, e belong to Λ such that $h \leq e$. Then, $W(h) \cap G(e) \subseteq W_\star(h)$ and $W(h) \cap D(e) \subseteq W_\star(h)$.*

Proof. Let w lie in $W(h) \cap G(e)$. We can write $w = w_1w_2 = w_2w_1$ where w_1 lies in $W_\star(h)$ and w_2 lies in $W^*(h)$. Since $h \leq e$, we have $\lambda^*(h) \subseteq \lambda^*(e)$ and $W^*(h) \subseteq W^*(e)$. Since w is assumed to belong to $G(e)$, this implies $w_2 = 1$. The proof of the second inclusion is similar. \square

Proposition 1.21. *Let e, f lie in Λ_\circ and w lie in $G(e) \cap D(f)$. There exists h in Λ_\circ with $h \leq e \wedge f$ such that w belongs to $W_\star(h)$ and $ewf = hw = h$.*

To prove the above proposition, we are going to use the existence of a normal decomposition in $R(M)$:

Proposition 1.22 ([14] Section 8.6). *For every w in $R(M)$ there exists a unique triple (w_1, e, w_2) with $e \in \Lambda$, $w_1 \in D_\star(e)$ and $w_2 \in G(e)$ such that $w = w_1ew_2$.*

Following [14], we call the triple (w_1, e, w_2) the *normal decomposition* of w .

Proof of Proposition 1.21. Consider the normal decomposition (w_1, h, w_2) of ewf . Then, w_1 belongs to $D_\star(h)$ and w_2 belongs to $G(h)$. The element $ewfw^{-1}$ is equal to $w_1hw_2w^{-1}$ and belongs to $E(R(M))$. Since w_1 lies in $D_\star(h)$, this implies that $w_3 = w_2w^{-1}w_1$ lies in $W_\star(h)$, and that $e \geq w_1hw_1^{-1}$. By Lemma 1.19, there exists w_4 in W and e_1, h_1 in Λ_\circ , with $e_1 \geq h_1$, such that $w_4e_1w_4^{-1} = e$ and $w_4h_1w_4^{-1} = w_1hw_1^{-1}$. Since Λ is a cross section for the action of W , we have $e_1 = e$ and $h_1 = h$. In particular, w_4 belongs to $W(e)$. Since w_1 belongs to $D_\star(h)$, we deduce that there exists r in $W_\star(h)$ such that $w_4 = w_1r$ with $\ell(w_4) = \ell(w_1) + \ell(r)$. Then, w_1 lies in $W(e)$. Now, write $w_2 = w_2''w_2'$ where w_2'' lies in $W^*(h)$ and w_2' belongs to $G_\star(h)$. One has $ewf = w_1w_2''hw_2'$, and w_1w_2'' lies in $D(h)$. By symmetry, we get that w_2' belongs to $W(f)$. Hence, $w_1^{-1}ww_2'^{-1}$ is equal to $w_3^{-1}w_2''$ and belongs to $W(h)$. By hypothesis w lies in $G(e) \cap D(f)$. Then we must have $\ell(w_3^{-1}w_2'') = \ell(w_1^{-1}) + \ell(w_2'^{-1}) + \ell(w)$. Since $w_3^{-1}w_2''$ belongs to $W(h)$, it follows that w_1 and w_2' belong to $W(h)$ too. This implies $w_1 = w_2' = 1$ and $w = w_3^{-1}w_2''$. Therefore, $ewf = hw_2'' = hw = wh$. Finally, w belongs to $W_\star(h)$ by Lemma 1.20. \square

1.3.4. *A presentation for $R(M)$.*

Notation 1.23. (i) For each w in W , we fix a reduced word representative \underline{w} .
(ii) We denote by $e \wedge_w f$ the unique letter in Λ that represents the element h in Proposition 1.21.

Note that for s in S , one has $\underline{s} = s$. We recall that Λ is a sub-lattice of $E(\overline{T})$ for the order \leq defined by $e \leq f$ if $ef = fe = e$. We are now ready to state a monoid presentation for $R(M)$:

Proposition 1.24. *The Renner monoid has the following monoid presentation whose generating set is $S \cup \Lambda_\circ$ and whose defining relations are:*

$$\begin{aligned} (\text{COX1}) \quad & s^2 = 1, & s \in S; \\ (\text{COX2}) \quad & |s, t|^m = |t, s|^m, & (\{s, t\}, m) \in \mathcal{E}(\Gamma); \\ (\text{TYM1}) \quad & se = es, & e \in \Lambda_\circ, s \in \lambda^*(e); \\ (\text{TYM2}) \quad & se = es = e, & e \in \Lambda_\circ, s \in \lambda_*(e); \\ (\text{TYM3}) \quad & \underline{ew}f = e \wedge_w f, & e, f \in \Lambda_\circ, w \in G(e) \cap D(f). \end{aligned}$$

Note that when $e \leq f$ and $w = 1$, then Relation (TYM3) becomes $ef = fe = e$. More generally, one has $e \wedge_1 f = e \wedge f$.

Proof. By remark 1.9 the submonoids $E(R)$ and W generate the monoid $R(M)$. As S is a generating set for W , it follows from the definition of Λ that the set $S \cup \Lambda_\circ$ generates $R(M)$ as a monoid. Clearly, Relations (COX1) and (COX2) hold in W , Relations (TYM1) and (TYM2) hold in $R(M)$. Relations (TYM3) hold in $R(M)$ by Proposition 1.21. It remains to prove that we obtain a presentation of the monoid $R(M)$. Let w belong to $R(M)$ with (w_1, e, w_2) as normal form. Consider any word ω on the alphabet $S \cup \Lambda_\circ$ that represents w . We claim that starting from ω , one can obtain the word $\underline{w_1 e w_2}$ using the relations of the above presentation only. This is almost obvious by induction on the number j of letters of the word ω that belong to Λ_\circ . The property holds for $j = 0$ (in this case $w = w_1$ and $e = w_2 = 1$) because (COX1) and (COX2) are the defining relations of the presentation of W . The case $j = 1$ is also clear, applying Relations (COX1), (COX2), (TYM1) and (TYM2). Now, for $j \geq 2$, the case j can be reduced to the case $j - 1$ using Relations (TYM3) (and the other relations). \square

The presentation in Proposition 1.24 is not minimal; some relations can be removed in order to obtain the presentation stated in Theorem 0.1. Let us introduce a notation used in this theorem:

Notation 1.25. *If e lies in Λ , we denote by $G^\dagger(e)$ the set $G(e) \cap \left(\bigcap_{f > e} W(f)\right)$. Similarly, we denote by $D^\dagger(e)$ the set $D(e) \cap \left(\bigcap_{f > e} W(f)\right)$.*

Remark 1.26. (i) $(\bigcap_{f > e} \lambda(f)) \cap \lambda_*(e) = \emptyset$ by Proposition 1.17.
(ii)

$$G^\dagger(e) = G(e) \cap W_{\bigcap_{f > e} \lambda(f)} \text{ and } D^\dagger(e) = D(e) \cap W_{\bigcap_{f > e} \lambda(f)}.$$

The reader may note that for $e \leq f$ one has $G^\dagger(e) \cap D^\dagger(f) = \{1\}$.

Proof of Theorem 0.1. We need to prove that every relations $\underline{ew}f = e \wedge_w f$ of type (TYM3) in Proposition 1.24 can be deduced from Relations (RBI) of Theorem 0.1, using the other common defining relations of type (COX1), (COX2), (TYM1) and (TYM2). We prove this by induction on the length of w . If $\ell(w) = 0$ then w is equal to 1 and therefore belongs to $G^\dagger(e) \cap D^\dagger(f)$. Assume $\ell(w) \geq 1$ and w does not belong to $G^\dagger(e) \cap D^\dagger(f)$. Assume furthermore w does not lie in $G^\dagger(e)$ (the other case is similar). Choose e_1 in Λ_\circ such that $e_1 > e$ and w does not lie in $W(e_1)$. Then, applying Relations (RIB), we can transform the word $\underline{ew}f$ into the word $\underline{ee_1 w}f$. Using relations (COX2), we can transform the

word \underline{w} into a word $\underline{w_1 w_2}$ where w_1 belongs to $W(e_1)$ and w_2 belongs to $G^\dagger(e_1)$. Then, applying Relations (COX2) and (TYM1), we can transform the word $ee_1\underline{w}f$ into the word $\underline{ew_1e_1w_2}f$. By hypothesis on w , we have $w_2 \neq 1$ and, therefore, $\ell(w_1) < \ell(w)$. Assume w_2 belongs to $D^\dagger(f)$. We can apply Relation (RIB) to transform $\underline{ew_1e_1w_2}f$ into $\underline{ew_1}(e_1 \wedge_{w_2} f)$. Using relations (COX2), we can transform $\underline{w_1}$ into a word $\underline{w_1' w_1'' w_1'''} with w_1''' in $W_*(e_1 \wedge_{w_2} f)$, w_1'' in $W^*(e_1 \wedge_{w_2} f)$ and w_1' in $D(e_1 \wedge_{w_2} f)$. Then $\underline{ew_1}(e_1 \wedge_{w_2} f)$ can be transformed into $\underline{ew_1'}(e_1 \wedge_{w_2} f)\underline{w_1''}$. Since $\ell(w_1') \leq \ell(w_1) < \ell(w)$, we can apply an induction argument to transform the word $\underline{ew_1}(e_1 \wedge_{w_2} f)$ into the word $e \wedge_{w_1} (e_1 \wedge_{w_2} f)\underline{w_1''}$. Now, by the unicity of the normal decomposition, w_1'' as to belong to $W_*(e \wedge_{w_1} (e_1 \wedge_{w_2} f))$. Therefore we can transform $e \wedge_{w_1} (e_1 \wedge_{w_2} f)\underline{w_1''}$ into $e \wedge_{w_1} (e_1 \wedge_{w_2} f)$ using Relations (TYM2). Note that the letters $e \wedge_{w_1} (e_1 \wedge_{w_2} f)$ and $e \wedge_w f$ has to be equal as they represent the same element in Λ . Assume finally that w_2 does not belong to $D^\dagger(f)$. By similar arguments we can, applying Relations (COX2) and (TYM1), transform the word $\underline{ew_1e_1w_2}f$ into a word $\underline{ew_1e_1w_3f_1w_4}$ where $f_1 > f$ in Λ_\circ and $w_2 = w_3w_4$ with w_3 in $D^\dagger(f_1)$ and w_4 in $W(f_1)$. At this stage we are in position to apply Relation (RBI). Thus, we can transform the word $\underline{ew_1e_1w_2}f$ into $\underline{ew_1}(e_1 \wedge_{w_3} f_1)\underline{w_4}f$. Since we have $\ell(w_1) + \ell(w_4) < \ell(w)$ we can proceed as in the first case to conclude. $\square$$

2. SOME PARTICULAR RENNER MONOIDS

Here we focus on some special Renner monoids considered in [6, 7, 8]. In each case, we provide an explicit monoid presentation using the general presentation obtained in Section 1.

2.1. The rook monoid. Consider $M = M_n$ and choose \mathbb{B} for Borel subgroup (see Example 1.14). In this case, the Weyl group is the symmetric group S_n . Its generating set S is $\{s_1, \dots, s_{n-1}\}$ where s_i is the transposition matrix corresponding to $(i, i+1)$. The cross section lattice $\Lambda = \{e_0, \dots, e_{n-1}, e_n\}$ is linear (we have $e_j \leq e_{j+1}$ for every j). For every j one has $\lambda_*(e_j) = \{s_i \mid j+1 \leq i\}$ and $\lambda^*(e_j) = \{s_i \mid i \leq j-1\}$. In particular, $G^\dagger(e_i) \cap D^\dagger(e_i) = \{1, s_i\}$, and for $i \neq j$ we have $G^\dagger(e_i) \cap D^\dagger(e_j) = \{1\}$.




FIGURE 1. Coxeter graph and Hasse diagram for M_n .

Therefore, we recover the monoid presentation of the rook monoid $R(M)$ stated in [3]: the generating set is $\{s_1, \dots, s_{n-1}, e_0, \dots, e_{n-1}\}$ and the defining relations

are

$$\begin{aligned}
s_i^2 &= 1, & 1 \leq i \leq n-1; \\
s_i s_j &= s_j s_i, & 1 \leq i, j \leq n-1 \text{ and } |i-j| \geq 2; \\
s_i s_{i+1} s_i &= s_{i+1} s_i s_{i+1}, & 1 \leq i \leq n-1; \\
e_j s_i &= s_i e_j & 1 \leq i < j \leq n-1; \\
e_j s_i &= s_i e_j = e_j & 0 \leq j < i \leq n-1; \\
e_i e_j &= e_j e_i = e_{\min(i,j)} & 0 \leq i, j \leq n-1; \\
e_i s_i e_i &= e_{i-1} & 1 \leq i \leq n-1.
\end{aligned}$$

2.2. The Symplectic Algebraic Monoid. Let n be a positive even integer and Sp_n be the *Symplectic Algebraic Group* [4, page 52]: assume ℓ lies in \mathbb{N} , and consider the matrix $J_\ell = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}$ in M_ℓ . Let $J = \begin{pmatrix} 0 & J_\ell \\ -J_\ell & 0 \end{pmatrix}$ in M_n , where $n = 2\ell$. Then, Sp_n is equal to $\{A \in M_n \mid A^t J A = J\}$, where A^t is the transpose matrix of A . We set $M = \overline{K^\times Sp_n}$. This monoid is a regular monoid with 0 whose associated reductive algebraic unit group is $K^\times Sp_n$. It is called the *Symplectic Algebraic Monoid* [6]. Let \mathbb{B} be the Borel subgroup of GL_n as defined in Example 1.14, and set $B = K^\times (\mathbb{B} \cap Sp_n)$. This is a Borel subgroup of the unit group of M . It is shown in [6] that the cross section lattice Λ of M is $\{e_0, e_1, \dots, e_\ell, e_n\}$ where the elements e_i correspond to the matrices of M_n defined in Example 1.14. In particular the cross section lattice Λ is linear. In this case, the Weyl group is a Coxeter group of type B_ℓ . In other words, the group W is isomorphic to the subgroup of S_n generated by the permutation matrices s_1, \dots, s_ℓ corresponding to $(1, 2)(n-1, n)$, $(2, 3)(n-2, n-1)$, \dots , $(\ell-1, \ell)(\ell+1, \ell+2)$, and $(\ell, \ell+1)$, respectively. One has $\lambda_*(e_i) = \{s_{i+1}, \dots, s_\ell\}$ and $\lambda^*(e_i) = \{s_1, \dots, s_{i-1}\}$. Therefore, $G^\uparrow(e_\ell) \cap D^\uparrow(e_\ell) = \{1, s_\ell, s_\ell s_{\ell-1} s_\ell\}$, and for i in $\{1, \dots, \ell-1\}$ one has $G^\uparrow(e_i) \cap D^\uparrow(e_i) = \{1, s_i\}$. A direct calculation proves that $e_i s_i e_i = s_i e_{i-1}$ for every i , and $e_\ell s_\ell s_{\ell-1} s_\ell e_\ell = e_{\ell-2}$. Hence, a monoid




FIGURE 2. Coxeter graph and Hasse diagram for Sp_n .

presentation of $R(M)$ is given by the generating set $\{s_1, \dots, s_\ell, e_0, \dots, e_\ell\}$ and the defining relations

$$\begin{aligned}
s_i^2 &= 1, & 1 \leq i \leq \ell; \\
s_i s_j &= s_j s_i, & 1 \leq i, j \leq \ell \text{ and } |i-j| \geq 2; \\
s_i s_{i+1} s_i &= s_{i+1} s_i s_{i+1}, & 1 \leq i \leq \ell-2; \\
s_\ell s_{\ell-1} s_\ell s_{\ell-1} &= s_{\ell-1} s_\ell s_{\ell-1} s_\ell; \\
e_j s_i &= s_i e_j & 1 \leq i < j \leq \ell; \\
e_j s_i &= s_i e_j = e_j & 0 \leq j < i \leq \ell; \\
e_i e_j &= e_j e_i = e_{\min(i,j)} & 0 \leq i, j \leq \ell; \\
e_i s_i e_i &= e_{i-1} & 1 \leq i \leq \ell; \\
e_\ell s_\ell s_{\ell-1} s_\ell e_\ell &= e_{\ell-2}.
\end{aligned}$$

2.3. The Special Orthogonal Algebraic Monoid. Let n be a positive integer and J_n be defined as in Section 2.2. The *Special Orthogonal Group* \mathbf{SO}_n is defined as $\mathbf{SO}_n = \{A \in SL_n \mid g^T J_n g = J_n\}$. The group $K^\times \mathbf{SO}_n$ is a connected reductive group. Following [7, 8], we define the *Special Orthogonal Algebraic Monoid* to be the Zariski closure $M = \overline{K^\times \mathbf{SO}_n}$ of $K^\times \mathbf{SO}_n$. This is an algebraic monoid [7, 8], and $B = \mathbb{B} \cap M$ is a Borel subgroup of its unit group. In this case, the cross section lattice depends on the parity of n . Furthermore, the Weyl group is a Coxeter group whose type depends on the parity of n too.




FIGURE 3. Coxeter graph and Hasse diagram for $\mathbf{SO}_{2\ell}$.

Assume $n = 2\ell$ is even. In this case, W is a Coxeter group of type D_{ℓ} . The standard generating set of W is $\{s_1, \dots, s_{\ell}\}$ where for $1 \leq i \leq \ell - 1$, the element s_i is the permutation matrix associated with $(i, i + 1)(n - i, n - i + 1)$, and s_{ℓ} is the permutation matrix associated with $(\ell - 1, \ell + 1)(\ell, \ell + 2)$. It is shown in [8] that the cross section Λ is equal to $\{e_0, e_1, \dots, e_{\ell}, f_{\ell}, e_n\}$. The elements e_i correspond to the matrices of M_n defined in Example 1.14; the element f_{ℓ} is the diagonal matrix $e_{\ell+1} + e_{\ell-1} - e_{\ell}$. The Hasse diagram of Λ is as represented in Figure 3. For j in $\{0, \dots, \ell - 2\}$ one has $\lambda_*(e_j) = \{s_i \mid j + 1 \leq i\}$ and $\lambda^*(e_j) = \{s_i \mid i \leq j - 1\}$. Furthermore, one can verified that

$$\lambda_*(e_{\ell-1}) = \lambda_*(f_{\ell}) = \lambda_*(e_{\ell}) = \emptyset,$$

$$\lambda^*(e_{\ell-1}) = \lambda^*(f_{\ell}) = \{s_i \mid i \leq \ell - 2\},$$

$$\lambda^*(e_{\ell}) = \{s_i \mid i \leq \ell - 1\},$$

$$\lambda^*(f_{\ell}) = \{s_i \mid i \neq \ell - 1\}.$$

Therefore, for i in $\{1, \dots, \ell - 2\}$ one has $G^\uparrow(e_i) \cap D^\uparrow(e_i) = \{1, s_i\}$. Furthermore,

$$G^\uparrow(e_{\ell-1}) \cap D^\uparrow(e_{\ell-1}) = \{1\} \text{ and}$$

$$G^\uparrow(f_{\ell}) \cap D^\uparrow(f_{\ell}) = \{1, s_{\ell-1}\} \quad ; \quad G^\uparrow(e_{\ell}) \cap D^\uparrow(e_{\ell}) = \{1, s_{\ell}\};$$

$$G^\uparrow(e_{\ell}) \cap D^\uparrow(f_{\ell}) = \{1, s_{\ell} s_{\ell-2} s_{\ell-1}\} \quad ; \quad G^\uparrow(f_{\ell}) \cap D^\uparrow(e_{\ell}) = \{1, s_{\ell-1} s_{\ell-2} s_{\ell}\}.$$

The monoid $R(M)$ has a presentation with $\{s_1, \dots, s_\ell, e_0, \dots, e_\ell, f_\ell\}$ for generating set and

$$\begin{aligned}
s_i^2 &= 1, & 1 \leq i \leq \ell; \\
s_i s_j &= s_j s_i, & 1 \leq i, j \leq \ell \text{ and } |i - j| \geq 2; \\
s_i s_{i+1} s_i &= s_{i+1} s_i s_{i+1}, & 1 \leq i \leq \ell - 2; \\
s_\ell s_{\ell-2} s_\ell &= s_{\ell-2} s_\ell s_{\ell-2}; \\
e_j s_i &= s_i e_j, & 1 \leq i < j \leq \ell; \\
e_j s_i &= s_i e_j = e_j, & 0 \leq j < i \leq \ell; \\
e_i e_j &= e_j e_i = e_{\min(i,j)}, & 0 \leq i, j \leq \ell; \\
f_\ell e_\ell &= e_\ell f_\ell = e_{\ell-1}; \\
e_i s_i e_i &= e_{i-1}, & 1 \leq i \leq \ell - 1; \\
e_\ell s_\ell e_\ell &= f_\ell s_{\ell-1} f_\ell = e_{\ell-2}; \\
e_\ell s_\ell s_{\ell-2} s_{\ell-1} f_\ell &= f_\ell s_{\ell-1} s_{\ell-2} s_\ell e_\ell = e_{\ell-3}.
\end{aligned}$$

for defining relations. Assume $n = 2\ell + 1$ is odd. In that case, W is a Coxeter group of type B_ℓ . It is shown in [7] that the cross section lattice is linear as in the case of the Symplectic Algebraic Monoid. It turns out that the Renner monoid of $\mathbf{SO}_{2\ell+1}$ is isomorphic to the Renner monoid of Symplectic Algebraic Monoid $\overline{K^\times Sp_{2\ell}}$, and that we obtain the same presentation than in the latter case.

2.4. More examples: Adjoint Representations. Let G be a simple algebraic group, and denote by \mathfrak{g} its Lie algebra. Let M be the algebraic monoid $\overline{K^\times Ad(G)}$ in $End(\mathfrak{g})$. The cross section lattice of M and the type map of M has been calculated for each Dynkin diagram (see [14, Sec. 7.4]). Therefore one can deduce a monoid presentation for each of the associated Renner monoid.

2.5. More examples: \mathcal{J} -irreducible algebraic monoids. In [13], Renner and Putcha consider among regular irreducible algebraic monoids those who are \mathcal{J} -irreducible, that is those whose cross section lattices have a unique minimal non-zero element. It is easy to see that the \mathcal{J} -irreducibility property is related to the existence of irreducible rational representations [13, Prop. 4.2]. Renner and Putcha determined the cross section lattice of those \mathcal{J} -irreducible that arise from special kind of dominant weights [13, Fig. 2,3]. Using [13, Theorem 4.13], one can deduce the associated type maps and therefore a monoid presentation of each corresponding Renner monoids.

3. A LENGTH FUNCTION ON $R(M)$

In this section we extend the length function defined in [3] to any Renner monoid. In all this section, we assume M is a regular irreducible algebraic monoid with a zero element. We denote by G the unit group of M . We fix a maximal torus T of G and we denote by W the Weyl group $N_G(T)/T$ of G . We denote by S the standard generating set associated with the canonical Coxeter structure of the Weyl group W . We fix a Borel subgroup B of G that contains T , and we denote by Λ the associated cross section lattice contained in $R(M)$. As before we set $\Lambda_\circ = \Lambda - \{1\}$.

3.1. Minimal word representatives. The definition of the length function on W and of a reduced word has been recalled in Section 1.2. If X is a set, we denote by X^* the set of finite words on X .

Definition 3.1. (i) We set $\ell(s) = 1$ for s in S and $\ell(e) = 0$ for e in Λ . Let x_1, \dots, x_k be in $S \cup \Lambda_\circ$ and consider the word $\omega = x_1 \cdots x_k$. Then, the *length* of the word ω is the integer $\ell(\omega)$ defined by $\ell(\omega) = \sum_{i=1}^k \ell(x_i)$.

(ii) The *length* of an element w which belongs to $R(M)$ is the integer $\ell(w)$ defined by

$$\ell(w) = \min \{ \ell(\omega), \omega \in (S \cup \Lambda_\circ)^* \mid \omega \text{ is a word representative of } w \}.$$

The following properties are direct consequences of the definition.

Proposition 3.2. *Let w belong to $R(M)$.*

(i) *The length function ℓ on $R(M)$ extends the length function ℓ defined on W .*

(ii) *If $\ell(w) = 0$ then w lies in Λ .*

(iii) *If s lies in S then $|\ell(sw) - \ell(w)| \leq 1$.*

(iv) *If w' belongs to $R(M)$, then $\ell(ww') \leq \ell(w) + \ell(w')$.*

Proof. Point (i) and (ii) are clear (the letters of every representative word of an element in W are in S). If w lie in $R(M)$ and s lie to S , then $\ell(sw) \leq \ell(w) + 1$. Since $w = s^2w = s(sw)$, the inequality $\ell(w) \leq \ell(sw) + 1$ holds too. Point (iii) follows, and Point (iv) is a direct consequence of (iii). \square

Proposition 3.3. *Let w belong to $R(M)$.*

(i) *If (w_1, e, w_2) is the normal decomposition of w , then $\ell(w) = \ell(w_1) + \ell(w_2)$.*

(ii) *If ω_1, ω_2 are two representative words of w on $S \cup \Lambda_\circ$ such that the equalities $\ell(w) = \ell(\omega_1) = \ell(\omega_2)$ hold, then using the defining relations of the presentation of $R(M)$ in Theorem 1.24, we can transform ω_1 into ω_2 without increasing the length.*

Proof. (i) Let ω be a representative word w on $S \cup \Lambda_\circ$ such that $\ell(w) = \ell(\omega)$. It is clear that we can repeat the argument of the proof of Theorem 1.24 without using the relation (BR1). Therefore $\ell(\omega) \geq \ell(w_1ew_2) = \ell(w_1) + \ell(w_2) \geq \ell(w)$.

(ii) is a direct consequence of the proof of (i). \square

Corollary 3.4. *Let w lie in $R(M)$ and e belongs to Λ_\circ . Denote by (w_1, f, w_2) the normal decomposition of w .*

(i) *One has $\ell(we) \leq \ell(w)$ and $\ell(ew) \leq \ell(w)$.*

(ii) *$\ell(we) = \ell(w)$ if and only if the normal decomposition of we is $(w_1, e \wedge f, w_2)$. Furthermore, in this case, w_2 lies in $W^*(e)$.*

Proof. (i) is a direct consequence of the definition of the length and of Proposition 3.3(i) : $\ell(we) = \ell(w_1fw_2e) \leq \ell(w_1) + 0 + \ell(w_2) + 0 = \ell(w)$. The same arguments prove that $\ell(ew) \leq \ell(w)$.

(ii) Decompose w_2 as a product $w'_2w''_2w'''_2$ where w'''_2 lies in $W_*(f)$, w''_2 lies in $W^*(f)$, w'_2 lies in $D(f)$ and $\ell(w_2) = \ell(w'_2) + \ell(w''_2) + \ell(w'''_2)$. Then, $we = w_1fw_2e = w_1fw'_2ew''_2 = w_1(f \wedge_{w'_2} e)w''_2$. In particular, $\ell(we) \leq \ell(w_1) + \ell(w''_2)$. Assume $\ell(we) = \ell(w)$. We must have $w'_2 = w'''_2 = 1$. The element w''_2 , that is w_2 , must belong to $D^*(f \wedge_1 e) = D^*(f \wedge e)$, and the element w_1 must belong to $G^*(f \wedge e)$. In particular, w_2 lies in $W^*(e)$. Furthermore, w_2 lies in $D(f \wedge e)$ since $\lambda^*(f \wedge e) \subseteq \lambda(f)$ by Proposition 1.17(i). Conversely, if the the normal decomposition of we is $(w_1, e \wedge f, w_2)$, then $\ell(we) = \ell(w_1) + \ell(w_2) = \ell(w)$. \square

3.2. Geometrical formula. In Proposition 3.6 below we provide a geometrical formula for the length function ℓ defined in the previous section. This formula extends naturally the geometrical definition of the length function on a Coxeter group. Another length function on Renner monoids has already been defined and investigated [16, 10, 14]. This length function has nice properties, which are similar to the ones in Propositions 3.2, 3.3(i) and 3.6. This alternative length function has been firstly introduced by Solomon [16] in the special case of rook monoids in order to verify a combinatorial formula that generalizes Rodrigues formula [15]. That is why we call this length function the *Solomon length function* in the sequel. We proved in [3] that our length function for the rook monoid verifies the same combinatorial formula. We also proved in [3] that in the case of the rook monoid, our presentation of $R(M)$ and our length function are related to the Hecke algebra. We believe this is still true in the general case.

Lemma 3.5. *Let w belong to $R(M)$ and denote by (w_1, e, w_2) its normal decomposition. Let s be in S .*

(i) *We have one of the two following cases:*

(a) *there exists t in $\lambda_*(e)$ such that $sw_1 = w_1t$. In this case, one has $sw = w$;*

(b) *the element sw_1 lies in $D_*(e)$ and (sw_1, e, w_2) is the normal decomposition of sw .*

(ii) *Denote by $\tilde{\ell}$ the Solomon length function on $R(M)$. Then,*

$$\ell(sw) - \ell(w) = \tilde{\ell}(sw) - \tilde{\ell}(w).$$

Proof. (i) If sw_1 lies in $D_*(e)$, then by Theorem 1.22, the triple (sw_1, e, w_2) is the normal decomposition of sw . Assume now that sw_1 does not belong to $D_*(e)$. In that case, e cannot be equal to 1. Since w_1 belongs to $D_*(e)$, by the exchange lemma, there exists t in $\lambda(e)$ such that $sw_1 = w_1t$. Therefore, $sw = sw_1ew_2 = w_1tew_2 = w_1ew_2 = w$.

(ii) The Solomon length $\tilde{\ell}(w)$ of an element w in $R(M)$ can be defined by the formula $\tilde{\ell}(w) = \ell(w_1) - \ell(w_2) + \tilde{\ell}_e$ where (w_1, e, w_2) is the normal decomposition of w and $\tilde{\ell}_e$ is a constant that depends on e only [10, Definition 4.1]. Therefore the result is a direct consequence of (i). \square

As a direct consequence of Lemma 3.5(ii) and [14, Theorem 8.18] we get Proposition 0.2.

Proposition 3.6. *Let w belong to $R(M)$, and (w_1, e, w_2) be its normal decomposition.*

$$\ell(w) = \dim(Bw_1eB) - \dim(Bew_2B).$$

When w lies in S_n , one has $e = w_2 = 1$, and we recover the well-known formula $\ell(w) = \dim(BwB) - \dim(B)$.

Proof. By [10, Section 4], for every normal decomposition (v_1, f, v_2) we have the equality $\dim(Bv_1fv_2B) = \ell(v_1) - \ell(v_2) + k_f$, where k_f is a constant that depends on f only. Therefore,

$$\dim(Bw_1eB) - \dim(Bew_2B) = \ell(w_1) + k_e - (-\ell(w_2) + k_e) = \ell(w).$$

\square

REFERENCES

- [1] BOURBAKI N. *éléments de mathématique. Fasc. XXXIV. Groupes et algèbres de Lie, Chapitre IV-V-VI*. Hermann, Paris 1968.
- [2] BRIESKORN E. Die Fundamentalgruppe des Raumes der regulären Orbits einer endlichen komplexen Spiegelungsgruppe. *Invent. Math.* 12 (1971) 57–61.
- [3] GODELLE E. The Braid rook monoid. *IJAC*. 18 (2008) 779–802.
- [4] HUMPHREYS J. *Linear algebraic groups*. Graduate Texts in Mathematics 21 Springer-Verlag, 1995.
- [5] HUMPHREYS J. *Reflection groups and Coxeter groups*. Cambridge Studies in Advanced Mathematics 29. Cambridge University Press, 1990.
- [6] LI Z. AND RENNER L. The Renner monoids and cell decompositions of the symplectic algebraic monoids. *IJAC* 13 (2003) 111–132.
- [7] LI Z. The Cross Section Lattices and Renner monoids of the odd Special Orthogonal algebraic monoids *semigroup forum* 66 (2003) 272–287.
- [8] LI Z. Idempotent lattices, Renner monoids and cross section lattices of the Special Orthogonal algebraic monoids. *J. Algebra* 270 (2003) 445–458.
- [9] LIPSCOMB S. *Symmetric inverse semigroups*. Mathematical Surveys and Monographs 46 American Mathematical Society, 1996.
- [10] PENNELL E. PUTCHA M. AND RENNER L. Analogue of the Bruhat-Chevalley Order for Reductive Monoids. *J. Algebra* 196 (1997) 339–368.
- [11] PUTCHA M. *Linear algebraic monoids*. London Mathematical Society Lecture Note Series 133 Cambridge University Press, Cambridge, 1988.
- [12] PUTCHA M. Monoid Hecke Algebras. *Trans. AMS* (1997) 3517–3534.
- [13] RENNER L. AND PUTCHA M. The system of idempotents and the Lattice of \mathcal{J} -classes of Reductive Algebraic Monoids. *J. Algebra* (1988) 385–399.
- [14] RENNER L. *Linear algebraic monoids*. Encyclopaedia of Mathematical Sciences 134, Invariant Theory and Algebraic Transformation Groups V. Springer-Verlag, 2005.
- [15] RODRIGUES, O. Note sur les inversions, ou dérangements produits dans les permutations. *J. de Math.* 4 (1839) 236–239.
- [16] SOLOMON L. The Bruhat decomposition, Tits system and Iwahori ring for the monoid of matrices over a finite field. *Geometrica Dedicata* (1990) 15–49.
- [17] SOLOMON L. An introduction to reductive monoids. *Semigroups, formal languages and groups*, J. Fountain, Ed., Klumer Academic Publishers (1995) 295–352.
- [18] SOLOMON L. Representations of the rook monoid. *J. Algebra* (2002) 309–242.
- [19] SOLOMON L. The Iwahori algebra of $\mathbf{M}_n(\mathbf{F}_q)$. A presentation and a representation on tensor space. *J. Algebra* (2004) 206–226.