Kinetics of the Oxidation of N-Aminopiperidine with Chloramine

Chaza Darwich, Mazen Elkhatib, Georg Steinhauser, Henri Delalu

► To cite this version:

Chaza Darwich, Mazen Elkhatib, Georg Steinhauser, Henri Delalu. Kinetics of the Oxidation of N-Aminopiperidine with Chloramine. *Kinetics and Catalysis*, 2009, 50 (1), pp.103-111. hal-00397533

HAL Id: hal-00397533

<https://hal.science/hal-00397533>

Submitted on 22 Jun 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Kinetics of the Oxidation of N-Aminopiperidine with Chloramine

Chaza Darwich^{1, *}, Mazen Elkhatab², Georg Steinhauser³ and Henri Delalu¹

¹ Laboratoire Hydrazines et Procédés, UMR 5179 CNRS-ISOCHEM (groupe SNPE), Université Claude Bernard Lyon 1, Bâtiment Berthollet, 22 Avenue Gaston Berger, F-69622 Villeurbanne Cedex, France. {e-mail: aromadervish@yahoo.com; delalu@univ-lyon1.fr, telephone : +33 4 72432664, fax : +33 4 72431291}.

² Laboratory of Applied Chemistry and Toxicology, Faculty of Sciences, Section 3, Department of Chemistry, P.O. Box 826, Tripoli, Lebanon {e-mail: mazen@ul.edu.lb, telephone : +961 3 490204, fax : +961 6 386365}.

³ Vienna University of Technology, Atominstitut der Österreichischen Universitäten, Stadionallee 2, A-1020 Vienna, Austria {e-mail: georg.steinhauser@ati.ac.at, telephone : +43 1 58 801 141 89, fax : +43 1 58 801 141 99}

* corresponding author

Abstract

The kinetics of the oxidation of N-aminopiperidine with chloramine was studied at different temperatures, with variable concentrations of the two reactants and at a pH ranging between 12 and 13.5. The reaction showed to be involving two steps: the first corresponded to the formation of a diazene intermediate, the second to the evolution of this intermediate into numerous compounds within a complex reactional chain. The rate law of the first step was determined by the Ostwald method and

found to be first order with respect to each reactant. The rate constant was determined at pH = 12.89 and T = 25°C:

$$k_2 = 1.15 \times 10^5 \exp(-39/RT) \text{ M}^{-1} \text{ s}^{-1} \quad (E_2 \text{ in kJ mol}^{-1})$$

With decreasing pH value, the first step exhibited acid catalysis phenomena, and diazene was converted into azopiperidine particularly faster. This created overlapping UV-absorptions between chloramine and azopiperidine, also observed in HPLC. GC/MS analyses were used to identify some of the numerous by-products formed. Their proportions are dependent of both pH and the reactants' concentrations ratio. A reaction mechanism taking this relationship into account, was suggested.

Keywords

Chloramine; Kinetics; N-aminopiperidine; Oxidation; Raschig process

Introduction

Heterocyclic compounds including an unsymmetrical hydrazine group are used in the pharmaceutical industry as precursors of medicinal drugs. In particular, N-aminopiperidine (**1**, NAPP, C₅H₁₂N₂) is a precursor of a selective CB1 endocannabinoid receptor antagonist, commercially known as *Rimonabant*, which is used for the treatment of obesity and for smoking cessation.

Presently, **1** is prepared by different methods, particularly in batch by the two following processes:

1. The Wright and Willette process [1], which is carried out in two steps (Scheme I):

- nitrosylation of 1-piperidine (PP) by addition of sodium nitrite to an acid solution of the amine (reaction 2)
- reduction of 1-nitrosopiperidine by a chemical or catalytic method (reaction 3)

Scheme I

This process leads to a high yield (75 to 92%) but, before the reduction step, the product of reaction 2 must be purified by distillation or recrystallization. Several methods are possible for its reduction [2-12]. Nevertheless, it must be handled with a lot of precaution because of its highly carcinogenic properties, which complicates synthesis on an industrial scale.

2. The urea method, which includes three steps: the first one is the preparation of 1-piperidylurea (reaction 4) followed by chlorination leading to 1-piperidyl-3-chlorourea (reaction 5). The latter is finally converted to N-aminopiperidine by addition of a concentrated solution of sodium hydroxide [13-15]. The reaction mechanism is a Hoffmann rearrangement and the yield obtained is about 82%.

However, this method would hardly be adapted to a continuous synthesis because it involves numerous steps (Scheme II).

Scheme II

Other methods reported in the literature were also used to prepare **1** [16-18], however many difficulties involving these methods were found. Therefore, to synthesize **1** in a high scale we have undertaken an aqueous amination of 1-piperidine by the Raschig process [19, 20]. This most environmentally-friendly route can be schematized by the following two reactions (Scheme III):

Scheme III

However, it presents the major drawback of leading to numerous by-products. This behavior is particularly due to the simultaneous oxidizing/aminating character of NH_2Cl . In particular, the reaction between **1** and chloramine is one of the principal side reactions observed during the synthesis of **1** by the Raschig process [21]. This reaction limits the yield and leads to the precipitation of by-products difficult to separate during the continuous extraction of **1** [21].

In this paper, we report a kinetic study of the monochloramine / N-aminopiperidine reaction.

Experimental

Reagents

All reagents and salts used were reagent grade products from ALDRICH[®] and PROLABO RP[®]. Water was passed through an ion-exchange resin, then distilled twice, deoxygenated and stored under nitrogen.

NH₂Cl is unstable in water, it was therefore prepared extemporaneously at -10°C by reacting 25 ml of sodium hypochlorite 2 M and 20 ml of NH₃/NH₄Cl aqueous solution ([NH₄Cl] = 2.3 M, [NH₃] = 3.6 M) in the presence of diethyl ether (40 ml). The organic layer (0.8 – 1 M of NH₂Cl) was shaken and washed several times with aliquots of distilled water. Aqueous solution of chloramine was obtained by re-extraction from the ethereal phase. Its concentration was determined by UV spectroscopy at $\lambda = 243$ nm ($\epsilon = 458 \text{ M}^{-1} \text{ cm}^{-1}$) [22]. N-aminopiperidine (97%) was delivered by ALDRICH[®].

Apparatus

The apparatus consisted of two thermo-stated vessels of borosilicate glass, one on the top of the other and joined by a conical fitting. The lower reactor (200 cm³) had inlets allowing the measurement of pH and temperature, influx of circulating nitrogen and removal of aliquots for analysis. Because of the sensitivity of hydrazines to oxidation upon exposure to air, the mixture was monitored by an

oxygen-sensitive electrode connected to a numerical indicator. The upper cylindrical vessel (100 cm³) was blocked at its base by a solid stopper (17 mm i.d.) fastened to a control rod. This set-up allows a quick introduction of the ampoule contents into the reactor and hence a precise definition of the start of the reaction. A slightly reduced pressure was maintained throughout the reaction mixture, and the reactor temperature was kept constant to $\pm 0.1^\circ\text{C}$ (thermocouple). A glass electrode (TACUSSEL TB/HS model) and a calomel reference electrode connected to a TACUSSEL ISIS 20000 pH meter were used for pH measurements.

Results and Discussion

Procedure and Analysis

Reactant solutions were prepared at the same pH (see Experimental): **1** was dissolved in deoxygenated water and introduced into the lower reactor. The pH value was adjusted by addition of sodium hydroxide and/or a buffer solution. When thermal equilibrium was reached, the aqueous solution of chloramine (prepared according to the above procedure) of identical pH was added from the upper vessel.

The concentration of chloramine was monitored by making use of its maximum ultraviolet absorption ($\epsilon_{\text{NH}_2\text{Cl}} = 458 \text{ M}^{-1} \text{ cm}^{-1}$ at $\lambda = 243 \text{ nm}$). It was analyzed either by UV spectrophotometry using a Cary 1E double-beam spectrophotometer or by HPLC using a HP 1100 chromatograph equipped with a Diode Array Detector. For experiments monitored by HPLC, the separation was done on a $150 \times 3 \text{ mm}$ ODS XDB-C₈ column ($d_p = 5 \mu\text{m}$) using MeOH/H₂O (70/30) as mobile phase (rate flow = 0.5 ml min^{-1}).

1 is transparent to UV, therefore reaction orders can be easily determined by analyzing the temporal chloramine evolution. In some experiments, aliquots were treated with formaldehyde (40-fold excess) in order to stop the reaction between NH_2Cl and **1** by converting the latter into its hydrazone (**2**, FNAPP, $\text{C}_6\text{H}_{12}\text{N}_2$), which has a maximum absorption in UV at 237 nm ($\epsilon_{\text{FNAPP}} = 4485 \text{ M}^{-1} \text{ cm}^{-1}$). This new method of NAPP derivatization with formaldehyde was recently developed in our laboratory (Scheme IV) [21].

Scheme IV

GC/MS analyses were carried out on a HP 5890 chromatograph coupled to a mass spectrometer HP 5970 equipped with a 30-m long CP-Sil C_{19} column (250 μm i.d., $d_f = 1.5 \mu\text{m}$). Methodological details on the apparatus and the experimental procedure have been published previously [23, 24].

Characterization of the reaction mixture

Figure 1 shows a UV spectrophotometric evolution of the mixture at different times of the reaction ($[\text{NAPP}]_0 = 20 \times 10^{-3} \text{ M}$, $[\text{NH}_2\text{Cl}]_0 = 2 \times 10^{-3} \text{ M}$, $\text{pH} = 12.89$ and $T = 25^\circ\text{C}$). In a first step, NH_2Cl absorption decreased at 243 nm, with the appearance of an isobestic point at 278 nm, which proved that the disappearance of chloramine is simultaneous to the formation of a new product **3** also absorbing in UV. Since **1** is

transparent in UV, **3** must result from the oxidation of **1** by chloramine (see Scheme V).

In a second step, the isobestic point disappeared while UV absorption increased and simultaneously deviated to lower wavelengths. At the end of the second step, the reaction mixture's evolution, monitored by UV spectrophotometry, led to a unique maximum at 237 nm, which is a characteristic wavelength for the absorption of hydrazones.

In order to understand the evolution of compounds in the reaction mixture with respect to the two steps described above, we compare the C₅H₁₀NNH₂/NH₂Cl interaction with the ones of CH₃NHNH₂/NH₂Cl, (CH₃)₂NNH₂/NH₂Cl, C₉H₉NNH₂/NH₂Cl and C₇H₁₂NNH₂/NH₂Cl which have been studied previously [23-25].

We found that the first elementary step of the NAPP oxidation leads transiently to an aminonitrene **3** (diazene intermediate, C₅H₁₀N₂) (Scheme V):

Scheme V

Figure 1. UV absorption spectra of the oxidation of *N*-aminopiperidine with chloramine (first step; $[C_5H_{10}NNH_2]_0 = 20 \times 10^{-3} M$, $[NH_2Cl]_0 = 2 \times 10^{-3} M$, $pH = 12.89$, $T = 25^\circ C$).

In the second step, **3** leads to different products (Scheme VI) depending on experimental conditions (pH, concentration, solvent):

Scheme VI

In order to identify these different products, GC/MS analyses were carried out on the reaction mixture (see chromatogram in Figure 2). Numerous peaks were observed, proving that **3** is converted to several products within a complex reactional chain. A

plausible reaction mechanism describing the formation of the most important by-products is reported later on.

Figure 2. Chromatogram obtained by GC/MS analysis for a NAPP - NH_2Cl mixture;

($[\text{NH}_2\text{Cl}]_0 = 2 \times 10^{-3} \text{ M}$, $[\text{C}_5\text{H}_{10}\text{NNH}_2]_0 = 20 \times 10^{-3} \text{ M}$, $\text{pH} = 12.89$, $T = 25^\circ\text{C}$).

The masses associated to the different peaks observed are as follows:

Peak	m/z
a	85
b	100
c	114
d	98
e	168
f	168
g	166
h	196
i	196
j	184
k	194

Taking previous phenomena into account, the kinetic study of the first step can be established using $[\text{NH}_2\text{Cl}] = f(t)$. For experiments monitored by UV spectrophotometry, only measurements registered in the first reaction instants were useful to determine $[\text{NH}_2\text{Cl}]$ because the UV absorption of **3** remains negligible in the beginning of the reaction.

Further experiments were carried out at lower pH values ($\text{pH} \leq 11.5$). Under these conditions, the first and second steps appeared to be accelerated. For instance, Figures 3 and 4 show UV absorptions of the reaction mixture with respect to time in two different experiments.

Figure 3. UV absorption spectra of NAPP – NH_2Cl mixture at $\text{pH} = 11.5$;

($[\text{C}_5\text{H}_{10}\text{NNH}_2]_0 = 20 \times 10^{-3} \text{ M}$, $[\text{NH}_2\text{Cl}]_0 = 2 \times 10^{-3} \text{ M}$, $T = 25^\circ\text{C}$).

Figure 4. UV absorption spectra of NAPP – NH_2Cl mixture at $\text{pH} = 11$;
 $([\text{C}_5\text{H}_{10}\text{NNH}_2]_0 = 20 \times 10^{-3} \text{ M}, [\text{NH}_2\text{Cl}]_0 = 2 \times 10^{-3} \text{ M}, T = 25^\circ\text{C})$.

Under these conditions, a kinetic study based on $[\text{NH}_2\text{Cl}] = f(t)$ and/or $[\text{FNAPP}] = f(t)$ turned out to be impossible because the reactants are completely consumed in less than 3 minutes. Thus, the HPLC method would face limitations for the same reason.

Kinetics of oxidation of N-aminopiperidine with chloramine

Reaction order and stoichiometry

Experiments were conducted at 25°C with pH values ranging between 12 and 13.5 (at pH values below 12, it became impossible to complete the kinetic study for the reasons mentioned above).

A first series of measurements was performed at 25°C and pH = 12.89. The kinetic parameters were determined by the Ostwald method and the rate of disappearance of NH₂Cl may be expressed as follows:

$$r = -d[\text{NH}_2\text{Cl}]/dt = k_2 [\text{NH}_2\text{Cl}]^\alpha [\text{C}_5\text{H}_{10}\text{NNH}_2]^\beta$$

The stoichiometry of the oxidation of hydrazine with chloramine is 1:1 with respect to each reactant [21].

To evaluate α , three series of measurements were carried out, using a constant concentration of **1** (30×10^{-3} M) and NH₂Cl concentrations ranging from 1×10^{-3} to 4×10^{-3} M (pH = 12.89, T = 25°C). In the first reaction instants, the curves Log [NH₂Cl] = f(t) came up to be straight lines with the slope $\psi = k_2 [\text{C}_5\text{H}_{10}\text{NNH}_2]_0^\beta$ indicating that $\alpha = 1$ (Table 1). Similarly, β was determined by the same method and under the same conditions by maintaining the concentration of NH₂Cl constant (1×10^{-3} M) and varying the concentration of **1** (10×10^{-3} to 100×10^{-3} M).

Partial orders were also determined from experiments monitored by HPLC (Table 1), and gave the same results as for experiments followed by UV spectrophotometry. Experiments were based on the following initial concentration ratios:

$$2 \leq [\text{C}_5\text{H}_{10}\text{NNH}_2]_0/[\text{NH}_2\text{Cl}]_0 \leq 6$$

In all cases the curves:

$$\frac{1}{[\text{C}_5\text{H}_{10}\text{NNH}_2]_0 - [\text{NH}_2\text{Cl}]_0} \text{Log} \frac{[\text{NH}_2\text{Cl}]_0 [\text{C}_5\text{H}_{10}\text{NNH}_2]}{[\text{C}_5\text{H}_{10}\text{NNH}_2]_0 [\text{NH}_2\text{Cl}]} = f(t)$$

showed to be straight lines with the slope k_2 , which confirms that $\alpha = \beta = 1$ (Figure 5). Consequently, the second order rate constant at $\text{pH} = 12.89$, $T = 25^\circ\text{C}$ was found to be equal to $k_2 = 18 \times 10^{-3} \pm 0.7 \times 10^{-3} \text{ M}^{-1} \text{ s}^{-1}$.

Figure 5. Determination of the kinetic parameters in the oxidation of *N*-aminopiperidine with chloramine (first step of the reaction monitored by HPLC; $[\text{C}_5\text{H}_{10}\text{NNH}_2]_0 = 10 \times 10^{-3} \text{ M}$, $[\text{NH}_2\text{Cl}]_0 = 4 \times 10^{-3} \text{ M}$, $\text{pH} = 12.89$, $T = 25^\circ\text{C}$).

$[\text{NH}_2\text{Cl}]_0$ (M)	$[\text{C}_5\text{H}_{10}\text{NNH}_2]_0$ (M)	ψ (s^{-1})	k_2 ($\text{M}^{-1} \text{s}^{-1}$)
1×10^{-3}	10×10^{-3}	1.86×10^{-4}	18.60×10^{-3}
1×10^{-3}	20×10^{-3}	3.29×10^{-4}	16.46×10^{-3}
1×10^{-3}	30×10^{-3}	5.25×10^{-4}	17.50×10^{-3}
1×10^{-3}	40×10^{-3}	7.20×10^{-4}	18.00×10^{-3}
1×10^{-3}	50×10^{-3}	8.79×10^{-4}	17.58×10^{-3}
1×10^{-3}	60×10^{-3}	1.10×10^{-3}	18.21×10^{-3}
1×10^{-3}	100×10^{-3}	1.83×10^{-3}	18.30×10^{-3}
2×10^{-3}	30×10^{-3}	5.27×10^{-4}	17.56×10^{-3}
3×10^{-3}	30×10^{-3}	5.22×10^{-4}	17.39×10^{-3}
4×10^{-3}	30×10^{-3}	5.35×10^{-4}	17.84×10^{-3}
5×10^{-3}	10×10^{-3}	*	18.30×10^{-3}
5×10^{-3}	15×10^{-3}	*	18.61×10^{-3}
5×10^{-3}	20×10^{-3}	*	18.22×10^{-3}
5×10^{-3}	30×10^{-3}	*	17.20×10^{-3}

Table 1. Determination of the partial orders and the rate constant in the oxidation of *N*-aminopiperidine with chloramine ($\text{pH} = 12.89$, $T = 25^\circ\text{C}$).

A second series of measurements was performed at 25°C in a pH interval ranging between 12.0 and 13.5 for $1 \leq [\text{C}_5\text{H}_{10}\text{NNH}_2]_0/[\text{NH}_2\text{Cl}]_0 \leq 10$ (Table 2). The established rate law (partial orders and stoichiometry) were preserved. Furthermore, k_2 remained quite constant at this pH range with a slight shift towards higher values when the pH started decreasing.

To interpret the results, it is important to distinguish between two domains where the pH is above or below 12.89. For $12.0 < \text{pH} < 12.89$, k_2 was determined at the first reaction instants in the same procedure described for the first series of measurements. Under these conditions, the rate constant value began to increase gradually as pH went below 12.89.

To determine k_2 for pH values above 12.89, it is necessary to take the alkaline hydrolysis of chloramine into account. This reaction has been studied by several authors [24-30]. The first elementary step corresponds to the formation of a hydroxylamine intermediate (Scheme VII), which immediately reacts producing several compounds (NO^- , N_2O , $\text{N}_2\text{O}_2^{2-}$, ONOO^-).

Scheme VII

This reaction follows a second order rate law:

$$-d[\text{NH}_2\text{Cl}]/dt = k_3 [\text{NH}_2\text{Cl}] [\text{OH}^-]$$

where $k_3 = 62 \times 10^{-6} \text{ M}^{-1} \text{ s}^{-1}$ at 25°C [27].

Therefore, we followed simultaneously the disappearances of both reactants based on $[\text{FNAPP}] = f(t)$ and $[\text{NH}_2\text{Cl}] = f(t)$. By combining the following equations:

$$-d[\text{NAPP}]/dt = k_2 [\text{NH}_2\text{Cl}] [\text{NAPP}]$$

$$-d[\text{NH}_2\text{Cl}]/dt = k_2 [\text{NH}_2\text{Cl}] [\text{NAPP}] + k_3 [\text{NH}_2\text{Cl}] [\text{OH}^-]$$

$$-d[\text{OH}^-]/dt = k_2 [\text{NH}_2\text{Cl}] [\text{NAPP}] + k_3 [\text{NH}_2\text{Cl}] [\text{OH}^-]$$

an implicit equation, which is a function of instantaneous concentrations of NAPP and NH_2Cl , is obtained and its resolution allowed us to calculate the constant k_2 :

$$[\text{NH}_2\text{Cl}] = [\text{NH}_2\text{Cl}]_0 - [\text{OH}^-]_0 [1 - ([\text{NAPP}]/[\text{NAPP}]_0)^{k_3/k_2}] - \{k_2 [\text{NAPP}] [1 - ([\text{NAPP}]/[\text{NAPP}]_0)^{k_3/k_2} - 1]\}/(k_3 - k_2)$$

The values of k_2 for the range $12.0 < \text{pH} < 13.53$ are given in Table 2.

$[\text{NH}_2\text{Cl}]_0 \text{ (M)}$	$[\text{C}_5\text{H}_{10}\text{NNH}_2]_0 \text{ (M)}$	pH	$k_2 \text{ (M}^{-1} \text{ s}^{-1}\text{)}$
1×10^{-3}	1×10^{-3}	12.03	30.21×10^{-3}
1×10^{-3}	5×10^{-3}	12.54	25.33×10^{-3}
2×10^{-3}	10×10^{-3}	12.70	22.75×10^{-3}
1×10^{-3}	10×10^{-3}	12.89	18.60×10^{-3}
1×10^{-3}	1×10^{-3}	13.35	18.48×10^{-3}
1×10^{-3}	1×10^{-3}	13.53	18.14×10^{-3}

Table 2. Kinetics of the NAPP - NH_2Cl interaction. Determination of the rate constant values for $12.0 < \text{pH} < 13.53$ ($T = 25^\circ\text{C}$).

Influence of temperature

The temperature effect was studied at pH 12.89 between 15 and 45°C . Concentrations for **1** and NH_2Cl used were equal to $20 \times 10^{-3} \text{ M}$ and $2 \times 10^{-3} \text{ M}$, respectively. The variation of k_2 with temperature was found to comply with the Arrhenius law. The curve $\text{Log } k_2 = f(1/T)$ is a straight line of slope = $-E_2/R$ and Y intercept = $\text{Log } A_2$ ($r^2 = 0.999$). A_2 and E_2/R represent the Arrhenius factor and activation energy, respectively.

$$k_2 = 1.15 \times 10^5 \exp(-39/RT) \text{ M}^{-1} \text{ s}^{-1} \quad (E_2 \text{ in kJ mol}^{-1})$$

The enthalpy and entropy of activation can be deduced from the following formulas:

$$\Delta H_2^\ddagger = E_2 - RT \text{ and } \Delta S_2^\ddagger = \text{Log } (A_2 h)/(e k_B T)$$

where k_B is Boltzmann's constant and h is Planck's constant

$$(k_B = 1.38033 \times 10^{-23} \text{ J K}^{-1}, h = 6.623 \times 10^{-23} \text{ J s}).$$

The calculated values are:

$$\Delta H_2^\ddagger = 36.52 \text{ kJ mol}^{-1} \quad \Delta S_2^\ddagger = -156.3 \text{ J mol}^{-1} \text{ K}^{-1}$$

Mechanism

The experimental results showed that the reaction between $C_5H_{10}NNH_2$ and NH_2Cl consists of two steps, in which the first follows a second order rate law and leads to an aminonitrene **3** according to the following redox mechanism [31, 32] (Scheme VIII).

Scheme VIII

In the second step, **3** is converted to different compounds depending on experimental conditions. However, GC/MS (Figure 2) analyses of different reaction mixtures (conducted under different conditions) revealed identical results with the only variation of the peaks' intensities ratios. From this observation, the following conclusions can be drawn:

- With increasing pH, peak **d** increases its intensity. Hence, **d** might correspond to a hydrazone (1,2-diazacyclohept-1-ene) obtained by an intramolecular rearrangement of aminonitrene. The latter is responsible for the absorption at 237 nm appearing at the end of the second step in the experiments conducted at $\text{pH} \geq 12$. This conclusion

agrees with previous studies on the rearrangement aminonitrene – hydrazone described in Scheme IX [33].

Scheme IX

- Two peaks **h** and **i** of same $m/z = 196$ were also observed and their intensities showed to be more important when the concentration of chloramine is higher (the ratio $\frac{[196] + [196]}{[98]}$ increases with $[\text{NH}_2\text{Cl}]_0$). **h** and **i** correspond to two tetrazine compounds, namely azopiperidine and dipyrindododecahydro-s-tetrazine both resulting from diazene according to Scheme X. The suggested mechanism agrees with Figures 3 and 4, showing UV absorptions of the reaction mixture for experiments conducted at lower pH: At pH = 11 (Figure 4), UV spectra are mainly due to the azopiperidine absorption.

Scheme X

Azopiperidine ($m/z = 196$)

Dipyridododecahydro-s-tetrazine ($m/z = 196$)

- By studying the possible evolution routes of the two reactants, it was proven that the other peaks observed cannot be related to diazene. They are consequent upon a succeeding reactional chain based on the evolution of 1-piperidine (peak **a**) present as an impurity in the starting material (peak **b**) and also resulting from the oxidation of NAPP with air: first it formed 1-nitrosopiperidine (peak **c**) which is converted into 1-piperidine. The latter is responsible for the numerous peaks observed in GC/MS. A complete characterization of the compounds obtained will be topic of further investigations.

Acknowledgements

We are grateful to ISOCEM (groupe SNPE) for financial support. C. D. would like to thank the Centre National Français de la Recherche Scientifique for the PhD scholarship and Monsieur Antoine OLLAGNIER for valuable help in Figures

editing. The Claude Bernard University of Lyon and the Lebanese University are also gratefully acknowledged.

References

- [1] Wright, J.B. and Willette, R.E., *J. Med. Pharm. Chem.*, 1962, vol. 5, p. 815.
- [2] Hanna, C. and Schueler, F.W., *J. Am. Chem. Soc.*, 1952, vol. 74, p. 3693.
- [3] Fr Patent 1400256.
- [4] Smith, P.A.S. and Pars, H.G., *J. Org. Chem.*, 1959, vol. 24, p. 1325.
- [5] Lunn, G., Sansone, E.B. and Keefer, L.K., *J. Org. Chem.*, 1984, vol. 49, p. 3470.
- [6] Podgornaya, I.V., Tayusheva, N.N. and Postovskii, I.Y., *Zhurnal Obshchei Khimii*, 1964, vol. 34, p. 2521.
- [7] Allen & Hanburys, NL Patent 6510107, 1966.
- [8] Laboratoire Lafon, Be Patent 812749, 1974.
- [9] Latourette, H. K. and Pianfetti, J. A., US Patent 3317607, 1967.
- [10] Tuemmler, W. B. and Winkler, H. L. S., US Patent 2979505, 1961.
- [11] Lima, D. A., US Patent 3154538, 1964.
- [12] Zimmer, H., Audrieth, L.F., Zimmer, M. and Rowe, R.V., *J. Am. Chem. Soc.*, 1955, vol. 77, p. 790.
- [13] Ohme, R. and Preuschhof, H., Ger (east) Patent 76520, 1970.
- [14] Ohme, R. and Preuschhof, H., *J. Prakt. Chem.*, 1970, vol. 312, p. 349.
- [15] Murakami, Y., Yokoyama, Y., Sasakura, C. and Tamagawa, M., *Chem. Pharm. Bull.*, 1983, vol. 31, p. 423.
- [16] Ohme, R. and Preuschhof, H., *Liebigs Ann. Chem.*, 1968, vol. 713, p. 74.
- [17] Hasegawa, Y., Hyoda, S., Fujita, H., Sawada, H. and Oki, Y., Eur. Patent 850930, 1998.

- [18] Sumitani, H. and Matsui, N., Jp Patent 183250, 2003.
- [19] Raschig, F., *Chem. Ztg.*, 1907, vol. 31, p. 926.
- [20] Raschig, F., *Ber. Dtsch. Chem. Ges.*, 1907, vol. 40, p. 4580.
- [21] Darwich, C., Dissertation, Lyon: Université Lyon I, 2005.
- [22] Ferriol, M., Gazet, J. and Rizk-Ouaini, R., *Anal. Chim. Acta*, 1990, vol. 231, p. 161.
- [23] Elkhatib, M., Peyrot, L., Scharff, J.P. and Delalu, H., *Int. J. Chem. Kinet.*, 1998, vol. 30, p. 129.
- [24] Elkhatib, M., Marchand, A., Counioux, J.J. and Delalu, H., *Int. J. Chem. Kinet.*, 1995, vol. 27, p. 757.
- [25] Delalu, H., Dissertation, Lyon : Université Lyon I, 1977.
- [26] Anbar, M. and Yagil, G., *J. Am. Chem. Soc.*, 1962, vol. 84, p. 1790.
- [27] Yagil, G. and Anbar, M., *J. Inorg. Nucl. Chem.*, 1964, vol. 26, p. 453.
- [28] McCoy, R.E., *J. Am. Chem. Soc.*, 1954, vol. 76, p. 1447.
- [29] Lenoble, W.J., *Tetrahedron Lett.*, 1966, vol. 7, p. 727.
- [30] Elkhatib, M., Duriche, C., Peyrot, L., Metz, R. and Delalu, H., *Int. J. Chem. Kinet.*, 2002, vol. 34, p. 515.
- [31] Lemal, D.M., *Nitrenes*. In: Nitrenes, New York: Lwowski edition, Interscience, 1970.
- [32] Schmidt, E.W., *Hydrazine and its derivatives, preparation, properties, application*, New York: Wiley, 1984.
- [33] Peyrot, L., Elkhatib, M., Vignalou, J.R., Metz, R., Elomar, F. and Delalu, H., *J. Het. Chem.*, 2001, vol. 38, p. 885.