Resolution of the finite Markov moment problem[★]

Laurent Gosse

*IAC-CNR “Mauro Picone” (sezione di Bari)
Via Amendola 122/D - 70126 Bari, Italy*

Olof Runborg^{*}

NADA, KTH, 10044 Stockholm, Sweden

Abstract

We expose in full detail a constructive procedure to invert the so-called “finite Markov moment problem”. The proofs rely on the general theory of Toeplitz matrices together with the classical Newton’s relations.

Key words: Inverse problems, Finite Markov’s moment problem, Toeplitz matrices.

1991 MSC: 65J22

Résumé

Nous présentons en détail une procédure constructive pour inverser le “probleme fini des moments de Markov”. Les preuves reposent sur la théorie générale des matrices de Toeplitz et les classiques relations de Newton.

[★] Support by the Wolfgang Pauli Institute (Wien) in the frame of the “Nanoscience” Programme and the European network HYKE, funded by the EC as contract HPRN-CT-2002-00282 is acknowledged.

^{*} Corresponding Author

Email addresses: `l.gosse@ba.iac.cnr.it` (Laurent Gosse),
`olofr@nada.kth.se` (Olof Runborg).

Afin d'inverser le système fini et mal conditionné (1), Koborov, Sklyar et Fardigola, [9,13] ont proposé un algorithme récursif non-linéaire. Dans [7] nous avons prouvé un lemme le réduisant à l'extraction de valeurs propres généralisées, voir (4). Cette Note vise à expliquer en détail les raisons pour lesquelles cette procédure simplifiée résout le problème des moments de Markov. Après avoir rappelé quelques éléments de la théorie des matrices de Toeplitz et les relations de Newton (Propositions 1 et 2), nous reformulons cet algorithme simplifié afin d'établir facilement certains lemmes techniques. Finalement, le Théorème 1 démontre le lien entre valeurs propres généralisées (4) et l'inversion de (1).

1 Introduction

We aim at inverting a moment system often associated with the prestigious name of Markov, as appearing in [2,3,6,11,12] in several fields of application; consult [5,10,14] for general background on moment problems. The original problem is the following. Given the *moments* m_k for $k = 1, \dots, K$, find a bounded measurable *density* function f and a real value $X > 0$ such that

- (1) $\int_0^X f(\xi) \xi^{k-1} d\xi = m_k, \quad k = 1, \dots, K,$
- (2) $|f(\xi)| = 1$ almost everywhere on $]0, X[$,
- (3) f has no more than $K - 1$ discontinuity points inside $]0, X[$.

The solution is a piecewise constant function taking values in $\{-1, 1\}$ a.e. on $]0, X[$ and changing sign in at most $K - 1$ points, which we denote $\{u_k\}$, ordered such that $0 \leq u_1 \leq \dots \leq u_K = X$. Finding $\{u_k\}$ from $\{m_k\}$ is an ill-conditioned problem when the u_k values come close to each other; its Jacobian is a Vandermonde matrix and iterative numerical resolution routines require extremely good starting guesses. For less than 4 moments, however, a direct method based on solving polynomial equations was presented in [12]. Here we are concerned with an arbitrary number of moments $K \in \mathbb{N}$.

We consider a slightly modified version of the problem which is more relevant for us. In this case f takes values in $\{1, 0\}$ instead of $\{-1, 1\}$ and the moments are scaled as $m_k \rightarrow km_k$. Moreover, to simplify the discussion we confine ourselves to the case when K is even, setting $K =: 2n$. The resulting problem can then be written as an algebraic system of

nonlinear equations: Given m_k find u_k such that

$$m_k = \sum_{j=1}^n u_{2j}^k - u_{2j-1}^k, \quad k = 1, \dots, K = 2n. \quad (1)$$

An algorithm for solving this problem was presented by Koborov, Sklyar and Fardigola in [9,13]. It requires solving a sequence of high degree polynomial equations, constructed through a rather complicated process with unclear stability properties. In [7] we showed that the algorithm can be put in a more simple form that makes it much more suitable for numerical implementation. The simplified algorithm reads

- (1) Construct the matrices A and B :

$$A = \begin{pmatrix} 1 & & & \\ -m_1 & 2 & & \\ \vdots & \ddots & \ddots & \\ -m_{2n-1} & \dots & -m_1 & 2n \end{pmatrix}, \quad B = \begin{pmatrix} 1 & & & \\ m_1 & 2 & & \\ \vdots & \ddots & \ddots & \\ m_{2n-1} & \dots & m_1 & 2n \end{pmatrix}. \quad (2)$$

- (2) Let $\mathbf{m} = (m_1, m_2, \dots, m_{2n})^T$ and solve the lower triangular Toeplitz linear systems

$$A\mathbf{a} = \mathbf{m}, \quad B\mathbf{b} = -\mathbf{m}, \quad (3)$$

to get $\mathbf{a}$ and $\mathbf{b}$.

- (3) Construct the matrices A_1, A_2 from $\mathbf{a} = (a_1, a_2, \dots, a_{2n})^T$ as

$$A_1 = \begin{pmatrix} a_1 & a_2 & \dots & a_n \\ a_2 & a_3 & \dots & a_{n+1} \\ \vdots & \vdots & \ddots & \vdots \\ a_n & a_{n+1} & \dots & a_{2n-1} \end{pmatrix}, \quad A_2 = \begin{pmatrix} a_2 & a_3 & \dots & a_{n+1} \\ a_3 & a_4 & \dots & a_{n+2} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n+1} & a_{n+2} & \dots & a_{2n} \end{pmatrix},$$

and the corresponding matrices B_1, B_2 from $\mathbf{b}$.

- (4) The values $\{u_k\}$ can then be computed as the generalized eigenvalues of the problems

$$A_2\mathbf{v} = uA_1\mathbf{v}, \quad B_2\mathbf{v} = uB_1\mathbf{v}. \quad (4)$$

The forthcoming section is devoted to a complete justification of this algorithm. We recall that these inversion routines have been shown to be numerically efficient in the paper [7].

2 Analysis of the algorithm

We begin by stating two classical results of prime importance for the analysis.

Let $\mathbb{L}^n \subset \mathbb{R}^{n \times n}$ be the set of lower triangular $n \times n$ real Toeplitz matrices. We define the diagonal scaling matrix and the mapping $\mathcal{T} : \mathbb{R}^n \rightarrow \mathbb{L}^n$ as

$$\Lambda = \begin{pmatrix} 0 & & & \\ & 1 & & \\ & & \ddots & \\ & & & n-1 \end{pmatrix}; \quad \mathcal{T}(\mathbf{x}) := \begin{pmatrix} x_1 & & & \\ x_2 & x_1 & & \\ \vdots & \ddots & \ddots & \\ x_n & \dots & x_2 & x_1 \end{pmatrix}, \quad \mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}.$$

The mapping $\mathcal{T}$ has the following properties, see *e.g.* [1]:

Proposition 1 *Lower triangular Toeplitz matrices commute and $\mathbb{L}^n$ is closed under matrix multiplication and (when the inverse exists) inversion,*

$$\mathcal{T}(\mathbf{x})\mathcal{T}(\mathbf{y}) = \mathcal{T}(\mathbf{y})\mathcal{T}(\mathbf{x}) \in \mathbb{L}^n, \quad \mathcal{T}(\mathbf{x})^{-1} \in \mathbb{L}^n. \quad (5)$$

Moreover, $\mathcal{T}$ is linear and

$$\mathcal{T}(\mathbf{x})\mathbf{y} = \mathcal{T}(\mathbf{y})\mathbf{x}, \quad \mathcal{T}(\mathbf{x})\mathcal{T}(\mathbf{y}) = \mathcal{T}(\mathcal{T}(\mathbf{x})\mathbf{y}). \quad (6)$$

The Λ matrix has the property

$$\mathcal{T}(\Lambda\mathbf{x}) = \Lambda\mathcal{T}(\mathbf{x}) - \mathcal{T}(\mathbf{x})\Lambda. \quad (7)$$

Another result that we rely heavily on is the classical Newton relations, see *e.g.* [8]:

Proposition 2 (Newton's relations) *Let P be the n -degree polynomial,*

$$P(x) = c_0 + c_1x + \dots + c_nx^n =: c_n(x - x_1) \dots (x - x_n).$$

Set $S_0 = n$ and define S_k for $k > 0$ as the sum of the roots of P taken to the power k , $S_k = \sum_{j=1}^n x_j^k$. Then, the $n+1$ following relations hold:

$$c_k S_0 + c_{k+1} S_1 + \dots + c_n S_{n-k} = k c_k, \quad k = 0, \dots, n. \quad (8)$$

2.1 Reformulation of the simplified algorithm

We want to write the equation $A\mathbf{a} = \mathbf{m}$ using the mapping $\mathcal{T}$: hence we augment the $\mathbf{m}$ and $\mathbf{a}$ -vectors with a zero and one element, respectively, to get $\tilde{\mathbf{m}} = (0, \mathbf{m})^T$ and $\tilde{\mathbf{a}} = (1, \mathbf{a})^T$, both in $\mathbb{R}^{K+1}$. We observe that the A -matrix in (2) is the lower right $K \times K$ block of $\Lambda - \mathcal{T}(\tilde{\mathbf{m}})$. Therefore,

$$(\Lambda - \mathcal{T}(\tilde{\mathbf{m}}))\tilde{\mathbf{a}} = -\tilde{\mathbf{m}} + \begin{pmatrix} 0 \\ A\mathbf{a} \end{pmatrix} = \begin{pmatrix} 0 \\ A\mathbf{a} - \mathbf{m} \end{pmatrix}.$$

Thus the equation $A\mathbf{a} = \mathbf{m}$ in (3) is equivalent to

$$\mathcal{T}(\tilde{\mathbf{m}})\tilde{\mathbf{a}} = \Lambda\tilde{\mathbf{a}}. \quad (9)$$

By the same argument, $B\mathbf{b} = -\mathbf{m}$ in (3) is equivalent to

$$\mathcal{T}(\tilde{\mathbf{m}})\tilde{\mathbf{b}} = -\Lambda\tilde{\mathbf{b}}. \quad (10)$$

with $\tilde{\mathbf{b}} = (1, \mathbf{b})^T$.

We can then directly also show that $\mathcal{T}(\tilde{\mathbf{a}})$ and $\mathcal{T}(\tilde{\mathbf{b}})$ are in fact each other's inverses.

Lemma 1 *When $\tilde{\mathbf{a}}$ are $\tilde{\mathbf{b}}$ given by (9, 10) then $\mathcal{T}(\tilde{\mathbf{a}})\mathcal{T}(\tilde{\mathbf{b}}) = I$.*

Proof: By Proposition 1 and (9, 10),

$$\Lambda\mathcal{T}(\tilde{\mathbf{a}})\tilde{\mathbf{b}} = \mathcal{T}(\Lambda\tilde{\mathbf{a}})\tilde{\mathbf{b}} + \mathcal{T}(\tilde{\mathbf{a}})\Lambda\tilde{\mathbf{b}} = \mathcal{T}(\mathcal{T}(\tilde{\mathbf{m}})\tilde{\mathbf{a}})\tilde{\mathbf{b}} + \mathcal{T}(\tilde{\mathbf{a}})\Lambda\tilde{\mathbf{b}} = \mathcal{T}(\tilde{\mathbf{a}}) \left[\mathcal{T}(\tilde{\mathbf{m}})\tilde{\mathbf{b}} + \Lambda\tilde{\mathbf{b}} \right] = 0.$$

Thus $\mathcal{T}(\tilde{\mathbf{a}})\tilde{\mathbf{b}}$ lies in the nullspace of Λ which is spanned by the vector $\mathbf{1} := (1, 0, \dots, 0)^T$. Moreover, $(\mathcal{T}(\tilde{\mathbf{a}})\tilde{\mathbf{b}})_1 = 1$ since the first elements of $\tilde{\mathbf{a}}$ and $\tilde{\mathbf{b}}$ are both one and we must in fact have $\mathcal{T}(\tilde{\mathbf{a}})\tilde{\mathbf{b}} = \mathbf{1}$. The lemma then follows from (6) and the definition of $\mathcal{T}$. $\square$

2.2 What lies beneath the algorithm

To understand the workings of the algorithm we need to introduce some new quantities and determine how they relate to $\tilde{\mathbf{a}}$, $\tilde{\mathbf{b}}$ and $\tilde{\mathbf{m}}$.

Let us start with some notation: we set $x_j = u_{2j-1}$ and $y_j = u_{2j}$ for $j = 1, \dots, n$. Furthermore, we introduce the sums

$$X_k = \sum_{j=1}^n x_j^k, \quad Y_k = \sum_{j=1}^n y_j^k, \quad k = 1, 2, \dots, K = 2n, \quad (11)$$

and define $X_0 = Y_0 = K$. In the even case, it then holds that

$$m_k = \sum_{j=1}^n x_j^k - \sum_{j=1}^n y_j^k = X_k - Y_k, \quad k = 1, \dots, K = 2n.$$

We also define the two polynomials

$$p(x) = (x - x_1) \cdots (x - x_n) =: c_0 + c_1 x + \cdots + c_{n-1} x^{n-1} + c_n x^n, \quad (12)$$

and

$$q(x) = (x - y_1) \cdots (x - y_n) =: d_0 + d_1 x + \cdots + d_{n-1} x^{n-1} + d_n x^n. \quad (13)$$

We note here that by construction $c_n = d_n = 1$.

By applying (8) to $x^n p(x)$ with $k = 0, \dots, K$ we get

$$\begin{pmatrix} X_0 \\ X_1 & X_0 \\ \vdots & \ddots & \ddots \\ X_K & \dots & X_1 & X_0 \end{pmatrix} \begin{pmatrix} c_n \\ c_{n-1} \\ \vdots \\ c_0 \\ 0 \\ \vdots \\ 0 \end{pmatrix} = \begin{pmatrix} K c_n \\ (K-1) c_{n-1} \\ \vdots \\ (K-n) c_0 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

The analogous system of equations holds also for Y_k and d_k . We introduce now some shorthand notation to write these equations in a concise form. First we set $\bar{\mathbf{c}} = (c_n, \dots, c_0)^T \in \mathbb{R}^{n+1}$ and $\bar{\mathbf{d}} = (d_n, \dots, d_0)^T \in \mathbb{R}^{n+1}$. We then construct the larger vectors, padded with zeros: $\mathbf{c} = (\bar{\mathbf{c}}, \mathbf{0})^T$ and $\mathbf{d} = (\bar{\mathbf{d}}, \mathbf{0})^T$, both in $\mathbb{R}^{K+1}$. Finally, we let $\mathbf{X} = (X_0, \dots, X_K)^T$ and $\mathbf{Y} = (Y_0, \dots, Y_K)^T$. Using $\mathcal{T}$ and Λ we can state the systems of equations above as follows:

$$\mathcal{T}(\mathbf{X})\mathbf{c} = (KI - \Lambda)\mathbf{c}, \quad \mathcal{T}(\mathbf{Y})\mathbf{d} = (KI - \Lambda)\mathbf{d}. \quad (14)$$

We also clearly have $\tilde{\mathbf{m}} = \mathbf{X} - \mathbf{Y}$.

Before we can relate $\mathbf{c}$ and $\mathbf{d}$ with $\tilde{\mathbf{a}}$ we need the following lemma:

Lemma 2 *Let $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be defined by $\mathbf{f}(\mathbf{x}) := \mathcal{T}(\mathbf{x})^{-1} \Lambda \mathbf{x}$ for $\mathbf{x}$ with a non-zero first element. Then $\mathbf{f}(\mathbf{x}_1) = \mathbf{f}(\mathbf{x}_2)$ implies that $\mathbf{x}_1 = \alpha \mathbf{x}_2$ for some non-zero $\alpha \in \mathbb{R}$.*

Proof: Suppose $\mathbf{f}(\mathbf{x}) = \mathbf{y}$. Then $\mathcal{T}(\mathbf{x})\mathbf{y} = \Lambda\mathbf{x}$ and by Proposition 1, $(\mathcal{T}(\mathbf{y}) - \Lambda)\mathbf{x} = 0$. Hence $\mathbf{f}(\mathbf{x}_1) = \mathbf{f}(\mathbf{x}_2)$ implies that $\mathbf{x}_1$ and $\mathbf{x}_2$ both lie in the nullspace of $\mathcal{T}(\mathbf{y}) - \Lambda$. Since the top left element of Λ is zero, it follows that the first element of $\mathbf{y}$ is zero and therefore the diagonal of $\mathcal{T}(\mathbf{y})$ is zero. Consequently, the nullspace of $\mathcal{T}(\mathbf{y}) - \Lambda$ has the same dimension as that of Λ , which is one. $\square$

We can now merge together and express the general structure from (12, 13) and (9, 10) in the most concise way.

Lemma 3 Suppose $\mathbf{c}, \mathbf{d}$ are defined by (12, 13) and $\tilde{\mathbf{a}}, \tilde{\mathbf{b}}$ by (9, 10). Then

$$\mathcal{T}(\tilde{\mathbf{a}})\mathbf{c} = \mathbf{d}, \quad \mathcal{T}(\tilde{\mathbf{b}})\mathbf{d} = \mathbf{c}.$$

Proof: We only need to prove the left equality. The right one follows immediately from Lemma 1. Let $\mathbf{v} = \mathcal{T}(\tilde{\mathbf{a}})\mathbf{c} = \mathcal{T}(\mathbf{c})\tilde{\mathbf{a}}$. We want to show that $\mathbf{v} = \mathbf{d}$. We note first that by (14)

$$\mathcal{T}(\mathbf{X})\mathbf{c} = (KI - \Lambda)\mathbf{c} \Rightarrow \mathcal{T}(\mathbf{c})\mathbf{X} = (KI - \Lambda)\mathbf{c} \Rightarrow \mathbf{X} = K\mathcal{T}(\mathbf{c})^{-1}\mathbf{c} - \mathcal{T}(\mathbf{c})^{-1}\Lambda\mathbf{c},$$

where $\mathcal{T}(\mathbf{c})$ is invertible since $c_n = 1$. Moreover, it is clear that $\mathcal{T}(\mathbf{y})\mathbf{1} = \mathbf{y}$ for all $\mathbf{y}$. Hence, $\mathbf{X} = K\mathbf{1} - \mathcal{T}(\mathbf{c})^{-1}\Lambda\mathbf{c}$. In the same way we also obtain $\mathbf{Y} = K\mathbf{1} - \mathcal{T}(\mathbf{d})^{-1}\Lambda\mathbf{d}$. Then,

$$\begin{aligned} \mathcal{T}(\tilde{\mathbf{m}})\tilde{\mathbf{a}} &= \mathcal{T}(\tilde{\mathbf{a}})\tilde{\mathbf{m}} = \mathcal{T}(\tilde{\mathbf{a}})(\mathbf{X} - \mathbf{Y}) \\ &= -\mathcal{T}(\tilde{\mathbf{a}})\mathcal{T}(\mathbf{c})^{-1}\Lambda\mathbf{c} + \mathcal{T}(\tilde{\mathbf{a}})\mathcal{T}(\mathbf{d})^{-1}\Lambda\mathbf{d}. \end{aligned}$$

We now note that by Proposition 1,

$$\begin{aligned} \mathcal{T}(\tilde{\mathbf{a}})\mathcal{T}(\mathbf{c})^{-1}\Lambda\mathbf{c} &= \mathcal{T}(\mathbf{c})^{-1}\mathcal{T}(\tilde{\mathbf{a}})\Lambda\mathbf{c} = \mathcal{T}(\mathbf{c})^{-1}\mathcal{T}(\Lambda\mathbf{c})\tilde{\mathbf{a}} = \mathcal{T}(\mathbf{c})^{-1}\Lambda\mathcal{T}(\mathbf{c})\tilde{\mathbf{a}} - \Lambda\tilde{\mathbf{a}} \\ &= \mathcal{T}(\mathbf{c})^{-1}\Lambda\mathbf{v} - \Lambda\tilde{\mathbf{a}}. \end{aligned}$$

Since also, $\mathcal{T}(\mathbf{c})\mathcal{T}(\tilde{\mathbf{a}}) = \mathcal{T}(\mathbf{v})$ we get

$$\begin{aligned} \mathcal{T}(\tilde{\mathbf{m}})\tilde{\mathbf{a}} &= -\mathcal{T}(\mathbf{c})^{-1}\Lambda\mathbf{v} + \Lambda\tilde{\mathbf{a}} + \mathcal{T}(\tilde{\mathbf{a}})\mathcal{T}(\mathbf{d})^{-1}\Lambda\mathbf{d} \\ &= \mathcal{T}(\tilde{\mathbf{a}})\left[\mathcal{T}(\mathbf{d})^{-1}\Lambda\mathbf{d} - \mathcal{T}(\mathbf{v})^{-1}\Lambda\mathbf{v}\right] + \Lambda\tilde{\mathbf{a}}. \end{aligned}$$

Consequently, by (9), $\mathcal{T}(\mathbf{d})^{-1}\Lambda\mathbf{d} = \mathcal{T}(\mathbf{v})^{-1}\Lambda\mathbf{v}$ and by Lemma 2, $\mathbf{v} = \alpha\mathbf{d}$; for some $\alpha \in \mathbb{R}$. But for the first element in $\mathbf{v}$ we then have $v_1 = c_n = \alpha d_n$ and we get $\alpha = 1$ since $c_n = d_n = 1$. $\square$

Finally, we also establish the following lemma.

Lemma 4 Let V and W be the Vandermonde matrices corresponding to the roots of $\tilde{p}(x) := xp(x)$ and $\tilde{q}(x) := xq(x)$ respectively. Then

$$V^T R \mathcal{T}(\tilde{\mathbf{c}}) V = \text{diag}(\{\tilde{p}'(x_k)\}), \quad W^T R \mathcal{T}(\tilde{\mathbf{d}}) W = \text{diag}(\{\tilde{q}'(x_k)\}),$$

where $R = \{\delta_{n+2-i-j}\} \in \mathbb{R}^{n+1 \times n+1}$ is the reversion matrix and $x_0 = y_0 = 0$.

Proof: We have

$$(V^T R T(\bar{\mathbf{c}}) V)_{ij} = \mathbf{x}_{i-1}^T R T(\bar{\mathbf{c}}) \mathbf{x}_{j-1} = \sum_{r=0}^n \sum_{\ell=0}^r c_r x_{i-1}^\ell x_{j-1}^{r-\ell} = \begin{cases} \frac{\tilde{p}(x_{i-1}) - \tilde{p}(x_{j-1})}{x_{i-1} - x_{j-1}}, & i \neq j, \\ \tilde{p}'(x_{i-1}), & i = j, \end{cases}$$

showing the left equality. The right equality follows in the same way. $\square$

2.3 Conclusion

We can now conclude and show that the unknown values u_j in (1) are indeed the generalized eigenvalues of (4).

Theorem 1 Suppose $K = 2n$; let $\mathbf{a}, \mathbf{b}$ be defined by (3). If all values $\{x_j\} \cup \{y_j\}$ are distinct, then $\{x_j\}, \{y_j\}$ are the generalized eigenvalues of (4).

Proof: Let $\tilde{\mathbf{a}}, \tilde{\mathbf{b}}$ be defined by (9, 10), which is equivalent to (3). Also define $\mathbf{c}$ and $\mathbf{d}$ as before by (12, 13). By Lemma 3 we have $\mathcal{T}(\tilde{\mathbf{a}})\mathbf{c} = \mathbf{d} \in \mathbb{R}^{2n+1}$, i.e.

$$\left(\begin{array}{cccc|cccc} 1 & & & & & & & \\ a_1 & 1 & & & & & & \\ \vdots & & \ddots & & & & & \\ a_n & \dots & a_1 & 1 & & & & \\ \hline a_{n+1} & a_n & \dots & a_1 & 1 & & & \\ a_{n+2} & a_{n+1} & \dots & a_2 & a_1 & 1 & & \\ \vdots & \vdots & & \ddots & \vdots & & \ddots & \\ a_{2n} & a_{2n-1} & \dots & a_n & a_{n-1} & \dots & a_1 & 1 \end{array} \right) \begin{pmatrix} \bar{\mathbf{c}} \\ - \\ 0 \end{pmatrix} = \begin{pmatrix} \bar{\mathbf{d}} \\ - \\ 0 \end{pmatrix}. \quad (15)$$

Clearly, the lower left block of the matrix multiplied by $\bar{\mathbf{c}}$ is zero, i.e. $\sum_{i=0}^n c_i a_{i+k} = 0$ for $k = 1, \dots, n$. Now, let v_i be the coefficients of the polynomial $v(x) := p(x)/(x - x_j)$ for some fixed j . Hence, by the special structure of (12),

$$c_0 + c_1 x + \dots + c_n x^n =: (v_1 + v_2 x + \dots + v_n x^{n-1})(x - x_j),$$

and, for $i = 0, \dots, n$,

$$c_i = \begin{cases} -x_j v_{i+1}, & i = 0, \\ v_i - x_j v_{i+1}, & 1 \leq i \leq n-1, \\ v_i, & i = n. \end{cases}$$

Thus we deduce,

$$0 = -x_j v_1 a_k + \sum_{i=1}^{n-1} (v_i - x_j v_{i+1}) a_{i+k} + v_n a_{n+k} = \sum_{i=1}^n v_i a_{i+k} - x_j \sum_{i=1}^n v_i a_{i+k-1},$$

which is the componentwise statement of $A_2 \mathbf{v} = x_j A_1 \mathbf{v}$. It remains to show that the rightmost sum is non-zero for at least some k , so that x_j is indeed a well-defined generalized eigenvalue. Let $\bar{\mathbf{a}} = (1, a_1, \dots, a_n)^T$ and $\bar{\mathbf{v}} = (0, v_1, \dots, v_n)^T$. Then (15) gives $\mathcal{T}(\bar{\mathbf{a}})\bar{\mathbf{c}} = \bar{\mathbf{d}}$ and, using Lemma 4 while taking $k = 1$ we have the sum

$$\sum_{i=1}^n v_i a_i = \bar{\mathbf{a}}^T \bar{\mathbf{v}} = (\mathcal{T}(\bar{\mathbf{c}})^{-1} \bar{\mathbf{d}})^T \bar{\mathbf{v}} = (V^T R \bar{\mathbf{d}})^T \text{diag}(\{\tilde{p}'(x_k)^{-1}\}) V^T \bar{\mathbf{v}} = q(x_j),$$

since $V^T R \bar{\mathbf{d}} = \{q(x_k)\}$ and $V^T \bar{\mathbf{v}} = \{x_k v(x_k)\} = \{\delta_{k-j} \tilde{p}'(x_k)\}$. Hence, the sum is non-zero because $x_j \neq y_i$ for all i, j . The same argument can be used for any j , which proves the theorem for $\{x_j\}$. The proof for $\{y_j\}$ is identical upon exchanging the roles of $\mathbf{c}$, $\bar{\mathbf{a}}$ and $\mathbf{d}$, $\tilde{\mathbf{b}}$. This leads to (4). $\square$

References

- [1] D. Bini, *Polynomial and matrix computations. I. Fundamental Algorithms*, Birkhäuser Verlag.
- [2] Y. Brenier, *Équations de moment et conditions d'entropie pour des modèles cinétiques* (French) [Moment equations and entropy conditions for kinetic models] Séminaire sur les Équations aux Dérivées Partielles, 1994–1995, Exp. No. XXII, 11 pp., École Polytech., Palaiseau, 1995.
- [3] Y. Brenier and L. Corrias, *A kinetic formulation for multibranch entropy solutions of scalar conservation laws*, Ann. I.H.P. Nonlinear Anal. **15** (1998), 169–190.
- [4] D. Czarkowski, D. & G. Chudnovsky, I.W. Selesnick, *Solving the optimal PWM problem for single-phase inverters*, IEEE Trans. Circ. Syst. I **49** (2002), 465–475.

- [5] P. Diaconis, D. Friedman, *The Markov moment problem and de Finetti's theorem*, to appear in Math. Zeitschrift
- [6] L. Gosse, *Using K-branch entropy solutions for multivalued geometric optics computations*, J. Comp. Phys. **180** (2002) 155–182.
- [7] L. Gosse & O. Runborg, *Finite moment problems and applications to multiphase computations in geometric optics*, Comm. Math. Sci. **3** (2005) 373–392.
- [8] I.N. Hepstein, *Topics in Algebra*, Ginn Waltham MA (1964) page 208.
- [9] V.I. Korobov, G. M. Sklyar, *Time-optimality and the power moment problem* (Russian) Mat. Sb. (N.S.) **134** (176) (1987) 186–206, 287; translation in Math. USSR-Sb. **62** (1989), no. 1, 185–206
- [10] M.G. Krein, A.A. Nudel'man, *The Markov moment problem and extremal problems*, A.M.S. translations (1977).
- [11] A.S. Lewis, *Superresolution in the Markov moment problem*, J. Math. Anal. Appl. **197** (1996) 774–780.
- [12] O. Runborg, *Some new results in multiphase geometrical optics*, Math. Mod. Numer. Anal. **34** (2000), 1203 – 1231.
- [13] G.M. Sklyar, L.V. Fardigola, *The Markov power moment problem in problems of controllability and frequency extinguishing for the wave equation on a half-axis*, J. Math. Anal. Appl. **276** (2002) 109–134.
- [14] G. Talenti, *Recovering a function from a finite number of moments*, Inverse Problems **3** (1987) 501–517.