



HAL
open science

The primitive idempotents of the p -permutation ring

Serge Bouc, Jacques Thévenaz

► **To cite this version:**

Serge Bouc, Jacques Thévenaz. The primitive idempotents of the p -permutation ring. 2009. hal-00430256

HAL Id: hal-00430256

<https://hal.science/hal-00430256>

Preprint submitted on 6 Nov 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The primitive idempotents of the p -permutation ring

Serge Bouc and Jacques Thévenaz

Abstract : Let G be a finite group, let p be a prime number, and let K be a field of characteristic 0 and k be a field of characteristic p , both large enough. In this note we state explicit formulae for the primitive idempotents of $K \otimes_{\mathbb{Z}} pp_k(G)$, where $pp_k(G)$ is the ring of p -permutation kG -modules.

AMS Subject Classification : 19A22, 20C20.

Key words : p -permutation, idempotent, trivial source.

1. Introduction

Let G be a finite group, let p be a prime number, and let K be a field of characteristic 0 and k be a field of characteristic p , both large enough. In this note we state explicit formulae for the primitive idempotents of $K \otimes_{\mathbb{Z}} pp_k(G)$, where $pp_k(G)$ is the ring of p -permutation kG -modules (also called the trivial source ring).

To obtain these formulae, we first use induction and restriction to reduce to the case where G is cyclic modulo p , i.e. G has a normal Sylow p -subgroup with cyclic quotient. Then we solve the easy and well known case where G is a cyclic p' -group. Finally we conclude by using the natural ring homomorphism from the Burnside ring $B(G)$ of G to $pp_k(G)$, and the classical formulae for the primitive idempotents of $K \otimes_{\mathbb{Z}} B(G)$.

Our formulae are an essential tool in [2], where Cartan matrices of Mackey algebras are considered, and some invariants of these matrices (determinant, rank) are explicitly computed.

2. p -permutation modules

2.1. Notation.

- Throughout the paper, G will be a fixed finite group and p a fixed prime number. We consider a field k of characteristic p and we denote by kG the group algebra of G over k . We assume that k is large enough in the sense that it is a splitting field for every group algebra $k(N_G(P)/P)$, where P runs through the set of all p -subgroups of G .

- We let K be a field of characteristic 0 and we assume that K is large enough in the sense that it contains the values of all the Brauer characters of the groups $N_G(P)/P$, where P runs through the set of all p -subgroups of G .

We recall quickly how Brauer characters are defined. We let \bar{k} be an extension of k containing all the n -th roots of unity, where n is the p' -part of the exponent of G . We choose an isomorphism $\theta : \mu_n(\bar{k}) \rightarrow \mu_n(\mathbb{C})$ from the group of n -th roots of unity in \bar{k} and the corresponding group in \mathbb{C} . If V is an r -dimensional kH -module for the group $H = N_G(P)/P$ and if s is an element of the set $H_{p'}$ of all p' -elements of H , the matrix of the action of s on V has eigenvalues $(\lambda_1, \dots, \lambda_r)$ in the group $\mu_n(\bar{k})$. The *Brauer character* ϕ_V of V is the central function defined on $H_{p'}$, with values in the field $\mathbb{Q}[\mu_n(\mathbb{C})]$, sending s to $\sum_{i=1}^r \theta(\lambda_i)$. The actual values of Brauer characters may lie in a subfield of $\mathbb{Q}[\mu_n(\mathbb{C})]$ and we simply require that K contains all these values.

2.2. Remark : Let V be as above and let W be a t -dimensional kH -module. If s has eigenvalues (μ_1, \dots, μ_t) on W , its eigenvalues for the diagonal action of H on $V \otimes_k W$ are $(\lambda_i \mu_j)_{1 \leq i \leq r, 1 \leq j \leq t}$. It follows that $\phi_{V \otimes_k W}(s) = \sum_{i=1}^r \sum_{j=1}^t \theta(\lambda_i \mu_j) = \phi_V(s) \phi_W(s)$.

- When H is a subgroup of G , and M is a kG -module, we denote by $\text{Res}_H^G M$ the kH -module obtained by restricting the action of G to H . When L is a kH -module, we denote by $\text{Ind}_H^G L$ the induced kG -module.

- When M is a kG -module, and P is a subgroup of G , the k -vector space of fixed points of P on M is denoted by M^P . When $Q \leq P$ are subgroups of G , the *relative trace* is the map $\text{tr}_Q^P : M^Q \rightarrow M^P$ defined by $\text{tr}_Q^P(m) = \sum_{x \in [P/Q]} x \cdot m$.

- When M is a kG -module, the *Brauer quotient* of M at P is the k -vector space

$$M[P] = M^P / \sum_{Q < P} \text{tr}_Q^P M^Q .$$

This k -vector space has a natural structure of $k\overline{N}_G(P)$ -module, where as usual $\overline{N}_G(P) = N_G(P)/P$. It is equal to zero if P is not a p -group.

- If P is a normal p -subgroup of G and M is a $k(G/P)$ -module, denote by $\text{Inf}_{G/P}^G M$ the kG -module obtained from M by inflation to G . Then there is an isomorphism

$$(\text{Inf}_{G/P}^G M)[P] \cong M$$

of $k(G/P)$ -modules.

- When G acts on a set X (on the left), and $x, y \in X$, we write $x =_G y$ if x and y are in the same G -orbit. We denote by $[G \backslash X]$ a set of representatives

of G -orbits on X , and by X^G the set of fixed points of G on X . For $x \in X$, we denote by G_x its stabilizer in G .

2.3. Review of p -permutation modules. We begin by recalling some definitions and basic results. We refer to [3], and to [1] Sections 3.11 and 5.5 for details :

2.4. Definition. A permutation kG -module is a kG -module admitting a G -invariant k -basis. A p -permutation kG -module M is a kG -module such that $\text{Res}_S^G M$ is a permutation kS -module, where S is a Sylow p -subgroup of G .

The p -permutation kG -modules are also called *trivial source modules*, because the indecomposable ones coincide with the indecomposable modules having a trivial source (see [3] 0.4). Moreover, the p -permutation modules also coincide with the direct summands of permutation modules (see [1], Lemma 3.11.2).

2.5. Proposition.

1. If H is a subgroup of G , and M is a p -permutation kG -module, then the restriction $\text{Res}_H^G M$ of M to H is a p -permutation kH -module.
2. If H is a subgroup of G , and L is a p -permutation kH -module, then the induced module $\text{Ind}_H^G L$ is a p -permutation kG -module.
3. If N is a normal subgroup of G , and L is a p -permutation $k(G/N)$ -module, the inflated module $\text{Inf}_{G/N}^G L$ is a p -permutation kG -module.
4. If P is a p -group, and M is a permutation kP -module with P -invariant basis X , then the image of the set X^P in $M[P]$ is a k -basis of $M[P]$.
5. If P is a p -subgroup of G , and M is a p -permutation kG -module, then the Brauer quotient $M[P]$ is a p -permutation $k\overline{N}_G(P)$ -module.
6. If M and N are p -permutation kG -modules, then their tensor product $M \otimes_k N$ is again a p -permutation kG -module.

Proof : Assertions 1,2,3, and 6 are straightforward consequences of the same assertions for *permutation* modules. For Assertion 4, see [3] 1.1.(3). Assertion 5 follows easily from Assertion 4 (see also [3] 3.1). \square

This leads to the following definition :

2.6. Definition. The p -permutation ring $pp_k(G)$ is the Grothendieck group of the category of p -permutation kG -modules, with relations corresponding to direct sum decompositions, i.e. $[M] + [N] = [M \oplus N]$. The ring structure

on $pp_k(G)$ is induced by the tensor product of modules over k . The identity element of $pp_k(G)$ is the class of the trivial kG -module k .

As the Krull-Schmidt theorem holds for kG -modules, the additive group $pp_k(G)$ is a free (abelian) group on the set of isomorphism classes of indecomposable p -permutation kG -modules. These modules have the following properties :

2.7. Theorem. [[3] Theorem 3.2]

1. The vertices of an indecomposable p -permutation kG -module M are the maximal p -subgroups P of G such that $M[P] \neq \{0\}$.
2. An indecomposable p -permutation kG -module has vertex P if and only if $M[P]$ is a non-zero projective $k\overline{N}_G(P)$ -module.
3. The correspondence $M \mapsto M[P]$ induces a bijection between the isomorphism classes of indecomposable p -permutation kG -modules with vertex P and the isomorphism classes of indecomposable projective $k\overline{N}_G(P)$ -modules.

2.8. Notation. Let $\mathcal{P}_{G,p}$ denote the set of pairs (P, E) , where P is a p -subgroup of G , and E is an indecomposable projective $k\overline{N}_G(P)$ -module. The group G acts on $\mathcal{P}_{G,p}$ by conjugation, and we denote by $[\mathcal{P}_{G,p}]$ a set of representatives of G -orbits on $\mathcal{P}_{G,p}$.

For $(P, E) \in \mathcal{P}_{G,p}$, let $M_{P,E}$ denote the (unique up to isomorphism) indecomposable p -permutation kG -module such that $M_{P,E}[P] \cong E$.

2.9. Corollary. The classes of the modules $M_{P,E}$, for $(P, E) \in [\mathcal{P}_{G,p}]$ form a \mathbb{Z} -basis of $pp_k(G)$.

2.10. Notation. The operations Res_H^G , Ind_H^G , $\text{Inf}_{G/N}^G$ extend linearly to maps between the corresponding p -permutations rings, denoted with the same symbol.

The maps Res_H^G and $\text{Inf}_{G/N}^G$ are ring homomorphisms, whereas Ind_H^G is not in general. Similarly :

2.11. Proposition. Let P be a p -subgroup of G . Then the correspondence $M \mapsto M[P]$ induces a ring homomorphism $\text{Br}_P^G : pp_k(G) \rightarrow pp_k(\overline{N}_G(P))$.

Proof : Let M and N be p -permutation kG -modules. The canonical bilinear map $M \times N \rightarrow M \otimes_k N$ is G -equivariant, hence it induces a bilinear map $\beta_P : M[P] \times N[P] \rightarrow (M \otimes_k N)[P]$ (see [3] 1.2), which is $\overline{N}_G(P)$ -equivariant. Now if X is a P -invariant k -basis of M , and Y a P -invariant k -basis of N ,

then $X \times Y$ is a P -invariant basis of $M \otimes_k N$. The images of the sets X^P, Y^P , and $(X \times Y)^P$ are bases of $M[P], N[P]$, and $(M \otimes_k N)[P]$, respectively, and the restriction of β_P to these bases is the canonical bijection $X^P \times Y^P \rightarrow (X \times Y)^P$. It follows that β_P induces an isomorphism $M[P] \otimes_k N[P] \rightarrow (M \otimes_k N)[P]$ of $k\overline{N}_G(P)$ -modules. Proposition 2.11 follows. \square

2.12. Notation. Let $\mathcal{Q}_{G,p}$ denote the set of pairs (P, s) , where P is a p -subgroup of G , and s is a p' -element of $\overline{N}_G(P)$. The group G acts on $\mathcal{Q}_{G,p}$, and we denote by $[\mathcal{Q}_{G,p}]$ a set of representatives of G -orbits on $\mathcal{Q}_{G,p}$.

If $(P, s) \in \mathcal{Q}_{G,p}$, we denote by $N_G(P, s)$ the stabilizer of (P, s) in G , and by $\langle Ps \rangle$ the subgroup of $N_G(P)$ generated by Ps (i.e. the inverse image in $N_G(P)$ of the cyclic group $\langle s \rangle$ of $\overline{N}_G(P)$).

2.13. Remarks :

- When H is a subgroup of G , there is a natural inclusion of $\mathcal{Q}_{H,p}$ into $\mathcal{Q}_{G,p}$, as $\overline{N}_H(P) \leq \overline{N}_G(P)$ for any p -subgroup P of H . We will consider $\mathcal{Q}_{H,p}$ as a subset of $\mathcal{Q}_{G,p}$.
- When $(P, s) \in \mathcal{Q}_{G,p}$, the group $N_G(P, s)$ is the set of elements g in $N_G(P)$ whose image in $\overline{N}_G(P)$ centralizes s . In other words, there is a short exact sequence of groups

$$(2.14) \quad \mathbf{1} \rightarrow P \rightarrow N_G(P, s) \rightarrow C_{\overline{N}_G(P)}(s) \rightarrow \mathbf{1} .$$

In particular $N_G(P, s)$ is a subgroup of $N_G(\langle Ps \rangle)$.

2.15. Notation. Let $(P, s) \in \mathcal{Q}_{G,p}$. Let $\tau_{P,s}^G$ denote the additive map from $pp_k(G)$ to K sending the class of a p -permutation kG -module M to the value at s of the Brauer character of the $\overline{N}_G(P)$ -module $M[P]$.

2.16. Remarks :

- It is clear that $\tau_{P,s}^G(M)$ only depends on the restriction of M to the group $\langle Ps \rangle$. In other words

$$\tau_{P,s}^G = \tau_{P,s}^{\langle Ps \rangle} \circ \text{Res}_{\langle Ps \rangle}^G .$$

Furthermore, it is clear from the definition that

$$(2.17) \quad \tau_{P,s}^G = \tau_{\mathbf{1},s}^{\langle Ps \rangle / P} \circ \text{Br}_P^{\langle Ps \rangle} \circ \text{Res}_{\langle Ps \rangle}^G .$$

- It is easy to see that $\tau_{P,s}^G$ only depends on the G -orbit of (P, s) , that is, $\tau_{P^g, s^g}^G = \tau_{P,s}^G$ for every $g \in G$.

The following proposition is Corollary 5.5.5 in [1], but our construction of the species is slightly different (but equivalent, of course). For this reason, we sketch an independent proof :

2.18. Proposition.

1. The map $\tau_{P,s}^G$ is a ring homomorphism $pp_k(G) \rightarrow K$ and extends to a K -algebra homomorphism (a species) $\tau_{P,s}^G : K \otimes_{\mathbb{Z}} pp_k(G) \rightarrow K$.
2. The set $\{\tau_{P,s}^G \mid (P, s) \in [\mathcal{Q}_{G,p}]\}$ is the set of all distinct species from $K \otimes_{\mathbb{Z}} pp_k(G)$ to K . These species induce a K -algebra isomorphism

$$T = \prod_{(P,s) \in [\mathcal{Q}_{G,p}]} \tau_{P,s}^G : K \otimes_{\mathbb{Z}} pp_k(G) \rightarrow \prod_{(P,s) \in [\mathcal{Q}_{G,p}]} K .$$

Proof : By 2.17, to prove Assertion 1, it suffices to prove that $\tau_{\mathbf{1},s}^{\langle Ps \rangle / P}$ is a ring homomorphism, since both $\text{Res}_{\langle Ps \rangle}^G$ and $\text{Br}_P^{\langle Ps \rangle}$ are ring homomorphisms. In other words, we can assume that $P = \mathbf{1}$. Now the value of $\tau_{\mathbf{1},s}^G$ on the class of a kG -module M is the value $\phi_M(s)$ of the Brauer character of M at s , so Assertion 1 follows from Remark 2.2.

For Assertion 2, it suffices to prove that T is an isomorphism. Since $[\mathcal{P}_{G,p}]$ and $[\mathcal{Q}_{G,p}]$ have the same cardinality, the matrix \mathcal{M} of T is a square matrix. Let $(P, E) \in \mathcal{P}_{G,p}$, and $(Q, s) \in \mathcal{Q}_{G,p}$. Then $\tau_{Q,s}(M_{P,E})$ is equal to zero if Q is not contained in P up to G -conjugation, because in this case $M_{P,E}(Q) = \{0\}$ by Theorem 2.7. It follows that \mathcal{M} is block triangular. As moreover $M_{P,E}[P] \cong E$, we have that $\tau_{P,s}(M_{P,E}) = \phi_E(s)$. This means that the diagonal block of \mathcal{M} corresponding to P is the matrix of Brauer characters of projective $k\overline{N}_G(P)$ -modules, and these are linearly independent by Lemma 5.3.1 of [1]. It follows that all the diagonal blocks of \mathcal{M} are non singular, so \mathcal{M} is invertible, and T is an isomorphism. \square

2.19. Corollary. *The algebra $K \otimes_{\mathbb{Z}} pp_k(G)$ is a split semisimple commutative K -algebra. Its primitive idempotents $F_{P,s}^G$ are indexed by $[\mathcal{Q}_{G,p}]$, and the idempotent $F_{P,s}^G$ is characterized by*

$$\forall (R, u) \in \mathcal{Q}_{G,p}, \quad \tau_{R,u}^G(F_{P,s}^G) = \begin{cases} 1 & \text{if } (R, u) =_G (P, s) \\ 0 & \text{otherwise.} \end{cases}$$

3. Restriction and induction

3.1. Proposition. *Let $H \leq G$, and $(P, s) \in \mathcal{Q}_{G,p}$. Then*

$$\text{Res}_H^G F_{P,s}^G = \sum_{(Q,t)} F_{Q,t}^H ,$$

where (Q, t) runs through a set of representatives of H -conjugacy classes of G -conjugates of (P, s) contained in H .

Proof : Indeed, as Res_H^G is an algebra homomorphism, the element $\text{Res}_H^G F_{P,s}^G$ is an idempotent of $K \otimes_{\mathbb{Z}} \text{pp}_k(H)$, hence it is equal to a sum of some distinct primitive idempotents $F_{Q,t}^H$. The idempotent $F_{Q,t}^H$ appears in this decomposition if and only if $\tau_{Q,t}^H(\text{Res}_H^G F_{P,s}^G) = 1$. By Remark 2.16

$$\begin{aligned} \tau_{Q,t}^H(\text{Res}_H^G F_{P,s}^G) &= \tau_{Q,t}^{\langle Qt \rangle}(\text{Res}_{\langle Qt \rangle}^H \text{Res}_H^G F_{P,s}^G) \\ &= \tau_{Q,t}^{\langle Qt \rangle}(\text{Res}_{\langle Qt \rangle}^G F_{P,s}^G) \\ &= \tau_{Q,t}^G(F_{P,s}^G) . \end{aligned}$$

Now $\tau_{Q,t}^G(F_{P,s}^G)$ is equal to 1 if and only if (Q, t) and (P, s) are G -conjugate. This completes the proof. \square

3.2. Proposition. *Let $H \leq G$, and $(Q, t) \in \mathcal{Q}_{H,p}$. Then*

$$\text{Ind}_H^G F_{Q,t}^H = |N_G(Q, t) : N_H(Q, t)| F_{Q,t}^G .$$

Proof : Since $K \otimes_{\mathbb{Z}} \text{pp}_k(G)$ is a split semisimple commutative K -algebra, any element X in $K \otimes_{\mathbb{Z}} \text{pp}_k(G)$ can be written

$$(3.3) \quad X = \sum_{(P,s) \in [\mathcal{Q}_{G,p}]} \tau_{P,s}^G(X) F_{P,s}^G ,$$

and moreover for any $(P, s) \in \mathcal{Q}_{G,p}$

$$\tau_{P,s}^G(X) F_{P,s}^G = X \cdot F_{P,s}^G .$$

Setting $X = \text{Ind}_H^G F_{Q,t}^H$ in this equation gives

$$\begin{aligned} \tau_{P,s}^G(\text{Ind}_H^G F_{Q,t}^H) F_{P,s}^G &= (\text{Ind}_H^G F_{Q,t}^H) \cdot F_{P,s}^G \\ &= \text{Ind}_H^G(F_{Q,t}^H \cdot \text{Res}_H^G F_{P,s}^G) . \end{aligned}$$

By Proposition 3.1, the element $\text{Res}_H^G F_{P,s}^G$ is equal to the sum of the distinct idempotents F_{P^y, s^y}^H associated to elements y of G such that $\langle Ps \rangle^y \leq H$. The product $F_{Q,t}^H \cdot F_{P^y, s^y}^H$ is equal to zero, unless (Q, t) is H -conjugate to (P^y, s^y) , which implies that (Q, t) and (P, s) are G -conjugate. It follows that the only non zero term in the right hand side of Equation 3.3 is the term corresponding to (Q, t) . Hence

$$\text{Ind}_H^G F_{Q,t}^H = \tau_{Q,t}^G(\text{Ind}_H^G F_{Q,t}^H) F_{Q,t}^G .$$

Now by Remark 2.16 and the Mackey formula

$$\begin{aligned}\tau_{Q,t}^G(\text{Ind}_H^G F_{Q,t}^H) &= \tau_{Q,t}^{\langle Qt \rangle}(\text{Res}_{\langle Qt \rangle}^G \text{Ind}_H^G F_{Q,t}^H) \\ &= \tau_{Q,t}^{\langle Qt \rangle} \left(\sum_{x \in \langle Qt \rangle \backslash G/H} \text{Ind}_{\langle Qt \rangle \cap xH}^{\langle Qt \rangle} {}^x \text{Res}_{\langle Qt \rangle \cap xH}^H F_{Q,t}^H \right).\end{aligned}$$

By Proposition 3.1, the element $\text{Res}_{\langle Qt \rangle \cap xH}^H F_{Q,t}^H$ is equal to the sum of the distinct idempotents $F_{Q^y, ty}^{\langle Qt \rangle \cap xH}$ corresponding to elements $y \in H$ such that $\langle Qt \rangle^y \leq \langle Qt \rangle \cap xH$. This implies $\langle Qt \rangle^y = \langle Qt \rangle^x$, i.e. $y \in N_G(\langle Qt \rangle)x$, thus $x \in N_G(\langle Qt \rangle) \cdot H$. This gives

$$\begin{aligned}\tau_{Q,t}^G(\text{Ind}_H^G F_{Q,t}^H) &= \tau_{Q,t}^{\langle Qt \rangle} \left(\sum_{\substack{x \in N_G(\langle Qt \rangle)H/H \\ y \in N_H(Q,t) \backslash N_G(\langle Qt \rangle)x}} {}^x F_{Q^y, ty}^{\langle Qt \rangle} \right) \\ &= \sum_{\substack{x \in N_G(\langle Qt \rangle)/N_H(\langle Qt \rangle) \\ y \in N_H(Q,t) \backslash N_G(\langle Qt \rangle)x}} \tau_{Q,t}^{\langle Qt \rangle} (F_{Q^{yx^{-1}}, tyx^{-1}}^{\langle Qt \rangle}) \\ &= \sum_{z \in N_H(Q,t) \backslash N_G(\langle Qt \rangle)} \tau_{Q,t}^{\langle Qt \rangle} (F_{Q^z, tz}^{\langle Qt \rangle}),\end{aligned}$$

where $z = yx^{-1}$. Finally $\tau_{Q,t}^{\langle Qt \rangle} (F_{Q^z, tz}^{\langle Qt \rangle})$ is equal to 1 if (Q^z, t^z) is conjugate to (Q, t) in $\langle Qt \rangle$, and to zero otherwise.

If $u \in \langle Qt \rangle$ is such that $(Q^z, t^z)^u = (Q, t)$, then $zu \in N_G(Q, t)$. But since $[\langle Qt \rangle, t] \leq Q$, we have $\langle Qt \rangle \leq N_G(Q, t)$, so $u \in N_G(Q, t)$, hence $z \in N_G(Q, t)$, and $(Q^z, t^z) = (Q, t)$. It follows that

$$\tau_{Q,t}^G(\text{Ind}_H^G F_{Q,t}^H) = |N_G(Q, t) : N_H(Q, t)|,$$

which completes the proof of the proposition. \square

3.4. Corollary. *Let $(P, s) \in \mathcal{Q}_{G,p}$. Then*

$$F_{P,s}^G = \frac{|s|}{|C_{\overline{N}_G(P)}(s)|} \text{Ind}_{\langle Ps \rangle}^G F_{P,s}^{\langle Ps \rangle}.$$

Proof : Apply Proposition 3.2 with $(Q, t) = (P, s)$ and $H = \langle Ps \rangle$. Then $N_H(Q, t) = \langle Ps \rangle$, thus by Exact sequence 2.14

$$|N_G(Q, t) : N_H(Q, t)| = \frac{|P| |C_{\overline{N}_G(P)}(s)|}{|P| |s|} = \frac{|C_{\overline{N}_G(P)}(s)|}{|s|},$$

and the corollary follows. \square

4. Idempotents

It follows from Corollary 3.4 that, in order to find formulae for the primitive idempotents $F_{P,s}^G$ of $K \otimes_{\mathbb{Z}} pp_k(G)$, it suffices to find the formula expressing the idempotent $F_{P,s}^{\langle Ps \rangle}$. In other words, we can assume that $G = \langle Ps \rangle$, i.e. that G has a normal Sylow p -subgroup P with cyclic quotient generated by s .

4.1. A morphism from the Burnside ring. When G is an arbitrary finite group, there is an obvious ring homomorphism \mathcal{L}_G from the Burnside ring $B(G)$ to $pp_k(G)$, induced by the *linearization* operation, sending a finite G -set X to the permutation module kX , which is obviously a p -permutation module. This morphism also commutes with restriction and induction : if $H \leq G$, then

$$(4.2) \quad \mathcal{L}_H \circ \text{Res}_H^G = \text{Res}_H^G \circ \mathcal{L}_G, \quad \mathcal{L}_G \circ \text{Ind}_H^G = \text{Ind}_H^G \circ \mathcal{L}_H.$$

Indeed, for any G -set X , the kH -modules $k\text{Res}_H^G X$ and $\text{Res}_H^G(kX)$ are isomorphic, and for any H -set Y , the kG -modules $k\text{Ind}_H^G Y$ and $\text{Ind}_H^G(kY)$ are isomorphic.

Similarly, when P is a p -subgroup of G , the ring homomorphism $\Phi_P : B(G) \rightarrow B(\overline{N}_G(P))$ induced by the operation $X \mapsto X^P$ on G -sets, is compatible with the Brauer morphism $\text{Br}_P^G : pp_k(G) \rightarrow pp_k(\overline{N}_G(P))$:

$$(4.3) \quad \mathcal{L}_{\overline{N}_G(P)} \circ \Phi_P = \text{Br}_P^G \circ \mathcal{L}_G.$$

This is because for any G -set X , the $k\overline{N}_G(P)$ -modules $k(X^P)$ and $(kX)[P]$ are isomorphic.

The ring homomorphism \mathcal{L}_G extends linearly to a K -algebra homomorphism $K \otimes_{\mathbb{Z}} B(G) \rightarrow K \otimes_{\mathbb{Z}} pp_k(G)$, still denoted by \mathcal{L}_G . The algebra $K \otimes_{\mathbb{Z}} B(G)$ is also a split semisimple commutative K -algebra. Its species are the K -algebra maps

$$K \otimes_{\mathbb{Z}} B(G) \rightarrow K, \quad X \mapsto |X^H|,$$

where H runs through the set of all subgroups of G up to conjugation. Here we denote by $|X^H|$ the number of H -fixed points of a G -set X and this notation is then extended K -linearly to any $X \in K \otimes_{\mathbb{Z}} B(G)$. The primitive idempotents e_H^G of $K \otimes_{\mathbb{Z}} B(G)$ are indexed by subgroups H of G , up to conjugation. They are given by the following formulae, found by Gluck ([4]) and later independently by Yoshida ([5]) :

$$(4.4) \quad e_H^G = \frac{1}{|N_G(H)|} \sum_{L \leq H} |L| \mu(L, H) G/L,$$

where μ denotes the Möbius function of the poset of subgroups of G . The idempotent e_H^G is characterized by the fact that for any $X \in K \otimes_{\mathbb{Z}} B(G)$

$$X \cdot e_H^G = |X^H| e_H^G .$$

4.5. Remark : Since $|X^H|$ only depends on $\text{Res}_H^G X$, it follows in particular that X is a scalar multiple of the “top” idempotent e_G^G if and only if $\text{Res}_H^G X = 0$ for any proper subgroup H of G . In particular, if N is a normal subgroup of G , then

$$(4.6) \quad (e_G^G)^N = e_{G/N}^{G/N} .$$

This is because for any proper subgroup H/N of G/N

$$\text{Res}_{H/N}^{G/N} (e_G^G)^N = (\text{Res}_H^G e_G^G)^N = 0 .$$

So $(e_G^G)^N$ is a scalar multiple of $e_{G/N}^{G/N}$. As it is also an idempotent, it is equal to 0 or $e_{G/N}^{G/N}$. Finally

$$|((e_G^G)^N)^{G/N}| = |(e_G^G)^G| = 1 ,$$

so $(e_G^G)^N$ is non zero.

4.7. The case of a cyclic p' -group. Suppose that G is a cyclic p' -group, of order n , generated by an element s . In this case, there are exactly n group homomorphisms from G to the multiplicative group k^\times of k . For each of these group homomorphisms φ , let k_φ denote the kG -module k on which the generator s acts by multiplication by $\varphi(s)$. As G is a p' -group, this module is simple and projective. The (classes of the) modules k_φ , for $\varphi \in \widehat{G} = \text{Hom}(G, k^\times)$, form a basis of $pp_k(G)$.

Since moreover for $\varphi, \psi \in \widehat{G}$, the modules $k_\varphi \otimes_k k_\psi$ and $k_{\varphi\psi}$ are isomorphic, the algebra $K \otimes_{\mathbb{Z}} pp_k(G)$ is isomorphic to the group algebra of the group \widehat{G} . This leads to the following classical formula :

4.8. Lemma. *Let G be a cyclic p' -group. Then for any $t \in G$,*

$$F_{1,t}^G = \frac{1}{n} \sum_{\varphi \in \widehat{G}} \tilde{\varphi}(t^{-1}) k_\varphi ,$$

where $\tilde{\varphi}$ is the Brauer character of k_φ .

Proof : Indeed for $s, t \in G$

$$\tau_{\mathbf{1},t}^G \left(\frac{1}{n} \sum_{\varphi \in \widehat{G}} \tilde{\varphi}(s^{-1}) k_{\varphi} \right) = \frac{1}{n} \sum_{\varphi \in \widehat{G}} \tilde{\varphi}(s^{-1}) \tilde{\varphi}(t) = \delta_{s,t} ,$$

where $\delta_{s,t}$ is the Kronecker symbol. □

4.9. The case $G = \langle Ps \rangle$. Suppose now more generally that $G = \langle Ps \rangle$, where P is a normal Sylow p -subgroup of G and s is a p' -element. In this case, by Proposition 3.1, the restriction of $F_{P,s}^G$ to any proper subgroup of G is equal to zero. Moreover, since $N_G(P, t) = G$ for any $t \in G/P$, the conjugacy class of the pair (P, t) reduces to $\{(P, t)\}$.

4.10. Lemma. *Suppose $G = \langle Ps \rangle$, and set $E_G^G = \mathcal{L}_G(e_G^G)$. Then*

$$E_G^G = \sum_{\langle t \rangle = \langle s \rangle} F_{P,t}^G .$$

Proof : By 4.2 and by Remark 4.5, the restriction of E_G^G to any proper subgroup of G is equal to zero. Let $(Q, t) \in \mathcal{Q}_{G,p}$, such that the group $L = \langle Qt \rangle$ is a proper subgroup of G . By Proposition 3.2, there is a rational number r such that

$$F_{Q,t}^G = r \operatorname{Ind}_L^G F_{Q,t}^L .$$

It follows that

$$E_G^G \cdot F_{Q,t}^G = r \operatorname{Ind}_L^G ((\operatorname{Res}_L^G E_G^G) \cdot F_{Q,t}^L) = 0 .$$

Now E_G^G is an idempotent of $K \otimes_{\mathbb{Z}} pp_k(G)$, hence is it a sum of some of the primitive idempotents $F_{Q,t}^G$ associated to pairs (Q, t) for which $\langle Qt \rangle = G$. This condition is equivalent to $Q = P$ and $\langle t \rangle = \langle s \rangle$.

It remains to show that all these idempotents $F_{P,t}^G$ appear in the decomposition of E_G^G , i.e. equivalently that $\tau_{P,t}^G(E_G^G) = 1$ for any generator t of $\langle s \rangle$. Now by 4.6 and Remark 2.16

$$\tau_{P,t}^G(E_G^G) = \tau_{\mathbf{1},t}^{G/P}(\operatorname{Br}_P^G(E_G^G)) = \tau_{\mathbf{1},t}^{\langle s \rangle}(E_{\langle s \rangle}^{\langle s \rangle}) .$$

Now the value at t of the Brauer character of a permutation module kX is equal to the number of fixed points of t on X . By K -linearity, this gives

$$\tau_{\mathbf{1},t}^{\langle s \rangle}(E_{\langle s \rangle}^{\langle s \rangle}) = |(e_{\langle s \rangle}^{\langle s \rangle})^t| ,$$

and this is equal to 1 if t generates $\langle s \rangle$, and to 0 otherwise, as was to be shown. □

4.11. Proposition. *Let $(P, s) \in \mathcal{Q}_{G,p}$, and suppose that $G = \langle Ps \rangle$. Then*

$$F_{P,s}^G = E_G^G \cdot \text{Inf}_{G/P}^G F_{1,s}^{G/P} .$$

Proof: Set $E_s = E_G^G \cdot \text{Inf}_{G/P}^G F_{1,s}^{G/P}$. Then E_s is an idempotent of $K \otimes_{\mathbb{Z}} \text{pp}_k(G)$, as it is the product of two (commuting) idempotents. Let $(Q, t) \in \mathcal{Q}_{G,p}$. If $\langle Qt \rangle \neq G$, then $\tau_{Q,t}^G(E_G^G) = 0$ by Lemma 4.10, thus $\tau_{Q,t}^G(E_s) = 0$. And if $\langle Qt \rangle = G$, then $Q = P$ and $\langle t \rangle = \langle s \rangle$. In this case

$$\tau_{Q,t}^G(E_s) = \tau_{P,t}^G(E_G^G) \cdot \tau_{P,t}^G(\text{Inf}_{G/P}^G F_{1,s}^{G/P}) .$$

By Lemma 4.10, the first factor in the right hand side is equal to 1. The second factor is equal to

$$\begin{aligned} \tau_{P,t}^G(\text{Inf}_{G/P}^G F_{1,s}^{G/P}) &= \tau_{1,t}^{G/P} \text{Br}_P^G(\text{Inf}_{G/P}^G F_{1,s}^{G/P}) \\ &= \tau_{1,t}^{G/P}(F_{1,s}^{G/P}) = \delta_{t,s} , \end{aligned}$$

where $\delta_{t,s}$ is the Kronecker symbol. Hence $\tau_{P,t}^G(E_s) = \delta_{t,s}$, and this completes the proof. \square

4.12. Theorem. *Let G be a finite group, and let $(P, s) \in \mathcal{Q}_{G,p}$. Then the primitive idempotent $F_{P,s}^G$ of the p -permutation algebra $K \otimes_{\mathbb{Z}} \text{pp}_k(G)$ is given by the following formula :*

$$F_{P,s}^G = \frac{1}{|P||s||C_{\overline{N}_G(P)}(s)|} \sum_{\substack{\varphi \in \langle s \rangle \\ L \leq \langle Ps \rangle \\ PL = \langle Ps \rangle}} \tilde{\varphi}(s^{-1}) |L| \mu(L, \langle Ps \rangle) \text{Ind}_L^G k_{L,\varphi}^{\langle Ps \rangle} ,$$

where $k_{L,\varphi}^{\langle Ps \rangle} = \text{Res}_L^{\langle Ps \rangle} \text{Inf}_{\langle s \rangle}^{\langle Ps \rangle} k_{\varphi}$.

Proof: By Corollary 3.4, and Proposition 4.11

$$F_{P,s}^G = \frac{|s|}{|C_{\overline{N}_G(P)}(s)|} \text{Ind}_{\langle Ps \rangle}^G (E_{\langle Ps \rangle}^{\langle Ps \rangle} \cdot \text{Inf}_{\langle s \rangle}^{\langle Ps \rangle} F_{1,s}^{\langle s \rangle}) .$$

By Equation 4.4, this gives

$$F_{P,s}^G = \frac{|s|}{|C_{\overline{N}_G(P)}(s)|} \text{Ind}_{\langle Ps \rangle}^G \frac{1}{|P||s|} \sum_{L \leq \langle Ps \rangle} |L| \mu(L, \langle Ps \rangle) \text{Ind}_L^{\langle Ps \rangle} k \cdot \text{Inf}_{\langle s \rangle}^{\langle Ps \rangle} F_{1,s}^{\langle s \rangle} .$$

Moreover for each $L \leq \langle Ps \rangle$

$$\begin{aligned} \text{Ind}_L^{\langle Ps \rangle} k \cdot \text{Inf}_{\langle s \rangle}^{\langle Ps \rangle} F_{1,s}^{\langle s \rangle} &\cong \text{Ind}_L^{\langle Ps \rangle} (\text{Res}_L^{\langle Ps \rangle} \text{Inf}_{\langle s \rangle}^{\langle Ps \rangle} F_{1,s}^{\langle s \rangle}) \\ &\cong \text{Ind}_L^{\langle Ps \rangle} \text{Inf}_{L/L \cap P}^L \text{Iso}_{LP/P}^{L/L \cap P} \text{Res}_{LP/P}^{\langle s \rangle} F_{1,s}^{\langle s \rangle} . \end{aligned}$$

Here we have used the fact that if L and P are subgroups of a group H , with $P \trianglelefteq H$, then there is an isomorphism of functors

$$\text{Res}_L^H \circ \text{Inf}_{H/P}^H \cong \text{Inf}_{L/L \cap P}^L \circ \text{Iso}_{LP/P}^{L/L \cap P} \circ \text{Res}_{LP/P}^{H/P} ,$$

which follows from the isomorphism of $(L, H/P)$ -bisets

$$H \times_H (H/P) \cong {}_L(H/P)_{H/P} \cong (L/L \cap P) \times_{L/L \cap P} (LP/P) \times_{LP/P} (H/P) .$$

Now Proposition 3.1 implies that $\text{Res}_{LP/P}^{\langle s \rangle} F_{1,s}^{\langle s \rangle} = 0$ if $LP/P \neq \langle s \rangle$, i.e. equivalently if $PL \neq \langle Ps \rangle$. It follows that

$$F_{P,s}^G = \frac{1}{|P| |C_{\overline{N}_G(P)}(s)|} \sum_{\substack{L \leq \langle Ps \rangle \\ PL = \langle Ps \rangle}} |L| \mu(L, \langle Ps \rangle) \text{Ind}_L^G (\text{Res}_L^{\langle Ps \rangle} \text{Inf}_{\langle s \rangle}^{\langle Ps \rangle} F_{1,s}^{\langle s \rangle}) .$$

By Lemma 4.8, this gives

$$F_{P,s}^G = \frac{1}{|P| |s| |C_{\overline{N}_G(P)}(s)|} \sum_{\substack{\varphi \in \widehat{\langle s \rangle} \\ L \leq \langle Ps \rangle \\ PL = \langle Ps \rangle}} \tilde{\varphi}(s^{-1}) |L| \mu(L, \langle Ps \rangle) \text{Ind}_L^G k_{L,\varphi}^{\langle Ps \rangle} ,$$

where $k_{L,\varphi}^{\langle Ps \rangle} = \text{Res}_L^{\langle Ps \rangle} \text{Inf}_{\langle s \rangle}^{\langle Ps \rangle} k_\varphi$, as was to be shown. \square

References

- [1] D. Benson. *Representations and cohomology I*, volume 30 of *Cambridge studies in advanced mathematics*. Cambridge University Press, 1991.
- [2] S. Bouc. On the Cartan matrix of Mackey algebras. Preprint, 2009.
- [3] M. Broué. On Scott modules and p -permutation modules: an approach through the Brauer morphism. *Proc.AMS*, 93(3):401–408, March 1985.
- [4] D. Gluck. Idempotent formula for the Burnside ring with applications to the p -subgroup simplicial complex. *Illinois J. Math.*, 25:63–67, 1981.

- [5] T. Yoshida. Idempotents in Burnside rings and Dress induction theorem.
J. Algebra, 80:90–105, 1983.

Serge Bouc, CNRS-LAMFA, Université de Picardie - Jules Verne,
33, rue St Leu, F-80039 Amiens Cedex 1, France.
`serge.bouc@u-picardie.fr`

Jacques Thévenaz, Institut de Géométrie, Algèbre et Topologie,
EPFL, Bâtiment BCH, CH-1015 Lausanne, Switzerland.
`Jacques.Thevenaz@epfl.ch`