A remark on the Schrödinger smoothing effect

Laurent Thomann

► To cite this version:

Laurent Thomann. A remark on the Schrödinger smoothing effect. *Asymptotic Analysis*, 2010, 69 (1-2), pp.117–123. hal-00422424v2

HAL Id: hal-00422424

<https://hal.science/hal-00422424v2>

Submitted on 25 Dec 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A REMARK ON THE SCHRÖDINGER SMOOTHING EFFECT

by

Laurent Thomann

Abstract. — We prove the equivalence between the smoothing effect for a Schrödinger operator and the decay of the associate spectral projectors. We give two applications to the Schrödinger operator in dimension one.

Résumé. — On donne une caractérisation de l'effet régularisant pour un opérateur de Schrödinger par la décroissance de ses projecteurs spectraux. On en déduit deux applications à l'opérateur de Schrödinger en dimension un.

1. Introduction

Let $d \geq 1$, and consider the linear Schrödinger equation

$$(1.1) \quad \begin{cases} i\partial_t u = Hu, & (t, x) \in \mathbb{R} \times \mathbb{R}^d, \\ u(0, x) = f(x) \in L^2(\mathbb{R}^d), \end{cases}$$

where H is a self-adjoint operator on $L^2(\mathbb{R}^d)$.

By the Hille-Yoshida theorem, the equation (1.1) admits a unique solution $u(t) = e^{-itH}f \in \mathcal{C}(\mathbb{R}; L^2(\mathbb{R}^d))$. Under suitable conditions on H , this solution enjoys a local gain of regularity (in the space variable) : For all $T > 0$ there exists $C > 0$ so that

$$\left(\int_0^T \|\Psi(x) \langle H \rangle^{\frac{\gamma}{2}} e^{-itH} f\|_{L^2(\mathbb{R}^d)}^2 dt \right)^{\frac{1}{2}} \leq C \|f\|_{L^2(\mathbb{R}^d)},$$

for some weight Ψ and exponent $\gamma > 0$.

2000 Mathematics Subject Classification. — 35-XX.

Key words and phrases. — Schrödinger equation, potential, smoothing effect.

The author was supported in part by the grant ANR-07-BLAN-0250.

This phenomenon has been discovered by T. Kato [7] in the context of KdV equations. For the Schrödinger equation in the case $H = -\Delta$, it has been proved by P. Constantin- J.-C. Saut [2], P. Sjölin [11], L. Vega [12] and K. Yajima [13]. The variable coefficients case has been obtained by S. Doi [3, 4, 5, 6].

The more general results are due to L. Robbiano-C. Zuily [9, 10] for equations with obstacles and potentials.

Let H be a self adjoint operator on $L^2(\mathbb{R}^d)$. It can be represented thanks to the spectral measure by

$$H = \int \lambda dE_\lambda.$$

In the sequel we moreover assume that $H \geq 0$. For $N \geq 0$, we can then define the spectral projector P_N associated to H by

$$(1.2) \quad P_N = \mathbf{1}_{[N, N+1[}(H) = \int \mathbf{1}_{[N, N+1[}(\lambda) dE_\lambda.$$

Our main result is a characterisation of the smoothing effect by the decay of the spectral projectors. Denote by $\langle H \rangle = (1 + H^2)^{\frac{1}{2}}$.

Theorem 1.1 (Smoothing effect vs. decay). —

Let $\gamma > 0$ and $\Psi \in \mathcal{C}(\mathbb{R}^d, \mathbb{R})$. Then the following conditions are equivalent

(i) There exists $C_1 > 0$ so that for all $f \in L^2(\mathbb{R}^d)$

$$(1.3) \quad \left(\int_0^{2\pi} \|\Psi(x) \langle H \rangle^{\frac{\gamma}{2}} e^{-itH} f\|_{L^2(\mathbb{R}^d)}^2 dt \right)^{\frac{1}{2}} \leq C_1 \|f\|_{L^2(\mathbb{R}^d)}.$$

(ii) There exists $C_2 > 0$ so that for all $N \geq 1$ and $f \in L^2(\mathbb{R}^d)$

$$(1.4) \quad \|\Psi P_N f\|_{L^2(\mathbb{R}^d)} \leq C_2 N^{-\frac{\gamma}{2}} \|P_N f\|_{L^2(\mathbb{R}^d)}.$$

The interesting point is that we can take the same function Ψ and exponent $\gamma > 0$ in both statements (1.3) and (1.4).

By the works cited in the introduction, in the case $H = -\Delta$ on $\mathbb{R}^d$, (1.3) is known to hold with $\gamma = \frac{1}{2}$ and $\Psi(x) = \langle x \rangle^{-\frac{1}{2}-\nu}$, for any $\nu > 0$.

There is also a class of operators H on $L^2(\mathbb{R}^d)$ for which (1.3) is well understood. Let $V \in \mathcal{C}^\infty(\mathbb{R}, \mathbb{R}_+)$, and assume that for $|x|$ large enough $V(x) \geq C \langle x \rangle^k$ and that for any $j \in \mathbb{N}^d$, there exists $C_j > 0$ so that $|\partial_x^j V(x)| \leq C_j \langle x \rangle^{k-|j|}$. Then L. Robbiano and C. Zuily [9] show that the smoothing effect (1.3) holds for the operator $H = -\Delta + V(x)$, with $\gamma = \frac{1}{k}$ and $\Psi(x) = \langle x \rangle^{-\frac{1}{2}-\nu}$, for any $\nu > 0$.

We now turn to the case of dimension $d = 1$, and consider the operator $H = -\Delta + V(x)$. We make the following assumption on V

Assumption 1. — *We suppose that $V \in C^\infty(\mathbb{R}, \mathbb{R}_+)$, and that there exist $2 < m \leq k$ so that for $|x|$ large enough*

(i) *There exists $C > 1$ so that $\frac{1}{C}\langle x \rangle^k \leq V(x) \leq C\langle x \rangle^k$.*

(ii) *$V''(x) > 0$ and $xV'(x) \geq mV(x) > 0$*

(iii) *For any $j \in \mathbb{N}$, there exists $C_j > 0$ so that $|\partial_x^j V(x)| \leq C_j \langle x \rangle^{k-|j|}$.*

For instance $V(x) = \langle x \rangle^k$ with $k > 2$ satisfies Assumption 1.

It is well known that under Assumption 1, the operator H has a self-adjoint extension on $L^2(\mathbb{R})$ (still denoted by H) and has eigenfunctions $(e_n)_{n \geq 1}$ which form an Hilbertian basis of $L^2(\mathbb{R})$ and satisfy

$$He_n = \lambda_n^2 e_n, \quad n \geq 1,$$

with $\lambda_n \rightarrow +\infty$, when $n \rightarrow +\infty$.

For $N \in \mathbb{N}$ the spectral projector P_N defined in (1.2) can be written in the following way. Let $f = \sum_{n \geq 1} \alpha_n e_n \in L^2(\mathbb{R})$, then

$$P_N f = \sum_{N \leq \lambda_n^2 < N+1} \alpha_n e_n.$$

Observe that we then have $f = \sum_{N \geq 0} P_N f$.

For such a potential, we can remove the spectral projectors in (1.4) and deduce from Theorem 1.1

Corollary 1.2. —

Let $\gamma > 0$ and $\Psi \in \mathcal{C}(\mathbb{R}, \mathbb{R})$. Let $H = \Delta + V(x)$ so that $V(x) = x^2$ or $V(x)$ satisfies Assumption 1. Then the following conditions are equivalent

(i) *There exists $C_1 > 0$ so that for all $f \in L^2(\mathbb{R})$*

$$(1.5) \quad \left(\int_0^{2\pi} \|\Psi(x) \langle H \rangle^{\frac{\gamma}{2}} e^{-itH} f\|_{L^2(\mathbb{R})}^2 dt \right)^{\frac{1}{2}} \leq C_1 \|f\|_{L^2(\mathbb{R})}.$$

(ii) *There exists $C_2 > 0$ so that for all $n \geq 1$*

$$(1.6) \quad \|\Psi e_n\|_{L^2(\mathbb{R})} \leq C_2 \lambda_n^{-\gamma}, \quad \forall n \geq 1.$$

The statements (1.5) and (1.6) were obtained by K. Yajima & G. Zhang in [16] when Ψ is the indicator of a compact $K \subset \mathbb{R}$ and with $\gamma = \frac{1}{k}$.

The statement (1.5) holds for $\Psi(x) = \langle x \rangle^{-\frac{1}{2}-\nu}$, by [9], but as far as we know, the bound (1.6) with this Ψ was unknown.

With Theorem 1.1 we are also able to prove the following smoothing effect for the usual Laplacian Δ on $\mathbb{R}$.

Proposition 1.3. — *Let $\Psi \in L^2(\mathbb{R})$. Then there exists $C > 0$ so that for all $f \in L^2(\mathbb{R})$*

$$\left(\int_0^{2\pi} \|\Psi(x) \langle \Delta \rangle^{\frac{1}{4}} e^{-it\Delta} f\|_{L^2(\mathbb{R})}^2 dt \right)^{\frac{1}{2}} \leq C \|\Psi\|_{L^2(\mathbb{R})} \|f\|_{L^2(\mathbb{R})}.$$

From the works cited in the introduction, we have

$$\left(\int_{\mathbb{R}} \|\Psi(x) \langle \Delta \rangle^{\frac{1}{4}} e^{-it\Delta} f\|_{L^2(\mathbb{R})}^2 dt \right)^{\frac{1}{2}} \leq C \|f\|_{L^2(\mathbb{R})},$$

for $\Psi(x) = \langle x \rangle^{\frac{1}{2}-\nu}$, for any $\nu > 0$. Hence Proposition 1.3 shows that we can extend the class of the weights, but we are only able to prove local integrability in time.

Notation. — *We use the notation $a \lesssim b$ if there exists a universal constant $C > 0$ so that $a \leq Cb$.*

2. Proof of the results

We define the self adjoint operator $A = [H]$ (entire part of H) by

$$A = \int [\lambda] dE_\lambda.$$

Notice that we immediately have that $A - H$ is bounded on $L^2(\mathbb{R}^d)$.

The first step in the proof of Theorem 1.1 is to show that we can replace e^{-itH} by e^{-itA} in (1.3)

Lemma 2.1. — *Let $\gamma > 0$ and $\Psi \in \mathcal{C}(\mathbb{R}^d, \mathbb{R})$. Then the following conditions are equivalent*

(i) *There exists $C_1 > 0$ so that for all $f \in L^2(\mathbb{R}^d)$*

$$(2.1) \quad \left(\int_0^{2\pi} \|\Psi(x) \langle H \rangle^{\frac{\gamma}{2}} e^{-itA} f\|_{L^2(\mathbb{R}^d)}^2 dt \right)^{\frac{1}{2}} \leq C_1 \|f\|_{L^2(\mathbb{R}^d)}.$$

(ii) *There exists $C_2 > 0$ so that for all $f \in L^2(\mathbb{R}^d)$*

$$(2.2) \quad \left(\int_0^{2\pi} \|\Psi(x) \langle H \rangle^{\frac{\gamma}{2}} e^{-itH} f\|_{L^2(\mathbb{R}^d)}^2 dt \right)^{\frac{1}{2}} \leq C_2 \|f\|_{L^2(\mathbb{R}^d)}.$$

Proof. — We assume (2.1) and we prove (2.2). Let $f \in L^2(\mathbb{R}^d)$ and Define $v = e^{-itH}f$. This function solves the problem

$$(i\partial_t - A)v = (H - A)v, \quad v(0, x) = f(x).$$

Then by the Duhamel formula

$$\begin{aligned} e^{-itH}f = v &= e^{-itA}f - i \int_0^t e^{-i(t-s)A}(H - A)v \, ds \\ &= e^{-itA}f - i \int_0^{2\pi} \mathbf{1}_{\{s < t\}} e^{-i(t-s)A}(H - A)v \, ds. \end{aligned}$$

Therefore by (2.1) and Minkowski

$$\begin{aligned} \|\Psi \langle H \rangle^{\frac{\gamma}{2}} e^{-itH}v\|_{L_{2\pi}^2 L^2} &\lesssim \|\Psi \langle H \rangle^{\frac{\gamma}{2}} e^{-itA}v\|_{L_{2\pi}^2 L^2} \\ &\quad + \int_0^{2\pi} \|\Psi \langle H \rangle^{\frac{\gamma}{2}} \mathbf{1}_{\{s < t\}} e^{-i(t-s)A}(H - A)v\|_{L_t^2 L_x^2} \, ds \\ (2.3) \quad &\lesssim \|f\|_{L^2} + \int_0^{2\pi} \|(H - A)v\|_{L^2} \, ds. \end{aligned}$$

Now use that the operator $(H - A) : L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)$ is bounded, and by (2.3) we obtain

$$\|\Psi \langle H \rangle^{\frac{\gamma}{2}} e^{-itH}v\|_{L_{2\pi}^2 L^2} \lesssim \|f\|_{L^2},$$

which is (2.2).

The proof of the converse implication is similar. $\square$

Proof of Theorem 1.1. — The proof is based on Fourier analysis in time. This idea comes from [8] and has also been used in [16], but this proof was inspired by [1].

(i) $\implies$ (ii) : To prove this implication, we use the characterisation (2.1). From (1.2) and the definition of A , $e^{-itA}P_N f = e^{-itN}P_N f$. Hence it suffices to replace f with $P_N f$ in (1.3) and (1.4) follows.

(ii) $\implies$ (i) : Again we will use Lemma 2.1. We assume (2.2) and we first prove that

$$(2.4) \quad \|\Psi \langle A \rangle^{\frac{\gamma}{2}} e^{-itA}f\|_{L^2(0, 2\pi; L^2(\mathbb{R}^d))} \lesssim \|f\|_{L^2(\mathbb{R}^d)}.$$

Write $f = \sum_{N \geq 0} P_N f$, then

$$\Psi \langle A \rangle^{\frac{\gamma}{2}} e^{-itA}f = \sum_{N \geq 0} e^{-iNt} \langle N \rangle^{\frac{\gamma}{2}} \Psi P_N f.$$

Now by Parseval in time

$$\|\Psi \langle A \rangle^{\frac{\gamma}{2}} e^{-itA} f\|_{L^2(0,2\pi)}^2 \lesssim \sum_{N \geq 0} \langle N \rangle^\gamma |\Psi P_N f|^2,$$

and by integration in the space variable and (1.4)

$$\begin{aligned} \|\Psi \langle A \rangle^{\frac{\gamma}{2}} e^{-itA} f\|_{L^2(0,2\pi; L^2(\mathbb{R}^d))}^2 &\lesssim \sum_{N \geq 0} \langle N \rangle^\gamma \|\Psi P_N f\|_{L^2(\mathbb{R}^d)}^2 \\ &\lesssim \sum_{N \geq 0} \|P_N f\|_{L^2(\mathbb{R}^d)}^2 = \|f\|_{L^2(\mathbb{R}^d)}^2, \end{aligned}$$

which yields (2.4).

Now since the operator $\langle A \rangle^{-\gamma/2} \langle H \rangle^{\gamma/2}$ is bounded on L^2 and commutes with e^{-itA} , we have by (2.4)

$$\begin{aligned} \|\Psi \langle H \rangle^{\frac{\gamma}{2}} e^{-itA} f\|_{L^2(0,2\pi; L^2(\mathbb{R}^d))} &= \\ &= \|\Psi \langle A \rangle^{\frac{\gamma}{2}} e^{-itA} (\langle A \rangle^{-\frac{\gamma}{2}} \langle H \rangle^{\frac{\gamma}{2}} f)\|_{L^2(0,2\pi; L^2(\mathbb{R}^d))} \\ &\lesssim \|\langle A \rangle^{-\frac{\gamma}{2}} \langle H \rangle^{\frac{\gamma}{2}} f\|_{L^2(\mathbb{R}^d)} \\ &\lesssim \|f\|_{L^2(\mathbb{R}^d)}, \end{aligned}$$

which is (2.1). □

Proof of Corollary 1.2. — Let V satisfy Assumption 1. Then by [14, Lemma 3.3] there exists $C > 0$ such that

$$|\lambda_{n+1}^2 - \lambda_n^2| \geq C \lambda_n^{1-\frac{2}{m}},$$

for n large enough. This implies that $[\lambda_n^2] < [\lambda_{n+1}^2]$ for n large enough, because $m > 2$ and $\lambda_n \rightarrow +\infty$. As a consequence

$$P_N f = \alpha_n e_n, \text{ with } n \text{ so that } N \leq \lambda_n^2 < N+1,$$

and this yields the result.

We now consider $V(x) = x^2$. In this case, the eigenvalues are the integers $\lambda_n^2 = 2n+1$, and the claim follows. □

Remark 2.2. — With this time Fourier analysis, we can prove the following smoothing estimate for H which satisfies Assumption 1

$$\|\langle H \rangle^{\frac{\theta(q,k)}{2}} e^{-itH} f\|_{L^p(\mathbb{R}; L^2(0,T))} \lesssim \|f\|_{L^2(\mathbb{R})},$$

where θ is defined by

$$\theta(q, k) = \begin{cases} \frac{2}{k}(\frac{1}{2} - \frac{1}{q}) & \text{if } 2 \leq q < 4, \\ \frac{1}{2k} - \eta \text{ for any } \eta > 0 & \text{if } q = 4, \\ \frac{1}{2} - \frac{2}{3}(1 - \frac{1}{q})(1 - \frac{1}{k}) & \text{if } 4 < q < \infty, \\ \frac{4-k}{6k} & \text{if } q = \infty. \end{cases}$$

This was done in [16] with a slightly different formulation.

Proof of Proposition 1.3. — By Theorem 1.1, we have to prove that the operator T defined by

$$Tf(x) = N^{\frac{1}{4}}\Psi(x)\mathbf{1}_{[N, N+1]}(-\Delta)f(x),$$

is continuous from $L^2(\mathbb{R})$ to $L^2(\mathbb{R})$ with norm independent of $N \geq 1$. By the usual TT^* argument, it is enough to show the result for TT^* .

The kernel of T is $K(x, y) = N^{\frac{1}{4}}\Psi(x)F_N(x - y)$ where

$$(2.5) \quad F_N(u) = \frac{1}{2\pi} \int e^{iu\xi} \mathbf{1}_{[\sqrt{N}, \sqrt{N+1}]}(|\xi|) d\xi = 4 \cos(D_N u) \frac{\sin(C_N u)}{u},$$

with $C_N = (\sqrt{N+1} - \sqrt{N})/2$ and $D_N = (\sqrt{N+1} + \sqrt{N})/2$.

The kernel of TT^* is given by

$$\Lambda(x, z) = \int K(x, y) \overline{K}(z, y) dy,$$

and by Parseval and (2.5)

$$\begin{aligned} \Lambda(x, z) &= N^{\frac{1}{2}}\Psi(x)\Psi(z) \int F_N(x - y) \overline{F_N}(z - y) dy \\ &= \frac{1}{4} N^{\frac{1}{2}}\Psi(x)\Psi(z) \int e^{i(x-z)\xi} \mathbf{1}_{[\sqrt{N}, \sqrt{N+1}]}(|\xi|) d\xi \\ &= \pi N^{\frac{1}{2}}\Psi(x)\Psi(z) \cos(D_N(x - z)) \frac{\sin(C_N(x - z))}{x - z}. \end{aligned}$$

Now, since $C_N \lesssim 1/\sqrt{N}$ and $|\sin(x)| \leq |x|$, we deduce that $|\Lambda(x, z)| \leq C|\Psi(x)||\Psi(z)|$ (independent of $N \geq 1$), and TT^* is continuous for $\Psi \in L^2(\mathbb{R})$. $\square$

Acknowledgements. — *The author would like to thank D. Robert for many enriching discussions and the anonymous referee for valuable suggestions which improved this paper.*

References

- [1] N. Burq, G. Lebeau and F. Planchon. Global existence for energy critical waves in 3-D domains. *Journal of the A.M.S.*, (21) 831–845, 2008.
- [2] P. Constantin and J.-C. Saut. Local smoothing properties of dispersive equations. *Journal American Mathematical Society* (1988) 413–439.
- [3] S. Doi. Smoothing effects of Schrödinger evolution group on Riemannian manifolds. *Duke Math. J.* 82 (1996) 679–706.
- [4] S. Doi. Smoothing effects for Schrödinger evolution equation and global behavior of geodesic flow. *Math. Ann.* 318 (2000) 355–389.
- [5] S. Doi. Remarks on the Cauchy problem for Schrödinger type equations. *Comm. in PDE*, 21 (1996) 163–178.
- [6] S. Doi. Smoothness of solutions for Schrödinger equations with unbounded potential. *Publ. Res. Inst. Math. Sci* 41 (2005), 1, 175–221.
- [7] T. Kato. On nonlinear Schrödinger equations, II. H^s -solutions and unconditional well-posedness, *J. Anal. Math.*, 87 (1995) 281–306.
- [8] G. Mockenhaupt, A. Seeger and C. Sogge. Local smoothing of Fourier integral operators and Carleson-Sjölin estimates. *J. Amer. Math. Soc.*, 6(1):65–130, 1993.
- [9] L. Robbiano, and C. Zuily. Remark on the Kato smoothing effect for Schrödinger equation with superquadratic potentials. *Comm. Partial Differential Equations*, 33 (2008), no. 4-6, 718–727.
- [10] L. Robbiano, and C. Zuily. The Kato Smoothing Effect for Schrödinger equations with unbounded potentials in exterior domains. *Int. Math. Res. Notices*, 15 (2009), 1636–1698.
- [11] P. Sjölin. Regularity of solution to the Schrödinger equation. *Duke Math. J.* 55 (1987) 699–715.
- [12] L. Vega. Schrödinger equations, pointwise convergence to the initial data. *Proc. Amer. Math. Soc.* 102 (1988) 874–878.
- [13] K. Yajima. On smoothing property of Schrödinger propagator. *Lectures notes in Math.* 1450 Springer Verlag (1990) 20–35.
- [14] K. Yajima. Smoothness and non-smoothness of the fundamental solution of time dependent Schrödinger equations. *Comm. Math. Phys.* 180 (1996) no. 3, 605–629.
- [15] K. Yajima, and G. Zhang. Local smoothing property and Strichartz inequality for Schrödinger equations with potentials superquadratic at infinity. *J. Differential Equations* (2004), no. 1, 81–110.
- [16] K. Yajima, and G. Zhang. Smoothing property for Schrödinger equations with potential superquadratic at infinity. *Comm. Math. Phys.* 221 (2001), no. 3, 573–590.

LAURENT THOMANN, Université de Nantes, Laboratoire de Mathématiques
 J. Leray, UMR CNRS 6629, 2, rue de la Houssinière, F-
 44322 Nantes Cedex 03, France. • E-mail : laurent.thomann@univ-nantes.fr
 Url : <http://www.math.sciences.univ-nantes.fr/~thomann/>