

A new proof of Delsarte, Goethals and Mac Williams theorem on minimal weight codewords of generalized Reed-Muller codes

Elodie LEDUCQ

January 13, 2010

Abstract

We give a new proof of Delsarte, Goethals and Mac Williams theorem on minimal weight codewords of generalized Reed-Muller codes published in 1970. To prove this theorem, we consider intersection of support of minimal weight codewords with affine hyperplanes and we proceed by recursion.

1 Introduction

In the appendix of [1], Delsarte, Goethals and Mac Williams prove the theorem 1 below. However, at the beginning of their proof, they point out that "it would be very desirable to find a more sophisticated and shorter proof".

In this paper, we give a new proof of this theorem that we hope is simpler.

Let $q = p^n$, p being prime number.

We identify the \mathbb{F}_q -algebra $B_m^q = \mathbb{F}_q[X_1, \dots, X_m]/(X_1^q - X_1, \dots, X_m^q - X_m)$ to the \mathbb{F}_q -algebra of the functions from \mathbb{F}_q^m to \mathbb{F}_q through the isomorphism $P \mapsto (x \mapsto P(x))$.

For $f \in B_m^q$, let $S_f = \{x \in \mathbb{F}_q^m, f(x) \neq 0\}$ the support of f , and $|f| = \text{Card}(S_f)$ the weight of f . The Hamming distance in B_m^q is denoted by $d(., .)$.

For $0 \leq r \leq m(q-1)$, the r th order generalized Reed-Muller code of length q^m is

$$R_q(r, m) = \{P \in B_m^q, \deg(P) \leq r\}$$

where $\deg(P)$ is the degree of the representative of P with degree at most $q-1$ in each variable.

The affine group $\text{GA}_m(\mathbb{F}_q)$ acts on $R_q(r, m)$ by its natural action. The minimum weight of $R_q(r, m)$ is $(q-s)q^{m-t-1}$, where $r = t(q-1) + s$, $0 \leq s \leq q-2$ (see [1]).

The following theorem gives the codeword of minimum weight of $R_q(r, m)$

Theorem 1 *Let $r = t(q-1) + s < m(q-1)$. The minimal weight codewords of $R_q(r, m)$ are codewords of $R_q(r, m)$ whose support is the union of $(q-s)$ distinct parallel affine subspaces of codimension $t+1$ included in an affine subspace of codimension t .*

Remark :

1. Clearly, codewords of this form are of minimal weight.
2. Using lemma 2 and corollary 3 below, this theorem means that codewords of minimal weight are equivalent, under the action of the affine group, to a codeword of the following form: $f(x) = c \prod_{i=1}^t (x_i^{q-1} - 1) \prod_{j=1}^s (x_{t+1} - b_j)$ where $c \in \mathbb{F}_q^*$, b_j are distinct elements of \mathbb{F}_q .

2 Proof of Theorem 1

In this paper, we use freely the two following lemmas and their corollary proved in [1] p. 435.

Lemma 2 *If $P(x) = 0$ whenever $x_1 = a$, then $P(x) = (x_1 - a)Q(x)$ where $\deg_{x_1}(Q) \leq \deg_{x_1}(P) - 1$.*

Remark : In [1], the lemma A1.1 says that the exponent of x_1 in Q is at most $q-2$, but they actually prove the above lemma.

Corollary 3 *If $P(x) = 0$ unless $x_1 = b$, then $P(x) = (1 - (x_1 - b)^{q-1})Q(x_2, \dots, x_m)$.*

Lemma 4 *Let S be a subset of \mathbb{F}_q^m , such that $\text{Card}(S) = tq^n < q^m$, $0 < t < q$. Assume that for any hyperplane of \mathbb{F}_q^m , either $\text{Card}(S \cap H) = 0$ or $\text{Card}(S \cap H) \geq tq^{n-1}$. Then there exists an affine hyperplane of \mathbb{F}_q^m which does not meet S .*

Now, we prove a lemma that will be crucial in our proof of theorem 1 :

Lemma 5 *Let $r = t(q-1) + s$, $0 \leq s \leq q-2$, f be a minimal weight codeword of $R_q(r, m)$ and $S = S_f$. If H is an hyperplane of \mathbb{F}_q^m , such that $S \cap H \neq \emptyset$ and $S \cap H \neq S$, then either S meets all hyperplanes parallel to H or S meets $q-s$ hyperplanes parallel to H in q^{m-t-1} points.*

Proof : Since an affine transformation does not change weight, we can assume that $H = \{x, x_1 = 0\}$. Now assume that S does not meet k hyperplanes parallel to H , $k \geq 1$. As $S \cap H \neq \emptyset$ and $S \cap H \neq H$, we have $k \leq q-2$. By lemma 2, we can write

$$f(x) = (x_1 - b_1)^{\alpha_1} \dots (x_1 - b_k)^{\alpha_k} P(x)$$

where S_P meets all hyperplanes parallel to H .

Let $d = \sum_{i=1}^k \alpha_i \leq q-1$. We want to prove that $d = s$.

- First, assume that $d > s$. Then the degree of P is $(t-1)(q-1) + q-1 + s-d$ and $0 \leq q-1 + s-d \leq q-2$. For $c \notin \{b_1, \dots, b_k\}$, we consider $Q_c = (1 - (x_1 - c)^{q-1})P$. The degree of Q_c is $t(q-1) + q-1 + s-d$. So

$$\begin{aligned} (q-s)q^{m-t-1} = \text{Card}(S) &= \sum_{c \notin \{b_1, \dots, b_k\}} \text{Card}(S_P \cap \{x_1 = c\}) \\ &\geq (q-k)(q - (q-1 + s-d))q^{m-t-1} \\ &= (q-k)(d-s+1)q^{m-t-1} \end{aligned}$$

and we obtain, since $d \geq k$,

$$(q-1-k)(d-s) \leq 0,$$

which is impossible, since $d > s$ and $k < q-1$.

- Now we have $d \leq s$. So $\deg(P) = t(q-1) + s-d$ and we get

$$\begin{aligned} (q-s)q^{m-t-1} = \text{Card}(S) &= \sum_{c \notin \{b_1, \dots, b_k\}} \text{Card}(S_P \cap \{x_1 = c\}) \\ &\geq (q-k)(q-s+d)q^{m-t-2} \end{aligned}$$

which gives

$$(d-k)q + k(s-d) \leq 0.$$

Hence, since $d \geq k$, $k \geq 1$ and $s \geq d$, necessarily $d = k = s$ and $\text{Card}(S_P \cap \{x_1 = c\}) = q^{m-t-1}$ for $c \notin \{b_1, \dots, b_k\}$.

□

Now, we are able to prove theorem 1.

Proof : We prove first the case where $t = 0$ and $t = m-1$.

- $t = 0$.

If $s = 0$, then $\deg(f) = 0$. Thus, since $f \neq 0$, we have $f = c$, for $c \in \mathbb{F}_q^*$ and $S_f = \mathbb{F}_q^m$.

Otherwise, let H be an affine hyperplane of \mathbb{F}_q^m , then $\text{Card}(S_f \cap H) = 0$ or $\text{Card}(S_f \cap H) \geq (q-s)q^{m-2}$.

Hence, by lemma 4, there exists an affine hyperplane H_0 , such that $S_f \cap H_0 = \emptyset$.

However, \mathbb{F}_q^m is the union of the q hyperplanes parallel to H_0 , so there exists H_1 , parallel to H_0 , such that $H_1 \cap S_f \neq \emptyset$.

Furthermore, since $|f| = (q-s)q^{m-1} \geq 2q^{m-1}$, $S_f \cap H_1 \neq S_f$. So, by lemma 5,

since $S_f \cap H_0 = \emptyset$, S_f meets $(q-s)$ hyperplanes parallel to H_1 , say H_1, \dots, H_{q-s} , in q^{m-1} points, this means that $S_f = \bigcup_{i=1}^{q-s} H_i$.

- $t = m - 1$.

Let $f \in R_q((m-1)(q-1) + s, m)$, $0 \leq s \leq q-2$, such that $|f| = q-s$. We put $S = S_f$. Let $\omega_1, \omega_2 \in S$ and H be an hyperplane, such that $\omega_1, \omega_2 \in H$. Assume that $S \cap H \neq S$ then, by lemma 5, either S meets all hyperplanes parallel to H (which is possible only if $s = 0$) or S meets $(q-s)$ hyperplanes parallel to H in one point. In both cases we get a contradiction, since in both cases S meets each hyperplane in exactly one point and $\omega_1, \omega_2 \in H$. So S is included in all hyperplanes H , such that $\omega_1, \omega_2 \in H$, this means that S is included in the line through ω_1 and ω_2 .

Now we prove the theorem for general t by recursion.

- Assume that for a fixed t , $1 \leq t \leq m-2$ (we have already proved the case where $t = 0$) and for all $0 \leq s \leq q-2$, the support of a codeword of minimal weight in $R_q((t+1)(q-1) + s, m)$ is the union of $(q-s)$ distinct parallel affine subspaces of codimension $t+2$ included in an affine subspace of codimension $t+1$.

Let $f \in R_q(t(q-1) + s, m)$, such that $|f| = (q-s)q^{m-t-1}$.

We put $S = S_f$. Let $a \in S$ and $F = \{\overrightarrow{ab}, b \in S\}$. We have :

$$\text{Card}(F) = (q-s)q^{m-t-1} \leq q^{rg(F)}.$$

Thus, since $0 \leq s \leq q-2$, we have $rg(F) \geq m-t$.

Let $\overrightarrow{v_1}, \dots, \overrightarrow{v_{m-t}}$ be $m-t$ independent vectors of F and \overrightarrow{u} , such that $\overrightarrow{u} \notin \text{Vect}(\overrightarrow{v_1}, \dots, \overrightarrow{v_{m-t}})$.

Since $t \geq 1$, there exists an affine hyperplane, say H , such that $a + \overrightarrow{v_1}, \dots, a + \overrightarrow{v_{m-t}} \in H$ and $a + \overrightarrow{u} \notin H$.

Assume that $S \cap H \neq S$. Then by lemma 5, either S meets all hyperplanes parallel to H or S meets $(q-s)$ hyperplanes parallel to H in q^{m-t-1} points.

- *1st case* : S meets $(q-s)$ hyperplanes.

By applying an affine transformation, we can assume that the $q-s$ hyperplanes are $H_i = \{x, x_1 = a_i\}$, $a_i \in \mathbb{F}_q$. Without loss of generality, we can assume that $H = H_1$.

$$\text{Let } P = \prod_{i=2}^{q-s} (x_1 - a_i) f(x),$$

$$\deg(P) \leq t(q-1) + s + q - 1 - s = (t+1)(q-1) \text{ and}$$

$|P| = \text{Card}(S \cap \{x_1 = a_1\}) = q^{m-t-1}$. So P is a codeword of minimal weight in $R_q((t+1)(q-1), m)$, and, by recursion hypothesis, $S_p = S \cap H$ is an affine subspace of codimension $t+1$.

- *2nd case* : S meets all the hyperplanes.

For all G_a hyperplane of equation $(z = a)$, $a \in \mathbb{F}_q$, parallel to $H = G_0$,

$g_a = f \cdot (1 - (z - a)^{q-1}) \in R_q((t+1)(q-1) + s, m)$ and $g_a \neq 0$. So
 $\text{Card}(S_{g_a}) \geq (q-s)q^{m-t-2}$.
 Since $\text{Card}(S) = (q-s)q^{m-t-1}$ and $S_{g_a} = S \cap G_a$,
 $\text{Card}(S_{g_a}) = (q-s)q^{m-t-2}$.
 By recursion hypothesis, $S_{g_0} = S \cap H$ is included in an affine subspace of
 codimension $t+1$.

In both cases, $S \cap H$ is included in an affine subspace of codimension $t+1$ which
 is impossible since $a + \vec{v}_1, \dots, a + \vec{v}_{m-t} \in S \cap H$.
 So $S \cap H = S$ and $a + \vec{u} \notin S$, which means that $\vec{u} \notin F$.
 Hence, $F \subset \text{Vect}(\vec{v}_1, \dots, \vec{v}_{m-t})$, i.e. S is included in an affine subspace of codi-
 mension t , say A .

By applying an affine transformation, we can assume that
 $A = \{x, x_1 = 0, \dots, x_t = 0\}$. Then by corollary 3, we can write

$$f(x) = \prod_{i=1}^t (x_i^{q-1} - 1) P(x_{t+1}, \dots, x_m)$$

$P \in R_q(s, m-t)$ and $|P| = |f| = (q-s)q^{m-t-1}$, thus, by the case where $t=0$,
 S_P is the union of $(q-s)$ parallel hyperplanes of A which gives the result.

□

Acknowledgements : I want to thank my supervisor, Jean-François Mestre,
 for his very helpful remarks.

References

- [1] P. Delsarte, J.M. Goethals, F.J. Mac Williams, On generalized Reed-Muller codes and their relatives, *Information and Control*, 16, 403-442 (1970)