

1 Earthworms (*Millsonia anomala*, Megascolecidae) do not increase rice growth
2 through enhanced nitrogen mineralization.

3

4 M. Blouin, S. Barot, P. Lavelle

5

6 *Laboratoire d'Ecologie des Sols Tropicaux (LEST), UMR 137, IRD, Bondy, 93143,*
7 *France.*

8

9 **Abstract**

10

11 Earthworms have been shown to increase plant growth in 75% of the experiments
12 that have compared plant growth in their presence and absence. However, the
13 relative importance of the different mechanisms advanced to explain such a
14 stimulatory effect has never been tested. In a laboratory experiment we observed
15 increased growth of rice plants in the presence of earthworms (*Millsonia anomala*,
16 Megascolecidae) and demonstrated that enhanced nitrogen release (generally
17 considered as the principal mechanism involved in earthworm positive effect on
18 plants) was not responsible for this result: earthworms had the same stimulatory
19 effect on plant growth (+20 %) irrespective of whether the soil (provided with different
20 amounts of mineral-N fertilizer) was either N-limited or N-saturated. We discuss
21 alternative explanations for the observed variations in rice production

22

23

24 *Keywords:* Earthworm; Mechanisms affecting plant growth; Mineralization; Nitrogen
25 gradient

26

27

28 **1. Introduction**

29

30 Among the mechanisms by which earthworms modify plant growth at the
31 individual or community levels (Scheu, 2003; Brown et al., 2004), five have been
32 claimed to be responsible for the positive effect noted on plant production: (i)
33 increased mineralization of soil organic matter, which increases nutrient availability
34 (Barois et al., 1987; Knight et al., 1989; James, 1991; Curry and Byrne, 1992; Lavelle
35 et al., 1992; Subler et al., 1997), especially for nitrogen (N), the major limiting nutrient
36 in terrestrial ecosystems; (ii) modification of soil porosity and aggregation (Blanchart
37 et al., 1999; Shipitalo and Le Bayon, 2004), which induces changes in water and
38 oxygen availability for plants (Doube et al., 1997; Allaire-Leung et al., 2000); (iii)
39 production of plant growth regulators via the stimulation of microbial activity
40 (Frankenberger and Arshad, 1995; Muscolo et al., 1998; Nardi et al., 2002;
41 Quaggiotti et al., 2004); (iv) biocontrol of pests and parasites (Stephens et al., 1994;
42 Clapperton et al., 2001; Blouin et al., 2005); (v) stimulation of symbionts (Gange,
43 1993; Pedersen and Hendriksen, 1993; Furlong et al., 2002).

44

45 Enhanced N mineralization is the best documented mechanism and is generally
46 thought to be the most important. However, despite 313 studies on earthworm effects
47 on plant growth, (Brown et al., 1999; Scheu, 2003), no attempts have been made to
48 assess the relative importance of each of these potential mechanisms (Scheu, 2003;
49 Brown et al., 2004). Here, in a particular experimental case, we evaluate the
50 importance of enhanced mineralization by growing rice fertilised with different levels

51 of mineral N. If enhanced N mineralization is the main mechanism involved in the
52 stimulatory effect of earthworms on plant growth, earthworm effect should be most
53 important when the N availability in soil is low since earthworm activities provide the
54 major amounts of mineral N. Under the same hypothesis, the stimulatory effect of
55 earthworm activities should disappear when the availability of mineral N in the soil is
56 high, the N surplus due to earthworms becoming negligible.

57

58

59 **2. Materials and Methods**

60

61 *2.1 Experimental units*

62

63 Young rice seedlings (*Oryza sativa*, cv. Moroberekan) were grown for three
64 months under a $600 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ artificial light source, at 28 °C/day and 24
65 °C/night temperatures and at 75 % +/- 5 % air moisture. Pots (10 cm in diameter)
66 were filled with 1 kg of a sandy ultisol from Lamto savannah (Ivory Coast). Nitrogen is
67 particularly limiting in the soil of the Lamto savannah : 500 mg kg^{-1} total N (Abbadie
68 and Lensi, 1990; Lensi et al., 1992; Gilot, 1997; Lata et al., 1999), $1.5 \text{ to } 7.5 \text{ mg kg}^{-1}$
69 NH_4^+ and $< 1 \text{ mg kg}^{-1} \text{NO}_3^-$ (Martin, 1990).

70

71 *2.2 Earthworms*

72

73 The original soil fauna was eliminated by sieving (2 mm mesh) and freezing the
74 soil (Blouin et al., 2005). The earthworm *Millsonia anomala*, (Omodeo and Vaillaud,
75 1967), endemic to this region, is a large mesohumic compacting species (Blanchart

76 et al., 1999), whereas the other species, *Chuniodrilus zielae*, (Omodeo and Vaillaud,
77 1967) has a somewhat larger geographical distribution and is a thin polyhumic
78 decompacting worm (Blanchart et al., 1999). These two endogeic earthworms feed
79 on soil organic matter; consequently, the soil was not amended with organic matter.

80 Three immature *M. anomala* (0.85 ± 0.20 g biomass) and three *C. zielae* ($0.07 \pm$
81 0.02 g) were added in each pot according to the treatments. The population density
82 of *M. anomala* was similar to that of natural populations, whereas that of *C. zielae*
83 was slightly lower (Lavelle, 1978). There were four treatments: *M. anomala* alone
84 (M), *C. zielae* alone (Ch), both species present (MCh), and a control without
85 earthworms (C).

86 Some earthworms died during the experiment but the mortality of *M. anomala* and
87 *C. zielae* had no significant effect on total ($P = 0.24$ and 0.14 respectively),
88 aboveground ($P = 0.34$ and 0.12 respectively) or belowground ($P = 0.14$ and 0.21
89 respectively) dry biomasses. The mortality was quite similar in all treatments and not
90 related to the N concentrations.

91

92 2.3 Fertilizer

93

94 Soil mineral N content was increased by watering daily rice plants with a complete
95 fertilizing solution in which all major mineral nutrients and oligo-elements were kept at
96 constant concentrations except N. Since NH_4^+ is the preferred form of nitrogen taken
97 up by rice (Fried et al., 1965; Sasakawa and Yamamoto, 1978), we modified the
98 Hoagland-Arnon solution by replacing $\text{Ca}(\text{NO}_3)_2$ with CaCl_2 and KNO_3 with
99 $\text{NH}_4(\text{SO}_4)_2$ and adding silicium which is necessary for many grasses. Therefore, the
100 composition of the fertilizer was KH_2PO_4 : $2939 \mu\text{mol l}^{-1}$; CaCl_2 : $2495 \mu\text{mol l}^{-1}$; MgSO_4 :

101 3950 $\mu\text{mol l}^{-1}$; Na_2SiO_3 : 996 $\mu\text{mol l}^{-1}$; Fe-EDTA (13 % of Fe): 5 mg l^{-1} ; Oligo-
102 elements : H_3BO_3 : 55 $\mu\text{mol l}^{-1}$; MnSO_4 : 20 $\mu\text{mol l}^{-1}$; ZnSO_4 : 0,6 $\mu\text{mol l}^{-1}$; Na_2MoO_4 : 0,4
103 $\mu\text{mol l}^{-1}$; CuSO_4 : 0,6 $\mu\text{mol l}^{-1}$. NH_4^+ was supplied at five different concentrations (0,
104 25, 100, 400, 1600 $\mu\text{mol l}^{-1}$) in each of the four fauna treatments, resulting altogether
105 in 20 treatments; each was replicated 3 times.

106

107

108 *2.4 The NH_4^+ gradient*

109

110 The test of our hypothesis requires a range of N concentrations, from deficient to
111 excess N availability. To ensure N-limitation for plant production in the 0 $\mu\text{mol l}^{-1}$ NH_4^+
112 treatment, we used a soil that only had 0.05 % total N content (Martin, 1990; Abbadie
113 and Lensi, 1990; Lensi et al., 1992; Gilot, 1997; Lata et al., 1999) and we added all
114 the other macro- and micronutrients at adequate concentrations.

115 NH_4^+ is known to be limiting to rice seedlings growth below 500 $\mu\text{mol l}^{-1}$ in
116 hydroponical conditions (Wang et al., 1993; Kirk, 2001). In our experiments, the 1600
117 treatment showed clear evidence of NH_4^+ toxicity : excessive N concentration in plant
118 tissues (fig. 1 a) and deficit in root production (Britto and Kronzucker, 2002) (fig. 1 b).
119 This argues in favour of an alleviation of the N-limitation in the 1600 $\mu\text{mol l}^{-1}$
120 treatment.

121

122 Fig. 1

123

124

125 *2.4 Statistical analysis*

126

127 Two complementary analyses were conducted on our data set: an ANOVA and
128 an ANCOVA. We checked the homogeneity of variances (Bartlett's test) and
129 normality of the residues (Shapiro-Wilk's test) for total, aboveground and
130 belowground biomasses. Residuals met the conditions of homoscedasticity ($P =$
131 0.43, 0.69, 0.20 respectively) and normality ($P = 0.80, 0.67, 0.22$ respectively). With
132 full ANOVA model, the effects of nitrogen, *M. anomala*, *C. zitelae* and the interactions
133 between these factors were tested (Table 1). In this analysis, nitrogen was
134 considered as a discrete factor. To visualize our results more easily, an ANCOVA
135 was conducted considering nitrogen as a continuous variable. The effect of nitrogen
136 was modelled using a polynom whose order is statistically determined. The
137 significant factor(s) of the ANOVA (earthworm species) may affect significantly some
138 of the parameters of the polynom, modifying the regression equation as compared
139 with the control one.

140

141

142 3. Results

143

144 As expected, the ANOVA showed that the N gradient had a significant effect on
145 plant dry total and above-ground biomasses (Table 1). Comparisons between means
146 (Tukey's test for multiple comparisons) showed that total and above-ground
147 biomasses were not significantly different from the 0 $\mu\text{mol l}^{-1}$ to the 100 $\mu\text{mol l}^{-1}$
148 treatments, but increased significantly from the 100 to the 400 $\mu\text{mol l}^{-1}$ treatments
149 (respectively +18 and 19.5 %) and from the 400 to the 1600 $\mu\text{mol l}^{-1}$ treatments
150 (respectively +19 and 31 %), showing that plant growth was N-limited at the lowest

151 NH_4^+ concentrations. No significant effect of the N gradient was observed on below-
152 ground biomass (Fig. 1c). *M. anomala* had a significant effect on total, above-ground
153 and below-ground biomasses. However, its interaction with the N concentration was
154 not significant (Table 1). The positive effect of *M. anomala* on plant production did not
155 depend on the N concentration. Nor *C. zieleae* neither its interactions with the N
156 concentration or *M. anomala* had a significant effect on plant production (Table 1).

157

158 Table 1

159

160 As *C. zieleae* had no significant effect in the ANOVA, we realize the ANCOVA to
161 test the effect of *M. anomala*, independently of the presence of *C. zieleae* : control (C)
162 and *C. zieleae* treatments, and *M. anomala* and *M. anomala-C. zieleae* treatments
163 respectively were pooled together.

164

165 Table 2

166

167 The relationship between plant total and above-ground biomasses with the N-
168 gradient fitted on a second order polynom ($y = a + bx + cx^2$) (Table 2). Below-ground
169 biomass did not vary significantly along the N-gradient ($y = a$). For total and above-
170 ground biomasses *M. anomala* had a significant positive effect on the parameter a,
171 but not on b or c (Table 2; Fig. 2a and 2b). *M. anomala* also had a significant effect
172 on the parameter a for below-ground biomass (Fig. 2c). Taken together, *M. anomala*
173 had a constant positive effect on plant production along the N-gradient : + 20, 16 and
174 35 % for the total, above-ground and below-ground biomasses respectively.

175

176 Fig. 2

177

178 4. Discussion

179

180 We postulated that if enhanced N mineralization is an important mechanism
181 involved in the positive effect of earthworms on plant growth, earthworms should
182 have a positive effect in a N-limited environment but not in a N-saturated
183 environment, where the enhanced mineralization of N is negligible. Both an ANOVA,
184 taking into account the presence of *C. zitelae*, and an ANCOVA, where the presence
185 of *M. anomala* was considered independently of the presence of *C. zitelae*, showed
186 that the effect of *M. anomala* was constant whatever the N concentrations. Thus, we
187 can reject the hypothesis that the main effect of *M. anomala* on plant production was
188 due to increased N mineralization. If *M. anomala* would increase plant production
189 through an enhanced mineralization of other nutrients (such as P, K... etc), this
190 would have produced no effect on plant in a N-limited environment, but an increase
191 in plant growth in situations where plant is no longer limited by N. This was not
192 observed in our experiment (Fig. 2) and we therefore reject this hypothesis.

193

194 In contrast to *M. anomala*, *C. zitelae* had no significant effect on plant biomass.
195 Only a slight positive effect of *C. zitelae* on root biomass had been observed
196 previously (Derouard et al., 1997). In our experiment the soil was probably not
197 compacted enough to allow the decompacting effect of *C. zitelae* to influence plant
198 growth significantly.

199

200 The increase in plant aboveground biomass simultaneously with the absence of
201 an increase in belowground biomass along the N gradient indicates that the rice
202 plants allocated less resource to the root system as N availability increased
203 (Thornley, 1972; Wilson, 1988; Andrews, 1993). In contrast, the presence of *M.*
204 *anomala* increased both above- and belowground biomasses. Two separate
205 mechanisms were probably responsible for the differing plant responses to the N
206 gradient and to the presence or *M. anomala* (see below).

207 Since *M. anomala* effect was constant over the N-gradient, our experiment does
208 not support the widespread belief that enhanced mineralization by earthworms is the
209 main causal mechanism to explain the increased rice growth in presence of *M.*
210 *anomala*. Nevertheless, this mechanism could be important in other plant-earthworm
211 associations. Moreover, our experiment allows the hypothesis of an enhanced
212 nitrogen mineralization to be excluded at high levels of N, but not at low levels:
213 different mechanisms could in fact be involved at different N concentrations. The
214 mechanism(s) responsible for the positive effect on plant growth at high N
215 concentration cannot be clearly identified from our results. Nevertheless, an
216 improvement of soil water or oxygen availabilities by *M. anomala* can probably be
217 ruled out because soil was maintained at an optimal value of 80 % field capacity by
218 daily watering. Control of specialist parasites or the stimulation of symbionts likely did
219 not occur in our experiment because the soil originated from a non-cultivated
220 savannah with no crop-specific parasites or symbionts, and the original soil fauna
221 had been eliminated. We cannot, however, discard the hypothesis that microbial
222 generalist parasites or symbionts may have been controlled or stimulated by *M.*
223 *anomala*. This leaves the production of plant growth regulators (Frankenberger and
224 Arshad, 1995; Muscolo et al., 1998; Nardi et al., 2002; Quaggiotti et al., 2004) as the

225 probable explanation of the stimulatory effect of *M. anomala* on rice in our
226 experiment. The possibility and the detailed mechanisms of the control of plant
227 physiology via phytohormones secreted into soils by the bacteria associated with
228 earthworms activities should be studied thoroughly.

229

230

231 **Acknowledgements**

232

233 We thank Jérôme Tondoh and Souleymane Konaté for supplying soil and
234 animals, Anne Pando and Jocelyne Roman for their technical help, Daniel Chessel
235 for his advice on statistical analyses and Aram Marks for his help in English
236 improvement.

237

238

239

240

241 **References**

242

243 Abbadie, L. and Lensi, R., 1990. Carbon and nitrogen mineralization and
244 denitrification in a humid savanna of West Africa (Lamto, Côte d'Ivoire). *Acta*
245 *Oecologica* 11, 717–728.

246 Allaire-Leung, S.E., Gupta, S.C. and Moncrief, J.F., 2000. Water and solute
247 movement in soil as influenced by macropore characteristics. 1. Macropore
248 continuity. *Journal of Contaminant Hydrology* 41, 283–301.

- 249 Andrews, M., 1993. Nitrogen effect on the partitioning of dry matter between shoot
250 and root of higher plants. *Current Topics in Plant Physiology* 1, 119-126.
- 251 Barois, I., Verdier, B., Kaiser, P., Mariotti, A., Rangel, P. and Lavelle, P., 1987.
252 Influence of the tropical earthworm *Pontoscolex corethrurus* (Glossoscolecidae)
253 on the fixation and mineralization of nitrogen. In: Bonvicini, A.M. and Omodeo, P.
254 (Eds), *On Earthworms*. Mucchi, Bologna, Italy, pp. 151-158.
- 255 Blanchart, E., Albrecht, A., Alegre, J., Duboisset, A., Gilot, C., Pashanasi, B., Lavelle,
256 P. and Brussaard, L., 1999. Effects of earthworms on soil structure and physical
257 properties. In: Lavelle, P., Brussaard, L. and Hendrix, P. (Eds), *Earthworm*
258 *management in tropical agroecosystems*. CAB International, Wallingford, pp.
259 149–172.
- 260 Blouin, M., Zuily-Fodil, Y., Pham-Thi, A.T., Laffray, D., Reversat, G., Pando, A.,
261 Tondoh, J. and Lavelle, P., 2005. Belowground organism activities affect plant
262 aboveground phenotype, inducing plant tolerance to parasites. *Ecology Letters* 8,
263 202-208.
- 264 Britto, D.T. and Kronzucker, H.J., 2002. NH_4^+ toxicity in higher plants: a critical
265 review. *Journal of Plant Physiology* 159, 567-584.
- 266 Brown, G.G., Edwards, C.A. and Brussaard, L., 2004. How earthworms affect plant
267 growth: burrowing into the mechanisms. In: Edwards, C.A. (Eds), *Earthworm*
268 *ecology*. CRC Press, Boca Raton, USA, pp. 13-49.
- 269 Brown, G.G., Pashanasi, B., Villenave, C., Patron, J.C., Senapati, B.K., Giri, S.,
270 Barois, I., Lavelle, P., Blanchart, E., Blakemore, R.J., Spain, A.V. and Boyer, J.,
271 1999. Effects of earthworms on plant production in the tropics. In: Lavelle, P.,
272 Brussaard, L. and Hendrix, P. (Eds), *The management of earthworms in tropical*
273 *agroecosystems*. CAB International, Wallingford, pp. 87-148.

- 274 Clapperton, M.J., Lee, N.O., Binet, F. and Conner, R.L., 2001. Earthworms indirectly
275 reduce the effects of take-all (*Gaeumannomyces graminis* var. *tritici*) on soft white
276 spring wheat (*Triticum aestivum* cv. Fielder). *Soil Biology and Biochemistry* 33,
277 1531-1538.
- 278 Curry, J.P. and Byrne, D., 1992. The role of earthworms in straw decomposition and
279 nitrogen turnover in arable land in Ireland. *Soil Biology and Biochemistry* 24,
280 1409-1412.
- 281 Derouard, L., Tondoh, J., Vilcosqui, L. and Lavelle, P., 1997. Effects of earthworm
282 introduction on soil processes and plant growth. *Soil Biology and Biochemistry* 29,
283 541-545.
- 284 Doube, B.M., Williams, P.M.L. and Willmott, P.J., 1997. The influence of two species
285 of earthworm (*Aporrectodea trapezoides* and *Aporrectodea rosea*) on the growth
286 of wheat, barley and faba beans in three soil types in the greenhouse. *Soil
287 Biology and Biochemistry* 29, 503-509.
- 288 Frankenberger, W.T. and Arshad, M., 1995. *Phytohormones in soils : Microbial
289 production and function*. Marcel Dekker, New York, 503 pp.
- 290 Fried, M.F., Zsoldos, F., Vose, P.B. and Shatokhin, I.L., 1965. Characterizing the
291 NO_3^- and NH_4^+ uptake process of rice roots by use of ^{15}N labelled NH_4NO_3 .
292 *Physiologia Plantarum* 18, 313-320.
- 293 Furlong, M.A., Singleton, D.R., Coleman, D.C. and Whitman, W.B., 2002. Molecular
294 and culture-based analyses of prokaryotic communities from an agricultural soil
295 and the burrows and casts of the earthworm *Lumbricus rubellus*. *Applied and
296 Environmental Microbiology* 68, 1265-1279.
- 297 Gange, A.C., 1993. Translocation of mycorrhizal fungi by earthworms during early
298 succession. *Soil Biology and Biochemistry* 25, 1021-1026.

- 299 Gilot, C., 1997. Effects of a tropical geophageous earthworm, *Millsonia anomala*
300 (Megascolecidae), on soil characteristics and production of a yam crop in Ivory
301 Coast. *Soil Biology and Biochemistry* 29, 353-359.
- 302 James, S.W., 1991. Soil nitrogen, phosphorus, and organic matter processing by
303 earthworms in tallgrass prairie. *Ecology* 72, 2101-2109.
- 304 Kirk, G.J.D., 2001. Plant-mediated processes to acquire nutrients: nitrogen uptake by
305 rice plants. *Plant and Soil* 232, 129-134.
- 306 Knight, D., Elliott, P.W. and Anderson, J.M., 1989. Effects of earthworms upon
307 transformations and movement of nitrogen from organic matter applied to
308 agricultural soils. In: Hansen, J.A. and Henriksen, K. (Eds), *Nitrogen in organic*
309 *wastes applied to soils*. Academic Press, London, pp. 59-80.
- 310 Lata, J.C., Durand, J., Lensi, R. and Abbadie, L., 1999. Stable coexistence of
311 contrasted nitrification statuses in a wet tropical savanna ecosystem. *Functional*
312 *Ecology* 13, 762–768.
- 313 Lavelle, P., 1978. Les vers de terre de la savane de Lamto (Côte d'Ivoire):
314 peuplements, populations et fonctions dans l'écosystème. Thèse d'état,
315 Laboratoire de zoologie de l'ENS, Paris 6, Paris, pp.
- 316 Lavelle, P., Melendez, G., Pashanasi, B. and Schaefer, R., 1992. Nitrogen
317 mineralization and reorganization in casts of the geophagous tropical earthworm
318 *Pontoscolex corethrurus* (Glossoscolecidae). *Biology and Fertility of Soils* 14, 49-
319 53.
- 320 Lensi, R., Domenach, A.M. and Abbadie, L., 1992. Field study of nitrification and
321 denitrification in a wet savanna of West Africa (Lamto, Côte d'Ivoire). *Plant and*
322 *Soil* 147, 107–113.

- 323 Martin, S., 1990. Modélisation de la dynamique et du rôle d'une population de vers
324 de terre *Millsonia anomala* dans les savanes humides de Côte d'Ivoire. Université
325 Paris 6, Paris, 206 pp.
- 326 Muscolo, A., Cutrupi, S. and Nardi, S., 1998. IAA detection in humic substances. Soil
327 Biology and Biochemistry 30, 1199-1201.
- 328 Nardi, S., Pizzeghello, D., Muscolo, A. and Vianello, A., 2002. Physiological effects of
329 humic substances on higher plants. Soil Biology and Biochemistry 34, 1527-1536.
- 330 Omodeo, P. and Vaillaud, M., 1967. Les Oligochètes de la savane de Gpakobo en
331 Côte d'Ivoire. Bulletin de l'Institut Français d'Afrique Noire 29, 925-944.
- 332 Pedersen, J.C. and Hendriksen, N.B., 1993. Effect of passage through the intestinal
333 tract of detritivore earthworms (*Lumbricus spp.*) on the number of selected Gram-
334 negative and total bacteria. Biology and Fertility of Soils 16, 227-232.
- 335 Quaggiotti, S., Ruperti, B., Pizzeghello, D., Francioso, O., Tugnoli, V. and Nardi, S.,
336 2004. Effect of low molecular size humic substances on nitrate uptake and
337 expression of genes involved in nitrate transport in maize (*Zea mays* L.). Journal
338 of Experimental Botany 55, 803-813.
- 339 Sasakawa, H. and Yamamoto, Y., 1978. Comparison of the uptake of nitrate and
340 ammonium by rice seedlings. Influences of light, temperature, oxygen
341 concentration, exogenous sucrose and metabolic inhibitors. Plant Physiology 62,
342 649-669.
- 343 Scheu, S., 2003. Effects of earthworms on plant growth: patterns and perspectives.
344 Pedobiologia 47, 846-856.
- 345 Shipitalo, M.J. and Le Bayon, R.C., 2004. Quantifying the effects of earthworms on
346 soil aggregation and porosity. In: Edwards, C.A. (Eds), Earthworm ecology. CRC
347 Press, Boca Raton, pp. 441.

- 348 Stephens, P.M., Davoren, C.W., Doube, B.M. and Ryder, M.H., 1994. Ability of the
349 lumbricid earthworms *Aporrectodea rosea* and *Aporrectodea trapezoides* to
350 reduce the severity of take-all under greenhouse and field conditions. *Soil Biology*
351 *and Biochemistry* 26, 1291-1297.
- 352 Subler, S., Baranski, C.M. and Edwards, C.A., 1997. Earthworm additions increased
353 short-term nitrogen availability and leaching in two grain-crop agroecosystems.
354 *Soil Biology and Biochemistry* 29, 413-421.
- 355 Thornley, J.H.M., 1972. A balanced quantitative model for root:shoot ratios in
356 vegetative plants. *Annals of Botany* 68, 211-216.
- 357 Wang, M.Y., Siddiqi, M.Y., Ruth, T.J. and Glass, A.D.M., 1993. Ammonium uptake by
358 rice roots. II. Kinetics of $^{13}\text{NH}_4^+$ influx across the plasmalemma. *Plant Physiology*
359 103, 1259-1267.
- 360 Wilson, J.B., 1988. A review of evidence on the control of shoot:root ratio, in relation
361 to models. *Annals of Botany* 61, 433-449.
- 362

363 Table 1

364 Factors affecting total, shoot and root biomasses from a three-way ANOVA

	df	biomass								
		total			aboveground		belowground			
		F	P		F	P	F	P		
Nitrogen	4	8.5	<0.0001	***	18.3	<0.0001	***	0.8	0.53	
<i>M. anomala</i>	1	9.1	0.004	**	8.4	0.006	**	9.4	0.004	**
<i>C. zielae</i>	1	1.0	0.33		1.1	0.29		0.7	0.39	
Nitrogen * <i>M. anomala</i>	4	0.9	0.50		0.9	0.45		0.7	0.63	
Nitrogen * <i>C. zielae</i>	4	0.9	0.49		0.4	0.78		1.8	0.16	
<i>M. anomala</i> * <i>C. zielae</i>	1	0.2	0.64		0.2	0.64		0.2	0.66	
Nitrogen * <i>M.a.</i> * <i>C.z.</i>	4	1.3	0.28		1.1	0.37		1.7	0.17	

365 ***: p < 0.001, **: p < 0.01, *: p < 0.1, n = 60

366

367 Table 2

368 Polynomial equations determined with the ANCOVA

model: $y = a + bN + cN^2 + dN^3$ coefficients of	df	biomass								
		total			aboveground		belowground			
		F	P		F	P	F	P		
N (b)	1	18.0	<0.0001	***	42.0	<0.0001	***	0.2	0.68	
N ² (c)	1	14.5	0.0004	***	33.1	<0.0001	***	0.2	0.64	
N ³ (d)	1	0.03	0.87		0.98	0.33		1.3	0.25	
<i>M. anomala</i> (a)	4	8.7	0.005	**	8.27	0.006	**	8.5	0.005	**
N * <i>M. anomala</i> (b)	4	0.3	0.57		0.26	0.61		0.4	0.53	
N ² * <i>M. anomala</i> (c)	1	0.001	0.98		0.0004	0.99		0.01	0.91	
N ³ * <i>M. anomala</i> (d)	1	0.8	0.37		1.19	0.28		0.3	0.58	

369 ***: p < 0.001, **: p < 0.01, *: p < 0.1, n = 60

370 Fig. 1

371 Rice responses to the N-gradient according to the presence of earthworms. (a)
372 Percentage of N in plants. (b) Shoot:root ratio of plants. Without lines: control without
373 fauna; Horizontal lines: *M. anomala*; Vertical lines: *C. zitelae*; Horizontal and vertical
374 lines: *C. zitelae* and *M. anomala*. Means \pm s.e., n = 60.

375

376 Fig. 2

377 Response of plant dry biomasses to earthworms along a nitrogen gradient. (a)
378 Total biomass, (b) Shoot biomass, (c) Root biomass. Nitrogen concentration was
379 expressed as $\log(1+[N])$ for more clarity. The polynomial equation of curves was
380 obtained by regression analysis, where the significance of the different coefficients
381 was tested. O: control without fauna; Δ : *C. zitelae*; +: *M. anomala*; x: *C. zitelae* and *M.*
382 *anomala*. The different regression lines are represented only for the significant
383 factors; solid line: without *M. anomala*, dashed line: with *M. anomala*. n = 60.

384

385