



**HAL**  
open science

# Geometric Reproducing Kernels for Signal Reconstruction

Eli Appleboim, Emil Saucan, Yehoshua Y. Zeevi

► **To cite this version:**

Eli Appleboim, Emil Saucan, Yehoshua Y. Zeevi. Geometric Reproducing Kernels for Signal Reconstruction. SAMPTA'09, May 2009, Marseille, France. pp.General session. hal-00453451

**HAL Id: hal-00453451**

**<https://hal.science/hal-00453451>**

Submitted on 4 Feb 2010

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Geometric Reproducing Kernels for Signal Reconstruction

Eli Appleboim <sup>(1)</sup>, Emil Saucan <sup>(2)</sup> and Yehoshua Y. Zeevi <sup>(1)</sup>

(1) Technion, Dept. of Electrical Engineering

(2) Technion, Dept. of Mathematics

eliap@ee.technion.ac.il, semil@ee.technion.ac.il, zeevi@ee.technion.ac.il

## Abstract:

In this paper we propose a smoothing method for non smooth signals, which control the geometry of a sampled signal. The signal is considered as a geometric object and the smoothing is done using a smoothing kernel function that controls the curvature of the obtained smooth signal in a close neighborhood of a metric curvature measure of the original signal.

## 1. Introduction

In [11], [12], a sampling scheme for signals that posses Riemannian geometric structure was introduced. It turns out that a variety of signals fall in this setting while gray scale images is just one such example. Rather than some Nyquist rate, the sampling scheme presented in [11], [12], is based on geometric characteristics of the sampled signals. Being precise, the following sampling theorem was proved.

**Theorem 1** *Let  $\Sigma^n, n \geq 2$  be a connected, not necessarily compact, smooth manifold, with finitely many compact boundary components. Then there exists a sampling scheme of  $\Sigma^n$ , with a proper density  $\mathcal{D} = \mathcal{D}(p) = \mathcal{D}\left(\frac{1}{k(p)}\right)$ , where  $k(p) = \max\{|k_1|, \dots, |k_{2n}|\}$ , and where  $k_1, \dots, k_{2n}$  are the principal (normal) curvatures of  $\Sigma^n$ , at the point  $p \in \Sigma^n$ .*

While the assumed Riemannian structure relies on the assumption that the signal satisfies  $\mathcal{C}^2$  smoothness criteria, the authors presented in [11], an extended version of Theorem 1 also for non smooth geometric signals, where the proposed strategy uses smoothing of the original signal. The following theorem was proved.

**Theorem 2** *Let  $\Sigma$  be a connected, non-necessarily compact surface of class  $\mathcal{C}^0$ . Then, for any  $\delta > 0$ , there exists a  $\delta$ -sampling of  $\Sigma$ , such that if  $\Sigma_\delta \rightarrow \Sigma$ , then  $\mathcal{D}_\delta \rightarrow \mathcal{D}$ , where  $\mathcal{D}_\delta$  and  $\mathcal{D}$  denote the densities of  $\Sigma_\delta$  and  $\Sigma$ , respectively.*

In the above Theorem 2  $\Sigma_\delta$  is a smoothing of  $\Sigma$  obtained by a convolution of  $\Sigma$  with a partition of unity kernel. Such a kernel being very common for manifolds smoothing indeed guarantees that the resultant manifold is as smooth as we wish however, in this process we do not have any control on the curvature of the obtained manifold.

Some natural question raise in this context,

1. To what extent can we smooth the original signal, using such a reproducing kernel while assuming a predefined bounds on the curvature of the resultant manifold?
2. Can the reproducing kernel be made local, namely, can we have different kernel characteristics for different areas along the sampled signals, while being able to glue the smoothed signal along common boundaries?
3. In what way if at any, we can give affirmative answers to 1 and 2 that are adaptive to the signal? Meaning, how can we have good prior estimates for the desired curvature bounds?

This paper aims at answering the above questions. Note that answering question 1 is analogous to smoothen a signal to have a predefined frequency band-pass, using a band-pass filter as commonly done in signal processing for decades. Answering 1, 2, 3 is equivalent to the use of filter banks with different band-pass characteristics. In all, giving affirmative answers to all above questions give rise to an adaptive non uniform sampling scheme for a variety of signals.

We will focus along the paper on signals that are do not admit a Riemannian structure but rather have a more general geometric structure of the so called Alexandrov spaces. We will term such signals as geometric-signals.

## 2. Preliminaries

In this section we will give some basic preliminary definitions and notations.

### 2.1 Alexandrov spaces

**Definition 3 (Alexandrov - Toponogov)** [ [9]] *A complete metric space  $X$ , satisfies the **triangle comparison condition** w.r.t  $\kappa \in \mathbb{R}$  if for every geodesic triangle  $\Delta_{pqr} \in X$ , there exists a **comparison triangle**, i.e. a triangle,  $\Delta_{p'q'r'} \in \mathbb{M}_\kappa^2$ , such that*

$$pq = p'q'; \quad qr = q'r'; \quad rp = r'p'$$

so that, for every point  $s \in pr$  we have that

$$d_X(s, q) > d_{\mathbb{M}_\kappa^2}(s', q')$$

where  $s' \in p'r'$  such that

$$ps = p's'; \quad sr = s'r'$$

Where  $\mathbb{M}_\kappa^2$  is a complete simply connected surface of constant curvature  $\kappa$ .




Figure 1: Comparison triangle.

**Definition 4** A complete metric space  $X$ , is an **Alexandrov space** of curvature  $> \kappa$  iff

1. For all  $x, y, \in X$  there exists a length minimizing curve  $\gamma$  joining  $x$  and  $y$  such that,

$$L(\gamma) = d_X(x, y);$$

where  $L$  denotes the arc length of curves in  $X$  and  $d_X$  stands for the metric given on  $X$ .  $\gamma$  is called a **minimal geodesic**.

2.  $X$  satisfies the triangle comparison condition for  $\kappa$ .
- 3.

$$\dim_H X < \infty;$$

$\dim_H =$  Hausdorff dimension.

**Remark 5** In a similar way, while reversing the direction of inequalities, one can define Alexandrov space of curvature  $< \kappa$ . For instance, in the comparison triangle condition, we will demand,

$$d_X(s, q) < d_{\mathbb{M}_\kappa^2}(s', q')$$

**Definition 6 (Gromov)** If  $X$  is an Alexandrov space of curvature  $< \kappa$  and  $\kappa \leq 0$  then  $X$  is called **CAT( $\kappa$ )-space**. CAT = Cartan-Alexandrov-Toponogov.

### 2.1.1 Examples:

1. Every complete Riemannian manifold of bounded sectional curvature.
2. The boundary of convex set in  $\mathbb{R}^n$  is an Alexandrov space of curvature  $\geq 0$ .
3. If  $X_i$  is a sequence of  $n$ -dimensional Alexandrov spaces of curv.  $\geq \kappa$  then their Gromov-Hausdorff limit, if exists, is an Alexandrov space of curv.  $\geq \kappa$  and dimension  $\leq n$ .

If the limit of the above sequence is of dimension  $< n$  we say the sequence **collapses**.

If  $X$  is an Alexandrov space then there exists a self-adjoint operator  $\Delta$ , called the **Laplacian** defined on  $L^2(X)$  so that,

$$\int_X \langle \nabla u, \nabla v \rangle d\mathcal{H}^n = \int_X v \nabla u d\mathcal{H}^n$$

where  $\mathcal{H}^n$  is the  $n^{\text{th}}$  Hausdorff measure of  $X$ ,  $u \in \mathcal{D}(\Delta)$ ,  $v \in W^{1,2}(X)$ .

**Theorem 7 ([6])** 1. If  $X$  is compact then the spectrum of  $\Delta$  is discrete.

2. There exists a continuous heat kernel  $h_t(x, y)$  on  $X$  so that,

$$e^{-t\Delta}u(x) = \int_X h_t(x, y)u(y)d\mathcal{H}^n(y)$$

## 2.2 Approximations of manifolds

Let  $M$  be a complete Riemannian manifold of bounded sectional curvature. Let  $p \in M$  be some point and let  $\phi_i$  be some  $C^\infty$  kernel function supported on some  $\epsilon_i$ -neighborhood of  $p$ . For example one can take  $\phi$  to be partition of unity, heat kernel and others. Let  $M_i$  be the manifold obtained by convolution,

$$M_i = \int_M \phi_i * M d\mu;$$

Note that  $M_i$  is smooth in a  $\delta_i$  neighborhood of  $p$  even if  $M$  fails to be smooth at  $p$ . Well known results (see for instance, [7]) in differential topology assert that,

$$\epsilon_i \rightarrow 0 \Rightarrow M_j \rightarrow M;$$

where convergence of manifolds is considered in the Gromov-Hausdorff topology. While the above result concerns the convergence on a topological level, in order to have curvature control we have to account for geometric convergence as well. This is guaranteed from the studies in [3], [4] and [10]. In [3], [4] it is proved that similar convergence to the above also exist for Betti numbers which are generalizations of Euler characteristic to all dimensions and are related to curvature through higher dimensional of Gauss-Bonnet type theorems [2]. In [10] the question of proper gluing of approximations in adjacent neighborhoods is addressed. It is shown that one can obtain geometric convergence in different neighborhoods  $V, U$  of the points  $p, q$  resp. so that, on the common boundary  $\partial V \cap \partial U$  the approximations coincide. In addition, if we write the heat operator on a manifold,  $\mathcal{N}$ , as

$$e^{-t\Delta_{\mathcal{N}}}f(x),$$

where  $f \in L^2(\mathcal{N})$  and  $t > 0, x \in \mathcal{N}$ , and  $\Delta_{\mathcal{N}}$ , denotes the Laplace-Beltrami operator associated with  $\mathcal{N}$ , then there is a smooth kernel function  $K_{\mathcal{N}}$ , such that,

$$e^{-t\Delta_{\mathcal{N}}}f(x) = \int_{\mathcal{N}} K_{\mathcal{N}}(t, x, y)f(y)dy;$$

In [3] convergence of the heat kernel is also achieved,

$$e^{-t\Delta_{M_i}} \rightarrow e^{-t\Delta_M}$$

### 3. Smoothing geometric signals with curvature control

In this section we present the results concerning questions 1, 2 and 3 posed in the introduction. These results give us the ability to smoothen a geometric signal while having an adaptive control on obtained curvatures.

**Definition 8** We say that a signal is a **geometric signal** iff it admits a structure of an Alexandrov space for some  $\kappa \in \mathbb{R}$ .

Let  $\Sigma$  be a geometric signal of sectional curvature bounded from below (above). Let  $p \in \Sigma$  be a point, and  $U(p) \subset \Sigma$  some compact neighborhood of  $p$ . Let

$$\kappa = \limsup K$$

such that  $U(p)$  is an Alexandrov space of curvature  $> K$ .

#### 3.1 Approximations of geometric signals

**Theorem 9 ([1])** Given a point  $p$  on  $\Sigma$ , there exists smooth local kernel  $\phi_i$  as above, yielding a sequence of manifolds  $M_i$ , smooth inside an  $\epsilon_i$  neighborhoods of  $p$ , such that

1.

$$M_i = \int_{\Sigma} \phi_i * \Sigma d\mu \rightarrow \Sigma,$$

as  $\epsilon \rightarrow 0$ .

2. If we further assume that while the Riemannian manifolds  $M_i$  converge to  $\Sigma$ , **no collapse occurs** i.e. the Hausdorff dimension of  $\Sigma$  is the same as of  $M_i$ , then, the sectional curvature  $K_i(p)$  of  $M_i$  at  $p$  satisfies,

$$\lim_{\epsilon \rightarrow 0} K_i(p) = \kappa;$$

The theorem above answers both questions 1 and 2. We can control the curvature of the obtained smooth signals in an adaptive way by making it converge to the lim sup of Alexandrov curvature of the signal  $\Sigma$ .

#### 3.2 Gluing

By arguments similar to those in [10] we have,

**Theorem 10 ([1])** Let the above smooth approximations of  $\Sigma$  be given in neighborhoods of two points  $p, q$ . Then they coincide as well as their sectional curvatures  $K_{i,V_i}, K_{i,U_i}$  on the common boundary, if non empty.

### 4. Sampling of geometric signals

We propose the following scheme for sampling of a geometric signals.

1. Consider the signal as an Alexandrov space. This requires the representation of the signal as a tame metric space in a meaningful manner.
2. Assess the appropriate Alexandrov curvature bound. This can be done by the use of discrete metric curvature measures.

3. Smooth the signal while controlling the curvature of the smoothed signal to suitably approximate the estimated curvature.
4. Sample the smoothed signal according to Theorem 1

#### 4.1 Special case - images

It is common to regard images as surfaces embedded in some  $\mathbb{R}^n$ . For gray scale images  $\mathbb{R}^3$  is considered while for color images it is usual to take  $\mathbb{R}^5$ . Figure 2 shows image re-sampled according to the geometric sampling proposed in Theorem1. In this example no smoothing was applied prior to sampling and artifacts of this can be seen in the reconstructed image. ‘‘Flat areas’’ of the image have 20 times reduced sampling resolution with respect to the original resolution.






Figure 2: Geometric sampling of a gray scale image. **Top to bottom** - original Lena; Lena resampled. The white dots are the new sampling points. One can see the sparseness w.r.t the original; Lena reconstructed. Reconstruction using linear interpolation over the sampling points. No smoothing was done.

In order to estimate the curvature of an image as an Alexandrov space we can take the set of discrete curvature measures proposed in [5] where such measures are suggested for very general cell-complexes. It is shown in [5]

that the one-dimensional curvature measure resembles the **Ricci** curvature of a cell-complex which, in the case of images (since they are 2-dimensional manifolds) coincides with the Gaussian curvature. Figure 3 shows the combinatorial Ricci (= Gauss) curvature of the image in Figure 2, see [13] for details about the adoption of the curvature measures introduced in [5] to images.



Figure 3: Discrete Ricci curvature of Lena. Apart from giving an assessment for the curvature of the image as an Alexandrov space, it also serves as an excellent edge detector as itself.

## 5. Further study

Current and future studies of geometric sampling of images and signals, focus on two aspects. First we wish to modify the smoothing process introduced herein so it will be done in the Fourier domain rather than the spatial domain. Namely, we wish to smooth the Fourier transform of the signal while considering curvature in the Fourier plane. This is inspired by the Nash embedding Theorem [8] while the Fourier transform of a manifold is smoothed prior to its embedding thus achieving a higher degree of smoothness with respect to smoothing in the spatial domain.

Another direction of study is devoted to the development of a geometric theory of sparse representations and geometric compress sensing.

## References:

- [1] Appleboim, E., Saucan, E. and Zeevi, Y. Y. *Geometric reproducing kernels for signals*, preprint.
- [2] Bochner, S. and Yano, K., *Curvature and Betti numbers*, Ann. Math. Stud. 32, 1953.
- [3] Cheeger, J. and Gromov, M., *On the characteristic numbers of complete manifolds of bounded curvature and finite volume*, Diff. Geom. and Com. Anal. Chavel Farkas Ed., Springer, 1985.
- [4] Cheeger, J. and Gromov, M., *Bounds on the Von Neumann dimension of  $L^2$ -cohomology and the Gauss-Bonnet theorem for open manifolds*, J. Diff. Geom. 21, 1985.
- [5] Forman, R., *Bochner's method for cell-complexes and combinatorial Ricci curvature*, Disc. Comp. Geom., 29, 2003.
- [6] Kuwae, K. Machigashira, Y. and Shioya, T., *Sobolev spaces, Laplacian and heat kernel on Alexandrov spaces*, Math. Z. 238, 2001.
- [7] Munkres, J. *Elementary Differential Topology*, Ann. Math. Stud. 54, 1966.
- [8] Nash, J., *The Imbedding problem for Riemannian manifolds*, Ann. Math. 63, 1956.
- [9] Otsu, Y. and Shioya, T., *The Riemannian structure of Alexandrov Spaces*, J. Diff. Geom., 39, 1994.
- [10] Petersen, P., Wei, G. and Ye, R., *Controlled geometry via smoothing*, Comm. Math. Helv., 74, 1999.
- [11] Saucan, E., Appleboim, E. and Zeevi, Y. Y. *Sampling and Reconstruction of Surfaces and Higher Dimensional Manifolds*, J. Math. Imaging. Vis., 30, 2008.
- [12] Saucan, E., Appleboim, E. and Zeevi, Y. Y. *Geometric Sampling of Manifolds for Image Representation and Processing* LNCS, 4485, 2007.
- [13] Saucan, E. Appleboim, E., Wolansky G. and Zeevi, Y. Y., *Combinatorial Ricci curvature for image processing*, Midas Jour. Proc. MICCAI 2008