
LONG NON-CROSSING CONFIGURATIONS IN THE PLANE

ADRIAN DUMITRESCU¹ AND CSABA D. TÓTH²

¹ Department of Computer Science, University of Wisconsin–Milwaukee, WI 53201-0784, USA
E-mail address: ad@cs.uwm.edu

² Department of Mathematics and Statistics, University of Calgary, AB, Canada
E-mail address: cdtoth@ucalgary.ca

ABSTRACT. We revisit several maximization problems for geometric networks design under the non-crossing constraint, first studied by Alon, Rajagopalan and Suri (ACM Symposium on Computational Geometry, 1993). Given a set of n points in the plane in general position (no three points collinear), compute a longest non-crossing configuration composed of straight line segments that is: (a) a matching (b) a Hamiltonian path (c) a spanning tree. Here we obtain new results for (b) and (c), as well as for the Hamiltonian cycle problem:

(i) For the longest non-crossing Hamiltonian path problem, we give an approximation algorithm with ratio $\frac{2}{\pi+1} \approx 0.4829$. The previous best ratio, due to Alon et al., was $1/\pi \approx 0.3183$. Moreover, the ratio of our algorithm is close to $2/\pi$ on a relatively broad class of instances: for point sets whose perimeter (or diameter) is much shorter than the maximum length matching. The algorithm runs in $O(n^{7/3} \log n)$ time.

(ii) For the longest non-crossing spanning tree problem, we give an approximation algorithm with ratio 0.502 which runs in $O(n \log n)$ time. The previous ratio, $1/2$, due to Alon et al., was achieved by a quadratic time algorithm. Along the way, we first re-derive the result of Alon et al. with a faster $O(n \log n)$ -time algorithm and a very simple analysis.

(iii) For the longest non-crossing Hamiltonian cycle problem, we give an approximation algorithm whose ratio is close to $2/\pi$ on a relatively broad class of instances: for point sets with the product (diameter \times convex hull size) much smaller than the maximum length matching. The algorithm runs in $O(n^{7/3} \log n)$ time. No previous approximation results were known for this problem.

1. Introduction

Self-crossing in planar configurations is typically an undesirable attribute. Many structures studied in computational geometry, in particular those involving a minimization condition, have the non-crossing attribute for free, for instance minimum spanning trees, minimum length matchings, Voronoi diagrams, etc. The non-crossing property usually follows

1998 ACM Subject Classification: F.2.2 Geometrical problems and computations.

Key words and phrases: Longest non-crossing Hamiltonian path, longest non-crossing Hamiltonian cycle, longest non-crossing spanning tree, approximation algorithm.

Adrian Dumitrescu was supported in part by NSF CAREER grant CCF-0444188. Part of the research by this author was done at Ecole Polytechnique Fédérale de Lausanne. Csaba D. Tóth was supported in part by NSERC grant RGPIN 35586. Part of the research by this author was done at Tufts University.

from the triangle inequality. Alon et al. [3] have considered the problems of computing (i) the longest non-crossing matching, (ii) the longest non-crossing Hamiltonian path and (iii) the longest non-crossing spanning tree, given n points in the plane. Although they were unable to prove it, they suspected that all these problems are NP -hard. The survey articles by Eppstein [8, pp. 439] and Mitchell [14, pp. 680] list these as open problems in the area of geometric network optimization. The problem of approximating the longest non-crossing Hamiltonian cycle is also of interest and wide open [4, pp. 338].

Without the non-crossing condition explicitly enforced, the problem of minimizing or maximizing the length of a spanning tree, Hamiltonian cycle or path, perfect matching, triangulation, etc. has a rich history. However if such structures are required to be non-crossing much less is known, in particular for the maximization variants. While for minimization problems, the non-crossing property comes usually for free via the triangle inequality, in contrast, for maximization problems, the non-crossing property conflicts directly with the length maximizing objective. This is another reason why these problems are interesting to study.

Related work. The existence of non-crossing Hamiltonian paths and cycles in geometric graphs has been studied in [2, 5]. Various Ramsey-type results for non-crossing spanning trees, paths and cycles have been obtained in [11] and [12]. The Euclidean MAX TSP, the problem of computing a longest straight-line tour of a set of points, has been proven NP -hard in dimensions three or higher [9], while its complexity in the Euclidean plane remains open [14]. In contrast, the shortest non-crossing matching and the shortest non-crossing spanning tree are both computable in polynomial time [8, 14], as they coincide with the shortest matching and the shortest spanning tree respectively.

Definitions and notations. A set S of points in the plane is said to be in *general position* if no three points are collinear. General position will be assumed throughout this paper. Given a set of n points in the plane, the results of Alon et al. are as follows: (i) A non-crossing matching whose total length is at least $2/\pi$ of the longest (possibly crossing) matching can be computed in $O(n^{7/3} \log n)$ time. (ii) A non-crossing Hamiltonian path whose total length is at least $1/\pi$ of the longest (possibly crossing) Hamiltonian path can be computed in $O(n^{7/3} \log n)$ time. (iii) A non-crossing spanning tree whose total length is at least $n/(2n-2) \geq 1/2$ of the longest (possibly crossing) spanning tree can be computed in $O(n^2)$ time. The running times have been adjusted to reflect the current best upper bound of $O(n^{4/3})$ on the number of halving lines as established by Dey [6].

A *geometric graph* G is a pair (V, E) where V is a finite set of points in general position in the plane, and E is a set set of segments (edges) connecting points in V . The *length* of G , denoted $L(G)$, is the sum of the Euclidean lengths of all edges in G . The graph G is said to be *non-crossing* if its edges have pairwise disjoint interiors (collinear triples of points are forbidden in order to avoid overlapping collinear edges).

For a point set S , let $\text{conv}(S)$ be the convex hull of S , and let $P = P(S)$ denote the perimeter of $\text{conv}(S)$. Denote by $D = D(S)$ the diameter of S and write $n = |S|$. Let M_{OPT} be a longest (possibly crossing) matching of S , and let M_{OPT}^* be a longest non-crossing matching of S ; observe that for odd n , M_{OPT} is a nearly perfect matching, with $(n-1)/2$ edges. Let H_{OPT} be a longest (possibly crossing) Hamiltonian path of S , and let H_{OPT}^* be a longest non-crossing Hamiltonian path of S . Let T_{OPT} be a longest (possibly crossing) spanning tree of S , and let T_{OPT}^* be a longest non-crossing spanning tree of S . Finally, let Q_{OPT} be a longest (possibly crossing) Hamiltonian cycle of S , and let Q_{OPT}^*

be a longest non-crossing Hamiltonian cycle of S . The following inequalities are obvious: $L(M_{OPT}) \leq L(H_{OPT}) \leq L(T_{OPT})$.

Given a set S of n points in the plane, a line ℓ going through two points of S is called a *halving line* if there are $\lfloor (n-2)/2 \rfloor$ points on one side and $\lceil (n-2)/2 \rceil$ points on the other side [13]. A *bisecting line* ℓ of S is any line that partitions the point set evenly, i. e., neither of the two open halfplanes defined by ℓ contains more than $n/2$ points of S [7]. Observe that any halving line of S is also a bisecting line of S . Any bisecting line of S yields (perhaps non-uniquely) a bipartition $S = R \cup B$, with $R \cap B = \emptyset$, $||R| - |B|| \leq 1$, with R contained in one of the closed halfplanes determined by ℓ , and B contained in the other. We call $S = R \cup B$ a linearly separable bipartition, or balanced partition of S . Observe that for any non-zero direction vector \vec{v} , there is a bisecting line orthogonal to \vec{v} , see [7, Lemma 4.4]. Two bisecting lines are called *equivalent* if they can yield the same balanced partition of S . It is well known that the number of non-equivalent bisecting lines of a set is of the same order as the number of halving lines of the set, and any balanced bipartition can be obtained from a halving line [7, pp. 67].

Our results are summarized in the following three theorems¹.

Theorem 1.1. (i) *For the longest non-crossing Hamiltonian path problem, there is an approximation algorithm with ratio $\frac{2}{\pi+1} \approx 0.4829$ that runs in $O(n^{7/3} \log n)$ time.*

(ii) *Given a set of n points in the plane, one can compute a non-crossing Hamiltonian path H in $O(n^{7/3} \log n)$ time such that $L(H) \geq \frac{2}{\pi} L(H_{OPT}) - \frac{P}{\pi}$. In particular, if the point set satisfies the condition $\frac{P}{\pi} \leq \delta L(H_{OPT})$ for some small $\delta > 0$, then $L(H) \geq (\frac{2}{\pi} - \delta) L(H_{OPT})$.*

(iii) *Alternatively, one can compute a non-crossing Hamiltonian path H in $O(n \log n / \sqrt{\varepsilon})$ time, such that $L(H) \geq (1 - \varepsilon) \frac{2}{\pi} L(H_{OPT}) - \frac{P}{\pi}$.*

Theorem 1.2. *For the longest non-crossing spanning tree problem for a given set of n points in the plane, there is an approximation algorithm with ratio 0.502 and $O(n \log n)$ running time. More precisely, the algorithm computes a non-crossing spanning tree T such that $L(T) \geq 0.502 \cdot L(T_{OPT})$.*

Theorem 1.3. *Given a set S of n points in the plane, with $|\text{conv}(S)| = h$:*

(i) *One can compute a non-crossing Hamiltonian cycle Q in $O(n^{7/3} \log n)$ time such that $L(Q) \geq \frac{2}{\pi} L(Q_{OPT}) - (2h - 1) \frac{P}{\pi}$. In particular, if the point set satisfies the condition $(2h - 1) \frac{P}{\pi} \leq \delta L(Q_{OPT})$ for some small $\delta > 0$, then $L(Q) \geq (\frac{2}{\pi} - \delta) L(Q_{OPT})$.*

(ii) *Alternatively, one can compute a non-crossing Hamiltonian cycle Q in $O(n^3 \log n)$ time such that $L(Q) \geq \frac{2}{\pi} L(Q_{OPT}) - (h + 2) \frac{P}{\pi}$.*

(iii) *Alternatively, one can compute a non-crossing Hamiltonian cycle Q in $O(n \log n / \sqrt{\varepsilon})$ time, such that $L(Q) \geq (1 - \varepsilon) \frac{2}{\pi} L(Q_{OPT}) - (2h - 1) \frac{P}{\pi}$.*

2. The Hamiltonian path

In this section we prove Theorem 1.1. Let $S = \{p_1, \dots, p_n\}$. We follow an approach similar to that of Alon et al. using projections and an averaging argument, in conjunction with a result on bipartite embeddings of spanning paths in the plane. Abellanas et al. [1, Theorem 3.1] showed that every linearly separable bipartition $S = R \cup B$ with $||R| - |B|| \leq 1$, admits an alternating non-crossing spanning path such that the edges cross any separating

¹Due to space limitations, some proofs are omitted.

line ℓ at points ordered monotonically along ℓ . Such a Hamiltonian path can be computed in $O(n \log n)$ time. Their algorithm computes the same Hamiltonian path for any two equivalent halving lines, that is, the alternating path depends on the bipartition only rather than the separating line.

We now recall the algorithm of Abellanas et al. [1]; see Fig. 4 for an example. Let $S = R \cup B$ with $||R| - |B|| \leq 1$ be the red-blue bipartition given by a vertical line ℓ : R on the left, B on the right. Their algorithm constructs an alternating path A in the following way: Let rb be the top red-blue edge of the convex hull $\text{conv}(S)$, called the *top bridge*. If $|R| > |B|$, set $A := \{r\}$, if $|R| < |B|$, set $A := \{b\}$, else set A to $\{r\}$ or $\{b\}$ arbitrarily. At every step, recompute the top bridge rb of $S \setminus A$, and add r to A if the last point in A was blue, or add b to A if the last point in A was red. As pointed out by the authors, the resulting path A is non-crossing because A is disjoint from the convex hull of $S \setminus A$ at each step.

We improve the lower bound of Alon et al. by computing the longest Hamiltonian path corresponding to a bipartition and a Hamiltonian path of length at least the perimeter of the convex hull, and returning the longest of the two.

Lemma 2.1. *For a point set S , $|S| = n \geq 31$, a non-crossing Hamiltonian path of length at least $P(S)$ can be computed in $O(n \log n)$ time. The bound on the length is best possible.*

Consider a geometric graph $G = (V, E)$, and a point $q \notin V$, so that $V \cup \{q\}$ is in general position. We say that q sees a vertex $v \in V$ if the segment qv does not intersect any edge of G . Similarly, we say that q sees an edge $e \in E$, if the triangle formed by v and e does not intersect any other edge of G . We make use of the fact that if n is even then the two endpoints of an alternating path are on opposite sides of the separating line ℓ . If n is odd, we first construct an alternating path for a specific subset of $n - 1$ points, and then augment it to a Hamiltonian path on all n points using the following lemma.

Lemma 2.2. *Let $S = R \cup B$ with $||R| - |B|| \leq 1$, be a linearly separable bipartition given by line ℓ . Let $q \in S$, and A' be a non-crossing alternating path on $S \setminus \{q\}$ such that its (consecutive) edges cross ℓ at points ordered monotonically along ℓ . Then q sees one edge of A' and consequently, A' can be extended to a Hamiltonian path A on S , with $L(A') < L(A)$. The path A can be computed in $O(n)$ time, given A' .*

Fix a Cartesian coordinate system Γ . Let k be the number of halving lines of S , denote the angles they make with the x -axis of Γ by $0 \leq \alpha_1 < \dots < \alpha_k < \pi$. By relabeling the points assume that the optimal path is $H_{\text{OPT}} = p_1, p_2, \dots, p_n$. For two points $p_i, p_j \in S$, let β_{ij} be the angle in $[0, \pi)$ formed by the line through $p_i p_j$ and the x -axis. If n is odd, then a bisecting line of direction α (for any α) must be incident to at least one point of S , and denote an arbitrary such point by q_α .

Algorithm A1:

STEP 1. Compute a non-crossing Hamiltonian path H_1 of length at least $P(S)$, by Lemma 2.1.

STEP 2. If n is even, then for all non-equivalent bisections of S (i.e., for all balanced bipartitions of S), compute a non-crossing alternating path using the algorithm of Abellanas et al. [1], and let the longest such path be H_2 . If n is odd, then for all non-equivalent bisections of S , compute a non-crossing alternating path of the even point set $S \setminus \{q_\alpha\}$ using the algorithm of [1] and let the longest such path be H'_2 . Augment H'_2 with vertex q_α by Lemma 2.2 to a Hamiltonian path H_2 .

STEP 3. Output the longest of the two paths H_1 and H_2 .

By Lemma 2.1, the running time of STEP 1 is $O(n \log n)$. Since the number of halving lines of an n -element point set is $O(n^{4/3})$ and all can be generated within this time [6], the running time of STEP 2 is $O(n^{7/3} \log n)$, consequently the total running time of **A1** is also $O(n^{7/3} \log n)$.

We proceed with the analysis of the approximation ratio. For simplicity, we assume that n is even. The case of n odd is slightly different. For each $\alpha \in [0, \pi)$, let Γ_α be a (rotated) coordinate system, obtained from Γ via a counterclockwise rotation by α , and with the y -axis dividing evenly the point set S . Let x_i be the x -coordinate of point p_i with respect to Γ_α . For a given α , let H_α be a non-crossing alternating path with respect to a balanced bipartition induced by the y -axis of Γ_α , as computed by the algorithm. There are $O(1)$ balanced bipartitions given by any halving line of S . Recall that H_α does not depend continuously on α ; it depends only on the discrete bipartition. However, the coordinates of the points depend continuously on α . Assume that $H_\alpha = p_{\sigma(1)}, p_{\sigma(2)}, \dots, p_{\sigma(n)}$, where σ is a permutation of $[n]$; here σ depends on the bipartition (hence also on α). Let W_α denote the *width* of S in direction α , that is, the width of the smallest parallel strip of direction α that contains S . By projecting on the x -axis of Γ_α , we get

$$\begin{aligned} L(H_\alpha) &\geq |x_{\sigma(1)}| + 2|x_{\sigma(2)}| + \dots + 2|x_{\sigma(n-1)}| + |x_{\sigma(n)}| = 2 \sum_{i=1}^n |x_i| - |x_{\sigma(1)}| - |x_{\sigma(n)}| \\ &= \sum_{j=1}^{n-1} (|x_j| + |x_{j+1}|) + |x_1| + |x_n| - |x_{\sigma(1)}| - |x_{\sigma(n)}| \geq \sum_{j=1}^{n-1} (|x_j| + |x_{j+1}|) - W_\alpha \\ &\geq \sum_{j=1}^{n-1} |p_j p_{j+1}| |\cos(\beta_{jj+1} - \alpha)| - W_\alpha \end{aligned} \quad (2.1)$$

In the 2nd line of the above chain of inequalities, we use the fact that $p_{\sigma(1)}$ and $p_{\sigma(n)}$ lie on opposite sides of ℓ , since n is even, hence $|x_{\sigma(1)}| + |x_{\sigma(n)}| \leq |p_{\sigma(1)} p_{\sigma(n)}| \leq W_\alpha$. In the 3rd line, we make use of the following inequality: for any two points $p_i, p_j \in S$, $|p_i p_j| |\cos(\beta_{ij} - \alpha)| \leq |x_i| + |x_j|$, with equality if and only if the two points lie on opposite sides of the y -axis of Γ_α . Recall: for even n , H_2 is the longest of the $O(k)$ Hamiltonian non-crossing paths H_{α_i} over all $O(k)$ balanced bipartitions of S . (A given angle α_i yields $O(1)$ balanced partitions, and corresponding alternating paths denoted here H_{α_i} .) We thus have for each $\alpha \in [0, \pi)$:

$$L(H_2) \geq \sum_{j=1}^{n-1} |p_j p_{j+1}| |\cos(\beta_{jj+1} - \alpha)| - W_\alpha.$$

Note that

$$\int_0^\pi |\cos(\beta_{jj+1} - \alpha)| \, d\alpha = \int_0^\pi |\cos \alpha| \, d\alpha = 2,$$

and according to Cauchy's surface area formula, we have $\int_0^\pi W_\alpha \, d\alpha = P(S)$. By integrating both sides of the previous inequality over the α -interval $[0, \pi]$, we obtain

$$\begin{aligned} \pi L(H_2) &\geq 2 \sum_{j=1}^{n-1} |p_j p_{j+1}| - P(S) = 2L(H_{\text{OPT}}) - P(S), \\ L(H_2) &\geq \frac{2}{\pi} L(H_{\text{OPT}}) - \frac{P(S)}{\pi}. \end{aligned} \quad (2.2)$$

We now improve the old approximation ratio of $\frac{1}{\pi} \approx 0.3183$ to $\frac{2}{\pi+1} \approx 0.4829$, by balancing the lengths of the two paths computed in STEP 1 and STEP 2. Set $c = \frac{\pi+1}{2}$.

Case 1: $L(H_{\text{OPT}}) \leq cP(S)$. By considering the path computed in STEP 1, we get a ratio of at least

$$\frac{L(H_1)}{L(H_{\text{OPT}})} \geq \frac{P(S)}{L(H_{\text{OPT}})} \geq \frac{P(S)}{cP(S)} = \frac{2}{\pi+1}.$$

Case 2: $L(H_{\text{OPT}}) \geq cP(S)$. By considering the path computed in STEP 2 (inequality (2.2)), we get a ratio of at least

$$\frac{L(H_2)}{L(H_{\text{OPT}})} \geq \frac{\frac{2}{\pi}L(H_{\text{OPT}}) - \frac{1}{\pi}P(S)}{L(H_{\text{OPT}})} \geq \frac{2}{\pi} - \frac{1}{c\pi} = \frac{2}{\pi} \left(1 - \frac{1}{\pi+1}\right) = \frac{2}{\pi+1}.$$

Observe that if the point set satisfies the condition $\frac{P(S)}{\pi} \leq \delta L(H_{\text{OPT}})$, then by (2.2), we have

$$L(H) \geq \frac{2}{\pi}L(H_{\text{OPT}}) - \delta L(H_{\text{OPT}}) = \left(\frac{2}{\pi} - \delta\right)L(H_{\text{OPT}}).$$

This concludes the proofs of parts (i) and (ii) of Theorem 1.1.

(iii) With the same approach as in [3], a Hamiltonian path of length at least $(1 - \varepsilon)\frac{2}{\pi}L(H_{\text{OPT}}) - \frac{P(S)}{\pi}$ can be found by considering only $b/\sqrt{\varepsilon}$ angles $\theta_i = \frac{i\pi\sqrt{\varepsilon}}{b}$, for $i = 0, 1, \dots, \lfloor b/\sqrt{\varepsilon} \rfloor$, where b is a suitable absolute constant. The resulting running time is $O(n \log n/\sqrt{\varepsilon})$. This concludes the proof of Theorem 1.1.

3. The spanning tree

In this section we prove Theorem 1.2. Let $S = \{p_1, \dots, p_n\}$, where $p_i = (x_i, y_i)$. Given a point $p \in S$, the *star centered at p* , denoted S_p , is the spanning tree on S whose edges join p to all the other points. Since S is in general position, S_p is non-crossing for any $p \in S$. An *extended star centered at p* is a spanning tree of S consisting of paths of length 1 or 2 (edges) connecting p to all the other points. See Fig. 1. While the star centered at a point is unique, there may be many extended stars centered at the same point, and some of them may be self-crossing. In particular S_p is also an extended star.




Figure 1: A star (left) and a non-crossing extended star (right) on a same point set, both centered at the same point p .

The algorithm of Alon et al. computes the n stars centered at each of the points, and then outputs the longest one. The algorithm takes quadratic time, and the analysis shows a ratio of $\frac{n}{2n-2}$ (which tends to $1/2$ in the limit). Their algorithm works in any metric space.

As pointed out by Alon et al., the ratio $1/2$ is best possible (in the limit) for this specific algorithm. We first re-establish the $1/2$ approximation ratio using a faster algorithm, and also with a simpler analysis. Our algorithm works also in any metric space; however in this general setting, the running time remains quadratic.

Algorithm A2: Compute a diameter of the point set, and output the longest of the two stars centered at one of its endpoints.

Obviously the algorithm runs in $O(n \log n)$ time, with bottleneck being the diameter computation [15]. Let ab be a diameter pair, and assume w.l.o.g. that $|ab| = 1$. The ratio $1/2$ (or even $\frac{n}{2n-2}$) follows from the next lemma in conjunction with the obvious upper bound $L(T_{\text{OPT}}) \leq n$ (or $L(T_{\text{OPT}}) \leq n - 1$).

Lemma 3.1. *Let S_a and S_b be the stars centered at the points a and b , respectively. Then $L(S_a) + L(S_b) \geq n$.*

Proof. Assume that $a = p_1$, $b = p_2$. For each $i = 3, \dots, n$, the triangle inequality for the triple a, b, p_i gives

$$|ap_i| + |bp_i| \geq |ab| = 1.$$

By summing up we have

$$L(S_a) + L(S_b) = \sum_{i=3}^n (|ap_i| + |bp_i|) + 2|ab| \geq (n-2) + 2 = n. \quad \blacksquare$$

We now continue with the new algorithm that achieves a (provable) $\frac{1}{2} + \frac{1}{500}$ approximation ratio within the same running time $O(n \log n)$.

Algorithm A3: Compute a diameter ab of the point set, and output the longest of the 5 non-crossing structures S_a, S_b, S_h, E_a, E_b , described below.

Assume w.l.o.g. that the ab is a horizontal unit segment, where $a = (0, 0)$ and $b = (1, 0)$. Let $h = (x_h, y_h)$ be a point in S with a largest value of $|y|$. By symmetry, we can assume that $y_h \geq 0$. S_a, S_b , and S_h are the 3 stars centered at a, b , and h respectively. E_a , resp. E_b , are two non-crossing extended stars centered at a , resp. b ; details to follow. Each of the five structures can be computed in $O(n \log n)$ time, so the total execution time is also $O(n \log n)$.

Set $\delta = 0.05$, $w = 0.6$, $t = 0.6$ and $z = 0.48$, and refer to Fig. 2. Let ℓ_1, ℓ_2, ℓ_3 , and ℓ_4 , be four parallel vertical lines: $\ell_1 : x = 0$, $\ell_2 : x = 0.2$, $\ell_3 : x = 0.8$, $\ell_4 : x = 1$. Obviously, all points in S lie in the strip bounded by ℓ_1 and ℓ_4 . Let V_m be the vertical parallel strip symmetric about the midpoint of ab and of width w . We refer to V_m as the middle strip; V_m is bounded by the vertical lines ℓ_2 and ℓ_3 . Let V_a and V_b be the two vertical strips of width 0.2 bounded by ℓ_1 and ℓ_2 , and by ℓ_3 and ℓ_4 respectively. Let $c = (x_c, y_c)$ be the intersection point between ℓ_3 and the circular arc γ_a of unit radius centered at a and subtending an angle of 60° . We have $x_c = 0.8$ and

$$y_c = \sqrt{1 - 0.8^2} = 0.6 = t.$$

We now describe the two extended star structures E_a and E_b . See also Fig. 3 for an example. To construct E_a , first compute the order of visibility of the points in V_b from point a by sorting. Then connect a with each point in the right strip V_b . Note that $b \in V_b$, thus $V_b \neq \emptyset$. Call S'_a the resulting star. The edges of this star together with the vertical line ℓ_3 divide $V_a \cup V_m$ into convex regions (wedges with a common apex a) ordered top-down. The subset of points in each wedge can be computed using binary search in overall $O(n \log n)$




Figure 2: A diameter pair a, b at unit distance, and the three vertical strips V_a, V_m , and V_b . The two circular arcs γ_a and γ_b of unit radius centered at a and b intersect at the point $(1/2, \sqrt{3}/2)$. All points of S above ab lie in the region bounded by ab, γ_a and γ_b .

time (over all wedges). S'_a is extended (augmented) as follows. In each wedge, say paq , all points are connected either to a or to p , depending on the best (longest) overall connection cost. We denote the resulting *extended star* structure by E_a . The construction of E_b is analogous. It is clear by construction that both E_a and E_b are non-crossing.

Lemma 3.2. *For each $p \in S$, let $d_{\max}(p)$ denote the maximum distance from p to other points in S . Then*

$$L(T_{OPT}) \leq \left[\sum_{i=1}^n d_{\max}(p_i) \right] - 1.$$

Proof. Consider T_{OPT} rooted at a and drawn as an abstract tree with the root at the top in the usual manner. Let $\pi(v)$ denote the parent of a (non-root) vertex v . Uniquely assign each edge $\pi(v)v$ of T_{OPT} to vertex v . Obviously, $L(\pi(v)v) \leq d_{\max}(v)$ holds for each edge in the tree. By adding up the above inequalities, and taking into account that $d_{\max}(a) = |ab| = 1$, the lemma follows. ■

Lemma 3.3. *Assume that $\sum_{i=1}^n |y_i| \geq \delta n$ for some positive constant $\delta \leq 1$. Then*

$$L(S_a) + L(S_b) \geq 2n\sqrt{\frac{1}{4} + \delta^2}.$$

Lemma 3.4. *Let n_a and n_b denote the number of points in the left and right vertical strips V_a and V_b . Then $L(E_a) \geq \frac{1+w}{4}(n + n_b)$, and similarly $L(E_b) \geq \frac{1+w}{4}(n + n_a)$. Consequently $L(E_a) + L(E_b) \geq \frac{1+w}{4}(2n + n_a + n_b)$. E_a and E_b can be constructed in $O(n \log n)$ time.*

Proof. The distance between ℓ_1 and ℓ_3 is $\frac{1+w}{2}$. By an argument similar to that in the proof of Lemma 3.1, the connection cost for a wedge with m points is at least $\frac{1+w}{4}m$. Therefore the total length of E_a is

$$L(E_a) \geq \frac{1+w}{2}n_b + \frac{1+w}{4}(n - n_b) = \frac{1+w}{4}(n + n_b).$$

The estimation of $L(E_b)$ is analogous. The running time has been established previously. ■

Lemma 3.5. *Assume that $\sum_{i=1}^n |y_i| \leq \delta n$ and $y_h \geq t$. Then $L(S_h) \geq (t - \delta)n$.*

Proof.

$$L(S_h) \geq \sum_{i=1}^n (y_h - y_i) = ny_h - \sum_{i=1}^n y_i \geq ny_h - \sum_{i=1}^n |y_i| \geq ny_h - \delta n \geq (t - \delta)n. \quad \blacksquare$$

Lemma 3.6. *Assume that $|y_h| \leq t = 0.6$. Let $p \in S$ be a point in the middle strip V_m , with y -coordinate satisfying $|y| \leq 0.15$. Then $d_{\max}(p) \leq 0.9605$.*

Proof. It is straightforward to check that the maximum distance is attained for a point p on ℓ_2 with y -coordinate -0.15 . The furthest point from p in the allowed region is c . Hence

$$d_{\max}(p) \leq |pc| = \sqrt{w^2 + (0.15 + t)^2} = \sqrt{0.6^2 + 0.75^2} \leq 0.9605. \quad \blacksquare$$

We now distinguish the following four cases to complete our estimation of the approximation ratio.

Case 1: $\sum_{i=1}^n |y_i| \geq \delta n$. The algorithm outputs² S_a or S_b . By Lemma 3.3, the approximation ratio is at least

$$\frac{L(S_a) + L(S_b)}{2L(T_{\text{OPT}})} \geq \sqrt{\frac{1}{4} + \delta^2} \geq 0.502.$$

Case 2: $\sum_{i=1}^n |y_i| \leq \delta n$ and $y_h \geq t$. The algorithm outputs S_h . By Lemma 3.5, the approximation ratio is at least $t - \delta = 0.55$.

Case 3: $\sum_{i=1}^n |y_i| \leq \delta n$ and $y_h \leq t$ and $n_a + n_b \geq (1 - z)n$. The algorithm outputs E_a or E_b . We only need the last inequality in estimating the length. By Lemma 3.4, the approximation ratio is at least

$$\frac{L(E_a) + L(E_b)}{2L(T_{\text{OPT}})} \geq \frac{1 + w}{4} \cdot \frac{2n + n_a + n_b}{2n} \geq \frac{(1 + w)(3 - z)}{8} = \frac{1.6 \cdot 2.52}{8} = 0.504.$$

Case 4: $\sum_{i=1}^n |y_i| \leq \delta n$ and $y_h \leq t$ and $n_a + n_b \leq (1 - z)n$. The algorithm outputs S_a or S_b . There are at least $zn = 0.48n$ points in the middle strip V_m . Observe that at most $n/3$ points in V_m have $|y_i| \geq 0.15$; otherwise we would have

$$\sum_{i=1}^n |y_i| \geq \sum_{V_m} |y_i| > 0.15 \cdot \frac{n}{3} = 0.05n = \delta n,$$

a contradiction. It follows that at least $12n/25 - n/3 = 11n/75$ points in the middle strip have $|y_i| \leq 0.15$. By Lemma 3.2 and Lemma 3.6,

$$L(T_{\text{OPT}}) \leq \frac{64n}{75} + 0.9605 \cdot \frac{11n}{75} \leq 0.9943n.$$

The approximation ratio is at least

$$\frac{L(S_a) + L(S_b)}{2L(T_{\text{OPT}})} \geq \frac{n}{2 \cdot 0.9943n} \geq 0.502.$$

This completes the list of cases and thereby the proof of Theorem 1.2.

Remark. The example in Fig. 3 with n points (n even) equally spaced along a circle shows that the constant 0.502 measuring the approximation ratio achieved by our algorithm **A3**

²Here and in other instances it is meant that the algorithm outputs a structure at least as long as these.

cannot be improved to anything larger than $2/\pi$. Indeed the lengths of the five structures computed by the algorithm are $L(S_a) = L(S_b) = L(S_h) = L(E_a) = L(E_b) = (1 - o(1))\frac{2}{\pi}n$, while $L(T_{\text{OPT}}) \geq L(H_{\text{OPT}}) = (1 - o(1))n$.




Figure 3: The non-crossing structure E_a for an example with $n = 16$ points on the circle. The middle strip V_m is bounded by the two dashed vertical lines.

4. The Hamiltonian cycle

In this section we present the proof of Theorem 1.3, which is similar (including notation) to that of Theorem 1.1. The rotated coordinate system Γ_α , and the x -coordinates x_i with respect to this system are denoted in the same way. By relabeling the points assume that the optimal cycle is $Q_{\text{OPT}} = p_1, p_2, \dots, p_n$ (with the convention that $p_{n+1} = p_1$). We approximate Q_{OPT} by constructing a non-crossing alternating path A on a subset of S , and then completing it to a non-crossing cycle using convex hull vertices. We need to observe that the alternating path A on the subset I of interior (non-hull) vertices of S produced by the algorithm of Abellanas et al. [1] is *not* good enough for *this* strategy: even though one endpoint of A (the first computed by the algorithm) is always on the convex hull of I , the other endpoint might be blocked by edges of A , so that A might not be extendible to a non-crossing Hamiltonian cycle (an example is shown in Fig. 4). Here, we give a stronger result that fits our purpose (for an even number of points).

Lemma 4.1. *Let $S = R \cup B$ with $|R| = |B|$, be a linearly separable bipartition given by line ℓ . Then S admits an alternating non-crossing spanning path A such that (1) the edges of A cross ℓ at points ordered monotonically along ℓ ; and (2) the two endpoints of A are incident to the two distinct edges of the convex hull that connect R and B (the two red-blue bridges). Such a Hamiltonian path can be computed in $O(n \log n)$ time. We refer to the underlying procedure as the two-endpoint path construction algorithm.*

Proof. We modify the algorithm of Abellanas et al. for path construction, so that the path is grown from the two endpoints and the two sub-paths merge "in the middle". Recall that $S = R \cup B$, and $|R| = |B|$, thus $|S|$ is even. Let $r_1 b_1$ and $r_2 b_2$ be the top and bottom red-blue edges of the convex hull $\text{conv}(S)$, respectively, called *top* and *bottom* bridges; it is possible that $r_1 = r_2$ or $b_1 = b_2$ but not both. One endpoint of A is an endpoint of the top bridge, and the other endpoint of A is an endpoint of the bottom bridge, and they




Figure 4: A non-crossing alternating path obtained by the algorithm of Abellanas et al. For the purpose of cycle construction, the path is non-extendible from its 2nd endpoint, vertex 7.

are chosen of opposite colors. Let $A = \{r_1, b_2\}$ or $A = \{b_1, r_2\}$ arbitrarily, containing two endpoints of the path. At every step, recompute the top and bottom bridges of $S \setminus A$, and append either the red or the blue vertex of each bridge to A such that the appended edges cross the separating line ℓ . In the last step, the convex hull of $S \setminus A$ is a red-blue segment that merges the two sub-paths. The two new edges added simultaneously at each step cannot cross each other; and they cannot cross previous edges, since they are separated from them by the convex hull of $S \setminus A$. Finally, they cannot extend the two sub-paths by the same point either, because $|S|$ is even. ■

The next lemma follows from [10, Lemma 2.1]; we will only need its corollary, Lemma 4.3.

Lemma 4.2. ([10]). *Let $P = p_1, p_2, \dots, p_n$ be a simple polygon (with the convention that $p_{n+1} = p_1$) and q be a point in the exterior of the convex hull of P , where $P \cup \{q\}$ is in general position. Then q sees one edge $p_i p_{i+1}$ of P . Such an edge can be found in $O(n)$ time.*

Lemma 4.3. *Let $P = p_1, p_2, \dots, p_n$ be a simple polygon (with the convention that $p_{n+1} = p_1$) and q be a point in the exterior of the convex hull of P , where $P \cup \{q\}$ is in general position. Then the polygonal cycle P can be extended to include q so that $P \cup \{q\}$ is still a simple polygon. More precisely, there exists $i \in [n]$, so that $Q = p_1, \dots, p_i, q, p_{i+1}, \dots, p_n$ is a simple polygon. Moreover, $L(Q) > L(P)$. The extension can be computed in $O(n)$ time.*

Proof. By Lemma 4.2, q sees one edge $p_i p_{i+1}$ of P . Replacing this edge of P by the two edges $p_i q$ and $q p_{i+1}$ results in a simple polygon $Q = p_1, \dots, p_i, q, p_{i+1}, \dots, p_n$. By the triangle inequality, $L(Q) > L(P)$. The extension can be computed in $O(n)$ time, as determined by the time needed to find a visible edge. ■

Note that the condition in the lemma that q lies in the exterior of the convex hull of P , is indeed necessary. Otherwise one cannot guarantee that q sees an edge of P .

(i) Let $S = S' \cup S''$, where S' is the set of convex hull vertices and S'' is the set of interior points. Let $S' = \{p_{j_1}, p_{j_2}, \dots, p_{j_h}\}$. Put $h = |S'|$, $m = |S''|$, thus $n = h + m$. Assume first for simplicity that m is even. An easy modification of the algorithm, explained below, is used for m odd.

Algorithm A4:

STEP 1. For all non-equivalent bisections of S'' (i.e., for all balanced bipartitions of S''):
 1. Compute a non-crossing alternating path A by using the two-endpoint path construction algorithm (Lemma 4.1). 2. Extend A to a cycle by connecting its endpoints to (one or two) convex hull vertices. 3. Further extend this cycle to include the remaining hull vertices, by repeated invocation of Lemma 4.3.

STEP 2. Output the longest such cycle (containing all points of S).

Observe that after STEP 1.1, the two endpoints of the path are vertices of $\text{conv}(S'')$, hence they can be connected to hull vertices to make a cycle. If m is odd, then there is a point $q \in S''$ on the line ℓ . Use the two-endpoint path construction algorithm for $S'' \setminus \{q\}$, and the same bisecting line ℓ . If q is in the interior of $\text{conv}(S'' \setminus \{q\})$, then extend the path with point q , using Lemma 2.2. Otherwise, q sees the top or bottom bridge of $\text{conv}(S'' \setminus \{q\})$, so the path can be extended by connecting q to the endpoint visible to q . The two endpoints of the extended path are on $\text{conv}(S'')$, hence they can be connected to hull vertices to make a cycle, as in the case of even m .

References

- [1] M. Abellanas, J. Garcia, G. Hernández, M. Noy, and P. Ramos: Bipartite embeddings of trees in the plane, *Discrete Applied Mathematics*, **93** (1999), 141–148.
- [2] O. Aichholzer, S. Cabello, R. Fabila-Monroy, D. Flores-Peñaloza, T. Hackl, C. Huemer, F. Hurtado, and D. R. Wood: Edge-removal and non-crossing configurations in geometric graphs, *Proc. 24th European Workshop on Computational Geometry*, Nancy 2008, pp. 119–122.
- [3] N. Alon, S. Rajagopalan and S. Suri: Long non-crossing configurations in the plane, *Fundamenta Informaticae* **22** (1995), 385–394. Also in *Proc. 9th ACM Sympos. on Comput. Geom.*, 1993, 257–263.
- [4] M. Bern and D. Eppstein: Approximation algorithms for geometric problems, in *Approximation Algorithms for NP-hard Problems* (D. S. Hochbaum, editor), PWS, Boston, 1997, pp. 296–345.
- [5] J. Černý, Z. Dvořák, V. Jelínek, and J. Kára: Noncrossing Hamiltonian paths in geometric graphs, *Discrete Applied Mathematics*, **155** (2007), 1096–1105.
- [6] T. K. Dey: Improved bounds on planar k -sets and related problems, *Discrete & Computational Geometry*, **19** (1998), 373–382.
- [7] H. Edelsbrunner: *Algorithms in Combinatorial Geometry*, Springer-Verlag, Heidelberg, 1987.
- [8] D. Eppstein: Spanning trees and spanners, in *Handbook of Computational Geometry* (J.-R. Sack and J. Urrutia, editors), Elsevier Science, Amsterdam, 2000, pp. 425–461.
- [9] S. P. Fekete: Simplicity and hardness of the maximum traveling salesman problem under geometric distances, *Proceedings of the 10th ACM-SIAM Symposium on Discrete Algorithms*, 1999, pp. 337–345.
- [10] F. Hurtado, M. Kano, D. Rappaport, and Cs. D. Tóth: Encompassing colored planar straight line graphs, *Computational Geometry: Theory and Applications*, **39** (1) (2008), 14–23.
- [11] G. Károlyi, J. Pach and G. Tóth: Ramsey-type results for geometric graphs. I, *Discrete and Computational Geometry* **18** (1997), 247–255.
- [12] G. Károlyi, J. Pach, G. Tóth and P. Valtr: Ramsey-type results for geometric graphs. II, *Discrete and Computational Geometry*, **20** (1998), 375–388.
- [13] L. Lovász: On the number of halving lines, *Ann. Univ. Sci. Budapest, Eötvös, Sec. Math.*, **14** (1971), 107–108.
- [14] J. S. B. Mitchell: Geometric shortest paths and network optimization, in *Handbook of Computational Geometry* (J.-R. Sack and J. Urrutia, editors), Elsevier Science, Amsterdam, 2000, pp. 633–701.
- [15] F. Preparata and M. Shamos: *Computational Geometry: An Introduction*, Springer, New York, 1985.