

**Validation study of a French version of the modified Telephone Interview for Cognitive Status (F-TICS-m) in elderly women**

*Running Title: Validation of TICS-m in French*

*Keywords: TICS-m; telephone screening; cognitive impairment; dementia; MMSE; validation; elderly*

Marie-Noël Vercambre<sup>1</sup>, Hélène Cuvelier<sup>1</sup>, Yasemin Atakuman Gayon<sup>1</sup>, Isabelle Hardy-Léger<sup>2</sup>, Claudine Berr<sup>3</sup>, Christophe Trivalle<sup>4</sup>, Marie-Christine Boutron-Ruault<sup>1</sup>, Françoise Clavel-Chapelon<sup>1</sup>

<sup>1</sup> INSERM (Institut National de la Santé et de la Recherche Médicale), ERI 20, Université Paris-Sud EA 4045, and Institut Gustave Roussy, Villejuif, F-94805 France

<sup>2</sup> Institut Gustave Roussy, Unité de Psycho-Oncologie, Villejuif, F-94805 France.

<sup>3</sup> INSERM, U888, Montpellier, F-34093 France; Université Montpellier1, Montpellier, F-34000 France

<sup>4</sup> Hôpital Paul Brousse, Assistance Publique-Hôpitaux de Paris, Pôle Vieillessement - Réadaptation - Accompagnement, Villejuif, F-94800 France.

**Correspondence** to Françoise Clavel-Chapelon (clavel@igr.fr)

INSERM ERI-20, E3N-EPIC Study, Institut Gustave Roussy, Espace Maurice Tubiana  
39 rue Camille Desmoulins  
94805 Villejuif Cedex  
France

Tel: (+33) 1 42 11 41 48; Fax: (+33) 1 42 11 40 00

## Key points

1. Characterization of cognitive status in large-scale epidemiologic studies is challenging.
2. Cognitive assessment by telephone represents an interesting cost-effective method.
3. In a sample of 120 community-dwelling French women, the Telephone Interview for Cognitive Status (F-TICS-m) showed satisfactory concurrent validity against the gold-standard classification from a neuropsychological examination.
4. Our results are consistent with previous evaluation studies in other languages and settings to support the validity of TICS-m to appraise global cognitive functioning.

**ACKNOWLEDGMENTS:** The authors report no conflict of interest. The E3N cohort is supported by the French League against Cancer, the European Community, the “Mutuelle Générale de l’Education Nationale”, the French Institute of Health and Medical Research, the Gustave Roussy Institute and several general councils in France. MN Vercambre is on a grant from the Statlife Company. None of the funding agencies played a role in the design and conducting of the study, analysis and interpretation of data, or preparation and approval of the manuscript. The authors are grateful to Laurence Lacoste (LL), whose preliminary work was useful for launching this study, and to the team of Bruno Vellas (INSERM U558, CHU de Toulouse) for helpful suggestions when adapting TICS-m in French. We also express our gratitude to Dominique Villebrun for help with the bibliography and Jerri Bram for assistance with English editing.

## **Word count:**

Abstract: 249; Text: 2,858

## ABSTRACT

**Objective:** To evaluate the performance of a French version of the modified Telephone Interview for Cognitive Status (F-TICS-m) in identifying cognitive decline among elderly women.

**Methods:** All women aged 72-86 participating in the “Etude Epidémiologique auprès de Femmes de l’Education Nationale” (E3N) cohort and living in or near Paris constituted the target population of the validation study. Volunteer women (n=120) underwent both a 20-min telephone interview and a face-to-face neuropsychological examination at an interval of few days (median interval: 10 days). The telephone interview included F-TICS-m, as well as a recall of key elements of a short story, arithmetic/verbal problems and two verbal fluency tests. Neuropsychological examination consisted of a standardized battery of cognitive tests (including the Mini-Mental State Examination (MMSE), the Free and Cued Selective Reminding Test-FCSRT, Trailmaking tests A and B, the similarities subtest of the Wechsler Adult Intelligence Scale-III...) and the Instrumental Activities of Daily Living questionnaire and the Geriatric Depression Scale. Neuropsychological examination led to classification of each subject as cognitively normal (n=92) or impaired/demented (n=28).

**Results:** F-TICS-m showed satisfactory internal consistency (Cronbach’s  $\alpha=0.69$ ). It correlated linearly with MMSE (Pearson’s  $r=0.72$ ). Concurrent validity against the gold-standard classification was satisfactory, with an area under the ROC curve (AUC) of 0.83. The combination of F-TICS-m and the other telephone tests had no additional effect on discrimination power in our sample (AUC=0.81).

**Conclusion:** F-TICS-m is a valid instrument for assessing the overall cognitive status of French elderly women. Its validity in men and its reproducibility warrant further studies.

## INTRODUCTION

Dementia is an important public health concern, with a prevalence that is rapidly increasing with population ageing (Kalaria *et al.*, 2008). In the absence of curative treatments, prevention has become a crucial challenge, and epidemiologic studies on risk factors in cognitive decline are warranted. However, progress in the field has been hampered by difficulties in evaluating cognitive status in large populations. Health care data are unable to exhaustively identify dementia cases due to underdiagnosis (Kurz *et al.*, 2001; Tyas *et al.*, 2006; Falagas *et al.*, 2007). Consequently, epidemiologic studies on cognitive decline/dementia must develop active strategies for characterizing cognitive status.

Conventional in-person neuropsychological assessment techniques are the gold-standard for identifying the cognitively impaired, but this approach is costly, especially in geographically dispersed studied populations. Telephone interviews provide an interesting alternative, since they are inexpensive and well accepted by elderly subjects. Thus, several telephone-administered cognitive tests have been developed during the last two decades (Ball and McLaren, 1997). In particular, the Telephone Interview for Cognitive Status (TICS) (Brandt *et al.*, 1988) has been broadly modelled on the Mini-Mental State examination (MMSE) (Folstein *et al.*, 1975), currently the most widely used screening instrument for cognitive impairment (Shulman *et al.*, 2006), though initially designed for face-to-face administration. The modified TICS-m (Brandt *et al.*, 1993; Welsh *et al.*, 1993) is a further refinement of the TICS instrument: a delayed recall task was added in order to broaden cognitive scope and accuracy.

The English version of TICS-m has demonstrated its value as a screening instrument in various populations (Welsh *et al.*, 1993; Plassman *et al.*, 1994; Gallo and Breitner, 1995; de Jager *et al.*, 2003; Lines *et al.*, 2003) and has been used in several Anglo-Saxon

epidemiologic studies (Grodstein *et al.*, 2003; Barber and Stott, 2004; Crooks *et al.*, 2005).

Furthermore, TICS and/or TICS-m have been translated into many languages and evaluated:

Spanish (Gude *et al.*, 1994), Finnish (Jarvenpaa *et al.*, 2002), Hebrew (Beeri *et al.*, 2003),

Italian (Ferrucci *et al.*, 1998; Dal Forno *et al.*, 2006), Japanese (Konagaya *et al.*, 2007) and

German (Loerbroks *et al.*, 2008), but not yet French.

The objective of the present study was to evaluate the performance of a French version of

TICS-m (F-TICS-m) as compared with conclusions from a complete neuropsychological

examination. The validation study was conducted in the framework of the “Etude

Epidémiologique de Femmes de l’Education Nationale” (E3N) study (Vercambre *et al.*,

2009).

## METHODS

### Subjects and procedure

The target population for the present study included all E3N cohort participants born between 1925 and 1930, alive at the beginning of the validation study and living in or near Paris. All eligible women were contacted by mail or telephone and informed of the purpose of the research and procedure. Volunteers able to speak/understand French and who had no hearing impairment underwent a telephone interview and were invited to undergo a standard neuropsychological examination at Paul Brousse Hospital (Villejuif, France) within two weeks. Those who participated in this second step were included in the validation study. At the end of the examination, results were communicated to the participant and a report was sent to her physician.

Telephone interviews and face-to-face examinations were conducted between September 2007 and September 2008. All study subjects signed an informed consent form in compliance with the rules of the French National Commission for Computed Data and Individual Freedom (Commission Nationale Informatique et Libertés) from which we obtained approval.

### The F-TICS-m instrument

The French version of TICS-m (available on [www.idf.inserm.fr/site/eri20/page.asp?page=785](http://www.idf.inserm.fr/site/eri20/page.asp?page=785)) compiles all translated items from the English version of either the TICS or TICS-m questionnaires. Minor modifications of culture-dependent questions were made (e.g. the question concerning the present American vice-president was changed to a question about the last French president). Moreover, we chose to follow the BEC96 method (Signoret, 1989) for the 10-word list memory task: 10 words from the Alzheimer's Disease Assessment Scale (Mohs, 1996) were given three times (each followed by a free recall); then, following other questions, there was a delayed recall without

prior presentation. In the total, only scores from the first and last recalls were taken into account.

Ultimately, the F-TICS-m instrument included consecutive questions evaluating temporal and spatial orientation (scored on 5 and 3 points, respectively); the immediate recall task of the 10-word list (scored on 10 points); the attention and calculation tasks of the MMSE (serial 7s and spell backwards) (scored on 6 points); six questions evaluating semantic memory (scored on 6 points); a delayed recall of the 10-word list (scored on 10 points); and two tasks involving language (repetition of two complex sentences and comprehension of a simple command) (scored on 3 points). Thus, F-TICS-m had a maximum score of 43.

#### Telephone interview

Each telephone interview was carried out by one of the two neuropsychologists involved in the project (HC and LL). Before beginning cognitive assessment, the neuropsychologist asked the respondent if she was alone in a quiet room with no pen or pencil and no temporal information within easy reach (newspaper, calendar, agenda). F-TICS-m administration was followed by four other cognitive tests (feasible by telephone) that were added to increase sensitivity to non-Alzheimer dementia. These additional tests were proposed in the following order: a recall of key elements of a short story (subtest of Wechsler's clinical memory scale, third edition (Wechsler, 2001)); arithmetic /verbal problems; and two verbal fluency tests (name as many words starting with "s" as possible within one minute; name as many fruits as possible within one minute). The latter three tests were chosen from the BEC96 battery (Signoret, 1989). At the end of the telephone interview, self-rated morale was appraised through a simple question ("How was your mood over the last two weeks: good, average, or poor?"). The telephone interview lasted no more than 20 minutes (10 minutes for F-TICS-m and 10 minutes for other tests).

### Neuropsychological examination

Within two weeks after the telephone interview (median delay: 10 days), each participant underwent a complete standardized examination by a neuropsychologist (HC or LL). This included anamnesis, a cognitive test battery and other questionnaires, as cited below. The cognitive domains evaluated were, by order: (1) global cognitive functioning: French version of MMSE (Derouesne *et al.*, 1999); (2) memory: 16 words of the Free and Cued Selective Reminding test or FCSRT (Van der Linden *et al.*, 2004); (3) executive function: Trail Making tests A and B (Tombaugh, 2004); (4) visual-constructive capacities: Clock Drawing test (Shulman, 2000); (5) autonomy: four Instrumental Activities of Daily Living (Barberger-Gateau *et al.*, 1999); (6) depression: Geriatric Depression Scale-15 items (Sheikh and Yesavage, 1986); (7) language: a French picture-naming test (DO80) (Deloche and Hannequin, 1997) and lexical evocations (Cardebat *et al.*, 1990); and (8) reasoning: similarities subtest of the Wechsler Adult Intelligence Scale-III (Wechsler, 2000).

### Gold-standard classification

After all examinations had been carried out, two neuropsychologists blinded to telephone scores (HC and IHL) classified participants according to results from the neuropsychological examination. Criteria were approved by an experienced geriatrician (CT). Four groups were observed: (1) normal cognitive status (cognitive scores within age- and education-adjusted norms), (2) mild cognitive impairment (according to Ravaglia *et al.* (Ravaglia *et al.*, 2008): deficit in one cognitive domain, but no evidence of functional dependency), (3) possible dementia (poor scores in many cognitive domains, but not so pathological and/or possibly related to depression), and (4) probable dementia (abnormal scores in many cognitive

domains with impact on autonomy). To evaluate discriminative ability of F-TICS-m, we contrasted scores of cognitively healthy women (1) with cognitively impaired women (2+3+4).

#### Statistical analysis

In order to assess the performance of the French version of TICS-m, we consecutively evaluated its internal consistency, construct validity and concurrent validity (i.e. discriminative ability) (Gifford and Cummings, 1999), but first, we evaluated selection biases in our study, in comparing women included in the analysis sample with the rest of the female target population and with women who participated only in the telephone interview. We described then distribution and covariates of F-TICS-m. Cronbach's alpha was calculated to evaluate the performance of the set of F-TICS-m items at measuring a single unidimensional latent construct (internal consistency); Pearson's correlation was used to quantify the linear relationship between F-TICS-m and MMSE scores (construct validity). To assess concurrent validity, we compared F-TICS-m distribution according to the gold-standard cognitive status as defined through the neuropsychological examination. We also carried out receiver operating curve (ROC) analysis (Swets, 1979), which summarizes the discriminative ability of the instrument as a whole. We computed the area under the curve (AUC statistics) for different cognitive tests and compared them with each other: F-TICS-m, F-TICS (F-TICS-m from which the delayed recall score had been post-hoc subtracted) and face-to-face MMSE. In addition, in order to determine whether F-TICS-m performance could be improved by adding other telephone tests, we compared discrimination statistics obtained with F-TICS-m alone to those obtained with the overall telephone score, constructed as the average of the five Z-scores to the different cognitive tests comprising the telephone interview.

Since sensitivity and specificity do not indicate whether the test will provide the correct screening result, we also calculated F-TICS-m positive and negative predictive values for different cut-off points.

All statistical analyses were performed using SAS software, version 9.1 (SAS Institute, Inc., Cary, NC). All results were considered significant at the 5 % level. All statistical tests were two-sided.

## RESULTS

### Selection and subject characteristics

Among 726 eligible E3N participants, 310 responded to our mailing/telephone contact.

Among them, 121 women refused to participate and 4 no longer fulfilled the inclusion criteria (e.g. they had moved out of the Paris area). Among the 185 women who took part in the telephone interview, 65 did not undergo the neuropsychological examination, mainly due to motor impairment. Finally, the validation sample included 120 women. Among them, face-to-face neuropsychological examination identified 28 who were cognitively impaired (6 probable dementia, 4 possible dementia and 18 mild cognitive impairment). The remaining 92 women were considered to have no cognitive impairment.

Compared to non-included women, those who had participated in both the telephone interview and face-to-face examination were somewhat younger (mean age: 78.8 vs. 79.2 years old, p-value in the Wilcoxon-Mann-Whitney test =0.01) and more likely to have responded to all E3N questionnaires (73.3% vs. 49.5%, p-value in Chi<sup>2</sup> test <0.001). No significant difference was found in education level (p-value to the Chi<sup>2</sup> test = 0.53).

Compared to included women, those who were interviewed only by telephone did not differ significantly in terms of F-TICS-m score, age, education level or self-reported morale, but they were less likely to have responded to all E3N questionnaires than those who underwent neuropsychological examination (58.5% vs. 73.3%, p-value to the Chi<sup>2</sup> test = 0.04).

### Distribution and covariates of F-TICS-m

In the 120-woman validation sample, observed F-TICS-m scores ranged between 16 and 40, with a mean (SD) of 32.6 (4.2). Cut-off points for first, second and third quartiles were, respectively, 31, 33 and 35.

F-TICS-m statistics among the 92 cognitively normal women are presented in Table 1 by age group, level of education, E3N compliance status and self-reported morale. On average, women in the younger age group had higher F-TICS-m score ( $p < 0.05$ ). More educated women tended also to score higher ( $p = 0.06$ ). The F-TICS-m score in intellectually intact women was associated neither with E3N compliance nor with self-reported morale.

#### Internal consistency and construct validity

In our data, Cronbach's alpha of F-TICS-m equalled 0.69. F-TICS-m correlated linearly with face-to-face MMSE score, with a Pearson's correlation coefficient of 0.72.

#### Discriminative ability

Compared to women in the normal group, both groups of women with mild cognitive impairment and possible or probable dementia had a significantly lower mean F-TICS-m score (p-values in Wilcoxon-Mann-Whitney test  $\leq 0.001$ ) (Table 2).

AUC statistics for F-TICS-m, F-TICS, face-to-face MMSE and overall telephone score equalled, respectively, 0.83, 0.78, 0.72 and 0.81 (Figure 1). F-TICS-m outperformed both F-TICS and MMSE, with sensitivity and specificity values higher for the majority of score cut-offs. As compared with overall telephone score, F-TICS-m alone did not present less satisfactory overall discriminative capacity.

Discrimination statistics for cognitive impairment are presented in Table 3 for selected cut-points of the F-TICS-m score. For instance, specificity, sensitivity and positive and negative predictive values were 0.89, 0.68, 0.66, and 0.90, respectively, for a cut-off of 30, and 0.60, 0.86, 0.39, and 0.93, respectively, for a cut-off of 33.

## DISCUSSION

In our sample of elderly French women participating in the E3N cohort, F-TICS-m proved to be efficient at identifying cognitive impairment. In addition, combining F-TICS-m with four other cognitive telephone tests did not [appreciably](#) improve its discriminative ability, thus supporting the validity of F-TICS-m alone to appraise global cognitive functioning. Our results are consistent with previous evaluation studies of TICS (Dal Forno *et al.*, 2006; Konagaya *et al.*, 2007) and TICS-m (Welsh *et al.*, 1993; de Jager *et al.*, 2003) in other languages, which demonstrated the widespread usefulness of this instrument. More specifically, the 10-word delayed recall of the TICS-m provides valuable information on cognitive assessment, since use of this task appears to improve discriminative accuracy in our sample of highly functioning women.

Some limitations of the present study have to be discussed. First, the rather low response and completion rates may have resulted in biases when non-responders differed substantially from responders in terms of the studied characteristics. Besides, the number of cognitively-impaired women was small compared with the number of non-impaired women, as expected in a population setting. Although statistics are less robust than with larger numbers, our sample size is consistent with previously published validation studies of cognitive tests by telephone (de Jager *et al.*, 2003; Jarvenpaa *et al.*, 2002; Konagaya *et al.*, 2007). On the whole, women participating in the validation study showed largely intact cognitive ability (only 7 scored below 25 points on MMSE in face-to-face examination). Thus, the prevalence of impairment in our study may well have been lower than that of the general population and positive predictive value may have been underestimated. Second, the study sample comprised only women, precluding extending our findings to male populations; moreover, participants were mainly former teachers and thus highly educated. Nevertheless, the English version of TICS-m had previously shown no difference between men and women, nor between low and

high education groups (de Jager *et al.*, 2003), suggesting that the F-TICS-m may be used at the population level. Third, we cannot rule out the possibility that subtle variations in ratings may have occurred among the raters, although the telephone interview and neuropsychological examination were highly standardized. Our data did not enable calculation of test-retest and inter-rater reliability. However, studies involving other language versions have suggested good reproducibility (Dal Forno *et al.*, 2006; Loerbroks *et al.*, 2008). Indeed, in a sample of 35 educated, highly functioning older women which was similar to our sample, a test-retest study of two TICS assessments at 31-day intervals showed a correlation of 0.7 (Kang *et al.*, 2007).

Our study has several strengths. It is the first to evaluate a French version of TICS-m. In addition, all participants underwent the same type of confirmatory assessment (regardless of telephone screening results), thus excluding verification bias (Donald and Van Til, 2001). Another strong feature of our study was that it involved a sample from the general population (i.e. a population with a relevant basal rate of impairment) rather than using a case-control design (that would artificially exclude subjects with intermediate cognitive decline or specific dementia etiology). Although performance statistics are generally lower in the former context (because of a more heterogeneous range of cognitive abilities), it provides more reliable cut-off points for screening purposes (Ritchie and Fuhrer, 1996) and is essential for demonstrating accuracy at detecting any type of impairment, rather than just one specific type (Cullen *et al.*, 2007).

Our data, along with previous studies, plead for the usefulness of a simple test such as TICS-m to efficiently evaluate cognitive function by telephone. Although large batteries of neuropsychological tests are essential in the psychogeriatrics field to make an accurate and reliable diagnosis, TICS-m is particularly suitable for large-scale community screening and/or

epidemiological follow-up: in addition to its low cost, it can be rapidly administered, which ensures better acceptability, especially among the cognitively-impaired and the physically-impaired elderly, who may not be strongly motivated to participate. Despite its advantages, however, TICS-m, like other telephone tests, is not adapted to detecting cognitive difficulties involving praxies and visual-spatial capacities. Moreover, its use is limited to subjects without hearing problems and those with non-reduced social contact.

Interestingly, TICS-m distribution was shown, in previous reports, and in our highly educated population, more symmetric and normal-shaped than MMSE distribution, suggesting that TICS-m is less subject to the ceiling effect that habitually limits MMSE. This statistical characteristic, which is probably due to the difficulty involved in the 10-word memory task (de Jager *et al.*, 2003), makes TICS-m more sensitive for detecting early cognitive impairment than TICS or MMSE alone. This difference between TICS and TICS-m has been reported previously (Welsh *et al.*, 1993), but it appeared more clear-cut in our sample of highly educated women.

In conclusion, our results support the reliability of F-TICS-m for assessing cognitive function in a French female population. Its use in men and its reproducibility warrant further studies, but our results already suggest promising perspectives in the field of geriatric epidemiology.

## REFERENCES

- Ball C, McLaren P. 1997. The tele-assessment of cognitive state: a review. *J Telemed Telecare* 3(3): 126-31.
- Barber M, Stott DJ. 2004. Validity of the Telephone Interview for Cognitive Status (TICS) in post-stroke subjects. *Int J Geriatr Psychiatry* 19(1): 75-9.
- Barberger-Gateau P, Fabrigoule C, Helmer C, Rouch I, Dartigues JF. 1999. Functional impairment in instrumental activities of daily living: an early clinical sign of dementia? *J Am Geriatr Soc* 47(4): 456-62.
- Beeri MS, Werner P, Davidson M, Schmidler J, Silverman J. 2003. Validation of the modified telephone interview for cognitive status (TICS-m) in Hebrew. *Int J Geriatr Psychiatry* 18(5): 381-6.
- Brandt J, Spencer M, Folstein M. 1988. The telephone Interview for Cognitive Status. *Neuropsych Neuropsychol and Behavioral Neurology* 1(2): 111-7.
- Brandt J, Welsh KA, Breitner JC, Folstein MF, Helms M, Christian JC. 1993. Hereditary influences on cognitive functioning in older men. A study of 4000 twin pairs. *Arch Neurol* 50(6): 599-603.
- Cardebat D, Doyon B, Puel M, Goulet P, Joannette Y. 1990. [Formal and semantic lexical evocation in normal subjects. Performance and dynamics of production as a function of sex, age and educational level]. *Acta Neurol Belg* 90(4): 207-17.
- Crooks VC, Clark L, Petitti DB, Chui H, Chiu V. 2005. Validation of multi-stage telephone-based identification of cognitive impairment and dementia. *BMC Neurol* 5(1): 8.
- Cullen B, O'Neill B, Evans JJ, Coen RF, Lawlor BA. 2007. A review of screening tests for cognitive impairment. *J Neurol Neurosurg Psychiatry* 78(8): 790-9.

Dal Forno G, Chiovenda P, Bressi F, Ferreri F, Grossi E, Brandt J, Rossini PM, Pasqualetti P. 2006. Use of an Italian version of the telephone interview for cognitive status in Alzheimer's disease. *Int J Geriatr Psychiatry* 21(2): 126-33.

de Jager CA, Budge MM, Clarke R. 2003. Utility of TICS-M for the assessment of cognitive function in older adults. *Int J Geriatr Psychiatry* 18(4): 318-24.

Deloche G, Hannequin D. 1997. Test de dénomination orale d'images (DO 80). ECPA, Paris.

Derouesne C, Poitreneau J, Hugonot L, Kalafat M, Dubois B, Laurent B. 1999. [Mini-Mental State Examination: a useful method for the evaluation of the cognitive status of patients by the clinician. Consensual French version]. *Presse Med* 28(21): 1141-8.

Donald A, Van Til L. 2001. Evaluating screening tests for dementia and cognitive impairment in a heterogeneous population in the presence of verification bias. *Int Psychogeriatr* 13 Supp 1: 203-14.

Falagas ME, Vardakas KZ, Vergidis PI. 2007. Under-diagnosis of common chronic diseases: prevalence and impact on human health. *Int J Clin Pract* 61(9): 1569-79.

Ferrucci L, Del L, I, Guralnik JM, Bandinelli S, Benvenuti E, Salani B, Lamponi M, Ubezio C, Benvenuti F, Baroni A. 1998. Is the telephone interview for cognitive status a valid alternative in persons who cannot be evaluated by the Mini Mental State Examination? *Aging (Milano)* 10(4): 332-8.

Folstein MF, Folstein SE, McHugh PR. 1975. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 12(3): 189-98.

Gallo JJ, Breitner JC. 1995. Alzheimer's disease in the NAS-NRC Registry of aging twin veterans, IV. Performance characteristics of a two-stage telephone screening procedure for Alzheimer's dementia. *Psychol Med* 25(6): 1211-9.

Gifford DR, Cummings JL. 1999. Evaluating dementia screening tests: methodologic standards to rate their performance. *Neurology* 52(2): 224-7.

Grodstein F, Chen J, Willett WC. 2003. High-dose antioxidant supplements and cognitive function in community-dwelling elderly women. *Am J Clin Nutr* 77(4): 975-84.

Gude RR, Calvo Mauri JF, Carrasco Lopez FJ. 1994. [The Spanish version and pilot study of a telephone test of cognitive status for evaluation and screening in dementia assessment and follow-up]. *Aten Primaria* 13(2): 61-6.

Jarvenpaa T, Rinne JO, Raiha I, Koskenvuo M, Lopponen M, Hinkka S, Kaprio J. 2002. Characteristics of two telephone screens for cognitive impairment. *Dement Geriatr Cogn Disord* 13(3): 149-55.

Kalaria RN, Maestre GE, Arizaga R, Friedland RP, Galasko D, Hall K, Luchsinger JA, Ogunniyi A, Perry EK, Potocnik F, Prince M, Stewart R, Wimo A, Zhang ZX, Antuono P. 2008. Alzheimer's disease and vascular dementia in developing countries: prevalence, management, and risk factors. *Lancet Neurol* 7(9): 812-26.

Kang JH, Cook N, Manson J, Buring JE, Grodstein F. 2007. Low dose aspirin and cognitive function in the women's health study cognitive cohort. *BMJ* 334(7601): 987.

Katz S. 1983. Assessing self-maintenance: activities of daily living, mobility, and instrumental activities of daily living. *J Am Geriatr Soc* 31(12): 721-7.

Konagaya Y, Washimi Y, Hattori H, Takeda A, Watanabe T, Ohta T. 2007. Validation of the Telephone Interview for Cognitive Status (TICS) in Japanese. *Int J Geriatr Psychiatry* 22(7): 695-700.

Kurz X, Scuvee-Moreau J, Salmon E, Pepin JL, Ventura M, Dresse A. 2001. Dementia in Belgium: prevalence in aged patients consulting in general practice (french). *Rev Med Liege* 56(12): 835-9.

Lines CR, McCarroll KA, Lipton RB, Block GA. 2003. Telephone screening for amnesic mild cognitive impairment. *Neurology* 60(2): 261-6.

Loerbroks A, Amelang M, Sturmer T. 2008. Reproducibility of a telephone interview assessing cognitive function and depressive symptoms in older adults in Germany. *Int J Geriatr Psychiatry* 23(10):1098-101.

Mohs RC. 1996. The Alzheimer's Disease Assessment Scale. *Int Psychogeriatr* 8(2): 195-203.

Plassman BL, Newman TT, Welsh KA, Helms M, Breitner JC. 1994. Properties of the telephone interview for cognitive status. *Neuropsych, Neuropsychol, and Behavioral Neurology* 7: 235-41.

Ravaglia G, Forti P, Montesi F, Lucicesare A, Pisacane N, Rietti E, Dalmonte E, Bianchin M, Mecocci P. 2008 Mild cognitive impairment: epidemiology and dementia risk in an elderly Italian population. *J Am Geriatr Soc* 56(1): 51-8.

Ritchie K, Fuhrer R. 1996. The validation of an informant screening test for irreversible cognitive decline in the elderly: performance characteristics within a general population sample. *Int J Geriatr Psychiatry* 11(2): 149-56.

Sheikh JI, Yesavage JA. 1986. Geriatric depression scale (GDS): recent findings and development of a shorter version. In: *Clinical gerontology: a guide to assessment and intervention*, Brink TL (ed). Howarth Press: New York, 165-172.

Shulman KI. 2000. Clock-drawing: is it the ideal cognitive screening test? *Int J Geriatr Psychiatry* 15(6): 548-61.

Shulman KI, Herrmann N, Brodaty H, Chiu H, Lawlor B, Ritchie K, Scanlan JM. 2006. IPA survey of brief cognitive screening instruments. *Int Psychogeriatr* 18(2): 281-94.

Signoret JL. 1989. [Evaluation of memory troubles and associated cognitive disorders. BEC96]. IPSEN, Paris.

Swets JA. 1979. ROC analysis applied to the evaluation of medical imaging techniques. *Invest Radiol* 14(2): 109-21.

- Tombaugh TN. 2004. Trail Making Test A and B: normative data stratified by age and education. *Arch Clin Neuropsychol* 19(2): 203-14.
- Tyas SL, Tate RB, Wooldrage K, Manfreda J, Strain LA. 2006. Estimating the incidence of dementia: the impact of adjusting for subject attrition using health care utilization data. *Ann Epidemiol* 16(6): 477-84.
- Van der Linden M, Coyette F, Poitrenaud J, membres du GRENEM. 2004. L'épreuve de rappel libre/rappel indicé à 16 items (RL/RI-16). In: *L'évaluation des troubles de la mémoire. Présentation de quatre tests de mémoire épisodique (avec leur étalonnage)* Solal, Van der Linden M, Adam S, Agniel A, membres du GRENEM (eds). Marseille, 25-47.
- Wechsler D. 2000. WAIS III. Echelle d'Intelligence de WECHSLER pour adultes Troisième édition. ECPA, Paris.
- Vercambre MN, Boutron-Ruault MC, Ritchie K, Clavel-Chapelon F, Berr C. 2009. Long-term association of food and nutrient intakes with cognitive and functional decline: a 13-year follow-up study of elderly French women. *Br J Nutr* in press 1-9.
- Wechsler D. 2001. Echelle clinique de Mémoire de Wechsler troisième édition. ECPA, Paris.
- Welsh KA, Breitner JC, Magruder-Habib K. 1993. Detection of dementia in the elderly using telephone screening of cognitive status. *Neuropsych, Neuropsychol, and Behavioral Neurology* 6: 103-10.

**Table 1.** F-TICS-m statistics according to selected characteristics of the 92 cognitively normal women included in the validation study sample

	n	Mean score (SD)	Range	P-value*
<b>Age</b>				
<80 years	63	34.3 (2.8)	22-40	
+80 years	29	33.2 (2.7)	28-38	0.046
<b>Education level</b>				
< 12 years	16	32.4 (4.1)	22-38	
≥ 12 years	76	34.2 (2.4)	28-40	0.064
<b>E3N compliance status</b>				
At least one missing questionnaire	24	33.9 (2.6)	28-38	
No missing questionnaire	68	33.9 (2.9)	22-40	0.89
<b>Self-rated morale</b>				
Good	47	34.1 (2.5)	28-40	
Average	28	34.3 (2.5)	29-38	0.12
Poor	17	32.6 (3.7)	22-37	0.16

F-TICS-m: French version of the modified TICS; SD: standard deviation

\* Wilcoxon-Mann-Whitney Test

**Table 2.** F-TICS-m statistics according to cognitive status as assessed through face-to-face neuropsychological examination

Cognitive status	n	Mean score (SD)	Min	Q1	Median	Q3	Max	P-value*
Normal	92	33.9 (2.8)	22**	32	34	36	40	
Mild cognitive impairment	18	30.9 (3.6)	25	28	30/31	33	37	0.001
Possible or probable dementia	10	23.7 (4.5)	16	23	25	26	30	<0.001

F-TICS-m: French version of the modified TICS; SD: standard deviation; Q1, Q3: first and third quartiles

\*Wilcoxon-Mann-Whitney Test

\*\* One woman in the “normal” group, with a 22 F-TICS-m score was highly depressed (the next low score observed in this group was 29). Although she presented slight executive troubles associated with depressive symptoms, she was classified as normal from the dementia point of view, considering her overall test performing

**Table 3.** Discrimination statistics for selected cut-off points applied to F-TICS-m score to identify cognitive impairment.

Cut-off point	Specificity	Sensitivity	Positive predictive value	Negative predictive value
34	0.47	0.86	0.33	0.91
33	0.60	0.86	0.39	0.93
32	0.71	0.75	0.44	0.90
31	0.83	0.71	0.56	0.90
30	0.89	0.68	0.66	0.90
29	0.95	0.54	0.75	0.87
28	0.98	0.50	0.88	0.87
27	0.99	0.46	0.93	0.86

**Figure 1.** ROC analysis for cognitive impairment screening\*: comparison of F-TICS-m, F-TICS, face-to-face MMSE and the overall telephone score

F-TICS-m: French version of the modified TICS; F-TICS: F-TICS-m from which the 10-word delayed recall task was post-hoc removed; MMSE: Mini Mental State Examination; overall telephone score: score constructed as the average of the five Z-scores on the different cognitive tests comprising the telephone interview;

ROC: receiver operating curve; AUC: area under the curve

\*Gold-standard assessment based on face-to-face neuropsychological examination