

ON CERTAIN CONSTRUCTIONS OF p -ADIC FAMILIES OF SIEGEL MODULAR FORMS OF EVEN GENUS

HISA-AKI KAWAMURA

To Professor Hiroshi Saito in memoriam

ABSTRACT. Suppose that $p > 5$ is a rational prime. Starting from a well-known p -adic analytic family of ordinary elliptic cusp forms of level p due to Hida, we construct a certain p -adic analytic family of holomorphic Siegel cusp forms of arbitrary even genus and of level p associated with Hida's p -adic analytic family via the Duke-Imamoğlu lifting provided by Ikeda. Moreover, we also derive a similar result for the Siegel Eisenstein series of even genus with trivial Nebentypus.

CONTENTS

1. Introduction	1
2. Preliminaries	8
2.1. Classical Duke-Imamoğlu lifting	8
2.2. Λ -adic Siegel modular forms	14
3. Cuspidal Λ -adic modular forms of genus 1	16
3.1. Hida's universal ordinary p -stabilized newforms	16
3.2. Λ -adic Shintani lifting	20
4. Main results	27
5. Appendix	36
5.1. Λ -adic Siegel Eisenstein series of even genus	36
5.2. Numerical evidences	38
References	39

1. INTRODUCTION

For a given rational prime $p > 5$, the study of p -adic analytic families of modular forms was initiated by Kummer and Eisenstein for non-cuspidal modular forms on the elliptic modular group $SL_2(\mathbb{Z})$, and afterwards was developed from various points of view by Iwasawa, Kubota-Leopoldt, Serre,

Date: 19 September, 2010.

This research was partially supported by the JSPS Core-to-Core Program “New Developments of Arithmetic Geometry, Motive, Galois Theory and Their Practical Applications”. It is also supported by the JSPS International Training Program (ITP).

Swinnerton-Dyer, Katz, Deligne-Ribet, Hida, Wiles and others. In particular, Hida [H3] and Wiles [Wi], among others, introduced the notion of Λ -adic modular forms, where $\Lambda = \mathbb{Z}_p[[1 + p\mathbb{Z}_p]]$ is the Iwasawa algebra, as formal q -expansions with coefficients in a finite flat Λ -algebra R whose specialization at each arithmetic point in the Λ -adic analytic space $\mathfrak{X}(R) = \text{Hom}_{\text{cont}}(R, \overline{\mathbb{Q}}_p)$ gives rise to the q -expansion of an elliptic modular form. In this context, a p -adic analytic family of modular forms can be regarded as an infinite collection of modular forms parametrized by varying weights whose components are interpolated by a Λ -adic modular form simultaneously.

In fact, a specific use of Hida's theory allows us to construct a Λ -adic modular form such that every specialization gives rise to a p -ordinary elliptic Hecke eigenform (i.e. a simultaneous eigenfunction of all Hecke operators such that the eigenvalue of the Atkin-Lehner U_p -operator is a p -adic unit), which is called the universal ordinary p -stabilized newform of tame level 1:

Fact 1.1 (cf. §3.1 below). Let $\Lambda_1 = \mathbb{Z}_p[[\mathbb{Z}_p^\times]]$ be the completed group ring on \mathbb{Z}_p^\times over \mathbb{Z}_p , and $\omega = \omega_p$ the Teichmüller character. There exist a Λ_1 -algebra R^{ord} finite flat over Λ and a formal q -expansion $\mathbf{f}_{\text{ord}} \in R^{\text{ord}}[[q]]$ such that for each arithmetic point $P \in \mathfrak{X}(R^{\text{ord}})$ of weight $2k > 2$ with trivial Nebentypus (i.e. P lies over the $\overline{\mathbb{Q}}_p^\times$ -valued continuous character $y \mapsto y^{2k}\omega(y)^{2k}$ on \mathbb{Z}_p^\times), the specialization $\mathbf{f}_{\text{ord}}(P)$ coincides with an ordinary p -stabilized newform f_{2k}^* of weight $2k$ on the congruence subgroup $\Gamma_0(p) \subset \text{SL}_2(\mathbb{Z})$ of level p with trivial Nebentypus. Namely, there exists a p -ordinary normalized cuspidal Hecke eigenform $f_{2k} = \sum_{m=1}^{\infty} a_m(f_{2k})q^m$ of weight $2k$ on $\text{SL}_2(\mathbb{Z})$ such that

$$f_{2k}^*(z) = f_{2k}(z) - \beta_p(f_{2k})f_{2k}(pz)$$

for all $z \in \mathfrak{H}_1 := \{z \in \mathbb{C} \mid \text{Im}(z) > 0\}$, and f_{2k}^* possesses the U_p -eigenvalue $\alpha_p(f_{2k})$, where $\alpha_p(f_{2k})$ and $\beta_p(f_{2k})$ denote the unit and non-unit p -adic roots of the equation

$$(1) \quad X^2 - a_p(f_{2k})X + p^{2k-1} = 0,$$

respectively.

In this setting, we pick an integer $k_0 \geq 6$ as small as possible to have an arithmetic point $P_0 \in \mathfrak{X}(R^{\text{ord}})$ of weight $2k_0$ with trivial Nebentypus corresponding to an actual modular form $\mathbf{f}_{\text{ord}}(P_0) = f_{2k_0}^*$, and fix an analytic neighborhood \mathfrak{U}_0 of P_0 in $\mathfrak{X}(R^{\text{ord}})$ on which every coefficient of \mathbf{f}_{ord} can be regarded as a p -adic analytic function. Then, by varying the weights $2k$ of arithmetic points $P \in \mathfrak{U}_0$ with trivial Nebentypus, we obtain an ordinary p -adic analytic family $\{f_{2k}^*\}$ parametrized by weights $2k \geq 2k_0$ with

$2k \equiv 2k_0 \pmod{(p-1)p^{m-1}}$ for some sufficiently large integer $m \geq 1$. Hereinafter, we refer to it as the Hida family of level p .

On the other hand, as an affirmative answer to the Duke-Imamoğlu conjecture concerning on a generalization of the Saito-Kurokawa lifting towards higher genus, Ikeda [I1] established the following:

Fact 1.2 (cf. §2.1 below). For each integer $n \geq 1$, let f be a normalized cuspidal Hecke eigenform of weight $2k$ on $\mathrm{SL}_2(\mathbb{Z})$ with $k \equiv n \pmod{2}$. Then there exists a non-zero cuspidal Hecke eigenform $\mathrm{Lift}^{(2n)}(f)$ of weight $k+n$ on the Siegel modular group $\mathrm{Sp}_{4n}(\mathbb{Z}) \subset \mathrm{GL}_{4n}(\mathbb{Z})$ of genus $2n$ such that the standard L -function $L(s, \mathrm{Lift}^{(2n)}(f), \mathrm{st})$ is taken of the form

$$L(s, \mathrm{Lift}^{(2n)}(f), \mathrm{st}) = \zeta(s) \prod_{i=1}^{2n} L(s+k+n-i, f),$$

where $\zeta(s)$ and $L(s, f)$ denote Riemann’s zeta function and Hecke’s L -function associated with f , respectively.

When $n = 1$, $\mathrm{Lift}^{(2)}(f)$ coincides with the Saito-Kurokawa lifting of f , whose existence was firstly conjectured by Saito and Kurokawa [Ku], and afterwards was shown by Maaß [M], Andrianov [An] and Zagier [Z]. In accordance with the tradition, we refer to $\mathrm{Lift}^{(2n)}(f)$ as the Duke-Imamoğlu lifting of f throughout the present article. We should mention that such particular objects obtained by means of the lifting process from lower genus are, of course, not “genuine” Siegel cusp forms of higher genus in the strict sense. Indeed, the above functoriality equation yields that the associated Satake parameter $(\psi_0(p), \psi_1(p), \dots, \psi_{2n}(p)) \in (\overline{\mathbb{Q}}_p^\times)^{2n+1}$ at p is taken as

$$\psi_i(p) = \begin{cases} p^{nk-n(n+1)/2} & \text{if } i = 0, \\ \alpha_p(f)p^{-k+i} & \text{if } 1 \leq i \leq n, \\ \beta_p(f)p^{-k-n+i} & \text{if } n+1 \leq i \leq 2n, \end{cases}$$

uniquely up to the action of the Weyl group $W_{2n} \simeq \mathfrak{S}_{2n} \times \{\pm 1\}^{2n}$, where $(\alpha_p(f), \beta_p(f))$ denotes the ordered pair of the roots of the same equation as (1) with the inequality of p -adic valuations $v_p(\alpha_p(f)) \leq v_p(\beta_p(f))$. Therefore the famous Ramanujan-Petersson conjecture for f implies the fact that $\mathrm{Lift}^{(2n)}(f)$ generates a non-tempered cuspidal automorphic representations of the group $\mathrm{GSp}_{4n}(\mathbb{A}_{\mathbb{Q}})$ of symplectic similitudes, where $\mathbb{A}_{\mathbb{Q}}$ denotes the ring of adèles of \mathbb{Q} . However, it has also been observed that some significant properties of $\mathrm{Lift}^{(2n)}(f)$ can be derived from corresponding properties of f . For instance, as will be explained more precisely in the sequel, the Fourier expansion of $\mathrm{Lift}^{(2n)}(f)$ can be written explicitly in terms of those of f and

a cuspidal Hecke eigenform of half-integral weight corresponding to f via the Shimura correspondence.

Now, let us explain our results. Let $\{f_{2k}\}$ be the infinite collection of p -ordinary normalized cuspidal Hecke eigenforms on $\mathrm{SL}_2(\mathbb{Z})$ corresponding to the Hida family $\{f_{2k}^*\}$ via the ordinary p -stabilization. The aim of the present article is to construct a p -adic analytic family of Siegel cusp forms on the congruence subgroup $\Gamma_0(p) \subset \mathrm{Sp}_{4n}(\mathbb{Z})$ of level p corresponding to the collection $\{\mathrm{Lift}^{(2n)}(f_{2k})\}$ under a suitable p -stabilization process. Namely, our main results are summarized as follows:

Theorem 1.3. *For each integer $n \geq 1$, let k_0 be a positive integer with $k_0 > n + 1$ and $k_0 \equiv n \pmod{2}$, $P_0 \in \mathfrak{X}(R^{\mathrm{ord}})$ an arithmetic point of weight $2k_0$ with trivial Nebentypus, and \mathfrak{U}_0 a fixed analytic neighborhood of P_0 . For each integer $k \geq k_0$ with $k \equiv k_0 \pmod{2}$, put*

$$\Phi_p^*(Y) := (Y - \alpha_p(f_{2k})^n)^{-1} \prod_{r=1}^{2n} \prod_{1 \leq i_1 < \dots < i_r \leq 2n} (Y - \psi_0(p)\psi_{i_1}(p) \cdots \psi_{i_r}(p)),$$

$$\Psi_p^*(Y) := (Y - \alpha_p(f_{2k})^{n-1} p^{k+n-1}) \prod_{i=1}^n (Y - \alpha_p(f_{2k})^{n-1} \beta_p(f_{2k}) p^{2i-2}),$$

and

$$\mathrm{Lift}^{(2n)}(f_{2k})^* := \frac{\Psi_p^*(\alpha_p(f_{2k})^n)}{\Phi_p^*(\alpha_p(f_{2k})^n)} \cdot \mathrm{Lift}^{(2n)}(f_{2k})|_{k+n} \Phi_p^*(U_{p,0}),$$

where $(\psi_0(p), \psi_1(p), \dots, \psi_{2n}(p))$ is the Satake parameter of $\mathrm{Lift}^{(2n)}(f_{2k})$ taken as above, and $U_{p,0}$ is the Hecke operator corresponding to the double coset

$$I_{4n} \mathrm{diag}(\underbrace{1, \dots, 1}_{2n}, \underbrace{p, \dots, p}_{2n}) I_{4n}$$

with respect to the standard Iwahori subgroup I_{4n} of $\mathrm{GSp}_{4n}(\mathbb{Z}_p)$. Then we have

- (i) $\mathrm{Lift}^{(2n)}(f_{2k})^*$ is a cuspidal Hecke eigenform of weight $k+n$ on $\Gamma_0(p) \subset \mathrm{Sp}_{4n}(\mathbb{Z})$ with trivial Nebentypus such that the eigenvalues agree with those of $\mathrm{Lift}^{(2n)}(f_{2k})$ for each prime $l \neq p$, and we have

$$\mathrm{Lift}^{(2n)}(f_{2k})^*|_{k+n} U_{p,0} = \alpha_p(f_{2k})^n \cdot \mathrm{Lift}^{(2n)}(f_{2k})^*.$$

- (ii) Let $\sigma : \Lambda_1 \rightarrow \Lambda_1$ be the ring homomorphism induced from the group homomorphism $y \mapsto y^2$ on \mathbb{Z}_p^\times , and

$$\tilde{R}^{\mathrm{ord}} := R^{\mathrm{ord}} \otimes_{\Lambda_1, \sigma} \Lambda_1,$$

a finite Λ_1 -algebra with the structure homomorphism $\lambda \mapsto 1 \otimes \lambda$ on Λ_1 . If $\tilde{P}_0 \in \mathfrak{X}(\tilde{R}^{\mathrm{ord}})$ is an arithmetic point lying over P_0 , there exist a formal Fourier expansion \mathbf{F} with coefficients in the localization $(\tilde{R}^{\mathrm{ord}})_{(\tilde{P}_0)}$ of \tilde{R}^{ord} at \tilde{P}_0 , and a choice of p -adic periods $\Omega_P \in \overline{\mathbb{Q}}_p$ for $P \in \mathfrak{U}_0$ in the sense of Greenberg-Stevens [GS] satisfying the following properties:

- $\Omega_{P_0} \neq 0$.
- There exists a normalization of $\text{Lift}^{(2n)}(f_{2k})$ such that for each arithmetic point $\tilde{P} \in \tilde{\mathfrak{X}}(\tilde{R}^{\text{ord}})$ lying over some arithmetic point $P \in \mathfrak{U}_0$ of weight $2k$ with trivial Nebentypus, we have

$$\mathbf{F}(\tilde{P}) = \frac{\Omega_P}{\Omega^\epsilon(P)} \text{Lift}^{(2n)}(f_{2k})^* \neq 0,$$

where $\Omega^\epsilon(P) \in \mathbb{C}^\times$ is the non-zero complex period of f_{2k} with signature $\epsilon \in \{\pm\}$ in the sense of Manin [Ma] and Shimura [S2].

Therefore we obtain a p -adic analytic family of non-zero Siegel cusp forms $\{\frac{\Omega_P}{\Omega^\epsilon(P)} \text{Lift}^{(2n)}(f_{2k})^*\}$ parametrized by varying weights $2k \geq 2k_0$ with

$$k \equiv k_0 \pmod{2} \text{ and } 2k \equiv 2k_0 \pmod{(p-1)p^{m-1}}$$

for some sufficiently large $m \geq 1$.

We note that the existence of such p -adic analytic families have been suggested by Guerzhoy [Gu] for $n = 1$, and conjectured by Panchishkin [P2] for arbitrary $n > 1$. In particular, Guerzhoy [Gu] derived a similar p -adic interpolation property of an essential part of the Fourier expansion of $\text{Lift}^{(2)}(f_{2k})$ under mild conditions. The above theorem can be regarded as a generalization of his result reformulated in the way that seems most appropriate for the study of p -adic properties of the Duke-Imamoğlu lifting.

For the proof of Theorem 1.3, we will give an explicit form of the Fourier expansion of $\text{Lift}^{(2n)}(f_{2k})^*$ (cf. Theorem 4.1 below). By combining this with the Λ -adic Shintani lifting due to Stevens [St], we will give a Λ -adic analogue of the classical Duke-Imamoğlu lifting for the universal ordinary p -stabilized newform \mathbf{f}_{ord} , which allows us to resolve the p -adic interpolation problem for the whole Fourier expansion of $\text{Lift}^{(2n)}(f_{2k})^*$ (cf. Theorem 4.4 below).

Remark 1.4. From a geometric point of view, a p -adic deformation theory for Siegel modular forms of arbitrary genus has been established by Hida in the ordinary case (cf. [H4, H5], see also [TU]). Unfortunately, $\text{Lift}^{(2n)}(f_{2k})$ does not admit the ordinary p -stabilization in the sense of Hida. However, it turns out that a slight weaker version $\text{Lift}^{(2n)}(f_{2k})^*$ is sufficient to adapt the ordinary theory. In the same spirit as Skinner-Urban [SU], we refer to it as the “semi-ordinary” p -stabilization of $\text{Lift}^{(2n)}(f_{2k})$. For further details on the topic will be discussed in §4 below.

When $n = 1$, more generally in the same direction, Skinner-Urban [SU] produced a p -adic deformation of the cuspidal automorphic representation of $\text{GSp}_4(\mathbb{A}_{\mathbb{Q}})$ generated by $\text{Lift}^{(2)}(f)$ within the framework of a significant study of the Selmer group $H_f^1(\mathbb{Q}, V_f(k))$ defined by Bloch-Kato [BK], where V_f denotes the p -adic Galois representation associated with f in the sense of Eichler-Shimura and Deligne (cf. [De]). Although they must be not entirely smooth (e.g. we cannot associate Siegel modular forms of genus $2n \geq 4$

with Galois representations so far), similar arithmetic applications of p -adic analytic families would be stimulated by the recent progress on Ikeda's generalization of the Duke-Imamoğlu lifting towards a Langlands functorial lifting of cuspidal automorphic representations of $\mathrm{PGL}_2(\mathbb{A}_K)$ to $\mathrm{Sp}_{4n}(\mathbb{A}_K)$ over a totally real field extension K/\mathbb{Q} (cf. [I3]). Indeed, as a consequence of the period relation for $\mathrm{Lift}^{(2n)}(f)$ that was conjectured by Ikeda [I2] and afterwards was shown by Katsurada and the author [KaK], we may produce a certain kind of congruence properties occurring between $\mathrm{Lift}^{(2n)}(f)$ and some genuine Siegel cusp forms of genus $2n$ under mild conditions, which is very similar to those in [SU] (cf. [Ka2, Ka3], see also Brown [Br] for $n = 1$). This type of congruence properties and their applications were firstly conjectured by Harder [Ha] for the Saito-Kurokawa lifting and by Doi-Hida-Ishii [DHI] for the Doi-Naganuma lifting.

It should be emphasized that our approach based on the description of Fourier expansions is more explicit than the method using the theory of automorphic representations, and hence yields some practical benefit. For instance, the semi-ordinary p -stabilized form $\mathrm{Lift}^{(2n)}(f_{2k})^*$ can be regarded naturally as the Duke-Imamoğlu lifting of f_{2k}^* , although the p -local component of the associated cuspidal automorphic representation of $\mathrm{Sp}_{4n}(\mathbb{A}_{\mathbb{Q}})$ is a quadratic twist of the Steinberg representation in general (cf. [I3]).

Finally, as will be explained in §5, we note that the method we use for the Duke-Imamoğlu lifting is adaptable to the Eisenstein series as well. Indeed, by taking Serre's p -adic analytic family of ordinary p -stabilized Eisenstein series instead of the Hida family, we also obtain a similar result for the Siegel Eisenstein series of genus $2n$, which is closely related to the results due to Takemori [Ta] and Panchishkin [P1]. In addition, as conjectured in [P2], our constructions of p -adic analytic families are also extendable to those of families of Siegel cusp forms of odd genus by means of the Miyawaki lifting (cf. [I2]). The corresponding result will appear elsewhere.

Acknowledgements. The author is deeply grateful to Professor Alexei Panchishkin and Professor Haruzo Hida for their valuable suggestions and comments. Moreover, discussions with Professor Jacques Tilouine, Professor Hidenori Katsurada, Professor Siegfried Böcherer, Professor Tamotsu Ikeda, Professor Marc-Hubert Nicole and Dr. Tomokazu Kashio have been illuminating for him.

Notation. We denote by \mathbb{Z} , \mathbb{Q} , \mathbb{R} , and \mathbb{C} the ring of integers, fields of rational numbers, real numbers and complex numbers, respectively. We put $e(x) = \exp(2\pi\sqrt{-1}x)$ for $x \in \mathbb{C}$. For each rational prime l , we denote by \mathbb{Q}_l , \mathbb{Z}_l and \mathbb{Z}_l^\times the field of l -adic numbers, the ring of l -adic integers and the group of units of \mathbb{Z}_l , respectively. Hereinafter, we fix an algebraic closure $\overline{\mathbb{Q}}_l$ of \mathbb{Q}_l . Let $v_l(*)$ denote the l -adic valuation normalized as $v_l(l) = 1$, and $e_l(*)$ the continuous additive character on $\overline{\mathbb{Q}}_l$ such that $e_l(y) = e(y)$ for all $y \in \mathbb{Q}$.

Throughout the article, we fix an odd prime $p > 5$. From now on, we take the algebraic closure $\overline{\mathbb{Q}}$ of \mathbb{Q} inside \mathbb{C} , and identify it with the image under a fixed embedding $\overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}}_p$ once for all.

For each integer $g \geq 1$, let GSp_{2g} and Sp_{2g} be the \mathbb{Q} -linear algebraic groups introduced as follows:

$$\begin{aligned} \mathrm{GSp}_{2g} &:= \{ M \in \mathrm{GL}_{2g} \mid {}^t M J M = \mu(M) J \text{ for some } \mu(M) \in \mathrm{GL}_1 \}, \\ \mathrm{Sp}_{2g} &:= \{ M \in \mathrm{GSp}_{2g} \mid \mu(M) = 1 \}, \end{aligned}$$

where $J = J_{2g} = \begin{pmatrix} 0_g & 1_g \\ -1_g & 0_g \end{pmatrix}$ with the $g \times g$ unit (resp. zero) matrix 1_g (resp. 0_g). We denote by B_{2g} the standard Borel subgroup of GSp_{2g} , and by P_{2g} the associated Siegel parabolic subgroup, that is, $\mathrm{P}_{2g} = \{ M = \begin{pmatrix} * & * \\ 0_g & * \end{pmatrix} \in \mathrm{GSp}_{2g} \}$. Each real point $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathrm{GSp}_{2g}(\mathbb{R})$ with $A, B, C, D \in \mathrm{Mat}_{g \times g}(\mathbb{R})$ and $\mu(M) > 0$ acts on the Siegel upper-half space

$$\mathfrak{H}_g := \{ Z = X + \sqrt{-1}Y \in \mathrm{Mat}_{g \times g}(\mathbb{C}) \mid {}^t Z = Z, Y > 0 \text{ (positive definite)} \}$$

of genus g via the linear transformation $Z \mapsto M(Z) = (AZ + B)(CZ + D)^{-1}$. Then for a given $\kappa \in \mathbb{Z}$ and a function F on \mathfrak{H}_n , we define an action of M on F by

$$(F|_{\kappa} M)(Z) := \mu(M)^{g\kappa - g(g+1)/2} \det(CZ + D)^{-\kappa} F(M(Z)).$$

For handling Siegel modular forms of genus g , we consider the following congruence subgroups of the Siegel modular group $\mathrm{Sp}_{2g}(\mathbb{Z})$: for each integer $N \geq 1$, put

$$\begin{aligned} \Gamma_0(N) &:= \left\{ M \in \mathrm{Sp}_{2g}(\mathbb{Z}) \mid M \equiv \begin{pmatrix} * & * \\ 0_g & * \end{pmatrix} \pmod{N} \right\}, \\ \Gamma_1(N) &:= \left\{ M \in \mathrm{Sp}_{2g}(\mathbb{Z}) \mid M \equiv \begin{pmatrix} * & * \\ 0_g & 1_g \end{pmatrix} \pmod{N} \right\}. \end{aligned}$$

For each $\kappa \in \mathbb{Z}$, the space $\mathcal{M}_{\kappa}(\Gamma_1(N))^{(g)}$ of (*holomorphic*) *Siegel modular forms* of weight κ on $\Gamma_1(N) \subseteq \mathrm{Sp}_{2g}(\mathbb{Z})$, consists of \mathbb{C} -valued holomorphic functions F on \mathfrak{H}_g satisfying the following conditions:

- (i) $F|_{\kappa} M = F$ for any $M \in \Gamma_1(N)$;
- (ii) For each $M \in \mathrm{Sp}_{2g}(\mathbb{Z})$, the function $F|_{\kappa} M$ possesses a Fourier expansion of the form

$$(F|_{\kappa} M)(Z) = \sum_{T \in \mathrm{Sym}_g^*(\mathbb{Z})} A_T(F|_{\kappa} M) e(\mathrm{trace}(TZ)),$$

where we denote by $\mathrm{Sym}_g^*(\mathbb{Z})$ the dual lattice of $\mathrm{Sym}_g(\mathbb{Z})$, that is, consisting of all half-integral symmetric matrices:

$$\mathrm{Sym}_g^*(\mathbb{Z}) = \{ T = (t_{ij}) \in \mathrm{Sym}_g(\mathbb{Q}) \mid t_{ii}, 2t_{ij} \in \mathbb{Z} \ (1 \leq i < j \leq g) \}.$$

It satisfies that $A_T(F|_{\kappa} M) = 0$ unless $T \geq 0$ (semi positive definite) for all $M \in \mathrm{Sp}_{2g}(\mathbb{Z})$.

A modular form $F \in \mathcal{M}_{\kappa}(\Gamma_1(N))^{(g)}$ is said to be *cuspidal* (or a *cusp form*) if it satisfies a stronger condition $A_T(F|_{\kappa} M) = 0$ unless $T > 0$ for all $M \in$

$\mathrm{Sp}_{2g}(\mathbb{Z})$. We denote by $\mathcal{S}_\kappa(\Gamma_1(N))^{(g)}$ the subspace of $\mathcal{M}_\kappa(\Gamma_1(N))^{(g)}$ consisting of all cusp forms. When $N = 1$, we subsequently write $\mathcal{M}_\kappa(\mathrm{Sp}_{2g}(\mathbb{Z}))$ and $\mathcal{S}_\kappa(\mathrm{Sp}_{2g}(\mathbb{Z}))$ instead of $\mathcal{M}_\kappa(\Gamma_1(1))^{(g)}$ and $\mathcal{S}_\kappa(\Gamma_1(1))^{(g)}$, respectively. For each Dirichlet character χ modulo N , we denote by $\mathcal{M}_\kappa(\Gamma_0(N), \chi)^{(g)}$ the subspace of $\mathcal{M}_\kappa(\Gamma_1(N))^{(g)}$ consisting of all forms F with Nebentypus χ , that is, satisfying the condition

$$F|_\kappa M = \chi(\det D)F \quad \text{for all } M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma_0(N),$$

and put $\mathcal{S}_\kappa(\Gamma_0(N), \chi)^{(g)} := \mathcal{M}_\kappa(\Gamma_0(N), \chi)^{(g)} \cap \mathcal{S}_\kappa(\Gamma_1(N))^{(g)}$. In particular, if $\chi = \chi_0$ is the principal character, we naturally write $\mathcal{M}_\kappa(\Gamma_0(N))^{(g)} = \mathcal{M}_\kappa(\Gamma_0(N), \chi_0)^{(g)}$ and $\mathcal{S}_\kappa(\Gamma_0(N))^{(g)} = \mathcal{S}_\kappa(\Gamma_0(N), \chi_0)^{(g)}$, respectively.

For each $T = (t_{ij}) \in \mathrm{Sym}_g^*(\mathbb{Z})$ and $Z = (z_{ij}) \in \mathfrak{H}_g$, we write

$$q^T := e(\mathrm{trace}(TZ)) = \prod_{i=1}^g q_{ii}^{t_{ii}} \prod_{i < j \leq g} q_{ij}^{2t_{ij}},$$

where $q_{ij} = e(z_{ij})$ ($1 \leq i \leq j \leq n$). Then it follows from the definition that each $F \in \mathcal{M}_\kappa(\Gamma_1(N))^{(g)}$ possesses the usual Fourier expansion

$$F(Z) = \sum_{\substack{T \in \mathrm{Sym}_g^*(\mathbb{Z}), \\ T \geq 0}} A_T(F) q^T \in \mathbb{C}[q_{ij}^{\pm 1} \mid 1 \leq i < j \leq g][[q_{11}, \dots, q_{gg}]].$$

For each ring R , we write $R[[q]]^{(g)} := R[q_{ij}^{\pm 1} \mid 1 \leq i < j \leq g][[q_{11}, \dots, q_{gg}]]$, which is a generalization of Serre's ring $R[[q]]^{(1)} = R[[q]]$ consisting of all formal q -expansions with coefficients in R . In particular, if $F \in \mathcal{M}_\kappa(\Gamma_1(N))^{(g)}$ is a Hecke eigenform (i.e. a simultaneous eigenfunction of all Hecke operators with similitude prime to N), then it is well-known that the field K_F obtained by adjoining all Fourier coefficients of F to \mathbb{Q} is a totally real algebraic field of finite degree, to which we refer as the *Hecke field* of F . Hence we have

$$F \in K_F[[q]]^{(g)} \subset \overline{\mathbb{Q}}[[q]]^{(g)} \hookrightarrow \overline{\mathbb{Q}_p}[[q]]^{(g)}.$$

For further details on Siegel modular forms set out above, see [AnZ] or [Fr].

2. PRELIMINARIES

2.1. Classical Duke-Imamoğlu lifting. In this subsection, we review Ikeda's construction of the Duke-Imamoğlu lifting for elliptic cusp forms (cf. [I1]) and Kohlen's description of its Fourier expansion (cf. [Ko5]).

To begin with, we recall some basic facts on elliptic modular forms of half-integral weight which were initiated by Shimura. For each $M \in \Gamma_0(4) \subset \mathrm{SL}_2(\mathbb{Z})$ and $z \in \mathfrak{H}_1$, put

$$j(M, z) := \frac{\theta_{1/2}(M(z))}{\theta_{1/2}(z)},$$

where $\theta_{1/2}(z) = \sum_{m \in \mathbb{Z}} \mathbf{e}(m^2 z)$ is the standard theta function. It is well-known that $j(M, z)$ for $M \in \Gamma_0(4)$ satisfies a usual 1-cocycle relation, and hence defines a factor of automorphy. Then for each integers $k, N \geq 1$, a \mathbb{C} -valued holomorphic function h on \mathfrak{H}_1 is called a modular form of weight $k + 1/2$ on $\Gamma_0(4N)$ if it satisfies the following conditions similar to those in the integral weight case:

- (i) $(h|_{k+1/2} M)(z) := j(M, z)^{-2k-1} h(M(z)) = h(z)$ for any $M \in \Gamma_0(4N)$;
- (ii) For each $M \in \mathrm{SL}_2(\mathbb{Z})$, the form $h|_{k+1/2} M$ has a Fourier expansion

$$(h|_{k+1/2} M)(z) = \sum_{m=0}^{\infty} c_m(h|_{k+1/2} M) q^m,$$

where $q = e(z)$.

In particular, a modular form h is said to be cuspidal (or a cusp form) if it satisfies the condition $c_0(h|_{k+1/2} M) = 0$ for all $M \in \mathrm{SL}_2(\mathbb{Z})$. We denote by $\mathcal{M}_{k+1/2}(\Gamma_0(4N))^{(1)}$ and $\mathcal{S}_{k+1/2}(\Gamma_0(4N))^{(1)}$ the space consisting of all modular forms of weight $k + 1/2$ on $\Gamma_0(4N)$ and its cuspidal subspace, respectively.

As one of the most significant properties of such forms of half-integral weight, Shimura [S1] established that there exists a Hecke equivariant linear correspondence between $\mathcal{M}_{k+1/2}(\Gamma_0(4N))^{(1)}$ and $\mathcal{M}_{2k}(\Gamma_0(N))^{(1)}$, to which we refer as the *Shimura correspondence*. More precisely, Kohnen introduced the plus spaces $\mathcal{M}_{k+1/2}^+(\Gamma_0(4N))^{(1)}$ and $\mathcal{S}_{k+1/2}^+(\Gamma_0(4N))^{(1)}$ respectively to be the subspaces of $\mathcal{M}_{k+1/2}(\Gamma_0(4N))^{(1)}$ and $\mathcal{S}_{k+1/2}(\Gamma_0(4N))^{(1)}$ consisting of all forms h with

$$c_m(h) = 0 \text{ unless } (-1)^k m \equiv 0 \text{ or } 1 \pmod{4},$$

and showed that if either $k \geq 2$ or $k = 1$ and N is cubefree, the Shimura correspondence gives the diagram of linear isomorphisms

$$\begin{array}{ccc} \mathcal{M}_{k+1/2}^+(\Gamma_0(4N))^{(1)} & \xrightarrow{\cong} & \mathcal{M}_{2k}(\Gamma_0(N))^{(1)} \\ \cup & & \cup \\ \mathcal{S}_{k+1/2}^+(\Gamma_0(4N))^{(1)} & \xrightarrow{\cong} & \mathcal{S}_{2k}(\Gamma_0(N))^{(1)}, \end{array}$$

which is commutative with the actions of Hecke operators (cf. [Ko1, Ko2, Ko3]). When $N = 1$, the Shimura correspondence can be characterized explicitly in terms of Fourier expansions as follows: If $f = \sum_{m=1}^{\infty} a_m(f) q^m \in \mathcal{S}_{2k}(\mathrm{SL}_2(\mathbb{Z}))$ is a Hecke eigenform normalized as $a_1(f) = 1$, and

$$h = \sum_{\substack{m \geq 1, \\ (-1)^k m \equiv 0, 1 \pmod{4}}} c_m(h) q^m \in \mathcal{S}_{k+1/2}^+(\Gamma_0(4))^{(1)}$$

corresponds to f via the Shimura correspondence, then for each fundamental discriminant \mathfrak{d} (i.e. \mathfrak{d} is either 1 or the discriminant of a quadratic field) with

$(-1)^k \mathfrak{d} > 0$, and $1 \leq m \in \mathbb{Z}$, we have

$$(2) \quad c_{|\mathfrak{d}|m^2}(h) = c_{|\mathfrak{d}|}(h) \sum_{d|m} \mu(d) \left(\frac{\mathfrak{d}}{d} \right) d^{k-1} a_{(m/d)}(f),$$

where $\mu(d)$ is the Möbius function, and $\left(\frac{\mathfrak{d}}{d} \right)$ the Kronecker character corresponding to \mathfrak{d} . We note that the inverse correspondence of the Shimura correspondence is determined uniquely up to scalar multiplication. It is because unlike the integral weight case, there is no canonical normalization of half-integral weight forms. Hence we should choose a suitable normalization of it in accordance with the intended use.

Remark 2.1. As will be explained more precisely in §§3.2 below, for integers $N \geq 1$, $k \geq 2$ and a fundamental discriminant \mathfrak{d} with $(-1)^k \mathfrak{d} > 0$, Shintani [Sh] and Kohnen [Ko2] constructed a theta lifting

$$\vartheta_{\mathfrak{d}} : \mathcal{S}_{2k}(\Gamma_0(N))^{(1)} \longrightarrow \mathcal{S}_{k+1/2}^+(\Gamma_0(4N))^{(1)},$$

which gives an inverse correspondence of the Shimura correspondence admitting an algebraic normalization with respect to $(-1)^k \mathfrak{d}$.

On the other hand, for each integers $n, k \geq 1$ with $k > n + 1$ and $n \equiv k \pmod{2}$, we define the (*holomorphic*) *Siegel Eisenstein series* of weight $k+n$ on $\mathrm{Sp}_{4n}(\mathbb{Z})$ as follows: for each $Z \in \mathfrak{H}_{2n}$, put

$$\begin{aligned} E_{k+n}^{(2n)}(Z) &:= 2^{-n} \zeta(1-k-n) \prod_{i=1}^n \zeta(1-2k-2n+2i) \\ &\quad \times \sum_{M=\begin{pmatrix} * & * \\ C & D \end{pmatrix} \in \mathrm{P}_{4n} \cap \mathrm{Sp}_{4n}(\mathbb{Z}) \setminus \mathrm{Sp}_4(\mathbb{Z})} \det(CZ + D)^{-k-n}. \end{aligned}$$

It is well-known that $E_{k+n}^{(2n)}$ is a non-cuspidal Hecke eigenform in $\mathcal{M}_{k+n}(\mathrm{Sp}_{4n}(\mathbb{Z}))$. In addition, for each $T \in \mathrm{Sym}_{2n}^*(\mathbb{Z})$ with $T > 0$, we decompose the associated discriminant $\mathfrak{D}_T := (-1)^n \det(2T)$ into the form

$$\mathfrak{D}_T = \mathfrak{d}_T \mathfrak{f}_T^2$$

with the fundamental discriminant \mathfrak{d}_T corresponding to the quadratic field extension $\mathbb{Q}(\sqrt{\mathfrak{d}_T})/\mathbb{Q}$ and an integer $\mathfrak{f}_T \geq 1$. Then the Fourier coefficient $A_T(E_{k+n}^{(2n)})$ is taken of the following form:

$$(3) \quad A_T(E_{k+n}^{(2n)}) = L\left(1-k, \left(\frac{\mathfrak{d}_T}{*} \right)\right) \prod_{l|\mathfrak{f}_T} F_l(T; l^{k-n-1}),$$

where $L(s, \left(\frac{\mathfrak{d}_T}{*} \right)) := \sum_{m=1}^{\infty} \left(\frac{\mathfrak{d}_T}{m} \right) m^{-s}$, and for each prime l , $F_l(T; X) = F_l^{(2n)}(T; X)$ denotes the polynomial in one variable X with coefficients in \mathbb{Z} appearing in the factorization of the formal power series

$$b_l(T; X) = b_l^{(2n)}(T; X) := \sum_{R \in \mathrm{Sym}_{2n}(\mathbb{Q}_l)/\mathrm{Sym}_{2n}(\mathbb{Z}_l)} e_l(\mathrm{trace}(TR)) X^{v_l(\nu_R)},$$

where $\nu_R = [\mathbb{Z}_l^{2n} + \mathbb{Z}_l^{2n}R : \mathbb{Z}_l^{2n}]$, that is,

$$(4) \quad b_l(T; X) = \frac{(1 - X) \prod_{i=1}^n (1 - l^{2i} X^2)}{1 - \left(\frac{\mathfrak{d}_T}{l}\right) l^n X} F_l(T; X)$$

(cf. [S1, S4, Kil, Fe]). Moreover, it is known that $F_l(T; X)$ is the polynomial of degree $2v_l(\mathfrak{f}_T)$ with $F_l(T; 0) = 1$ and satisfies the functional equation

$$(5) \quad F_l(T; l^{-2n-1} X^{-1}) = (l^{2n+1} X^2)^{-v_l(\mathfrak{f}_T)} F_l(T; X)$$

(cf. [Ka1]). In particular, we have $F_l(T; X) = 1$ if $v_l(\mathfrak{f}_T) = 0$. We easily see that $F_l(uT; X) = F_l(T; X)$ for each $u \in \mathbb{Z}_l^\times$.

Remark 2.2. For each prime l , and for each nondegenerate $T \in \text{Sym}_{2n}(\mathbb{Z}_l)$, the formal power series $b_l(T; X)$ gives rise to the local Siegel series $b_l(T; s) := b_l(T; l^{-s})$ for $s \in \mathbb{C}$ with $\text{Re}(s) > 0$. In particular, for each even integer $\kappa > 2n + 1$, then the value $b_l(T; \kappa)$ coincides with the local density

$$\alpha_l(T, H_{2\kappa}) := \lim_{r \rightarrow \infty} (l^r)^{-4n\kappa + n(2n+1)} \times \#\{U \in \text{Mat}_{2\kappa \times 2n}(\mathbb{Z}_l/l^r \mathbb{Z}_l) \mid {}^t U H_{2\kappa} U - T \in l^r \text{Sym}_{2n}^*(\mathbb{Z}_l)\},$$

where $H_{2\kappa} = \frac{1}{2} \begin{pmatrix} 0_\kappa & 1_\kappa \\ 1_\kappa & 0_\kappa \end{pmatrix}$. In this connection, there have been numerous papers focusing on the local densities of quadratic forms.

Following [I1], we construct the the Duke-Imamoglu lifting as follows:

Theorem I (Theorems 3.2 and 3.3 in [I1]). *Suppose that n, k are positive integers with $n \equiv k \pmod{2}$. Let $f = \sum_{m=1}^\infty a_m(f) q^m \in \mathcal{S}_{2k}(\text{SL}_2(\mathbb{Z}))$ be a normalized Hecke eigenform, and $h = \sum_{m \geq 1} c_m(h) q^m \in \mathcal{S}_{k+1/2}^+(\Gamma_0(4))^{(1)}$ a corresponding Hecke eigenform as in (2). Then for each $0 < T \in \text{Sym}_{2n}^*(\mathbb{Z})$ with discriminant $\mathfrak{D}_T = \mathfrak{d}_T \mathfrak{f}_T^2$, put*

$$(6) \quad A_T(\text{Lift}^{(2n)}(f)) := c_{|\mathfrak{d}_T|}(h) \prod_{l \mid \mathfrak{f}_T} \alpha_l(f)^{v_l(\mathfrak{f}_T)} F_l(T; \beta_l(f) l^{-k-n}),$$

where for each prime l , we denote by $(\alpha_l(f), \beta_l(f))$ the ordered pair of the roots of $X^2 - \alpha_l(f)X + l^{2k-1} = 0$ with $v_l(\alpha_l(f)) \leq v_l(\beta_l(f))$. Then the Fourier expansion

$$\text{Lift}^{(2n)}(f) := \sum_{\substack{T \in \text{Sym}_{2n}^*(\mathbb{Z}), \\ T > 0}} A_T(\text{Lift}^{(2n)}(f)) q^T$$

gives rise to a Hecke eigenform in $\mathcal{S}_{k+n}(\text{Sp}_{4n}(\mathbb{Z}))$ such that

$$L(s, \text{Lift}^{(2n)}(f), \text{st}) = \zeta(s) \prod_{i=1}^{2n} L(s + k + n - i, f).$$

Remark 2.3. For a given Hecke eigenform $F \in \mathcal{M}_{k+n}(\text{Sp}_{4n}(\mathbb{Z}))$ with the Satake parameter $(\psi_0(l), \psi_1(l), \dots, \psi_{2n}(l)) \in (\mathbb{C}^\times)^{2n+1}/W_{2n}$ for each prime

l , then the spinor L -function $L(s, F, \text{spin})$ and the standard L -function $L(s, F, \text{st})$ associated with F are respectively defined as follows:

$$L(s, F, \text{spin}) := \prod_{l < \infty} \left\{ (1 - \psi_0(l)l^{-s}) \prod_{r=1}^{2n} \prod_{1 \leq i_1 < \dots < i_r \leq 2n} (1 - \psi_0(l)\psi_{i_1}(l) \cdots \psi_{i_r}(l)l^{-s}) \right\}^{-1},$$

$$L(s, F, \text{st}) := \prod_{l < \infty} \left\{ (1 - l^{-s}) \prod_{i=1}^{2n} (1 - \psi_i(l)l^{-s})(1 - \psi_i(l)^{-1}l^{-s}) \right\}^{-1},$$

Then it follows from the explicit form of $L(s, \text{Lift}^{(2n)}(f), \text{st})$ and the fundamental equation $\psi_0(l)^2\psi_1(l) \cdots \psi_{2n}(l) = l^{2n(k+n)-n(2n+1)}$ that the Satake parameter of $\text{Lift}^{(2n)}(f)$ is taken as

$$(7) \quad \psi_i(l) = \begin{cases} l^{k-n(n+1)/2} & \text{if } i = 0, \\ \alpha_l(f)l^{-k+i} & \text{if } 1 \leq i \leq n, \\ \beta_l(f)l^{-k-n+i} & \text{if } n+1 \leq i \leq 2n. \end{cases}$$

Hence the spinor L -function $L(s, \text{Lift}^{(2n)}(f), \text{spin})$ can be also written explicitly in terms of the symmetric power L -functions $L(s, f, \text{sym}^r)$ of f with some $0 \leq r \leq n$ (cf. [Mu, Sc2]).

According to the equation (3), we may formally look at the Siegel Eisenstein series $E_{k+n}^{(2n)}$ as the Duke-Imamoğlu lifting of the normalized elliptic Eisenstein series

$$E_{2k}^{(1)} = \frac{\zeta(1-2k)}{2} + \sum_{m=1}^{\infty} \sigma_{2k-1}(m)q^m \in \mathcal{M}_{2k}(\text{SL}_2(\mathbb{Z})),$$

where $\sigma_{2k-1}(m) = \sum_{0 < d|m} d^{2k-1}$. Indeed, we easily see that for each prime l ,

$$(\alpha_l(E_{2k}^{(1)}), \beta_l(E_{2k}^{(1)})) = (1, l^{2k-1}),$$

and it is well-known that Cohen's Eisenstein series $H_{k+1/2} \in \mathcal{M}_{k+1/2}^+(\Gamma_0(4))^{(1)}$ corresponds to $E_{2k}^{(1)}$ via the Shimura correspondence, and possesses the Fourier coefficient

$$c_{|\mathfrak{d}|}(H_{k+1/2}) = L(1-k, \left(\frac{\mathfrak{d}}{*} \right))$$

for each fundamental discriminant \mathfrak{d} with $(-1)^k \mathfrak{d} > 0$ (cf. [Co, EZ]).

We also note that the Duke-Imamoğlu lifting does not vanish identically. Indeed, for each $0 < T \in \text{Sym}_{2n}^*(\mathbb{Z})$ with $\mathfrak{D}_T = \mathfrak{d}_T$ (i.e. $\mathfrak{f}_T = 1$), the equation (6) yields the equation

$$A_T(\text{Lift}^{(2n)}(f)) = c_{|\mathfrak{d}_T|}(h).$$

Hence the non-vanishing of $A_T(\text{Lift}^{(2n)}(f))$ is guaranteed by the well-studied non-vanishing theorem for Fourier coefficients of h as follows:

Lemma 2.4. *For each integer $k \geq 6$, let $f \in \mathcal{S}_{2k}(\text{SL}_2(\mathbb{Z}))$ and $h \in \mathcal{S}_{k+1/2}^+(\Gamma_0(4))^{(1)}$ be taken as above. Then there exists a fundamental discriminant \mathfrak{d} such that $(-1)^k \mathfrak{d} > 0$ and*

$$c_{|\mathfrak{d}|}(h) \neq 0.$$

Moreover, for each prime p , a similar statement remains valid under the additional condition either $\mathfrak{d} \equiv 0 \pmod{p}$ or $\mathfrak{d} \not\equiv 0 \pmod{p}$.

Proof. For each fundamental discriminant \mathfrak{d} with $(-1)^k \mathfrak{d} > 0$, Kohnen-Zagier [KoZ] established the equation

$$(8) \quad \frac{c_{|\mathfrak{d}|}(h)^2}{\|h\|^2} = \frac{(k-1)!}{\pi^k} |\mathfrak{d}|^{k-1/2} \frac{L_{\mathfrak{d}}(k, f)}{\|f\|^2},$$

where $L_{\mathfrak{d}}(s, f) := \sum_{m=1}^{\infty} \left(\frac{\mathfrak{d}}{m}\right) a_m(f) m^{-s}$, and we denote by $\|f\|^2$ and $\|h\|^2$ the Petersson norms square of f and h respectively, that is,

$$\begin{aligned} \|f\|^2 &= \langle f, f \rangle := \int_{\text{SL}_2(\mathbb{Z}) \backslash \mathfrak{H}_1} |f(x + \sqrt{-1}y)|^2 y^{2k-2} dx dy, \\ \|h\|^2 &= \langle h, h \rangle := \frac{1}{6} \int_{\Gamma_0(4) \backslash \mathfrak{H}_1} |h(x + \sqrt{-1}y)|^2 y^{k-3/2} dx dy. \end{aligned}$$

Hence the existence of a fundamental discriminant \mathfrak{d} with desired properties follows immediately from the non-vanishing theorem for $L_{\mathfrak{d}_T}(k, f)$ (cf. [BFH, Wa2]). We complete the proof. \square

For the convenience in the sequel, we describe the Fourier expansion of $\text{Lift}^{(2n)}(f)$ a little more precisely. For each prime l dividing \mathfrak{f}_T , by virtue of the functional equation (5), we have

$$\alpha_l(f)^{v_l(\mathfrak{f}_T)} F_l(T; \beta_l(f) l^{-k-n}) = \beta_l(f)^{v_l(\mathfrak{f}_T)} F_l(T; \alpha_l(f) l^{-k-n}),$$

and hence $\alpha_l(f)^{v_l(\mathfrak{f}_T)} F_l(T; \beta_l(f) l^{-k-n})$ can be written in terms of $\alpha_l(f) + \beta_l(f) = a_l(f)$ and $l^{-k} \alpha_l(f) \beta_l(f) = l^{k-1}$. In fact, Kohnen showed that for each l , we have

$$(9) \quad \alpha_l(f)^{v_l(\mathfrak{f}_T)} F_l(T; \beta_l(f) l^{-k-n}) = \sum_{i=0}^{v_l(\mathfrak{f}_T)} \phi_T(l^{v_l(\mathfrak{f}_T)-i}) (l^{k-1})^{v_l(\mathfrak{f}_T)-i} a_{l^i}(f),$$

with some arithmetic function $\phi_T(d)$ with values in \mathbb{Z} defined for each integer $d \geq 1$ dividing \mathfrak{f}_T (cf. [Ko5]). Hence we obtain another explicit form

$$A_T(\text{Lift}^{(2n)}(f)) = c_{|\mathfrak{d}_T|}(h) \prod_{l|\mathfrak{f}_T} \sum_{i=0}^{v_l(\mathfrak{f}_T)} \phi_T(l^{v_l(\mathfrak{f}_T)-i}) (l^{k-1})^{v_l(\mathfrak{f}_T)-i} a_{l^i}(f).$$

We will make use of this equation as well as (6) in the sequel.

Remark 2.5. For a given $f \in \mathcal{S}_{2k}(\mathrm{SL}_2(\mathbb{Z}))$, Ikeda's construction of $\mathrm{Lift}^{(2n)}(f)$ obviously depends on the choice of $h \in \mathcal{S}_{k+1/2}^+(\Gamma_0(4))^{(1)}$. However, by combining the equation (2) with Kohnen's refinement of the Fourier expansion of $\mathrm{Lift}^{(2n)}(f)$, we may realize the Duke-Imamoğlu lifting as an explicit linear mapping $\mathcal{S}_{k+1/2}^+(\Gamma_0(4))^{(1)} \rightarrow \mathcal{S}_{k+n}(\mathrm{Sp}_{4n}(\mathbb{Z}))$. Moreover, Kohnen-Kojima [KoK] and Yamana [Y] characterized the image of the mapping in terms of a relation between Fourier coefficients, which can be regarded as a generalization of Maass' characterization of the Saito-Kurokawa lifting $\mathrm{Lift}^{(2)}(f)$.

2.2. Λ -adic Siegel modular forms. In this subsection, we introduce the notion of Λ -adic Siegel modular forms of arbitrary genus $g \geq 1$ from point of view of Fourier expansions.

Let $\Gamma = 1 + p\mathbb{Z}_p$ be the maximal torsion-free subgroup of \mathbb{Z}_p^\times . We choose and fix a topological generator $\gamma \in \Gamma$ such that $\Gamma = \gamma^{\mathbb{Z}_p}$. Let $\Lambda = \mathbb{Z}_p[[\Gamma]]$ and $\Lambda_1 = \mathbb{Z}_p[[\mathbb{Z}_p^\times]]$ be the completed group rings on Γ and on \mathbb{Z}_p^\times over \mathbb{Z}_p , respectively. We easily see that Λ_1 has a natural Λ -algebra structure induced from the isomorphism $\Lambda_1 \simeq \Lambda[\mu_{p-1}]$, where μ_{p-1} denotes the maximal torsion subgroup consisting of all $(p-1)$ -th roots of unity.

Remark 2.6. As is well-known, Λ is isomorphic to the power series ring $\mathbb{Z}_p[[X]]$ in one variable X with coefficients in \mathbb{Z}_p under $\gamma \mapsto 1 + X$. In addition, $\mathbb{Z}_p[[X]]$ is isomorphic to the ring $\mathrm{Dist}(\mathbb{Z}_p, \mathbb{Z}_p)$ consisting of all distributions on \mathbb{Z}_p with values in \mathbb{Z}_p . Indeed, every distribution $\mu \in \mathrm{Dist}(\mathbb{Z}_p, \mathbb{Z}_p)$ corresponds to the power series

$$A_\mu(X) = \int_{\mathbb{Z}_p} (1+X)^x d\mu(x) = \sum_{m=0}^{\infty} \int_{\mathbb{Z}_p} \binom{x}{m} d\mu(x) X^m \in \mathbb{Z}_p[[X]],$$

where $\binom{x}{m}$ is the binomial function. Therefore we obtain

$$\Lambda \simeq \mathbb{Z}_p[[X]] \simeq \mathrm{Dist}(\mathbb{Z}_p, \mathbb{Z}_p),$$

which allows us to consider the definition of Λ -adic Siegel modular forms below from a different point of view.

To begin with, we introduce the Λ -adic analytic spaces as an alternative notion for the weights of holomorphic Siegel modular forms in the following:

Definition 2.7 (Λ -adic analytic spaces). For each Λ_1 -algebra R finite flat over Λ , we define the Λ -adic analytic space $\mathfrak{X}(R)$ associated with R as

$$\mathfrak{X}(R) := \mathrm{Hom}_{\mathrm{cont}}(R, \overline{\mathbb{Q}}_p),$$

on which the following arithmetic data are introduced:

- (i) A point $P \in \mathfrak{X}(R)$ is said to be *arithmetic* if there exists an integer $\kappa \geq 2$ such that the restriction of P into $\mathfrak{X}(\Lambda) := \mathrm{Hom}_{\mathrm{cont}}(\Lambda, \overline{\mathbb{Q}}_p) \simeq \mathrm{Hom}_{\mathrm{cont}}(\Gamma, \overline{\mathbb{Q}}_p^\times)$ corresponds to a continuous character $P_\kappa : \Gamma \rightarrow \overline{\mathbb{Q}}_p^\times$

satisfying $P_\kappa(\gamma) = \gamma^\kappa$. We denote by $\mathfrak{X}_{\text{alg}}(R)$ the set consisting of all arithmetic points in $\mathfrak{X}(R)$.

- (ii) An arithmetic point $P \in \mathfrak{X}_{\text{alg}}(R)$ is said to be of *signature* (κ, ε) if there exist an integer $\kappa \geq 2$ and a finite character $\varepsilon : \mathbb{Z}_p^\times \rightarrow \overline{\mathbb{Q}}_p^\times$ such that P lies over the point $P_{\kappa, \varepsilon} \in \mathfrak{X}_{\text{alg}}(\Lambda_1) \simeq \text{Hom}_{\text{cont}}(\mathbb{Z}_p^\times, \overline{\mathbb{Q}}_p^\times)$ corresponding to the character $P_{\kappa, \varepsilon}(y) = y^\kappa \varepsilon(y)$ on \mathbb{Z}_p^\times . For simplicity, we denote such P by $P = (\kappa, \varepsilon)$ and often refer to it as the arithmetic point of *weight* κ with *Nebentypus* $\varepsilon\omega^{-\kappa}$ in the sequel.

We note that $\mathfrak{X}(\Lambda)$ has a natural analytic structure induced from the identification $\text{Hom}_{\text{cont}}(\Lambda, \overline{\mathbb{Q}}_p) \simeq \text{Hom}_{\text{cont}}(\Gamma, \overline{\mathbb{Q}}_p^\times)$. Moreover, for a given R , restrictions to Λ_1 and then to Λ induce a surjective finite-to-one mapping

$$\pi : \mathfrak{X}(R) \twoheadrightarrow \mathfrak{X}(\Lambda_1) \twoheadrightarrow \mathfrak{X}(\Lambda),$$

which allows us to define some analytic charts around all points of $\mathfrak{X}_{\text{alg}}(R)$. Indeed, it is established by Hida that each $P \in \mathfrak{X}_{\text{alg}}(R)$ is unramified over $\mathfrak{X}(\Lambda)$, and consequently there exists a natural local section of π

$$S_P : U_P \subseteq \mathfrak{X}(\Lambda) \rightarrow \mathfrak{X}(R)$$

defined on a neighborhood U_P of $\pi(P)$ such that $S_P(\pi(P)) = P$. These local sections endow $\mathfrak{X}(R)$ with analytic charts around points in $\mathfrak{X}_{\text{alg}}(R)$. For each $P \in \mathfrak{X}_{\text{alg}}(R)$, a function $\mathbf{f} : \mathfrak{U} \subseteq \mathfrak{X}(R) \rightarrow \overline{\mathbb{Q}}_p$ defined on $\mathfrak{U} = S_P(U_P)$ is said to be *analytic* if $\mathbf{f} \circ S_P : U_P \rightarrow \overline{\mathbb{Q}}_p$ is analytic. In parallel, an open subset $\mathfrak{U} \subseteq \mathfrak{X}(R)$ containing some $P \in \mathfrak{X}_{\text{alg}}(R)$ is called an *analytic neighborhood* of P if $\mathfrak{U} = S_P(U_P)$. For instance, we easily see that each element $\mathbf{a} \in R$ gives rise to an analytic function $\mathbf{a} : \mathfrak{X}_{\text{alg}}(R) \rightarrow \overline{\mathbb{Q}}_p$ defined by $\mathbf{a}(P) = P(\mathbf{a})$. In most generality, if $P \in \mathfrak{X}(R)$ is unramified over $\mathfrak{X}(\Lambda)$, then each element $\mathbf{a} \in R_{(P)}$ gives rise to an analytic function defined on some analytic neighborhood of P , where $R_{(P)}$ denotes the localization of R at P , and gives rise to a discrete valuation ring finite and unramified over Λ (cf. Corollary 1.4 in [H1]). According to the custom, we refer to the evaluation $\mathbf{a}(P)$ at $P \in \mathfrak{X}_{\text{alg}}(R)$ as the *specialization* of \mathbf{a} at P .

Following [GS, H3] and [P1], we define the Λ -adic Siegel modular forms as follows:

Definition 2.8 (Λ -adic Siegel modular forms). Let R be a Λ_1 -algebra finite flat over Λ . For each integer $g \geq 1$, pick $P_0 = (\kappa_0, \omega^{\kappa_0}) \in \mathfrak{X}_{\text{alg}}(R)$ with $\kappa_0 > g + 1$. A formal Fourier expansion

$$\mathbf{F} = \sum_{\substack{T \in \text{Sym}_g^*(\mathbb{Z}), \\ T \geq 0}} \mathbf{a}_T q^T \in R_{(P_0)}[[q]]^{(g)}$$

is called a Λ -adic Siegel modular form of genus g and of level 1 if there exists an analytic neighborhood \mathfrak{U}_0 of P_0 such that for each arithmetic point

$P = (\kappa, \omega^\kappa) \in \mathfrak{U}_0$ with $\kappa \geq \kappa_0$, the specialization

$$\mathbf{F}(P) := \sum_T \mathbf{a}_T(P) q^T \in \overline{\mathbb{Q}}_p[[q]]^{(g)}$$

gives rise to the Fourier expansion of a holomorphic Siegel modular form in $\mathcal{M}_\kappa(\Gamma_0(p))^{(g)}$. In particular, a Λ -adic Siegel modular form \mathbf{F} is said to be *cuspidal* (or a *cuspidal form*) if $\mathbf{F}(P) \in \mathcal{S}_\kappa(\Gamma_0(p))^{(g)}$ for almost all $P \in \mathfrak{U}_0$.

If there exists a Λ -adic Siegel modular form $\mathbf{F} \in R_{(P_0)}[[q]]^{(g)}$, then every coefficient $\mathbf{a}_T \in R_{(P_0)}$ of \mathbf{F} gives rise to an analytic function defined on \mathfrak{U}_0 . Hence every specialization $\mathbf{F}(P)$ gives a holomorphic Siegel modular form whose Fourier coefficients are p -adic analytic functions on \mathfrak{U}_0 . In this context, we mean by a *p -adic analytic family* the infinite collection of holomorphic Siegel modular forms $\{\mathbf{F}(P) \in \mathcal{M}_\kappa(\Gamma_0(p))^{(g)}\}$ parametrized by varying arithmetic points $P = (\kappa, \omega^\kappa) \in \mathfrak{U}_0$. In addition, by identifying such $P \in \mathfrak{U}_0$ with the element $(\kappa, \kappa \pmod{p-1})$ in Serre's p -adic weight space

$$\mathbb{Z}_p \times \mathbb{Z}/(p-1)\mathbb{Z} \simeq \varprojlim_{m \geq 1} \mathbb{Z}/(p-1)p^{m-1}\mathbb{Z},$$

we may also regard $\{\mathbf{F}(P)\}$ as a usual p -adic analytic family parametrized by varying integral weights $\kappa \geq \kappa_0$ with $\kappa \equiv \kappa_0 \pmod{(p-1)p^{m-1}}$ for some sufficiently large $m \geq 1$.

Remark 2.9. On purpose to construct a non-zero Λ -adic Siegel modular form \mathbf{F} , we should take a $P_0 = (\kappa_0, \omega^{\kappa_0}) \in \mathfrak{X}_{\text{alg}}(R)$ having a smallest possible $\kappa_0 \in \mathbb{Z}$ such that $\mathbf{F}(P_0)$ coincides with an actual holomorphic Siegel modular form $F_{\kappa_0} \in \mathcal{M}_{\kappa_0}(\Gamma_0(p))^{(g)}$. For this reason, the condition $\kappa_0 > g + 1$ will be practically required in the subsequent arguments for the Duke-Imamoğlu lifting and the holomorphic Siegel Eisenstein series. Indeed, this is neither more nor less than the condition of holomorphy of the Siegel Eisenstein series of genus g . However, even in the same context, it should better to assume a more general condition $\kappa_0 \geq g + 1$, which is evident from the fact that the smallest possible weight for holomorphic Siegel modular forms of genus g occurring in the de Rham cohomology is $g + 1$.

3. CUSPIDAL Λ -ADIC MODULAR FORMS OF GENUS 1

In this section, we review Hida's construction of a cuspidal Λ -adic modular form of genus 1 and of tame level 1, and then we apply the Stevens' Λ -adic Shintani lifting for it in order to get a p -adic analytic family of half-integral weight forms corresponding to Hida's family via the Shimura correspondence.

3.1. Hida's universal ordinary p -stabilized newforms. For each integer $r \geq 1$, let $X_1(p^r) = \Gamma_1(p^r) \backslash \mathfrak{H}_1 \cup \mathbb{P}^1(\mathbb{Q})$ be the compactified modular curve, and $V_r = H^1(X_1(p^r), \mathbb{Z}_p)$ the simplicial cohomology group of $X_1(p^r)$ with values in \mathbb{Z}_p . It is well-known that V_r is canonically isomorphic to the parabolic cohomology group $H_{\text{par}}^1(\Gamma_1(p^r), \mathbb{Z}_p) \subseteq H^1(\Gamma_1(p^r), \mathbb{Z}_p)$, which is

defined to be the image of the compact-support cohomology group under the natural map (cf. [S3]). We denote the *abstract Λ -adic Hecke algebra* of tame level 1 by the free polynomial algebra

$$\mathbb{T} := \Lambda_1[\mathbb{T}_m \mid 1 \leq m \in \mathbb{Z}]$$

generated by \mathbb{T}_m over Λ_1 . Since $\mathbb{T} \simeq \mathbb{Z}_p[\mathbb{T}_m, \mathbb{Z}_p^\times]$, a natural action of \mathbb{T} on V_r is defined by regarding the generator \mathbb{T}_m acts via the m -th Hecke correspondence and elements of \mathbb{Z}_p^\times act via the usual Nebentypus actions. For each pair of positive integers (r_1, r_2) with $r_1 \geq r_2$, the natural inclusion $\Gamma_1(p^{r_1}) \hookrightarrow \Gamma_1(p^{r_2})$ induces the corestriction $V_{r_1} \rightarrow V_{r_2}$, which commutes with the action of \mathbb{T} . Hence we may consider the projective limit

$$V_\infty := \varprojlim_{r \geq 1} V_r$$

with a \mathbb{T} -algebra structure. We denote by V_∞^{ord} the direct factor of V_∞ cut out by the ordinary idempotent $e_{\text{ord}} = \lim_{m \rightarrow \infty} \mathbb{T}_p^{m!}$, that is,

$$V_\infty^{\text{ord}} = e_{\text{ord}} \cdot V_\infty,$$

on which \mathbb{T}_p acts invertibly. We note that V_∞^{ord} is a Λ -algebra free of finite rank (cf. Theorem 3.1 in [H1]). Moreover, let $\mathcal{L} = \text{Frac}(\Lambda)$ be the fractional field of Λ , Hida constructed an idempotent e_{prim} in the image of $\mathbb{T} \otimes_\Lambda \mathcal{L}$ in $\text{End}_{\mathcal{L}}(V_\infty^{\text{ord}} \otimes_\Lambda \mathcal{L})$, which can be regarded as an analogue of the projection to the space of primitive Hecke eigenforms in Atkin-Lehner theory (cf. [H1], pp.250, 252). Then we define the *universal ordinary parabolic cohomology group* of tame level 1 by the \mathbb{T} -algebra

$$\mathbb{V}^{\text{ord}} := V_\infty^{\text{ord}} \cap e_{\text{prim}}(V_\infty^{\text{ord}} \otimes_\Lambda \mathcal{L}),$$

which is a reflexive Λ -algebra of finite rank and is consequently locally free of finite rank over Λ . Then the *universal p -ordinary Hecke algebra* of tame level 1 is defined to be the image R^{ord} of \mathbb{T} in $\text{End}_{\Lambda_1}(\mathbb{V}^{\text{ord}})$ under the homomorphism

$$h : \mathbb{T} \rightarrow \text{End}_{\Lambda_1}(\mathbb{V}^{\text{ord}}).$$

We note that R^{ord} is naturally equipped with a formal q -expansion

$$(10) \quad \mathbf{f}_{\text{ord}} = \sum_{m=1}^{\infty} \mathbf{a}_m q^m \in R^{\text{ord}}[[q]], \quad \mathbf{a}_m = h(\mathbb{T}_m),$$

which is called the *universal p -stabilized ordinary form* of tame level 1.

Next, we introduce the global data to be interpolated by \mathbf{f}_{ord} as follows:

Definition 3.1 (ordinary p -stabilized newforms). For given integers $\kappa \geq 2$ and $r \geq 1$, a Hecke eigenform $f_\kappa^* \in \mathcal{S}_\kappa(\Gamma_1(p^r))^{(1)}$ is called an *ordinary p -stabilized newform* if one of the following conditions holds true:

- (i) f_κ^* is a p -ordinary Hecke eigenform in $\mathcal{S}_\kappa^{\text{new}}(\Gamma_1(p^r))^{(1)}$, where we denote by $\mathcal{S}_\kappa^{\text{new}}(\Gamma_1(p^r))^{(1)}$ the subspace consisting of all newforms in $\mathcal{S}_\kappa(\Gamma_1(p^r))^{(1)}$.

(ii) If $r = 1$, then there exists a normalized ordinary Hecke eigenform $f_\kappa = \sum_{m=1}^{\infty} a_m(f_\kappa)q^m \in \mathcal{S}_\kappa(\mathrm{SL}_2(\mathbb{Z}))$ such that

$$f_\kappa^*(z) = f_\kappa(z) - \beta_p(f_\kappa)f_\kappa(pz) \quad (z \in \mathfrak{H}_1),$$

where $\beta_p(f_\kappa)$ is the non-unit root of $X^2 - a_p(f_\kappa)X + p^{\kappa-1} = 0$.

Remark 3.2. It follows from the definition that ordinary p -stabilized newforms are literally p -ordinary Hecke eigenforms. Indeed, the assertion is trivial in the case of (i). If $f_\kappa^* \in \mathcal{S}_\kappa(\Gamma_1(p))^{(1)}$ is taken as in (ii), then for each prime l , we have

$$a_l(f_\kappa^*) = \begin{cases} a_l(f_\kappa) & \text{if } l \neq p, \\ \alpha_p(f_\kappa) & \text{if } l = p, \end{cases}$$

where $\alpha_p(f_\kappa)$ denotes the p -adic unit appearing in the factorization

$$X^2 - a_p(f_\kappa)X + p^{\kappa-1} = (X - \alpha_p(f_\kappa))(X - \beta_p(f_\kappa)).$$

Hence we have

$$L(s, f_\kappa^*) = L^{(p)}(s, f_\kappa) \cdot (1 - \alpha_p(f_\kappa)p^{-s})^{-1},$$

where $L^{(p)}(s, f_\kappa)$ denotes Hecke's L -function of f_κ with the p -local Euler factor removed. For a given p -ordinary Hecke eigenform in $\mathcal{S}_\kappa(\mathrm{SL}_2(\mathbb{Z}))$, this type of p -adic normalization process selecting half the Euler factor at p is called the *ordinary p -stabilization*. However, we should note that each ordinary p -stabilized newform $f_\kappa^* \in \mathcal{S}_\kappa(\Gamma_1(p))^{(1)}$ is actually an oldform except for $\kappa = 2$. Indeed, if $f_\kappa^* \in \mathcal{S}_\kappa^{\mathrm{new}}(\Gamma_1(p))^{(1)}$, then we have

$$|a_p(f_\kappa^*)| = p^{\kappa/2-1}$$

(cf. Theorem 4.6.17 (ii) in [Mi]). If $\kappa > 2$, this contradicts the assumption that f_κ^* is ordinary at p . Hence we summarize that each ordinary p -stabilized newform is in fact a p -ordinary Hecke eigenform occurring in either $\mathcal{S}_2^{\mathrm{new}}(\Gamma_1(p))^{(1)}$, $\mathcal{S}_\kappa^{\mathrm{old}}(\Gamma_1(p))^{(1)} := \mathcal{S}_\kappa(\Gamma_1(p))^{(1)} - \mathcal{S}_\kappa^{\mathrm{new}}(\Gamma_1(p))^{(1)}$ with $\kappa > 2$, or $\mathcal{S}_\kappa^{\mathrm{new}}(\Gamma_1(p^r))^{(1)}$ with some $\kappa \geq 2$ and $r > 1$.

Then the following theorem has been established by Hida:

Theorem II (cf. Theorem 2.6 in [GS]). *Let r be a fixed positive integer, and $\mathbf{f}_{\mathrm{ord}} = \sum_{m=1}^{\infty} \mathbf{a}_m q^m \in R^{\mathrm{ord}}[[q]]$ the universal ordinary p -stabilized form of tame level 1 introduced above. Then for each $P \in \mathfrak{X}_{\mathrm{alg}}(R^{\mathrm{ord}})$, the specialization*

$$\mathbf{f}_{\mathrm{ord}}(P) = \sum_{m=1}^{\infty} \mathbf{a}_m(P)q^m \in \overline{\mathbb{Q}}_p[[q]]$$

induces a one-to-one correspondence

$$\left\{ P = (\kappa, \varepsilon) \in \mathfrak{X}_{\mathrm{alg}}(R^{\mathrm{ord}}) \mid 2 \leq \kappa \in \mathbb{Z}, \varepsilon : \mathbb{Z}_p^\times \rightarrow \overline{\mathbb{Q}}_p^\times \text{ (finite character)} \right\} \\ \xleftrightarrow{1:1} \left\{ f_\kappa^* \in \mathcal{S}_\kappa(\Gamma_0(p^r), \varepsilon\omega^{-\kappa})^{(1)} \mid \begin{array}{l} \text{ordinary } p\text{-stabilized newform} \\ \text{of tame level 1} \end{array} \right\}.$$

Remark 3.3. Obviously, Theorem II implies that \mathbf{f}_{ord} interpolates ordinary p -stabilized newforms of tame level 1, that is, of level p^r with some integer $r \geq 1$. In most generality, for each integer $N \geq 1$ prime to p , Hida constructed the universal ordinary p -stabilized newform of tame level N interpolating p -ordinary cuspidal Hecke eigenforms of level Np^r .

By applying Theorem II for $r = 1$, each $P = (2k, \omega^{2k}) \in \mathfrak{X}_{\text{alg}}(R^{\text{ord}})$ corresponds to an ordinary p -stabilized newform $\mathbf{f}_{\text{ord}}(P) = f_{2k}^* \in \mathcal{S}_{2k}(\Gamma_0(p))^{(1)}$ associated with a p -ordinary normalized Hecke eigenform $f_{2k} \in \mathcal{S}_{2k}(\text{SL}_2(\mathbb{Z}))$ via the ordinary p -stabilization. However, we note that $\dim_{\mathbb{C}} \mathcal{S}_{2k}(\text{SL}_2(\mathbb{Z})) = 0$ for $k < 6$, and hence $\mathbf{f}_{\text{ord}}(P)$ vanishes identically at $\{P = (2k, \omega^{2k}) \in \mathfrak{X}_{\text{alg}}(R^{\text{ord}}) \mid 1 < k < 6\}$. Therefore, for a fixed $P_0 = (2k_0, \omega^{2k_0}) \in \mathfrak{X}_{\text{alg}}(R^{\text{ord}})$ with $k_0 \geq 6$, we may regard $\mathbf{f}_{\text{ord}} \in R^{\text{ord}}[[q]]$ as a Λ -adic cusp form of genus 1, and we consequently obtain a p -adic analytic family of ordinary p -stabilized newforms $\{\mathbf{f}_{\text{ord}}(P) = f_{2k}^*\}$ parametrized by $P = (2k, \omega^{2k}) \in \mathfrak{X}_{\text{alg}}(R^{\text{ord}})$ with $k_0 \leq k \in \mathbb{Z}$.

In our setting, the choice of $P_0 = (2k_0, \omega^{2k_0})$ having the smallest possible weight $2k_0$ is obviously taken as $k_0 = 6$, that is, P_0 corresponds to Ramanujan’s Δ -function

$$f_{12} = q \prod_{m=1}^{\infty} (1 - q^m)^{24} = \sum_{m=1}^{\infty} \tau(m)q^m \in \mathcal{S}_{12}(\text{SL}_2(\mathbb{Z})).$$

In addition, we may choose any analytic neighborhood \mathfrak{U}_0 of such P_0 in $\mathfrak{X}(R^{\text{ord}})$. Since we will apply some lifting for \mathbf{f}_{ord} in the sequel, the choices of k_0 , P_0 and \mathfrak{U}_0 may vary depending on the intended use. For readers’ convenience, we present a list of ordinary primes with respect to the unique normalized Hecke eigenforms $f_{2k_0} \in \mathcal{S}_{2k_0}(\text{SL}_2(\mathbb{Z}))$ with $k_0 \in \{6, 8, 9, 10, 11, 13\}$, that is, rational primes at which f_{2k_0} is ordinary:

k_0	Ordinary primes for f_{2k_0}
6	$11 \leq p \leq 2399, 2417 \leq p \leq 19597$
8	$17 \leq p \leq 53, 61 \leq p \leq 15269, 15277 \leq p \leq 19597$
9	$17 \leq p \leq 14879$
10	$19 \leq p \leq 3361, 3373 \leq p \leq 9973$
11	$p = 11, 23 \leq p \leq 9973$
13	$29 \leq p \leq 9973$

For the smallest ordinary prime $p = 11$ with respect to f_{12} , we give a numerical example of another components of the Hida family:

Example 3.4. Since $6 + (11 - 1) = 16$, we focus on the 2-dimensional space $\mathcal{S}_{32}(\text{SL}_2(\mathbb{Z}))$. Then we may take a normalized Hecke eigenform $f_{32} \in$

$\mathcal{S}_{32}(\mathrm{SL}_2(\mathbb{Z}))$ determined uniquely up to Galois conjugation such that

$$\begin{aligned} f_{32} = & q + xq^2 + (432x + 50220)q^3 + (39960x + 87866368)q^4 \\ & - (1418560x - 18647219790)q^5 + (17312940x + 965671206912)q^6 \\ & - (71928864x - 16565902491320)q^7 - (462815680x - 89324586639360)q^8 \\ & + (7500885120x - 200500912849563)q^9 \\ & - (38038437810x + 3170978118696960)q^{10} \\ & + (29000909200x - 4470615038375388)q^{11} + \cdots \in K_{32}[[q]], \end{aligned}$$

where K_{32} denotes the real quadratic field

$$\mathbb{Q}[x]/(x^2 - 39960x - 2235350016).$$

We easily check that the norm of the difference $a_{11}(f_{12}) - a_{11}(f_{32})$ is factored into

$$2^8 \cdot 3^3 \cdot 5^4 \cdot 11 \cdot 368789 \cdot 99988481 \cdot 7376353157.$$

Therefore we obtain a congruence between f_{12} and f_{32} modulo a prime ideal of the ring of integers $\mathcal{O}_{K_{32}}$ of K_{32} lying over 11, which implies that their ordinary 11-stabilizations f_{12}^* and f_{32}^* reside both in the Hida family for $p = 11$.

3.2. Λ -adic Shintani lifting. As mentioned in the previous §§2.1, we review the construction of an inverse correspondence of the Shimura correspondence in the sense of Shintani [Sh] and Kohnen [Ko4]. Moreover, we introduce a similar lifting for the universal ordinary p -stabilized newform $\mathbf{f}_{\mathrm{ord}}$, which was constructed by Stevens [St].

For simplicity, suppose that $N \geq 1$ is odd squarefree and $k \geq 2$. Let \mathfrak{D} be an integer with $\mathfrak{D} \equiv 0, 1 \pmod{4}$ and $(-1)^k \mathfrak{D} > 0$. We denote by $\mathcal{L}(\mathfrak{D})$ the set of all primitive matrices $Q \in \mathrm{Sym}_2^*(\mathbb{Z})$ with discriminant $-\det(2Q) = \mathfrak{D}$. We may naturally identify each element $Q = \begin{pmatrix} a & b/2 \\ b/2 & c \end{pmatrix} \in \mathcal{L}(\mathfrak{D})$ with an integral binary quadratic form $Q(x, y) = ax^2 + bxy + cy^2$ with $\mathrm{gcd}(a, b, c) = 1$. For simplicity, we write $Q = [a, b, c]$ instead of $\begin{pmatrix} a & b/2 \\ b/2 & c \end{pmatrix}$ in the sequel. We also denote by $\mathcal{L}_N(\mathfrak{D})$ the subset of $\mathcal{L}(\mathfrak{D})$ consisting of all elements $[a, b, c]$ with $a \equiv 0 \pmod{N}$. We easily see that if $\mathfrak{D} \equiv 0 \pmod{N}$, then

$$\mathcal{L}_N(\mathfrak{D}) = \{ [a, b, c] \in \mathcal{L}(\mathfrak{D}) \mid a \equiv b \equiv 0 \pmod{N} \}.$$

We note that the congruence subgroup $\Gamma_0(N) \subseteq \mathrm{SL}_2(\mathbb{Z})$ acts on $\mathcal{L}_N(\mathfrak{D})$ as

$$\begin{aligned} \mathcal{L}_N(\mathfrak{D}) \times \Gamma_0(N) & \longrightarrow \mathcal{L}_N(\mathfrak{D}) \\ (Q, M) & \longmapsto Q \circ M := {}^tMQM, \end{aligned}$$

and we easily see that $\mathcal{L}_N(\mathfrak{D})/\Gamma_0(N)$ is finite. For each $Q = [a, b, c] \in \mathcal{L}_N(\mathfrak{D})$, we associate it with a geodesic cycle C_Q in $\Gamma_0(N) \backslash \mathfrak{H}_1$ that is defined

as the image of the semicircle

$$az^2 + b\operatorname{Re}(z) + c = 0$$

oriented either from left to right (resp. from right to left) if $a > 0$ (resp. $a < 0$) or from $-c/b$ to $\sqrt{-1}\infty$ according as $a \neq 0$ or $a = 0$. Then for each $f \in \mathcal{S}_{2k}(\Gamma_0(N))$, we define a cycle integral associated with f by

$$r_Q(f) := \int_{C_Q} f(z)Q(z, 1)^{k-1} dz.$$

Then the following theorem is given by Shintani and Kohnen:

Theorem III (cf. Theorem 2 in [Ko2]). *Let \mathfrak{d} be a fixed fundamental discriminant with $(-1)^k \mathfrak{d} > 0$. For each $f \in \mathcal{S}_{2k}(\Gamma_0(N))$, put*

$$\vartheta_{\mathfrak{d}}(f) := \sum_{\substack{m \geq 1, \\ (-1)^k m \equiv 0, 1 \pmod{4}}} \left\{ \sum_{0 < d | N} \mu(d) \left(\frac{\mathfrak{d}}{d} \right) d^{k-1} \times \sum_{Q \in \mathcal{L}_{(Nd)}(\mathfrak{d}md^2)/\Gamma_0(Nd)} \chi_{\mathfrak{d}}(Q) r_Q(f) \right\} q^m,$$

where $\chi_{\mathfrak{d}}$ denotes the generalized genus character associated with \mathfrak{d}_0 (cf. [GKZ]). Then we have $\vartheta_{\mathfrak{d}}(f) \in \mathcal{S}_{k+1/2}^+(\Gamma_0(4N))^{(1)}$. Moreover, the mapping

$$\vartheta_{\mathfrak{d}} : \mathcal{S}_{2k}(\Gamma_0(N))^{(1)} \longrightarrow \mathcal{S}_{k+1/2}^+(\Gamma_0(4N))^{(1)}$$

is Hecke equivariant in the sense of the Shimura correspondence.

This type of lifting from integral weight to half-integral weight was firstly introduced by Shintani [Sh], and afterwards was reformulated by Kohnen [Ko2]. According to the custom, we refer to $\vartheta_{\mathfrak{d}}$ as the \mathfrak{d} -th Shintani lifting.

For a given normalized Hecke eigenform $f \in \mathcal{S}_{2k}(\Gamma_0(N))$, we note that all of the Shintani lifting $\vartheta_{\mathfrak{d}}(f)$ give rise to Hecke eigenforms in $\mathcal{S}_{k+1/2}^+(\Gamma_0(4N))^{(1)}$ corresponding to f via the Shimura correspondence, however they differ from each other by the normalization of the Fourier coefficients depending on \mathfrak{d} . Indeed, suppose for simplicity that $N = 1$, $f \in \mathcal{S}_{2k}(\operatorname{SL}_2(\mathbb{Z}))$ is a normalized Hecke eigenform and $h \in \mathcal{S}_{k+1/2}^+(\Gamma_0(4))^{(1)}$ a corresponding Hecke eigenform via the Shimura correspondence as in (2). Then for each integer $m \geq 1$ with $(-1)^k m \equiv 0, 1 \pmod{4}$, we have

$$\begin{aligned} (11) \quad \frac{c_{|\mathfrak{d}|}(h)c_m(h)}{\|h\|^2} &= \frac{(-1)^{\lfloor k/2 \rfloor} 2^k}{\|f\|^2} \sum_{Q \in \mathcal{L}(\mathfrak{d}m)/\operatorname{SL}_2(\mathbb{Z})} \chi_{\mathfrak{d}}(Q) r_Q(f) \\ &= \frac{(-1)^{\lfloor k/2 \rfloor} 2^k}{\|f\|^2} c_m(\vartheta_{\mathfrak{d}}(f)) \end{aligned}$$

(cf. Theorem 3 in [Ko4]). Namely, the choice of \mathfrak{d} determines the normalization datum.

Remark 3.5. We note that the \mathfrak{d} -th Shintani lifting $\vartheta_{\mathfrak{d}}(f)$ admits a nice algebraic property. Indeed, by the equations (8) and (11), we have

$$c_{|\mathfrak{d}|}(\vartheta_{\mathfrak{d}}(f)) = (-1)^{\lfloor k/2 \rfloor} \frac{(k-1)!}{(2\pi)^k} \cdot |\mathfrak{d}|^{k-1/2} L_{\mathfrak{d}}(k, f).$$

On the other hand, by Manin [Ma] and Shimura [S2], we may associate f with two complex periods Ω^+ and Ω^- such that for each critical point $s \in \mathbb{Z}$ with $0 < s < 2k$, the special value $\pi^{-s} L_{\mathfrak{d}}(s, f) / \Omega^{\epsilon}$ resides in the field $K_f(\sqrt{|\mathfrak{d}|})$, where $\epsilon \in \{\pm\}$ is the signature of $(-1)^k \left(\frac{\mathfrak{d}}{-1}\right)$. Therefore we obtain $c_{|\mathfrak{d}|}(\vartheta_{\mathfrak{d}}(f)) / \Omega^{\epsilon} \in K_f(\sqrt{|\mathfrak{d}|})$. Moreover, let \mathcal{O}_f be the ring of integers in K_f . Then by combining Kohnen's theory and a result of Stevens (cf. Proposition 2.3.1 in [St]), we have

$$\frac{1}{\Omega^{\epsilon}} \vartheta_{\mathfrak{d}}(f) \in \mathcal{O}_f[[q]]$$

after taking a suitable normalization.

Now, let us consider a Λ -adic analogue of the Shintani lifting for the universal ordinary p -stabilized newform $\mathbf{f}_{\text{ord}} \in R^{\text{ord}}[[q]]$ (cf. (10) in §§3.1): We define the *metaplectic double covering* of the universal ordinary Hecke algebra R^{ord} by

$$\tilde{R}^{\text{ord}} := R^{\text{ord}} \otimes_{\Lambda_1, \sigma} \Lambda_1,$$

where the tensor product is taken with respect to the ring homomorphism $\sigma : \Lambda_1 \rightarrow \Lambda_1$ corresponding to the group homomorphism $y \mapsto y^2$ on \mathbb{Z}_p^{\times} . We note that \tilde{R}^{ord} has a natural Λ_1 -algebra structure induced from the homomorphism $\lambda \mapsto 1 \otimes \lambda$ on Λ_1 . Therefore we may define the associated Λ -adic analytic space $\mathfrak{X}(\tilde{R}^{\text{ord}})$ and its subset $\mathfrak{X}_{\text{alg}}(\tilde{R}^{\text{ord}})$ of arithmetic points as well as R^{ord} . However, we should mention that the ring homomorphism

$$\begin{array}{ccc} R^{\text{ord}} & \longrightarrow & \tilde{R}^{\text{ord}} \\ \mathbf{a} & \longmapsto & \mathbf{a} \otimes 1 \end{array}$$

is not a Λ_1 -algebra homomorphism. This causes the fact that the mapping induced by pullback on Λ -adic analytic spaces $\mathfrak{X}(\tilde{R}^{\text{ord}}) \rightarrow \mathfrak{X}(R^{\text{ord}})$ does not preserve the signatures of arithmetic points. Indeed, we easily see that if $\tilde{P} = (\kappa, \varepsilon) \in \mathfrak{X}_{\text{alg}}(\tilde{R}^{\text{ord}})$ lies over $P \in \mathfrak{X}_{\text{alg}}(R^{\text{ord}})$, then $P = (2\kappa, \varepsilon^2)$.

Then the following theorem is a refinement of Stevens' Λ -adic Shintani lifting for $\mathbf{f}_{\text{ord}} \in R^{\text{ord}}[[q]]$:

Theorem IV (cf. Theorem 3 in [St]). *For a fixed $P_0 = (2k_0, \omega^{2k_0}) \in \mathfrak{X}_{\text{alg}}(R^{\text{ord}})$ with $k_0 > 1$, let \mathfrak{d}_0 be a fundamental discriminant with $(-1)^{k_0} \mathfrak{d}_0 >$*

0 and $\mathfrak{d}_0 \equiv 0 \pmod{p}$. Then there exist a formal q -expansion

$$\Theta_{\mathfrak{d}_0} = \sum_{m \geq 1} \mathbf{b}(\mathfrak{d}_0; m) q^m \in \tilde{R}^{\text{ord}}[[q]]$$

and a choice of p -adic periods $\Omega_P \in \overline{\mathbb{Q}}_p$ for $P \in \mathfrak{X}_{\text{alg}}(R^{\text{ord}})$ (cf. [GS]) satisfying the following:

- (i) $\Omega_{P_0} \neq 0$.
- (ii) For each $\tilde{P} = (k, \omega^k) \in \mathfrak{X}_{\text{alg}}(\tilde{R}^{\text{ord}})$, the specialization

$$\Theta_{\mathfrak{d}_0}(\tilde{P}) = \sum_{m \geq 1} \mathbf{b}(\mathfrak{d}_0; m)(\tilde{P}) q^m \in \overline{\mathbb{Q}}_p[[q]]$$

gives rise to a holomorphic cusp form in $\mathcal{S}_{k+1/2}^+(\Gamma_0(4p))^{(1)}$. In particular, there exists an analytic neighborhood \mathfrak{U}_0 of P_0 such that for each \tilde{P} lying over $P = (2k, \omega^{2k}) \in \mathfrak{U}_0$, we have

$$\Theta_{\mathfrak{d}_0}(\tilde{P}) = \frac{\Omega_P}{\Omega^\epsilon(P)} \vartheta_{\mathfrak{d}_0}(\mathbf{f}_{\text{ord}}(P)),$$

where $\Omega^\epsilon(P)$ denotes the complex periods of $\mathbf{f}_{\text{ord}}(P)$ with signature $\epsilon \in \{\pm\}$.

We note that a non-vanishing property of $\Theta_{\mathfrak{d}_0}$ is naturally induced from the one of the classical Shintani lifting $\vartheta_{\mathfrak{d}_0}$. Indeed, if $\vartheta_{\mathfrak{d}_0}(\mathbf{f}_{\text{ord}}(P_0))$ is non-zero, then by virtue of the property (i), $\Theta_{\mathfrak{d}_0}$ does not vanish on an analytic neighborhood \mathfrak{U}_0 of P_0 . Hence a suitable choice of $(\mathfrak{d}_0, \mathfrak{U}_0)$ yields a p -adic analytic family of non-zero half-integral weight forms $\{\frac{\Omega_P}{\Omega^\epsilon(P)} \vartheta_{\mathfrak{d}_0}(\mathbf{f}_{\text{ord}}(P))\}$ parametrized by varying arithmetic points $P = (2k, \omega^{2k}) \in \mathfrak{U}_0$ with $k \geq k_0$. For further details on the non-vanishing properties of the classical Shintani lifting and its generalizations, see [Wa1, Wa2].

For each $P = (2k, \omega^{2k}) \in \mathfrak{U}_0$ with $k \geq k_0$, let $f_{2k} \in \mathcal{S}_{2k}(\text{SL}_2(\mathbb{Z}))$ be a p -ordinary normalized Hecke eigenform corresponding to $\mathbf{f}_{\text{ord}}(P) \in \mathcal{S}_{2k}(\Gamma_0(p))^{(1)}$ (i.e. $\mathbf{f}_{\text{ord}}(P) = f_{2k}^*$). Then by Theorem IV, we may also resolve the p -adic interpolation problem for some Fourier coefficients of the classical Shintani lifting $\vartheta_{\mathfrak{d}_0}(f_{2k}) \in \mathcal{S}_{k+1/2}^+(\Gamma_0(4))^{(1)}$.

For such occasions, we prepare the following:

Lemma 3.6. *Suppose that $\mathfrak{D} \equiv 0 \pmod{p}$. Then we have*

- (i) For each $Q \in \mathcal{L}(\mathfrak{D})$, there exists $Q' \in \mathcal{L}_p(\mathfrak{D})$ such that

$$Q' \equiv Q \pmod{\text{SL}_2(\mathbb{Z})}.$$

- (ii) If $\mathfrak{D} \equiv 0 \pmod{p^2}$ and $\left(\frac{\mathfrak{D}/p^2}{p}\right) = 1$, then for each $Q \in \mathcal{L}_p(\mathfrak{D})$, there exists $[a, b, c] \in \mathcal{L}_p(\mathfrak{D})$ such that $a \equiv 0 \pmod{p^3}$ and

$$Q \equiv [a, b, c] \pmod{\Gamma_0(p)}.$$

- (iii) *The identification $[a, b, c] \bmod \Gamma_0(p) \mapsto [a, b, c] \bmod \mathrm{SL}_2(\mathbb{Z})$ induces a one-to-one correspondence between $\mathcal{L}_p(\mathfrak{D})/\Gamma_0(p)$ and $\mathcal{L}_p(\mathfrak{D})/\mathrm{SL}_2(\mathbb{Z})$.*
- (iv) *If $\mathfrak{D} \not\equiv 0 \pmod{p^2}$, then the mapping $[a, b, c] \bmod \Gamma_0(p) \mapsto [p^{-1}a, b, pc] \bmod \mathrm{SL}_2(\mathbb{Z})$ induces a one-to-one correspondence between $\mathcal{L}_p(\mathfrak{D})/\Gamma_0(p)$ and $\mathcal{L}(\mathfrak{D})/\mathrm{SL}_2(\mathbb{Z})$.*

Proof. The assertions (i), (ii), (iii) and (iv) have appeared respectively as Lemmas 1, 3, 2 of [Gu] and Proposition in §I.1 of [GKZ]. For the readers' convenience, we present all of their proofs. For each $[a, b, c] \in \mathcal{L}(\mathfrak{D})$ with $b \not\equiv 0 \pmod{p}$, the assumption $\mathfrak{D} \equiv 0 \pmod{p}$ yields $a \not\equiv 0 \pmod{p}$. Then we put $\beta \equiv -b/(2a) \pmod{p}$ and

$$[a', b', c'] := [a, b, c] \circ \begin{pmatrix} 1 & \beta \\ 0 & 1 \end{pmatrix} \equiv [a, b, c] \pmod{\mathrm{SL}_2(\mathbb{Z})}.$$

We note that $b' = 2a\beta + b \equiv 0 \pmod{p}$. Hence we may assume that $b \equiv 0 \pmod{p}$. If $[a, b, c] \in \mathcal{L}(\mathfrak{D})$ satisfies $a \not\equiv 0 \pmod{p}$, then we easily see that $c \equiv 0 \pmod{p}$. Hence we have

$$[a, b, c] \equiv [a, b, c] \circ \begin{pmatrix} p & p-1 \\ 1 & 1 \end{pmatrix} \in \mathcal{L}_p(\mathfrak{D}) \pmod{\mathrm{SL}_2(\mathbb{Z})}$$

and we obtain the assertion (i). For each $Q = [a, b, c] \in \mathcal{L}_p(\mathfrak{D})$ with $\mathfrak{D} \equiv 0 \pmod{p^2}$, we easily see that $a \equiv 0 \pmod{p^2}$. If $Q' = [a', b', c'] \in \mathcal{L}_p(\mathfrak{D})$ satisfies $Q' = Q \circ \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$ with some $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \Gamma_0(p)$, then the condition $\left(\frac{\mathfrak{D}/p^2}{p}\right) = 1$ yields that the quadratic form $[p^{-2}a, p^{-1}b, c] \in \mathcal{L}_p(\mathfrak{D}/p^2)$ satisfies

$$[p^{-2}a, p^{-1}b, c](\alpha, p^{-1}\gamma) = (p^{-2}a)\alpha^2 + (p^{-1}b)\alpha(p^{-1}\gamma) + c(p^{-1}\gamma)^2 \equiv 0 \pmod{p}.$$

This equation implies the fact that Q' satisfies $p^{-2}a' \equiv 0 \pmod{p}$, and hence we obtain the assertion (ii). For the proof of the assertion (iii), it suffices to show the injectivity of the mapping $[a, b, c] \bmod \Gamma_0(p) \mapsto [a, b, c] \bmod \mathrm{SL}_2(\mathbb{Z})$.

Indeed, for $[a, b, c], [a', b', c'] \in \mathcal{L}_p(\mathfrak{D})$, if there exists $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$ such that

$$[a', b', c'] = [a, b, c] \circ \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix},$$

then we have $a' = a\alpha^2 + b\alpha\gamma + c\gamma^2$. Since $a, a', b \equiv 0 \pmod{p}$ and $\gcd(c, p) = 1$, we obtain $\gamma \equiv 0 \pmod{p}$. Namely, $[a, b, c] \equiv [a', b', c'] \pmod{\Gamma_0(p)}$. In order to prove the assertion (iv), it also suffices to show that the injectivity of the mapping $[a, b, c] \bmod \Gamma_0(p) \mapsto [p^{-1}a, b, pc] \bmod \mathrm{SL}_2(\mathbb{Z})$. Indeed, suppose that $[a, b, c], [a', b', c'] \in \mathcal{L}_p(\mathfrak{D})$ satisfy

$$[p^{-1}a', b', pc'] = [p^{-1}a, b, pc] \circ \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$$

with $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$. The assumption $\mathfrak{D} \not\equiv 0 \pmod{p^2}$ yields $a \not\equiv 0 \pmod{p^2}$. Since $pc' = p^{-1}a\beta^2 + b\beta\delta + pc\delta^2$, we have $\beta \equiv 0 \pmod{p}$. Then

we have

$$[a', b', c'] = [a, b, c] \circ \begin{pmatrix} \alpha & p^{-1}\beta \\ p\gamma & \delta \end{pmatrix},$$

and hence $[a', b', c'] \equiv [a, b, c] \pmod{\Gamma_0(p)}$. We complete the proof. \square

As a consequence of Theorem IV, we have the following:

Proposition 3.7. *Under the same notation and assumptions as above, let \mathfrak{d}_0 be a fixed fundamental discriminant with $(-1)^{k_0}\mathfrak{d}_0 > 0$, $\mathfrak{d}_0 \equiv 0 \pmod{p}$ and $c_{|\mathfrak{d}_0|}(\vartheta_{\mathfrak{d}_0}(f_{2k_0})) \neq 0$ (cf. Lemma 2.4). Then for each integer $k \geq k_0$ and each fundamental discriminant \mathfrak{d} with $(-1)^k\mathfrak{d} > 0$, there exist an analytic neighborhood \mathfrak{U}_0 of $P_0 = (2k_0, \omega^{2k_0}) \in \mathfrak{X}_{\text{alg}}(\tilde{R}^{\text{ord}})$ and an element $\mathbf{c}(\mathfrak{d}_0; |\mathfrak{d}|) \in (\tilde{R}^{\text{ord}})_{(\tilde{P}_0)}$ such that*

$$\mathbf{c}(\mathfrak{d}_0; |\mathfrak{d}|)(\tilde{P}) = \frac{\Omega_P}{\Omega^\epsilon(P)} \left(1 - \left(\frac{\mathfrak{d}}{p} \right) \beta_p(f_{2k}) p^{-k} \right) c_{|\mathfrak{d}|}(\vartheta_{\mathfrak{d}_0}(f_{2k}))$$

for each $\tilde{P} = (k, \omega^k) \in \mathfrak{X}_{\text{alg}}(\tilde{R}^{\text{ord}})$ lying over $P = (2k, \omega^{2k}) \in \mathfrak{U}_0$.

Proof. The following proof is essentially the same as the one of the main theorem in [Gu]: By virtue of Theorem IV, there exists an element $\mathbf{b}(\mathfrak{d}_0; |\mathfrak{d}|) \in \tilde{R}^{\text{ord}}$ such that for each $\tilde{P} \in \mathfrak{X}_{\text{alg}}(\tilde{R}^{\text{ord}})$ lying over $P = (2k, \omega^{2k}) \in \mathfrak{U}_0$, we have

$$\begin{aligned} \mathbf{b}(|\mathfrak{d}_0|; \mathfrak{d})(\tilde{P}) &= \frac{\Omega_P}{\Omega^\epsilon(P)} c_{|\mathfrak{d}|}(\vartheta_{\mathfrak{d}_0}(f_{2k}^*)) \\ &= \frac{\Omega_P}{\Omega^\epsilon(P)} \sum_{Q \in \mathcal{L}_p(\mathfrak{d}_0|\mathfrak{d}|)/\Gamma_0(p)} \chi_{\mathfrak{d}_0}(Q) r_Q(f_{2k}^*). \end{aligned}$$

(I) Suppose that $\mathfrak{d} \not\equiv 0 \pmod{p}$. Then by (i), (iii) of Lemma 3.6, we have

$$\begin{aligned} \sum_{Q \in \mathcal{L}_p(\mathfrak{d}_0|\mathfrak{d}|)/\Gamma_0(p)} \chi_{\mathfrak{d}_0}(Q) r_Q(f_{2k}^*) &= \sum_{Q \in \mathcal{L}(\mathfrak{d}_0|\mathfrak{d}|)/\text{SL}_2(\mathbb{Z})} \chi_{\mathfrak{d}_0}(Q) r_Q(f_{2k}) \\ &\quad - \beta_p(f_{2k}) \sum_{[a,b,c] \in \mathcal{L}_p(\mathfrak{d}_0|\mathfrak{d}|)/\Gamma_0(p)} \chi_{\mathfrak{d}_0}([a,b,c]) \int_{C_{[a,b,c]}} f_{2k}(pz)(az^2 + bz + c)^{k-1} dz. \end{aligned}$$

Here we easily see that

$$\chi_{\mathfrak{d}_0}([a, b, c]) = \left(\frac{\mathfrak{d}}{p} \right) \chi_{\mathfrak{d}_0}([p^{-1}a, b, pc])$$

for each $[a, b, c] \in \mathcal{L}_p(\mathfrak{d}_0|\mathfrak{d}|)/\Gamma_0(p)$, and hence it follows from (iv) of Lemma 3.6 that

$$\begin{aligned} & \sum_{[a,b,c] \in \mathcal{L}_p(\mathfrak{d}_0|\mathfrak{d}|)/\Gamma_0(p)} \chi_{\mathfrak{d}_0}([a, b, c]) \int_{C_{[a,b,c]}} f_{2k}(pz)(az^2 + bz + c)^{k-1} dz \\ &= \left(\frac{\mathfrak{d}}{p}\right) \sum_{[a,b,c] \in \mathcal{L}(\mathfrak{d}_0|\mathfrak{d}|)/\mathrm{SL}_2(\mathbb{Z})} \chi_{\mathfrak{d}_0}([a, b, c]) \\ & \quad \times \int_{C_{[a,b,c]}} f_{2k}(pz)\{a(pz)^2 + b(pz) + c\}^{k-1} \cdot p^{1-k} dz \\ &= \left(\frac{\mathfrak{d}}{p}\right) p^{-k} \sum_{Q=[a,b,c] \in \mathcal{L}(\mathfrak{d}_0|\mathfrak{d}|)/\mathrm{SL}_2(\mathbb{Z})} \chi_{\mathfrak{d}_0}(Q) r_Q(f_{2k}), \end{aligned}$$

where in the second equation we have made use of the transformation law with respect to $z \mapsto p^{-1}z$. Therefore we obtain that $\mathbf{c}(\mathfrak{d}_0; |\mathfrak{d}|) := \mathbf{b}(\mathfrak{d}_0; |\mathfrak{d}|) \in \tilde{R}^{\mathrm{ord}}$ satisfies the equation

$$\begin{aligned} & \mathbf{c}(\mathfrak{d}_0; |\mathfrak{d}|)(\tilde{P}) \\ &= \frac{\Omega_P}{\Omega^\epsilon(P)} \left(1 - \left(\frac{\mathfrak{d}}{p}\right) \beta_p(f_{2k}) p^{-k}\right) \sum_{Q \in \mathcal{L}(\mathfrak{d}_0|\mathfrak{d}|)/\mathrm{SL}_2(\mathbb{Z})} \chi_{\mathfrak{d}_0}(Q) r_Q(f_{2k}) \\ &= \frac{\Omega_P}{\Omega^\epsilon(P)} \left(1 - \left(\frac{\mathfrak{d}}{p}\right) \beta_p(f_{2k}) p^{-k}\right) c_{|\mathfrak{d}|}(\vartheta_{\mathfrak{d}_0}(f_{2k})). \end{aligned}$$

(II) Suppose that $\mathfrak{d} \equiv 0 \pmod{p}$ and $\left(\frac{\mathfrak{d}_0|\mathfrak{d}|/p^2}{p}\right) = 1$. Similarly to the case (I), in order to show that $\mathbf{c}(\mathfrak{d}_0; |\mathfrak{d}|) := \mathbf{b}(\mathfrak{d}_0; |\mathfrak{d}|)$ satisfies the desired property, it suffices to show the equation

(12)

$$\sum_{[a,b,c] \in \mathcal{L}_p(\mathfrak{d}_0|\mathfrak{d}|)/\Gamma_0(p)} \chi_{\mathfrak{d}_0}([a, b, c]) \int_{C_{[a,b,c]}} f_{2k}(pz)(az^2 + bz + c)^{k-1} dz = 0.$$

Indeed, we easily see that for each $s \in \mathbb{Z}/p\mathbb{Z}$, the cycle integrals on the left-hand side of the equation (12) are invariant under the translation $z \mapsto z + p^{-1}s$, and hence we have

$$\begin{aligned} & \sum_{[a,b,c] \in \mathcal{L}_p(\mathfrak{d}_0|\mathfrak{d}|)/\Gamma_0(p)} \chi_{\mathfrak{d}_0}([a, b, c]) \int_{C_{[a,b,c]}} f_{2k}(pz)(az^2 + bz + c)^{k-1} dz \\ &= p^{-1} \sum_{[a,b,c] \in \mathcal{L}_p(\mathfrak{d}_0|\mathfrak{d}|)/\Gamma_0(p)} \left(\int_{C_{[a,b,c]}} f_{2k}(pz)(az^2 + bz + c)^{k-1} dz \right) \\ & \quad \times \sum_{s \in \mathbb{Z}/p\mathbb{Z}} \chi_{\mathfrak{d}_0}([a, 2(p^{-1}a)s + b, (p^{-2}a)s^2 + (p^{-1}b)s + c]). \end{aligned}$$

Then it follows from (ii) of Lemma 3.6 and a simple calculation that

$$\begin{aligned} & \sum_{s \in \mathbb{Z}/p\mathbb{Z}} \chi_{\mathfrak{d}_0}([a, 2(p^{-1}a)s + b, (p^{-2}a)s^2 + (p^{-1}b)s + c]) \\ &= \left(\frac{\mathfrak{d}_0/p}{a}\right) \sum_{s \in \mathbb{Z}/p\mathbb{Z}} \left(\frac{p}{(p^{-2}a)s + (p^{-1}b)s + c}\right) \\ &= \left(\frac{\mathfrak{d}_0/p}{a}\right) \sum_{s \in \mathbb{Z}/p\mathbb{Z}} \left(\frac{p}{(p^{-1}b)s + c}\right) = 0. \end{aligned}$$

Hence we obtain the desired equation (12). (III) Suppose that $\mathfrak{d} \equiv 0 \pmod{p}$ and $\left(\frac{\mathfrak{d}_0|\mathfrak{d}|/p^2}{p}\right) = -1$. We note that we cannot prove the assertion along the same lines as (I) and (II). However, we may fortunately take an alternative route as follows: We note that by virtue of Lemma 2.4 and the equation (11), there exists a fundamental discriminant \mathfrak{d}_1 with $(-1)^{k_0}\mathfrak{d}_1 > 0$, $\mathfrak{d}_1 \not\equiv 0 \pmod{p}$ and $c_{|\mathfrak{d}_1|}(\vartheta_{\mathfrak{d}_0}(f_{2k_0})) \neq 0$. Then by taking the \mathfrak{d} -th Shintani lifting $\vartheta_{\mathfrak{d}}(f_{2k})$ as well as $\vartheta_{\mathfrak{d}_0}(f_{2k})$, the equation (9) yields

$$\begin{aligned} c_{|\mathfrak{d}|}(\vartheta_{\mathfrak{d}_0}(f_{2k})) &= \frac{c_{|\mathfrak{d}_1|}(\vartheta_{\mathfrak{d}}(f_{2k})) \cdot c_{|\mathfrak{d}_0|}(\vartheta_{\mathfrak{d}_0}(f_{2k}))}{c_{|\mathfrak{d}_1|}(\vartheta_{\mathfrak{d}_0}(f_{2k}))} \\ &= \frac{\left(1 - \left(\frac{\mathfrak{d}_1}{p}\right) \beta_p(f_{2k}) p^{-k}\right) c_{|\mathfrak{d}_1|}(\vartheta_{\mathfrak{d}}(f_{2k})) \cdot c_{|\mathfrak{d}_0|}(\vartheta_{\mathfrak{d}_0}(f_{2k}))}{\left(1 - \left(\frac{\mathfrak{d}_1}{p}\right) \beta_p(f_{2k}) p^{-k}\right) c_{|\mathfrak{d}_1|}(\vartheta_{\mathfrak{d}_0}(f_{2k}))}, \end{aligned}$$

where in the second equation of the above, we note that the p -adic interpolation properties of the numerator and the denominator have been already proved at the previous steps (I) and (II). Therefore, by taking a smaller analytic neighborhood \mathfrak{U}_0 of $P_0 \in \mathfrak{X}_{\text{alg}}(R^{\text{ord}})$ if it's necessary to avoid possible vanishing of the denominator, we obtain that the element

$$\mathbf{c}(\mathfrak{d}_0; |\mathfrak{d}|) := \frac{\mathbf{b}(\mathfrak{d}; |\mathfrak{d}_1|) \cdot \mathbf{b}(\mathfrak{d}_0; |\mathfrak{d}_0|)}{\mathbf{b}(\mathfrak{d}_0; |\mathfrak{d}_1|)} \in (\tilde{R}^{\text{ord}})_{(\tilde{P}_0)}$$

defines a quotient of analytic functions on \mathfrak{U}_0 and satisfies the desired interpolation property. We complete the proof. \square

4. MAIN RESULTS

In this section, we construct a Λ -adic Duke-Imamoğlu lifting for the universal ordinary p -stabilized newform $\mathbf{f}_{\text{ord}} \in R^{\text{ord}}[[q]]$. Therefore, we first introduce a suitable p -stabilization process for the Duke-Imamoğlu lifting of p -ordinary Hecke eigenforms in $\mathcal{S}_{2k}(\text{SL}_2(\mathbb{Z}))$. Moreover, to consider the p -adic interpolation problem in the sequel, we give an explicit form of its Fourier expansion.

Suppose that positive integers n and k are fixed as $n \equiv k \pmod{2}$. Let $f \in \mathcal{S}_{2k}(\text{SL}_2(\mathbb{Z}))$ be a normalized Hecke eigenform that is ordinary

at p , and $h \in \mathcal{S}_{k+1/2}^+(\Gamma_0(4))^{(1)}$ a corresponding Hecke eigenform via the Shimura correspondence. As mentioned in §§2.1, the Duke-Imamoğlu lifting $\text{Lift}^{(2n)}(f) \in \mathcal{S}_{k+n}(\text{Sp}_{4n}(\mathbb{Z}))$ is characterized as a Hecke eigenform such that for each prime l , the Satake parameter $(\psi_0(l), \psi_1(l), \dots, \psi_{2n}(l)) \in (\overline{\mathbb{Q}}^\times)^{2n+1}/W_{2n}$ is written explicitly in terms of the ordered pair $(\alpha_l(f), \beta_l(f))$ satisfying $X^2 - a_l(f)X + l^{2k-1} = (X - \alpha_l(f))(X - \beta_l(f))$ and $v_l(\alpha_l(f)) \leq v_l(\beta_l(f))$ (cf. Remark 2.4). In particular, if $l = p$, the ordinarity condition implies that $\alpha_p(f)$ is a p -adic unit, and hence $v_p(\beta_p(f)) = 2k - 1$. Then we recall that the Hecke polynomial $\Phi_p(Y) \in \overline{\mathbb{Q}}_p^\times[Y]$ associated with $\text{Lift}^{(2n)}(f)$ at p is defined as

$$\Phi_p(Y) := (Y - \psi_0(p)) \prod_{r=1}^{2n} \prod_{1 \leq i_1 < \dots < i_r \leq 2n} (Y - \psi_0(p)\psi_{i_1}(p) \cdots \psi_{i_r}(p)).$$

We note that the equation (7) yields that $\psi_0(p) = p^{nk-n(n+1)/2}$ and the product

$$(13) \quad \psi_0(p) \prod_{i=1}^n \psi_i(p) = \alpha_p(f)^n$$

is a unique unit p -adic root of $\Phi_p(Y)$. Put

$$\begin{aligned} \Phi_p^*(Y) &:= \Phi_p(Y) \cdot \{(Y - \psi_0(p))(Y - \psi_0(p) \prod_{i=1}^n \psi_i(p))\}^{-1} \\ &= \prod_{r=1}^{2n} \prod_{\substack{1 \leq i_1 < \dots < i_r \leq 2n, \\ (i_1, \dots, i_n) \neq (1, \dots, n)}} (Y - \psi_0(p)\psi_{i_1}(p) \cdots \psi_{i_r}(p)), \end{aligned}$$

and

$$\begin{aligned} \Psi_p^*(Y) &:= (Y - \psi_0(p) \left(\prod_{i=1}^n \psi_i(p) \right) \psi_{2n}(p)) \\ &\quad \times \prod_{j=1}^{n-1} (Y - \psi_0(p) \left(\prod_{i=1}^n \psi_i(p) \right) \psi_{2n-j}(p) \psi_{2n-j+1}(p)) \\ &\quad \times (Y - \psi_0(p) \left(\prod_{i=1}^{n-1} \psi_i(p) \right) \psi_{2n}(p)). \end{aligned}$$

Obviously, we have $\Psi_p^*(Y) \mid \Phi_p^*(Y) \mid \Phi_p(Y)$ and it follows from the equation (7) that $\Phi_p^*(Y)$ and $\Psi_p^*(Y)$ can be taken of the forms in Theorem 1.3.

On the other hand, the following Hecke operators at p are fundamental in the p -adic theory of Siegel modular forms of arbitrary genus $g \geq 1$:

$$U_{p,i} := \begin{cases} I_{2g} \operatorname{diag}(\underbrace{1, \dots, 1}_g, \underbrace{p, \dots, p}_g) I_{2g}, & \text{if } i = 0, \\ I_{2g} \operatorname{diag}(\underbrace{1, \dots, 1}_i, \underbrace{p, \dots, p}_{g-i}, \underbrace{p^2, \dots, p^2}_i, \underbrace{p, \dots, p}_{g-i}) I_{2g}, & \text{if } 1 \leq i \leq g-1, \end{cases}$$

where $I_{2g} := \{M \in \operatorname{GSp}_{2g}(\mathbb{Z}_p) \mid M \bmod p \in \operatorname{B}_{2g}(\mathbb{Z}/p\mathbb{Z})\}$. We note that these Hecke operators $U_{p,0}, U_{p,1}, \dots, U_{p,g-1}$ generate the Hecke algebra

$$\mathcal{H}_p(I_{2g}, S_{2g}) := \{I_{2g} M I_{2g} \mid M \in I_{2g} \backslash S_{2g} / I_{2g}\},$$

where $S_{2g} \subset \operatorname{GSp}_{2g}(\mathbb{Q}_p)$ is a semi-group such that

$$[I_{2g} \cap M^{-1} S_{2g} M : I_{2g}], [I_{2g} \cap M^{-1} S_{2g} M : M^{-1} S_{2g} M] < +\infty$$

for all $M \in S_{2g}$. In particular, we are interested in the operator $U_{p,0}$ which plays a central role in these operators. For instance, if $F = \sum_{T \geq 0} A_T(F) q^T \in \mathcal{M}_\kappa(\Gamma_0(N))^{(g)}$ with $\kappa \geq 0$ and $N \geq 1$, the action of $U_{p,0}$ on F admits the Fourier expansion

$$(14) \quad F|_\kappa U_{p,0} = \sum_{T \geq 0} A_{pT}(F) q^T,$$

which can be regarded as a generalization of the Atkin-Lehner U_p -operator acting on elliptic modular forms. Then we easily see that

$$F|_\kappa U_{p,0} \in \begin{cases} \mathcal{M}_\kappa(\Gamma_0(N))^{(g)}, & \text{if } p \mid N, \\ \mathcal{M}_\kappa(\Gamma_0(Np))^{(g)}, & \text{if } p \nmid N, \end{cases}$$

and the action of $U_{p,0}$ commutes with those of all Hecke operators at each prime $l \neq p$. For further details on the operator $U_{p,0}$, see [AnZ, Bö].

Now, we define a p -stabilization of $\operatorname{Lift}^{(2n)}(f)$ by

$$(15) \quad \operatorname{Lift}^{(2n)}(f)^* := \frac{\Psi_p^*(\alpha_p(f)^n)}{\Phi_p^*(\alpha_p(f)^n)} \cdot \operatorname{Lift}^{(2n)}(f)|_{k+n} \Phi_p^*(U_{p,0}).$$

Obviously, the equation (13) yields that

$$\frac{\Psi_p^*(\alpha_p(f)^n)}{\Phi_p^*(\alpha_p(f)^n)} \neq 0.$$

In particular, if $n = 1$, we have $\Psi_p^*(Y) = \Phi_p^*(Y) = (Y - p^k)(Y - \beta_p(f))$, and hence $\Psi_p^*(\alpha_p(f))/\Phi_p^*(\alpha_p(f)) = 1$. It also follows immediately from the above-mentioned properties of $U_{p,0}$ that $\operatorname{Lift}^{(2n)}(f)^* \in \mathcal{S}_{k+n}(\Gamma_0(p))^{(2n)}$ is

a Hecke eigenform such that for each prime $l \neq p$, the Hecke eigenvalues coincide with those of $\text{Lift}^{(2n)}(f)$.

Then the Fourier expansion of $\text{Lift}^{(2n)}(f)^*$ is written explicitly as follows:

Theorem 4.1. *Under the same notation and assumptions as above, if $0 < T \in \text{Sym}_{2n}^*(\mathbb{Z})$ satisfies $\mathfrak{D}_T = \mathfrak{d}_T \mathfrak{f}_T^2$, then we have*

$$\begin{aligned} A_T(\text{Lift}^{(2n)}(f)^*) &= \left(1 - \left(\frac{\mathfrak{d}_T}{p}\right) \beta_p(f) p^{-k}\right) c_{|\mathfrak{d}_T|}(h) \\ &\quad \times \alpha_p(f)^{v_p(\mathfrak{f}_T) + n(n+1)} \prod_{\substack{l|\mathfrak{f}_T, \\ l \neq p}} \alpha_l(f)^{v_l(\mathfrak{f}_T)} F_l(T; l^{-k-n} \beta_l(f)). \end{aligned}$$

Proof. By virtue of the equations (6) and (13), we obtain

$$\begin{aligned} &A_T(\text{Lift}^{(2n)}(f)^*) \\ &= \frac{\Psi_p^*(\alpha_p(f)^n)}{\Phi_p^*(\alpha_p(f)^n)} c_{|\mathfrak{d}_T|}(h) \prod_{\substack{l|\mathfrak{f}_T, \\ l \neq p}} \alpha_l(f)^{v_l(\mathfrak{f}_T)} F_l(T; l^{-k-n} \beta_l(f)) \\ &\quad \times \sum_{j=0}^{2^{2n}-2} (-1)^j s_j \left(\left\{ \psi_0(p) \psi_{i_1}(p) \cdots \psi_{i_r}(p) \mid \begin{array}{l} 1 \leq r \leq 2n, \\ 1 \leq i_1 < \cdots < i_r \leq 2n, \\ (i_1, \dots, i_n) \neq (1, \dots, n) \end{array} \right\} \right) \\ &\quad \times \alpha_p(f)^{v_p(\mathfrak{f}_T) + n(2^{2n}-2) - nj} F_p(p^{2^{2n}-2-j} T; p^{-k-n} \beta_p(f)) \\ &= c_{|\mathfrak{d}_T|}(h) \alpha_p(f)^{v_p(\mathfrak{f}_T) + n(n+1)} \prod_{\substack{l|\mathfrak{f}_T, \\ l \neq p}} \alpha_l(f)^{v_l(\mathfrak{f}_T)} F_l(T; l^{-k-n} \beta_l(f)) \\ &\quad \times \frac{\alpha_p(f)^{-n(n+1)} \Psi_p^*(\alpha_p(f)^n)}{\alpha_p(f)^{-n(2^{2n}-2)} \Phi_p^*(\alpha_p(f)^n)} \\ &\quad \times \sum_{j=0}^{2^{2n}-2} (-1)^j s_j \left(\left\{ \alpha_p(f)^{-n} \psi_0(p) \psi_{i_1}(p) \cdots \psi_{i_r}(p) \mid \begin{array}{l} 1 \leq r \leq 2n, \\ 1 \leq i_1 < \cdots < i_r \leq 2n, \\ (i_1, \dots, i_n) \neq (1, \dots, n) \end{array} \right\} \right) \\ &\quad \times F_p(p^{2^{2n}-2-j} T; p^{-k-n} \beta_p(f)), \end{aligned}$$

where $s_j(\{X_1, \dots, X_{2^{2n}-2}\})$ denotes the j -th elementary symmetric polynomial in variables $X_1, \dots, X_{2^{2n}-2}$. Here by the equation (7), we easily see that $\alpha_p(f)^{-n} \psi_0(p) = p^{n(n+1)/2} \{\beta_p(f) p^{-k-n}\}^n$,

$$\psi_i(p) = \begin{cases} p^{-i} \{\beta_p(f) p^{-k-n}\}^{-1}, & \text{if } 1 \leq i \leq n, \\ p^i \{\beta_p(f) p^{-k-n}\}, & \text{if } n+1 \leq i \leq 2n, \end{cases}$$

and

$$\begin{aligned} & \alpha_p(f)^{-n(n+1)} \Psi_p^*(\alpha_p(f)^n) \\ &= (1 - p^{2n} \{\beta_p(f) p^{-k-n}\}) \prod_{i=1}^n (1 - p^{2n+2i-1} \{\beta_p(f) p^{-k-n}\}^2). \end{aligned}$$

Put

$$\xi_{p,i}(X) := \begin{cases} p^{n(n+1)/2} X^n, & \text{if } i = 0, \\ p^{-i} X^{-1}, & \text{if } 1 \leq i \leq n, \\ p^i X, & \text{if } n+1 \leq i \leq 2n, \end{cases}$$

Hence it suffices to show that the equation

$$\begin{aligned} & (1 - p^{2n} X) \prod_{i=1}^n (1 - p^{2n+2i-1} X^2) \\ & \times \sum_{j=0}^{2^{2n}-2} (-1)^j s_j \left(\left\{ \xi_{p,0}(X) \xi_{p,i_1}(X) \cdots \xi_{p,i_r}(X) \mid \begin{array}{l} 1 \leq r \leq 2n, \\ 1 \leq i_1 < \cdots < i_r \leq 2n, \\ (i_1, \dots, i_r) \neq (1, \dots, r) \end{array} \right\} \right) \\ & \times F_p(p^{2^{2n}-2-j} T; X) \\ &= \left(1 - \left(\frac{\mathfrak{d}_T}{p} \right) p^n X \right) \prod_{r=1}^{2n} \prod_{\substack{1 \leq i_1 < \cdots < i_r \leq 2n, \\ (i_1, \dots, i_r) \neq (1, \dots, r)}} (1 - \xi_{p,0}(X) \xi_{p,i_1}(X) \cdots \xi_{p,i_r}(X)) \end{aligned}$$

holds for each $T \in \text{Sym}_{2n}^*(\mathbb{Z})$ with $T > 0$. First, we prove the assertion for $n = 1$. In that case, it suffices to show the equation

$$F_p(p^2 T; X) - (p^2 X + p^3 X^2) F_p(pT; X) + p^5 X^3 F_p(T; X) = 1 - \left(\frac{\mathfrak{d}_T}{p} \right) pX.$$

Indeed, for each $T \in \text{Sym}_2^*(\mathbb{Z})$ with $T > 0$, the polynomial $F_p(T; X)$ admits the explicit form

$$\begin{aligned} & F_p(T; X) \\ &= \sum_{i=0}^{v_p(\mathfrak{m}_T)} (p^2 X)^i \left\{ \sum_{j=0}^{v_p(\mathfrak{f}_T)-i} (p^3 X^2)^j - \left(\frac{\mathfrak{d}_T}{p} \right) pX \sum_{j=0}^{v_p(\mathfrak{f}_T)-i-1} (p^3 X^2)^j \right\}, \end{aligned}$$

where $\mathbf{m}_T = \max\{0 < m \in \mathbb{Z} \mid m^{-1}T \in \text{Sym}_2^*(\mathbb{Z})\}$ (cf. [K]). Therefore we have

$$\begin{aligned}
& F_p(p^2T; X) - (p^2X + p^3X^2)F_p(pT; X) + p^5X^3F_p(T; X) \\
&= \sum_{i=0}^{v_p(\mathbf{m}_T)+2} (p^2X)^i \left\{ \sum_{j=0}^{v_p(\mathfrak{f}_T)-i+2} (p^3X^2)^j - \left(\frac{\mathfrak{d}_T}{p}\right) pX \sum_{j=0}^{v_p(\mathfrak{f}_T)-i+1} (p^3X^2)^j \right\} \\
&\quad - \sum_{i=0}^{v_p(\mathbf{m}_T)+1} (p^2X)^{i+1} \left\{ \sum_{j=0}^{v_p(\mathfrak{f}_T)-i+1} (p^3X^2)^j - \left(\frac{\mathfrak{d}_T}{p}\right) pX \sum_{j=0}^{v_p(\mathfrak{f}_T)-i} (p^3X^2)^j \right\} \\
&\quad - \sum_{i=0}^{v_p(\mathbf{m}_T)+1} (p^2X)^i \left\{ \sum_{j=0}^{v_p(\mathfrak{f}_T)-i+1} (p^3X^2)^{j+1} - \left(\frac{\mathfrak{d}_T}{p}\right) pX \sum_{j=0}^{v_p(\mathfrak{f}_T)-i} (p^3X^2)^{j+1} \right\} \\
&\quad + \sum_{i=0}^{v_p(\mathbf{m}_T)} (p^2X)^{i+1} \left\{ \sum_{j=0}^{v_p(\mathfrak{f}_T)-i} (p^3X^2)^{j+1} - \left(\frac{\mathfrak{d}_T}{p}\right) pX \sum_{j=0}^{v_p(\mathfrak{f}_T)-i-1} (p^3X^2)^{j+1} \right\} \\
&= \sum_{i=0}^{v_p(\mathbf{m}_T)+2} (p^2X)^i \left\{ \sum_{j=0}^{v_p(\mathfrak{f}_T)-i+2} (p^3X^2)^j - \left(\frac{\mathfrak{d}_T}{p}\right) pX \sum_{j=0}^{v_p(\mathfrak{f}_T)-i+1} (p^3X^2)^j \right\} \\
&\quad - \sum_{i=1}^{v_p(\mathbf{m}_T)+2} (p^2X)^i \left\{ \sum_{j=0}^{v_p(\mathfrak{f}_T)-i+2} (p^3X^2)^j - \left(\frac{\mathfrak{d}_T}{p}\right) pX \sum_{j=0}^{v_p(\mathfrak{f}_T)-i+1} (p^3X^2)^j \right\} \\
&\quad - \sum_{i=0}^{v_p(\mathbf{m}_T)+1} (p^2X)^i \left\{ \sum_{j=0}^{v_p(\mathfrak{f}_T)-i+1} (p^3X^2)^{j+1} - \left(\frac{\mathfrak{d}_T}{p}\right) pX \sum_{j=0}^{v_p(\mathfrak{f}_T)-i} (p^3X^2)^{j+1} \right\} \\
&\quad + \sum_{i=1}^{v_p(\mathbf{m}_T)+1} (p^2X)^i \left\{ \sum_{j=0}^{v_p(\mathfrak{f}_T)-i+1} (p^3X^2)^{j+1} - \left(\frac{\mathfrak{d}_T}{p}\right) pX \sum_{j=0}^{v_p(\mathfrak{f}_T)-i} (p^3X^2)^{j+1} \right\} \\
&= \left\{ \sum_{j=0}^{v_p(\mathfrak{f}_T)+2} (p^3X^2)^j - \left(\frac{\mathfrak{d}_T}{p}\right) pX \sum_{j=0}^{v_p(\mathfrak{f}_T)+1} (p^3X^2)^j \right\} \\
&\quad - \left\{ \sum_{j=0}^{v_p(\mathfrak{f}_T)+1} (p^3X^2)^{j+1} - \left(\frac{\mathfrak{d}_T}{p}\right) pX \sum_{j=0}^{v_p(\mathfrak{f}_T)} (p^3X^2)^{j+1} \right\} \\
&= \left\{ \sum_{j=0}^{v_p(\mathfrak{f}_T)+2} (p^3X^2)^j - \left(\frac{\mathfrak{d}_T}{p}\right) pX \sum_{j=0}^{v_p(\mathfrak{f}_T)+1} (p^3X^2)^j \right\} \\
&\quad - \left\{ \sum_{j=1}^{v_p(\mathfrak{f}_T)+2} (p^3X^2)^j - \left(\frac{\mathfrak{d}_T}{p}\right) pX \sum_{j=1}^{v_p(\mathfrak{f}_T)+1} (p^3X^2)^j \right\} \\
&= 1 - \left(\frac{\mathfrak{d}_T}{p}\right) pX.
\end{aligned}$$

For each $n > 1$, it follows from the equation (4) that the desired equation is equivalent to the following:

$$\begin{aligned} & \sum_{j=0}^{2^{2n}-2} (-1)^j s_j \left(\left\{ \xi_{p,0}(X) \xi_{p,i_1}(X) \cdots \xi_{p,i_r}(X) \mid \begin{array}{l} 1 \leq r \leq 2n, \\ 1 \leq i_1 < \cdots < i_r \leq 2n, \\ (i_1, \dots, i_r) \neq (1, \dots, r) \end{array} \right\} \right) \\ & \quad \times b_p(p^{2^{2n}-2-j} T; X) \\ & = \frac{(1-X) \prod_{i=1}^n (1-p^{2i} X^2) \prod_{r=1}^{2n} \prod_{\substack{1 \leq i_1 < \cdots < i_r \leq 2n, \\ (i_1, \dots, i_r) \neq (1, \dots, r)}} (1 - \xi_{p,0}(X) \xi_{p,i_1}(X) \cdots \xi_{p,i_r}(X))}{(1-p^{2n} X) \prod_{i=1}^n (1-p^{2n+2i-1} X^2)}. \end{aligned}$$

This can be proved by making use of the same arguments as in [Zh] and [Ki2] (see also [BöS]). Now we complete the proof. \square

As a consequence of Theorem 4.1, we also have

Corollary 4.2. *Under the same assumption as above, we have*

$$\text{Lift}^{(2n)}(f)^*|_{k+n} U_{p,0} = \alpha_p(f)^n \cdot \text{Lift}^{(2n)}(f)^*.$$

Indeed, the desired equation follows immediately from Theorem 4.1. \square

We should mention that Courtieu-Panchishkin [CP] stated a general philosophy of the p -stabilization for Siegel modular forms. In accordance with it, we may also consider another p -stabilization

$$\begin{aligned} \text{Lift}^{(2n)}(f)^\dagger & := \text{Lift}^{(2n)}(f)|_{k+n} (U_{p,0} - \psi_0(p)) \cdot \Phi_p^*(U_{p,0}) \\ & = \text{Lift}^{(2n)}(f)|_{k+n} \Phi_p(U_{p,0}) \cdot (U_{p,0} - \psi_0(p) \prod_{i=1}^n \psi_i(p))^{-1}. \end{aligned}$$

Since $U_{p,0}$ annihilates the Hecke polynomial $\Phi_p(Y)$ (cf. Proposition 6.10 in [AnZ]), it follows from the equation (13) that

$$\begin{aligned} \text{Lift}^{(2n)}(f)^\dagger|_{k+n} U_{p,0} & = \psi_0(p) \prod_{i=1}^n \psi_i(p) \cdot \text{Lift}^{(2n)}(f)^\dagger \\ & = \alpha_p(f)^n \cdot \text{Lift}^{(2n)}(f)^\dagger. \end{aligned}$$

We easily verify that $U_{p,0}$ annihilates $\Phi_p^*(Y) \cdot (Y - \psi_0(p) \prod_{i=1}^n \psi_i(p))$ as well, and hence we may also prove Corollary 4.2 along the same line as above. Since we have

$$\frac{\Psi_p^*(\alpha_p(f)^n)}{\Phi_p^*(\alpha_p(f)^n)} \cdot \text{Lift}^{(2n)}(f)^\dagger = (1 - p^{nk-n(n+1)/2}) \cdot \text{Lift}^{(2n)}(f)^*,$$

$\text{Lift}^{(2n)}(f)^*$ and $\text{Lift}^{(2n)}(f)^\dagger$ are essentially the same, however, the former can be regarded as the principle p -stabilization for $\text{Lift}^{(2n)}(f)$ and the Siegel Eisenstein series $E_{k+n}^{(2n)}$ (cf. §§5.1 below).

Remark 4.3 (semi-ordinarity). Unfortunately, even if f is ordinary (and so is the associated Galois representation), $\text{Lift}^{(2n)}(f)$ does not admit the ordinary p -stabilization in the sense of Hida (cf. [H4, H5]). That is, $\text{Lift}^{(2n)}(f)$ may not be an eigenfunction of the Hecke operators $U_{p,0}, U_{p,1}, \dots, U_{p,2n-1}$ such that all the eigenvalues are p -adic units. It is because the fact that the p -local spherical representation associated with $\text{Lift}^{(2n)}(f)$ is not equal to any induced representation of an unramified character. On the other hand, Corollary 4.2 implies that there exists a p -stabilization $\text{Lift}^{(2n)}(f)^*$ so that the eigenvalue of $U_{p,0}$ is a p -adic unit. Following [SU], we refer to it as the *semi-ordinary* p -stabilization of $\text{Lift}^{(2n)}(f)$. In fact, it turns out that this condition is sufficient to adapt the ordinary theory. Since the operator $U_{p,0}$ is given by the trace of the Frobenius operator acting on the ordinary part of the cohomology of the Siegel variety, there is no need to use the overconvergence of canonical subgroup.

Now, for the universal ordinary p -stabilized newform $\mathbf{f}_{\text{ord}} = \sum_{m \geq 1} \mathbf{a}_m q^m \in R^{\text{ord}}[[q]]$ interpolating the Hida family $\{f_{2k}^*\}$, we construct its Λ -adic lifting interpolating the semi-ordinary p -stabilized Duke-Imamoglu lifting of f_{2k} :

Theorem 4.4. *For each integer $n \geq 1$, and a fixed integer $k_0 > n + 1$ with $k_0 \equiv n \pmod{2}$, let \mathfrak{d}_0 and \mathfrak{U}_0 be a pair of a fundamental discriminant and an analytic neighborhood of $P_0 \in \mathfrak{X}_{\text{alg}}(R^{\text{ord}})$ taken as in Proposition 3.7. For each integer $k \geq k_0$ with $k \equiv k_0 \pmod{2}$, we denote by $\text{Lift}_{\mathfrak{d}_0}^{(2n)}(f_{2k})$ the Duke-Imamoglu lifting of f_{2k} associated with the \mathfrak{d}_0 -th Shintani lifting $\vartheta_{\mathfrak{d}_0}(f_{2k})$. Then for $\tilde{P}_0 = (k_0, \omega^{k_0}) \in \mathfrak{X}_{\text{alg}}(\tilde{R}^{\text{ord}})$ lying over P_0 , there exist a formal Fourier expansion*

$$\mathbf{F} = \sum_{T > 0} \mathbf{a}_T q^T \in (\tilde{R}^{\text{ord}})_{(\tilde{P}_0)}[[q]]^{(2n)}$$

and a choice of p -adic period $\Omega_P \in \overline{\mathbb{Q}}_p$ for $P \in \mathfrak{U}_0$ satisfying the following:

- (i) $\Omega_{P_0} \neq 0$.
- (ii) For each $\tilde{P} = (k, \omega^k) \in \mathfrak{X}_{\text{alg}}(\tilde{R}^{\text{ord}})$ lying over $P = (2k, \omega^{2k}) \in \mathfrak{U}_0$ with $k \equiv k_0 \pmod{2}$, we have

$$\mathbf{F}(\tilde{P}) = \frac{\Omega_P}{\Omega^\epsilon(P)} \text{Lift}_{\mathfrak{d}_0}^{(2n)}(f_{2k})^*,$$

where $\Omega^\epsilon(P) \in \mathbb{C}^\times$ is the complex period of P with signature $\epsilon \in \{\pm\}$.

Proof. By combining Theorem 4.1 with the equation (9), we have

$$\begin{aligned}
 & A_T(\text{Lift}_{\mathfrak{d}_0}^{(2n)}(f_{2k})^*) \\
 &= \left(1 - \left(\frac{\mathfrak{d}_T}{p}\right) \beta_p(f_{2k})p^{-k}\right) c_{|\mathfrak{d}_T|}(\mathfrak{v}_{\mathfrak{d}_0}(f_{2k})) \\
 &\quad \times \alpha_p(f_{2k})^{v_p(\mathfrak{f}_T)+n(n+1)} \prod_{\substack{l|\mathfrak{f}_T, \\ l \neq p}} \alpha_l(f_{2k})^{v_l(\mathfrak{f}_T)} F_l(T; \beta_l(f_{2k})l^{-k-n}) \\
 &= \left(1 - \left(\frac{\mathfrak{d}_T}{p}\right) \beta_p(f_{2k})p^{-k}\right) c_{|\mathfrak{d}_T|}(\mathfrak{v}_{\mathfrak{d}_0}(f_{2k})) \\
 &\quad \times \alpha_p(f_{2k})^{n(n+1)} \prod_{\substack{l|\mathfrak{f}_T, \\ l \neq p}} \sum_{i=0}^{v_l(\mathfrak{f}_T)} \phi_T(l^{v_l(\mathfrak{f}_T)-i})(l^{v_l(\mathfrak{f}_T)-i})^{k-1} a_{li}(f).
 \end{aligned}$$

We easily see that for each prime $l \neq p$ and each integer $r \geq 0$, there exists an element $\mathbf{d}(l^r) \in \Lambda$ such that $\mathbf{d}(l^r)(P) = l^{r(k-1)}$ for each $P = (2k, \omega^{2k}) \in \mathfrak{X}_{\text{alg}}(R^{\text{ord}})$. Then we define an element $\mathbf{a}_T \in (\tilde{R}^{\text{ord}})_{(\tilde{P}_0)}$ by

$$(16) \quad \mathbf{a}_T := \mathbf{c}(\mathfrak{d}_0; |\mathfrak{d}|) \mathbf{a}_p^{n(n+1)} \prod_{\substack{l|\mathfrak{f}_T, \\ l \neq p}} \sum_{i=0}^{v_l(\mathfrak{f}_T)} \phi_T(l^{v_l(\mathfrak{f}_T)-i}) \mathbf{d}(l^{v_l(\mathfrak{f}_T)-i}) \mathbf{a}_{li},$$

where $\mathbf{c}(\mathfrak{d}_0; |\mathfrak{d}|) \in (\tilde{R}^{\text{ord}})_{(\tilde{P}_0)}$ is the element in Proposition 3.7. Hence the desired interpolation property follows immediately from Theorem II and Proposition 3.7, and we complete the proof. \square

According to the explicit form (16), the Λ -adic Duke-Imamoğlu lifting admits a non-vanishing property as well as the Λ -adic Shintani lifting that we have already established in §§3.2. Therefore we consequently obtain a semi-ordinary p -adic analytic family of non-zero cuspidal Hecke eigenforms $\{\frac{\Omega_P}{\Omega^\epsilon(P)} \text{Lift}_{\mathfrak{d}_0}^{(2n)}(f_{2k})^* \in \mathcal{S}_{k+n}(\Gamma_0(p))^{(2n)}\}$ parametrized by varying $P = (2k, \omega^{2k}) \in \mathfrak{U}_0$ with $k \equiv k_0 \equiv n \pmod{2}$.

Remark 4.5. Recently, Ikeda [I3] generalized the classical Duke-Imamoğlu lifting to a Langlands functorial lifting of cuspidal automorphic representations of $\text{PGL}_2(\mathbb{A}_K)$ to $\text{Sp}_{4n}(\mathbb{A}_K)$ over a totally real algebraic field extension K of \mathbb{Q} , which contains some generalizations of the classical Duke-Imamoğlu lifting for $\mathcal{S}_{2k}(\Gamma_0(N))^{(1)}$ with $N \geq 1$. Unfortunately, the case $N = p$ has been excluded from the framework so far. It is because resulting automorphic representations of $\text{Sp}_{4n}(\mathbb{A}_K)$ are likely to have the Steinberg representations as their p -local components. Therefore we cannot use the Duke-Imamoğlu lifting of $f_{2k}^* \in \mathcal{S}_{2k}(\Gamma_0(p))^{(1)}$ directly. However, by virtue of Theorem 4.1 and Corollary 4.2, we may regard the semi-ordinary p -stabilized

Duke-Imamoğlu lifting $\text{Lift}_{\mathfrak{d}_0}^{(2n)}(f_{2k})^* \in \mathcal{S}_{k+n}(\Gamma_0(p))^{(2n)}$ naturally as a possible generalized Duke-Imamoğlu lifting of $f_{2k}^* \in \mathcal{S}_{2k}^{\text{old}}(\Gamma_0(p))^{(1)}$. The author expects that the theory of the p -adic stabilization for Siegel modular forms is not only interesting in its own right but also useful in the study of classical Siegel modular forms and the associated automorphic representations.

5. APPENDIX

5.1. Λ -adic Siegel Eisenstein series of even genus. As applications of the methods we have used in the previous §4, we give a similar result for the Siegel Eisenstein series of even genus.

Recall that for the Eisenstein series $E_{2k}^{(1)} \in \mathcal{M}_{2k}(\text{SL}_2(\mathbb{Z}))$, the ordinary p -stabilization

$$(E_{2k}^{(1)})^*(z) := E_{2k}^{(1)}(z) - p^{2k-1}E_{2k}^{(1)}(pz) \in \mathcal{M}_{2k}(\Gamma_0(p))^{(1)} \quad (z \in \mathfrak{H}_1)$$

can be assembled into a p -adic analytic family (cf. [H3]). On the other hand, we have mentioned in §§2.1 that for each pair of positive integers n, k with $k > n + 1$ and $k \equiv n \pmod{2}$, the Siegel Eisenstein series $E_{k+n}^{(2n)} \in \mathcal{M}_{k+n}(\text{Sp}_{4n}(\mathbb{Z}))$ can be formally regarded as the Duke-Imamoğlu lifting of $E_{2k}^{(1)}$. Therefore by replacing the Hida family $\{f_{2k}^*\}$ with $\{(E_{2k}^{(1)})^*\}$ in the argument in §4, we may naturally deduce the following:

Theorem 5.1. *For each positive integers n and k with $k > n + 1$ and $k \equiv n \pmod{2}$, put*

$$(E_{k+n}^{(2n)})^* := \frac{\Psi_p^*(1)}{\Phi_p^*(1)} \cdot E_{k+n}^{(2n)}|_{k+n}\Phi_p^*(U_{p,0}),$$

where $\Phi_p^*(Y), \Psi_p^*(Y) \in \overline{\mathbb{Q}}_p^\times[Y]$ denote the polynomials defined in §4, but for $(\alpha_p(E_{2k}^{(1)}), \beta_p(E_{2k}^{(1)})) = (1, p^{2k-1})$. Then we have

- (i) $(E_{k+n}^{(2n)})^*$ is a non-cuspidal Hecke eigenform in $\mathcal{M}_{k+n}(\Gamma_0(p))^{(2n)}$ such that the Hecke eigenvalues agree with $E_{k+n}^{(2n)}$ for each $l \neq p$ and

$$(E_{k+n}^{(2n)})^*|_{k+n}U_{p,0} = (E_{k+n}^{(2n)})^*.$$

- (ii) Let $\mathcal{L} = \text{Frac}(\Lambda)$ be the field of fractions of Λ . There exists a formal Fourier expansion

$$\mathbf{E} = \sum_{T \geq 0} \mathbf{e}_T q^T \in \mathcal{L}[[q]]^{(2n)}$$

such that for each $P_{2k} \in \mathfrak{X}_{\text{alg}}(\Lambda)$ with $k > n + 1$ and $k \equiv n \pmod{2}$, we have

$$\mathbf{E}(P_{2k}) = (E_{k+n}^{(2n)})^*.$$

Proof. By making use of the fundamental properties of $U_{p,0}$ stated in §4, we easily obtain the assertion (i). For each $T \in \text{Sym}_{2n}^*(\mathbb{Z})$ with $T > 0$, by applying Theorem 4.1 for $E_{2k}^{(1)}$ instead of f_{2k} , we have

$$\begin{aligned} A_T((E_{k+n}^{(2n)})^*) &= \left(1 - \left(\frac{\mathfrak{d}_T}{p}\right) p^{k-1}\right) L\left(1 - k, \left(\frac{\mathfrak{d}_T}{*}\right)\right) \prod_{\substack{l \nmid T, \\ l \neq p}} F_l(T; l^{k-n-1}) \\ &= L^{(p)}\left(1 - k, \left(\frac{\mathfrak{d}_T}{*}\right)\right) \prod_{\substack{l \nmid T, \\ l \neq p}} F_l(T; l^{k-n-1}), \end{aligned}$$

where $L^{(p)}(s, \left(\frac{\mathfrak{d}_T}{*}\right)) := L(s, \left(\frac{\mathfrak{d}_T}{*}\right))(1 - \left(\frac{\mathfrak{d}_T}{p}\right) p^{-s})$. If $T \in \text{Sym}_{2n}^*(\mathbb{Z})$ has $\text{rank}(T) = 0$, then it follows from the definition of $E_{k+n}^{(2n)}$ that

$$\begin{aligned} A_T((E_{k+n}^{(2n)})^*) &= \frac{\Psi_p^*(1)}{\Phi_p^*(1)} \cdot \Phi_p^*(1) \cdot A_T(E_{k+n}^{(2n)}) \\ &= 2^{-n} \zeta^{(p)}(1 - k - n) \prod_{i=1}^n \zeta^{(p)}(1 - 2k - 2n + 2i), \end{aligned}$$

where $\zeta^{(p)}(s) := \zeta(s)(1 - p^{-s})$. Moreover, for each integer $0 < r < 2n$, if $T = \begin{pmatrix} 0_{2n-r} & \\ & T_1 \end{pmatrix}$ with some $0 < T_1 \in \text{Sym}_r^*(\mathbb{Z})$, then by virtue of Theorem 1 in [Ki1] (see also Theorem 4.4 in [Ka1]), the Fourier coefficient $A_T(E_{k+n}^{(2n)})$ can be written explicitly in terms of the product of special values of $\zeta(s)$ and the product of $F_l(T_1; l^{k+n-r-1}) = F_l^{(r)}(T_1; l^{k+n-r-1})$ taken over all prime l . Hence we may derive a similar explicit form of $A_T((E_{k+n}^{(2n)})^*)$ in terms of $\zeta^{(p)}(s)$ and $F_l(T_1; l^{k+n-r-1})$ with $l \neq p$. Therefore, the assertion (ii) follows immediately from the constructions of one-variable p -adic L -functions in the sense of Kubota-Leopoldt (cf. [Wi, H3]). \square

Remark 5.2. More generally in the same context, Panchishkin [P1] has already obtained a similar result for the Siegel Eisenstein series of arbitrary genus $g \geq 1$ that is twisted by a cyclotomic character. However, in that case, we may conduce the p -adic interpolation properties of the Fourier coefficients only for $0 < T \in \text{Sym}_g^*(\mathbb{Z})$ with $p \nmid \det(T)$. Fortunately, in our setting, the description of the semi-ordinary p -stabilization allows us to resolve the p -adic interpolation problem for the whole Fourier expansion.

When $n = 1$, Takemori [Ta] independently proved that for each integer $r \geq 0$ and each integer $N \geq 1$ prime to p , a similar result also holds for the Siegel Eisenstein series of genus 2 and of level Np^r with a primitive Nebentypus character. In the present article, we mainly dealt with the Duke-Imamoğlu lifting according to [I1], which requires the conditions $r = 0$ and $N = 1$. Therefore the Siegel Eisenstein series has been discussed under

the same conditions, but for arbitrary even genus $2n \geq 2$. We note that the method we use is also extendable to the Siegel Eisenstein series of even genus in a more general setting, at least, for $N > 1$.

5.2. Numerical evidences. Finally in this section, we present some numerical evidences of the p -adic analytic family $\{\text{Lift}^{(2n)}(f_{2k})^*\}$ which have been produced by using the Wolfram *Mathematica* 7.

Example 5.3. As mentioned in Example 3.4, Ramanujan's Δ -function $f_{12} \in \mathcal{S}_{12}(\text{SL}_2(\mathbb{Z}))$ and a normalized Hecke eigenform $f_{32} \in \mathcal{S}_{32}(\text{SL}_2(\mathbb{Z}))$ can be assembled into the same Hida family for $p = 11$. We easily see that

$$\begin{aligned} h_{13/2} &= q - 56q^4 + 120q^5 - 240q^8 + 9q^9 + 1440q^{12} + \cdots \in \mathbb{Z}[[q]], \\ h_{33/2} &= q + (x - 32768)q^4 + (2x - 65568)q^5 + (218x - 7116672)q^8 \\ &\quad + (432x - 14298687)q^9 - (2916x - 103037184)q^{12} + \cdots \in \mathcal{O}_{K_{32}}[[q]] \end{aligned}$$

give rise to Hecke eigenforms in $\mathcal{S}_{13/2}(\Gamma_0(4))^{(1)}$ and $\mathcal{S}_{33/2}(\Gamma_0(4))^{(1)}$ corresponding respectively to f_{12} and f_{32} via the Shimura correspondence, where $\mathcal{O}_{K_{32}}$ denotes the ring of integers in the real quadratic field K_{32} . By making use of the induction formulas of $F_l(T; X)$ (cf. [Ka1]), we computed Fourier coefficients of $\text{Lift}^{(4)}(f_{12}) \in \mathcal{S}_8(\text{Sp}_8(\mathbb{Z}))$ and $\text{Lift}^{(4)}(f_{32}) \in \mathcal{S}_{18}(\text{Sp}_8(\mathbb{Z}))$ for 4475 half-integral symmetric matrices in $\text{Sym}_4^*(\mathbb{Z})$ according to Nipp's table of equivalent classes of quaternary quadratic forms with discriminant up to 457 (cf. [N]). For simplicity, we denote by

$$[t_{11}, t_{22}, t_{33}, t_{44}, t_{12}, t_{13}, t_{23}, t_{14}, t_{24}, t_{34}]$$

the matrix

$$\begin{pmatrix} t_{11} & t_{12}/2 & t_{13}/2 & t_{14}/2 \\ t_{12}/2 & t_{22} & t_{23}/2 & t_{24}/2 \\ t_{13}/2 & t_{23}/2 & t_{33} & t_{34}/2 \\ t_{14}/2 & t_{24}/2 & t_{34}/2 & t_{44} \end{pmatrix} \in \text{Sym}_4^*(\mathbb{Z}).$$

For each $0 < T \in \text{Sym}_4^*(\mathbb{Z})$ with $\mathfrak{D}_T = \det(2T) = 121$, $\mathfrak{d}_T = 1$ and $\mathfrak{f}_T = 11$, is equivalent to one of the following three representatives:

$$\begin{aligned} T_1 &= [1, 1, 3, 3, 0, 1, 0, 0, 1, 0], & T_2 &= [1, 1, 4, 4, 1, 1, 0, 1, 1, 4], \\ T_3 &= [2, 2, 2, 2, 2, 1, 0, 1, 1, 2]. \end{aligned}$$

By virtue of Theorem 4.1 in [Ka1], we obtain

$$F_{11}(T_i; X) = 1 - 1452X + 161051X^2$$

for $i = 1, 2, 3$. Then we checked that the norm of the difference

$$A_{T_i}(\text{Lift}^{(4)}(f_{12})) - A_{T_i}(\text{Lift}^{(4)}(f_{32})) \quad (i = 1, 2, 3)$$

is factored into

$$2^8 \cdot 3^4 \cdot 5^5 \cdot 11 \cdot 171449 \cdot 680531 \cdot 35058959130397.$$

Moreover, for each $0 < T \in \text{Sym}_4^*(\mathbb{Z})$ with $4 \leq \mathfrak{D}_T \leq 457$ appearing in [N], the norm of $A_T(\text{Lift}^{(4)}(f_{12})) - A_T(\text{Lift}^{(4)}(f_{32}))$ is indeed divisible by $p = 11$.

This fact supports that $\text{Lift}^{(4)}(f_{12})$ is congruent to $\text{Lift}^{(4)}(f_{32})$ modulo a prime ideal of $\mathcal{O}_{K_{32}}$ lying over $p = 11$, and hence their semi-ordinary 11-stabilizations $\text{Lift}^{(4)}(f_{12})^*$ and $\text{Lift}^{(4)}(f_{32})^*$ reside both in the same 11-adic analytic family. Similarly, we also produced another examples of congruences occurring between Fourier coefficients of $\text{Lift}^{(4)}(f_{20}) \in \mathcal{S}_{12}(\text{Sp}_8(\mathbb{Z}))$ and $\text{Lift}^{(4)}(f_{56}) \in \mathcal{S}_{30}(\text{Sp}_8(\mathbb{Z}))$ for $p = 19$.

REFERENCES

- [An] A. Andrianov, *Modular descent and the Saito-Kurokawa conjecture*, Invent. Math. **53** (1979), 267–280.
- [AnZ] A. Andrianov and V. Zhuravlev, *Modular forms and Hecke operators*, Transl. of Math. Monogr., 145. AMS, Providence, RI, 1995.
- [BK] S. Bloch and K. Kato, *L-functions and Tamagawa numbers of motives*, *The Grothendieck Festschrift, Vol. I*, 333–400, Progr. Math., vol. 86, Birkhäuser Boston Inc., Boston, MA, 1990.
- [Bö] S. Böcherer, *On the Hecke operator $U(p)$. With an appendix by Ralf Schmidt*, J. Math. Kyoto Univ. **45** (2005), no.4, 807–829.
- [BöS] S. Böcherer and F. Sato, *Rationality of certain formal power series related to local densities*, Comment. Math. Univ. St.Paul. **36** (1987), No.1, 53–86.
- [Br] J. Brown, *Saito-Kurokawa lifts and applications to the Bloch-Kato conjecture*, Compos. Math. **143** (2007), no. 2, 290–322.
- [BFH] D. Bump, S. Friedberg and J. Hoffstein, *Nonvanishing theorems for L-functions of modular forms and their derivatives*, Invent. Math. **102** (1990), 543–618.
- [Co] H. Cohen, *Sums involving the values at negative integers of L-functions of quadratic characters*, Math. Ann. **217** (1975), 271–285.
- [CP] M. Courtieu and A. Panchishkin, *Non-Archimedean L-Functions and Arithmetical Siegel Modular Forms*, Lecture Notes in Math. vol. 1471, Springer-Verlag, Berlin, 2004.
- [De] P. Deligne, *Formes modulaires et représentations l-adiques*, Séminaire Bourbaki **355** (1969).
- [DHI] K. Doi, H. Hida and H. Ishii, *Discriminant of Hecke fields and twisted adjoint L-values for $GL(2)$* , Invent. Math. **134** (1998), 547–577.
- [EZ] M. Eichler and D. Zagier, *The theory of Jacobi forms*, Progr. Math., vol. 55, Birkhäuser Boston Inc., Boston, MA, 1985.
- [Fe] P. Feit, *Poles and residues of Eisenstein series for symplectic and unitary groups*, Mem. Amer. Math. Soc. **61** (1986), no. 346, iv+89 pp.
- [Fr] E. Freitag, *Siegelsche Modulfunktionen*, Grundlehren Math. Wiss., vol. 254, Springer-Verlag, Berlin, 1983.
- [GKZ] B. Gross, W. Kohnen and D. Zagier, *Heegner points and derivatives of L-series. II*, Math. Ann. **278** (1987), no.1–4, 497–562.
- [GS] R. Greenberg and G. Stevens, *p-adic L-functions and p-adic periods of modular forms*, Invent. Math. **111** (1993), 407–447.
- [Gu] P. Guerzhoy, *On p-adic families of Siegel cusp forms in the Masaß Spezialschar*, J. reine angew. Math. **523** (2000), 103–112.
- [Ha] G. Harder, *Eisensteinkohomologie und die Konstruktion gemischter Motive*, Lecture Notes in Math., vol. 1562, Springer-Verlag, Berlin, 1993.
- [H1] H. Hida, *Galois representations into $GL_2(\mathbb{Z}_p[[X]])$ attached to ordinary cusp forms*, Invent. Math. **85** (1986), no.3, 545–613.
- [H2] H. Hida, *Modules of congruences of Hecke algebras and L-functions associated with cusp forms*, Amer. J. Math. **110** (1988), no. 2, 323–382.

- [H3] H. Hida, *Elementary theory of L-functions and Eisenstein series*, London Math. Soc. Student Text **26** (1993).
- [H4] H. Hida, *Control theorems for coherent sheaves on Shimura varieties of PEL-type*, J. Inst. Math. Jussieu **1** (2002), 1–76.
- [H5] H. Hida, *p-adic automorphic forms on Shimura varieties*, Springer Monographs in Math., Springer-Verlag, New York, 2004.
- [I1] T. Ikeda, *On the lifting of elliptic modular forms to Siegel cusp forms of degree $2n$* , Ann. of Math. **154** (2001), no. 3, 641–681.
- [I2] T. Ikeda, *Pullback of the lifting of elliptic cusp forms and Miyawaki’s conjecture*, Duke Math. J. **131** (2006), no. 3, 469–497.
- [I3] T. Ikeda, *On the lifting of automorphic representations of $\mathrm{PGL}_2(\mathbb{A})$ to $\mathrm{Sp}_{2n}(\mathbb{A})$ or $\mathrm{Sp}_{2n+1}(\mathbb{A})$ over a totally real field*, preprint.
- [K] G. Kaufhold, *Dirichletsche Reihe mit Funktionalgleichung in der Theorie der Modulfunktion 2. Grades*, Math. Ann. **137** (1959), 454–476.
- [Ka1] H. Katsurada, *An explicit formula for Siegel series*, Amer. J. Math. **121** (1999), 415–452.
- [Ka2] H. Katsurada, *Congruence of Siegel modular forms and special values of their standard zeta functions*, Math. Z. **259** (2008), no.1, 97–111.
- [Ka3] H. Katsurada, *Congruence between Duke-Imamoglu-Ikeda lifts and non-Duke-Imamoglu-Ikeda lifts*, preprint.
- [KaK] H. Katsurada and H. Kawamura, *Ikeda’s conjecture on the period of the Duke-Imamoglu-Ikeda lift*, preprint.
- [Ki1] Y. Kitaoka, *Dirichlet series in the theory of quadratic forms*, Nagoya Math. J. **92** (1984), 73–84.
- [Ki2] Y. Kitaoka, *Local densities of quadratic forms and Fourier coefficients of Eisenstein series*, Nagoya Math. J. **103** (1986), 149–160.
- [Ko1] W. Kohnen, *Modular forms of half-integral weight on $\Gamma_0(4)$* , Math. Ann. **248** (1980), 249–266.
- [Ko2] W. Kohnen, *Beziehungen zwischen Modulformen halbganzen Gewichts und Modulformen ganzen Gewichts*, Dissertation Bonn. Math. Schr. **131** (1981).
- [Ko3] W. Kohnen, *Newforms of half-integral weight*, J. reine angew. Math. **333** (1982), 32–72.
- [Ko4] W. Kohnen, *Fourier coefficients of modular forms of half-integral weight*, Math. Ann. **271** (1985), no.2, 237–268.
- [Ko5] W. Kohnen, *Lifting modular forms of half-integral weight to Siegel modular forms of even genus*, Math. Ann. **322** (2002), 787–809.
- [KoK] W. Kohnen and H. Kojima, *A Maass space in higher genus*, Compositio. Math. **141** (2005), 313–322.
- [KoZ] W. Kohnen and D. Zagier, *Values of L-series of modular forms at the center of the critical strip*, Invent. Math. **64** (1981), 175–198.
- [Ku] N. Kurokawa, *Examples of eigenvalues of Hecke operators on Siegel cusp forms of degree two*, Invent. Math. **49** (1978), no. 2, 149–165.
- [M] H. Maass, *Über eine Spezialschar von Modulformen zweiten Grades I, II, III*, Invent. Math. **52** (1979), 95–104; Invent. Math. **53** (1979), 249–253; Invent. Math. **53** (1979), 255–265.
- [Ma] J. Manin, *Periods of cusp forms and p-adic Hecke series*, Mat. Sb. **92** (1973), 378–401; English transl. in Math. USSR-Sb. **21** (1973).
- [Mi] T. Miyake, *Modular forms*, Springer Monogr. in Math., Springer-Verlag, Berlin, 2006.
- [Mu] K. Murakawa, *Relations between symmetric power L-functions and spinor L-functions attached to Ikeda lifts*, Kodai Math. J. **25** (2002), 61–71.

- [N] G. Nipp, *Gordon Nipp's Tables of Quaternary Quadratic Forms* (website), <http://www2.research.att.com/~njas/lattices/nipp.html>
- [P1] A. Panchishkin, *On the Siegel-Eisenstein measure and its applications*, Israel J. Math. **120** (2000), Part B, 467–509.
- [P2] A. Panchishkin, *Two Modularity Lifting Conjectures for Families of Siegel Modular Forms*, Math. Notes **88** (2010), No.4, 25–32.
- [Sc1] R. Schmidt, *The Saito-Kurokawa lifting and functoriality*, Amer. J. Math. **127**(1) (2005), 209–240.
- [Sc2] R. Schmidt, *On the spin L -function of Ikeda's lifts*, preprint.
- [S1] G. Shimura, *On modular forms of half integral weight*, Ann. of Math. **97** (1973), 440–481.
- [S2] G. Shimura, *The periods of certain automorphic forms of arithmetic type*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **28** (1981), No.3, 605–632 (1982).
- [S3] G. Shimura, *Introduction to the arithmetic theory of automorphic functions*, Princeton Univ. Press, Princeton, NJ, 1994.
- [S4] G. Shimura, *Euler products and Fourier coefficients of automorphic forms on symplectic groups*, Invent. Math. **116** (1994), no.1-3, 531–576.
- [Sh] T. Shintani, *On construction of holomorphic cusp forms of half integral weight*, Nagoya Math. J. **58** (1975), 83–126.
- [SU] C. Skinner and E. Urban, *Sur les déformations p -adiques de certaines représentations automorphes*, J. Inst. Math. Jussieu **5** (2006), no. 4, 626–698.
- [St] G. Stevens, *Λ -adic modular forms of half-integral weight and a Λ -adic Shintani lifting*, Contemp. Math. **174** (1994), 129–151.
- [Ta] S. Takemori, *p -adic Siegel-Eisenstein series of degree two*, preprint.
- [TU] J. Tilouine and E. Urban, *Several-variable p -adic families of Siegel-Hilbert cusp eigensystems and their Galois representations*, Ann. Sci. École Norm. Sup. (4) **32** (1999), no.4, 499–574.
- [Wa1] J.-L. Waldspurger, *Correspondence de Shimura*, J. Math. Pures Appl. **59**(1) (1980), 1–132.
- [Wa2] J.-L. Waldspurger, *Correspondences de Shimura et quaternions*, Forum Math. **3** (1991), 219–307.
- [Wi] A. Wiles, *On ordinary λ -adic representations associated to modular forms*, Invent. Math. **94** (1988), No. 3, 529–573.
- [Y] S. Yamana, *Maass relations in higher genus*, Math. Z. **265** (2010), no. 2, 263–276.
- [Z] D. Zagier, *Sur la conjecture de Saito-Kurokawa (d'après H. Maass)*, *Seminar on Number Theory, Paris 1979–80*, Birkhäuser, 371–394, Progr. Math. vol. 12, Birkhäuser Boston, Mass., 1981.
- [Zh] N. A. Zharkovskaya, *The connection between the eigenvalues of Hecke operators and the Fourier coefficients of eigenfunctions for Siegel modular forms of genus n* (Russian), Mat. Sb. (N.S.) **96 (138)** (1975), no.4, 584–593, 646.

(PRIMARY) INSTITUT FOURIER, UFR DE MATHÉMATIQUES, UNIVERSITÉ GRENOBLE I, 100 RUES DES MATHS, BP 74, 38402 SAINT-MARTIN D'HÈRES CEDEX, FRANCE
E-mail address: hisa@fourier.ujf-grenoble.fr

(SECONDARY) DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, KITA 10 NISHI 8, KITA-KU, SAPPORO, 060-0810 HOKKAIDO, JAPAN
E-mail address: kawamura@math.sci.hokudai.ac.jp