



**HAL**  
open science

# Genetic structure of Indian scad mackerel *Decapterus russelli*: Pleistocene vicariance and secondary contact in the central Indo-West Pacific seas

Audrey Rohfritsch, Philippe Borsa

## ► To cite this version:

Audrey Rohfritsch, Philippe Borsa. Genetic structure of Indian scad mackerel *Decapterus russelli*: Pleistocene vicariance and secondary contact in the central Indo-West Pacific seas. *Heredity*, 2005, 95, pp.315-322. 10.1038/sj.hdy.6800727 . hal-00554479

**HAL Id: hal-00554479**

**<https://hal.science/hal-00554479>**

Submitted on 10 Jan 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

To be cited as:

Rohfritsch A., Borsa P. 2005. – Genetic structure of Indian scad mackerel *Decapterus russelli*: Pleistocene vicariance and secondary contact in the central Indo-West Pacific seas. *Heredity* 95, 315-322.

## **Genetic structure of Indian scad mackerel *Decapterus russelli*: Pleistocene vicariance and secondary contact in the central Indo-West Pacific seas**

A Rohfritsch<sup>1</sup> and P Borsa<sup>1,2</sup>

<sup>1</sup> *Laboratoire Génome Populations Interactions (CNRS UMR 5000, IFREMER URM16, IRD), Station Méditerranéenne de l'Environnement Littoral, UMII, 1 Quai de la Daurade, 34200 Sète, France*

<sup>2</sup> *Institut de Recherche pour le Développement, BP A5, 98848 Nouméa cedex, Nouvelle-Calédonie.*

*Correspondence: P. Borsa. Institut de Recherche pour le Développement, BPA5, 98848 Nouméa cx, Nouvelle-Calédonie. Fax +687 26 43 26; E-mail: Philippe.Borsa@noumea.ird.nc*

**Keywords:** cytochrome b; intron-length polymorphism; geographic structure; Indo-Malay archipelago

**Running title:** Genetic structure of Indian scad mackerel

Major genetic breaks between the Indian and Pacific oceans have been reported for marine fishes and invertebrates. The genetic structure and history of *Decapterus russelli* in the Indo-Malay archipelago were investigated using the cytochrome *b* gene sequence as mitochondrial marker and two length-polymorphic introns as nuclear markers. The existence of two major mitochondrial lineages separated by 2.2% average nucleotide divergence, and their heterogeneous geographical distributions, were confirmed. This indicated past geographic isolation, possibly caused by Pleistocene drop in sea level. The presence in sympatry of the two mitochondrial lineages was thought to result from secondary contact. Recent population bottleneck and subsequent rapid population expansion were indicated by low genetic diversities and strongly negative Tajima's *D*-values. This supports the hypothesis that the habitat available to *D. russelli* in the Pleistocene was restricted. Taking into account both mitochondrial and nuclear-DNA data, three geographically distinct populations were identified: one sampled in the Makassar Strait and in the Sulawesi Sea, one in the Arafura Sea, and the third from the entire western region of the Indo-Malay archipelago. Considering the high hydrological connectivity of this region of the Indo-Pacific, homing behaviour and, perhaps, the association of spawning with retention zones, are believed to be able to maintain such a structure in a species with pelagic life-history.

## Introduction

Many marine organisms have large population sizes, high fecundity and high dispersal capabilities as planktonic larvae and free-swimming adults. Because of the high levels of gene flow expected from such traits, one expects that genetic differentiation between geographic populations is limited and allopatric speciation occurs only when the populations have long been separated by geographical barriers (e.g. continents) or hydrological barriers (e.g. oceanographic fronts, divergences of water masses, salinity or temperature clines). Philopatric behaviour, gamete incompatibility, differences in spawning time or location, and habitat selection could also be factors involved in inducing, maintaining or enforcing genetic differentiation.

The Central Indo-West Pacific region is both geographically and hydrologically complex. This region is also reputed to have the highest marine biodiversity in the world (Mora *et al*, 2003, and references therein). Its shorelines have undergone important changes in the Pleistocene (Hewitt, 2000; Voris, 2000) with the shallow seas of the Sunda shelf (Java Sea) and Sahul shelf (Arafura Sea) left above present sea level, and the emergence of land barriers isolating the South China Sea from the Indian Ocean at its southern limit and from the Sulu Sea eastbound (Fig. 1). The Sulu Sea was itself isolated from the Sulawesi Sea (Fig. 1). Thus, repeated fluctuations of sea levels in this topographically complex region could have favoured the geographic isolation of several inland seas during the ice ages, leading to the genetic differentiation of the populations they harboured. This is thought to be the cause of a phylogeographic break between Indian and Pacific populations of several marine species, and also a factor contributing to the greater species diversity reported for the Indo-Malay region (Hewitt, 2000). Geographic discontinuities have been reported for several invertebrate species distributed widely in the Indo-West Pacific (Williams and Benzie, 1998; Barber *et al*, 2000; 2002; Benzie *et al*, 2002; Gopurenko and Hughes, 2002). While no genetic differences were evident for goatfish (*Mulloidichthys vanicolensis*) populations across the Indo-West Pacific (Stepien *et al*, 1994), two other surveys reported geographic disjunction between Indian and West Pacific mitochondrial lineages and populations of inshore fishes, which was thought to be reminiscent of Pleistocene isolation. A large phylogeographic discontinuity was reported for barramundi (*Lates calcarifer*) on either side of the Torres Strait (Chenoweth *et al*, 1998), which formed a land barrier connecting Australia and New Guinea during the Pleistocene (Voris, 2000). Admixture of haplotypes from two distinct mitochondrial lineages was interpreted as evidence for recent secondary introgression in barramundi (Chenoweth *et al*, 1998). The phylogeographic structure of false clown anemonefish (*Amphiprion ocellaris*), a coastal sedentary fish with short pelagic larval stage, was also explained by sea level changes during the Pleistocene, rather than by contemporary geography (Nelson *et al*, 2000). Significant differences in *Cytochrome b* haplotype frequencies were found between *A. ocellaris* populations from the western edge of the Sunda Shelf (West coast of Sumatra, Malacca Strait) and those from the rest of the archipelago, including South China Sea, Sunda Strait, Bali Strait, Sulu

Sea, and Sulawesi Sea. Gene flow in *A. ocellaris* was inferred to be highest between regions connected by the permanent current that flows through the Makassar Strait from the Sulawesi Sea to the Bali Strait (Indonesian throughflow: Fieux *et al*, 1994). Pronounced geographic structure was also inferred from restriction-fragment length polymorphisms of the mitochondrial control region among populations of snapper (*Pristipomoides multidens*), sampled from the seas between northern Australia, Indonesia and West Papua (Ovenden *et al*, 2004). This was believed to be caused by constraints to dispersal at all life stages.

In contrast with the foregoing examples of inshore, sedentary fishes and invertebrates, little genetic heterogeneity has been reported for pelagic fishes from the Indo-Pacific. In spite of extensive sampling, no significant genetic differences were found between Indian and Pacific-Ocean populations of bigeye tuna (*Thunnus obesus*) (Alvarado Bremer *et al*, 1998; Chow *et al*, 2000), and Indo-Pacific sailfish (*Istiophorus platypterus*) (Graves and McDowell, 1995). Although a preliminary survey hinted at the possible genetic distinctness of round scad mackerel (*Decapterus macrosoma*) from the Sunda Strait *vs.* populations of the Sunda Shelf (Arnaud *et al*, 1999), no genetic differences were evident at both mitochondrial and nuclear loci among round scad mackerel populations at the scale of the Indo-Malay archipelago (Borsa, 2003). The spatial heterogeneity reported for yellowfin tuna (*Thunnus albacares*) (Ward *et al*, 1997) is a noticeable exception to the general pattern of panmixia in pelagic fishes in the Indo-Pacific.

Here we address the case of a highly mobile, pelagic fish of the scad mackerel group that exhibits geographic structure across the Indo-Malay archipelago. The Indian scad mackerel, *Decapterus russelli* (Carangidae), is abundant and widely distributed in the Indo-West Pacific and is a major fishery resource in South-East Asia (Mansor and Abdullah, 1995; Smith-Vaniz, 1999). However, knowledge on the biology, ecology, distribution and stock structure of this and all other tropical Carangidae is still preliminary. First-hand information on the spawning areas and the reproductive periods of *D. russelli* is scarce (Delsman, 1926; De Jong, 1940). Fishery statistics show large fluctuations in the catches of *D. russelli* depending on the season (Sadhotomo, 1998). Salinity variation in relation with the monsoon cycle would be an important factor in the distribution and the migration of Indian scad mackerel in the Indo-Malay archipelago (Hardenberg, 1937; Potier and Boely, 1990). From the temporal patterns in fishery catches, Hardenberg (1937) suggested that at least two populations of scad mackerel seasonally enter the Java Sea. One population may originate from the seas east to the Java Sea and reach the central-western part of the Java Sea during the East monsoon; the other stock may originate in the Indian Ocean and reach the western part of the Java Sea, through the Sunda Strait, during the West monsoon. Hardenberg (1937) also suggested that a third population occupying the southern part of the South China Sea may reach the northwestern part of the Java Sea during the West monsoon.

Studying the geographical structure of *Decapterus russelli* populations using genetic markers will improve our knowledge of its biology and ecology. The identification of differentiated populations is also of interest to the marine biogeography of the Indo-Pacific as the geographic distribution of allele frequencies coupled with the gene phylogeny will allow insight into the evolutionary history of Indian scad mackerel (geographical barriers, bottleneck events, recolonization). In a preliminary survey based on the sequence polymorphism of the cytochrome *b* gene, Perrin and Borsa (2001) distinguished two putative lineages in *D. russelli* mitochondria in the Indo-Malay archipelago, which were separated by 2.2% average nucleotide divergence. These two mitochondrial lineages had a heterogeneous geographic distribution and apparently co-occur in the Sulawesi Sea only. The presence in some samples of these two distinct lineages could be explained by either recent secondary contact between formerly isolated populations, or admixture in the sample of cryptic species. Mitochondrial-DNA haplotype frequency data and sequence phylogeny were insufficient information to select among the potential scenarios of genetic differentiation or cryptic speciation in *D. russelli*. In the present study, the precision of the mitochondrial phylogeography of this species was increased by expanding the study area and the total sample size, and the population genetic structure of *D. russelli* in the Indo-Malay archipelago was investigated using novel nuclear markers. The question we addressed was that of the origin and maintenance of geographic differences in this highly vagile species.

## Materials and methods

### *DNA samples*

Four hundred and fifty-eight Indian scad mackerel [*Decapterus russelli* (Rüppell 1830)] were sampled in 1995-1998 from small-sized purse seiners operating on a daily basis in the vicinity of 8 landing places in the Indo-Malay archipelago (Kelang, Carita, Labuan, Pekalongan, Kota Kinabalu, Sandakan, Toli-Toli, and Makassar). Sample Tambelan, from around Tambelan Islands in the Indonesian part of the South China Sea was collected from the fish hold of a medium-sized purse seiner upon calling in Pontianak (Kalimantan Barat, Indonesia). Another sample, caught by trawling in the Arafura Sea off West Papua, was purchased frozen at the fish market in Ambon, South Maluku. Fig. 1 reports the sites and dates where Indian scad mackerel were sampled in Indonesia, Malaysia, and West Papua (see Table 1 for sample sizes). Individual length was available for one sample (Toli-Toli). Length-frequency data allowed us to define size classes and to test the null hypothesis of genetic homogeneity across cohorts.

For each individual, a piece of muscle or a pectoral fin was stored in 95% ethanol. DNA was extracted from ca. 5 mg dehydrated fish tissue placed in 500  $\mu$ L extraction buffer (25 mM EDTA; 0.05 M Tris HCl pH8; 1% SDS) with 12 U proteinase K overnight at 55 °C, followed by a phenol-chloroform-isoamylalcohol extraction cycle (Sambrook *et al.*, 1989). The DNA pellet was resuspended in 100  $\mu$ L deionized water and frozen at -20 °C for subsequent use as template DNA in polymerase chain reactions (PCR).

### *Cytochrome b gene: PCR/SSCP and sequencing*

A 355-base pair (bp) fragment of the mitochondrial (mt) DNA cytochrome *b* gene was amplified by PCR using universal primers *PAT* (5'> GCC CCT CAG AAT GAT ATT TGT CCT C 3'>) and *CTB* (5'> CCA ACA TTT CAG CAT GAT GAA 3'>) modified from, respectively, primers *CB1-L* and *CB2-H* of Palumbi *et al.* (1991). PCR amplifications were carried out in 25  $\mu$ L reaction mixture containing ~50 ng template DNA, 2.5 mM MgCl<sub>2</sub>, 0.48 mM dNTPs, 0.12  $\mu$ M each primer and 0.38 U *Taq* DNA polymerase (Promega, Madison WI, USA) in 1X specific *Taq* buffer (from 10X buffer supplied by Promega). PCR conditions were 40 seconds at 94 °C, followed by 35 cycles of denaturing at 94 °C for 20 s, hybridization at 45 °C for 12 s and polymerization at 72 °C for 12 s with a final elongation step at 72 °C for 3 minutes, in a Crocodile III thermocycler (Appligène, Strasbourg). Sequence polymorphism was detected by single strand conformation polymorphism (SSCP) analysis (Lessa and Applebaum, 1993). This technique allows the detection of ~90% sequences for fragments of size between 300 base pairs (bp) to 450 bp when these differ from one another by as little as one nucleotide change (Hayashi, 1991 in Lessa and Applebaum, 1993). For SSCP, 3  $\mu$ L PCR product was added to 6  $\mu$ L denaturing buffer containing 1 vol. 6X loading buffer (0.25% xylene cyanol, 0.25% bromophenol blue, 30% glycerol, 0.5 M EDTA) and 5 vol. 95% formamide, 10 mM NaOH and 20 mM EDTA, and was denatured at 95 °C for 5 min. The single-strand DNAs were run at 18 °C for 15 hrs 30 min at 2 W in 1X non-denaturing MDE polyacrylamide gel (FMC Corporation, Rockland ME, USA) in a vertical electrophoresis tank (Hoefer, San Francisco CA, USA). DNA bands were visualised by silver-nitrate staining (Wray *et al.*, 1981). All haplotypes that were detected by SSCP (Fig. 2A) were sequenced using the Thermo Sequenase Radiolabeled Terminator kit of Amersham Life Science (Cleveland OH, USA) after the purification of the PCR products [1 U exonuclease I and 1 U alkaline phosphatase (Amersham) added to 5  $\mu$ L PCR product, incubated at 37 °C for 15 min, and the enzymes inactivated at 80 °C for 15 min]. The DNA fragments produced by the sequencing reaction were separated by vertical electrophoresis in 6% denaturing polyacrylamide gel [6% 29:1 acrylamide/bis-acrylamide (BioRad Laboratories, Hercules, Canada) in 0.5X TBE buffer]. Our nomenclature for SSCP phenotypes followed that of Perrin and Borsa (2001).

### *Nuclear-DNA markers*

Two introns showing length polymorphism in preliminary tests were used as nuclear genetic markers in *Decapterus russelli*. The first intron of the gene encoding aldolase *B* (*Aldolase B-1*) was amplified using PCR primers *1.1F* (5'> GCT CCA GGA AAG GGA ATC CTG GC 3'>) and *1.2R* (5'> CTC GTG GAA GAA GAT GAT CCC GCC 3'>) designed in the flanking

exons by Lemaire (2001). Just before the PCR, primer 1.2R was radioactively labelled for 30 min at 37°C in a mixture of 2 µM primer, 1 U T4 polynucleotide-kinase (Eurogentec, Liège, Belgium), and 1.7 µM (1.85 MBq) [ $\gamma$ -<sup>33</sup>P]ATP (Isotopchim, Ganagobie, France). PCR amplifications were carried out in 10 µL reaction mixture containing 4 µL DNA solution, 2.5 mM MgCl<sub>2</sub>, 0.3 mM dNTP, 0.03 µM radioactively labeled primer, 0.4 µM primer 1.1F, 0.5 U *Taq* DNA polymerase (Promega) in 1X *Taq* buffer. The polymerase chain reaction was run in a Crocodile III thermocycler (Appligène, Strasbourg, France). PCR program parameters were 3 min at 95 °C, followed by 35 cycles of denaturing at 94 °C for 12 s, hybridization at 52 °C for 12 s and polymerization at 72 °C for 20 s and followed by a final elongation step at 72 °C for 3 min. Length polymorphism was detected after allowing the heat-denatured PCR products to migrate vertically for 3 hrs 30 min at 50 W in 6% denaturing polyacrylamide gel (Fig. 2B). The gel was transferred onto a sheet of Whatman paper, vacuum-dried for 1 hr at 80 °C, and autoradiographed against X-Omat film (Eastman-Kodak, Rochester NY, USA).

The second intron of the gonadotropin-releasing hormone gene (*GnRH3*) was amplified using PCR primers *GnRH2F* (5'> AGA AGT GTG GGA GAG CTA GAG GC 3'>) and *GnRH2R* (5'> AGA GAC ACC ACT TCT CCT GTA CCC 3'>) designed by Hassan *et al* (2002) from a sequence of the Cichlidae *Astatotilapia burtoni* (GenBank AF076963). PCR amplifications were carried out in 10 µL reaction mixture containing 1 µL DNA solution, 1.25 mM MgCl<sub>2</sub>, 0.296 mM dNTP, 0.3 µM of primer *GnRH2F* labelled with 6-FAM fluorescent dye (Sigma Genosys, London, UK; carboxylfluorescein; maximum absorbance at 494 nm, maximum emission at 525 nm), 0.3 µM of the other primer (*GnRH2R*), 0.25 U *Taq* polymerase (Promega) in 1X *Taq* buffer. The PCR program used was the same as for amplifying *Aldolase B* Intron 1. After 5 min denaturing at 95 °C in 2 vol. denaturing buffer, the PCR products were run for 6 hrs at 40 W in 8% denaturing polyacrylamide gel. Fluorescent DNA bands were revealed under blue laser light (488 nm) in an FMBIO II gel scanner (Hitachi Instruments, San José CA, USA) equipped with an emission filter specific to fluorescein (505±20 nm). Intron-length polymorphism was thus detected at intron locus *GnRH 3-2* (Fig. 2C).

#### Data analysis

A neighbour-joining phylogeny of *Decapterus russelli* mitochondria was constructed using Kimura's 2-parameter model (MEGA2; Kumar *et al*, 2001). Parsimony analysis (MP) was conducted in addition, using MEGA2. Homologous *Cytochrome b* sequences of *D. kurroides*, *D. macrosoma* and *D. tabl* from Ambon, South Maluku, *D. macarellus* from Tobelo, Maluku, and *Selar crumenophthalmus* from Biak, West Papua were used as outgroups. Support values for internal nodes of the tree were estimated using the bootstrap resampling procedure (Felsenstein, 1985). One thousand pseudoreplicates were carried out for the NJ and MP runs.

A minimum parsimony network was produced to visualize the intraspecific phylogeny of mtDNA haplotypes using the software TCS v. 1.13 (Clement *et al*, 2000). TCS estimates genealogies by implementing the statistical parsimony method described in Templeton *et al* (1992).

The gene diversities ( $h$ ), the nucleotide diversities ( $\pi$ ), and their standard deviations were estimated using ARLEQUIN (Schneider *et al*, 2000). The average nucleotide divergence between mitochondrial haplogroups *A* and *M* was estimated with Kimura's two-parameter model using MEGA2. ARLEQUIN was also used to test for departures from the equilibrium between mutation and genetic drift in the mtDNA sequence data, with Tajima's *D*-test (Tajima, 1989). For a selectively neutral marker, negative *D*-values indicate excesses of low-frequency haplotypes, relative to mutation / drift equilibrium, with one or very few alleles at high frequency and rare alleles that derive from the latter by very few mutations. This is generally ascribed to rapid population expansion following a severe reduction in effective size (bottleneck). Conversely, positive values point to secondary contact among previously differentiated lineages.

Correspondence analysis (CA) (Benzécri, 1982) was performed on the matrix of allelic frequencies per sample (AFC procedure in BIOMECO; Lebreton *et al*, 1990). This amounts to estimating the degree of genetic differentiation between populations because the eigenvalues of each axis of the CA are analogous to partial *F<sub>st</sub>* (Guinand, 1996).

Wright's  $F_{is}$  and global- and pairwise  $F_{st}$  were estimated, respectively, using Weir and Cockerham's (1984) multiallelic  $\hat{f}$  and  $\hat{\theta}$  estimators [procedure FSTATS of GENETIX (Belkhir *et al*, 2000)]. Estimator  $\hat{\theta}$  can be used for both haploid and diploid data and is independent of the sampling strategy (Weir and Cockerham, 1984). It is widely used, thus allowing meaningful comparisons with other surveys of genetic differentiation in natural populations. The null hypotheses  $f=0$  and  $\theta=0$  were tested using the permutation test procedures implemented in GENETIX, where pseudo-distributions of  $\hat{f}$  and  $\hat{\theta}$  were obtained by random permutations of alleles and genotypes, respectively, and used as the expected distributions under the null hypothesis. The probability of occurrence of a parameter value larger than or equal to the observed value was estimated as  $P=(n+1)/(N+1)$ , where  $n$  is the number of pseudo-values larger (or lower for negative  $\hat{f}$ ) than or equal to the observed value and  $N$  is the number of random permutations (Sokal and Rohlf, 1995).

Pairwise linkage disequilibria were tested using GENETIX (Belkhir *et al*, 2000). The average correlation coefficient between alleles at two loci ( $R_{ij}$ ) was estimated according to Garnier-Géré and Dillmann (1992). The null hypothesis ( $R_{ij} = 0$ ) was tested by permutations of genotypes at a locus.

## Results

### *Cytochrome b* sequences

Nineteen different SSCP phenotypes at the *Cytochrome b* locus were detected in the total sample (Fig. 2A), including three novel SSCP phenotypes that were not present in the samples formerly surveyed by Perrin and Borsa (2001). One of the SSCP phenotypes ( $b$ ) proved after sequencing to consist of two haplotypes ( $b1$ ,  $b2$ ) differing by one nucleotide substitution (Perrin and Borsa, 2001). Phenotype  $A$  was the most common in all samples (frequency > 0.82) except Toli-Toli and Makassar, where the most common SSCP phenotype was  $M$  (frequency > 0.63). Phenotype  $M$  was absent from all the other samples, but some other haplotypes ( $b1$ ,  $b2$ ,  $i$ ) of Perrin and Borsa's (2001) haplogroup  $M$  were present at low frequencies in samples Carita, Labuan and Pekalongan. The frequencies in a sample of all SSCP phenotypes except  $A$  and  $M$  were lower than 0.04 in a sample, and lower than 0.01 in the total sample.

Among the 307 bp sequenced for each of the 20 *Cytochrome b* haplotypes represented by the above 19 SSCP phenotypes so far detected in the total sample, 20 sites were variable, 11 of which were phylogenetically informative. Sequences have been deposited in GENBANK (<http://www.ncbi.nlm.nih.gov>) (accession numbers nos. AF307494-AF307510 and AF515757-AF515759). Eighteen of the 20 nucleotide substitutions in the total sequence dataset occurred at third-nucleotide positions.

Both NJ and MP trees of partial nucleotide sequences of the *Cytochrome b* gene (data not shown) featured the two phylogenetically distinct mitochondrial lineages, haplogroup  $A$  and haplogroup  $M$ , formerly identified by Perrin and Borsa (2001). The average nucleotidic distance between the two haplogroups was  $d \pm SE = 2.2 \pm 0.7\%$ . Haplogroups  $A$  and  $M$  were supported by bootstrap values of 72% and 56-64%, respectively. This is statistically weak, but the geographic distribution of haplotypes also pointed to the delineation between haplotypes  $A$  and  $M$  which were the most frequent haplotypes of haplogroups  $A$  and  $M$ , respectively. While the inclusion of haplotype  $M$  into haplogroup  $M$  together with haplotypes  $b1$ ,  $b2$  and  $i$  was not strongly supported, the clade formed by the three latter haplotypes was more substantiated (bootstrap score = 78-79%). The three novel haplotypes detected in the samples ( $ii$ ,  $bb'$  and  $d$ ) clustered with haplogroup- $A$  haplotypes.

The haplotype network (TCS diagram) derived from cytochrome  $b$  gene partial sequences is presented in Fig. 3. Three of these rarer haplotypes ( $b1$ ,  $b2$  and  $i$ ) previously assigned to haplogroup  $M$  actually appeared to form a distinct cluster, separated from  $A$  by 4 mutational steps. Haplotype  $M$  was separated from  $A$  by 5 mutational steps and from the  $b1/b2/i$  cluster by 3 or 4 mutational steps only. The reticulations in the haplotype network, including the apparently intermediate position of  $M$ , may be explained by homoplasy, resulting from multiple substitutions at a given nucleotide site, or by recombination (see Hoarau *et al*, 2002 for such an example in flounder, *Platichthys flesus*).

Samples Toli-Toli and Makassar showed high gene ( $\hat{h} \geq 0.40$ ) and nucleotide ( $\hat{\pi} \geq 0.007$ ) diversities at the *Cytochrome b* locus (Table 1). Both samples possessed the *M* haplotype at high frequency, together with a substantial proportion of clade-*A* haplotypes (Table 1). All the other samples had  $\hat{h} \leq 0.32$  and  $\hat{\pi} \leq 0.002$  (Table 1) and the Arafura Sea sample was totally monomorphic for haplotype *A* (Table 1).

Tajima's *D*-values were strongly negative for all samples except Makassar and Toli-Toli, for which they were strongly positive (Table 1).

#### *Intron-length polymorphism*

The size of allele  $r_0$  at the *Aldolase B-1* locus was 265 bp. All the other alleles differed from allele  $r_0$  by a few base pairs. Allele  $r_0$  was the most common (frequency  $> 0.75$ ) in all samples except in the Arafura Sea sample, which was dominated by allele  $r_1$  (frequency = 0.67). Five other alleles were detected at locus *Aldolase B-1*, with frequencies  $\leq 0.06$  (Fig. 2B; Table 1).

Allele sizes (including primers) at the *GnRH 3-2* locus were centered around 300 pb, estimated by linear interpolation against the CXR 60-400bp ladder (Promega, Madison WI, USA). Because of difficulty in distinguishing some alleles differing of one another by tiny size differences, these were grouped as compound alleles. Compound alleles 30, 40, 90, and 110 each included two or three alleles (see footnote to Table 1) that were distinct on some gels, but not on other gels. The three alleles dominant in frequency were compound alleles 40 (0.54 > frequency per sample > 0.16), 50 (0.54 > frequency > 0.27), and 110 (0.33 > frequency > 0.09). The four other alleles that were detected, including compound allele 90 had frequencies  $\leq 0.15$  (Table 1).

#### *Population genetic structure*

Fixation index values ( $\hat{f}$ ) estimated from each sample at the *Aldolase B-1* and *GnRH 3-2* loci were not significantly different from zero ( $0.04 < P < 0.83$ ; 2-tailed test). Therefore, the hypothesis of panmixia was not rejected for any of the populations sampled.

The  $\hat{\theta}$ -values on the total set of samples significantly departed from 0 at all three loci (*Cytochrome b*:  $\hat{\theta} = 0.370$ ,  $P < 0.001$ ; *Aldolase B-1*:  $\hat{\theta} = 0.165$ ,  $P < 0.001$ ; *GnRH 3-2*:  $\hat{\theta} = 0.016$ ,  $P < 0.01$ ; 1000 random permutations using GENETIX), thereby demonstrating strong geographic structure of *Decapterus russelli* populations across the Indo-Malay archipelago. The null hypothesis of homogeneity could not be rejected at any one locus (*Cytochrome b*:  $\hat{\theta} = -0.006$ ; *Aldolase B-1*:  $\hat{\theta} = -0.005$ ; *GnRH 3-2*:  $\hat{\theta} = -0.001$ ; 1000 random permutations using GENETIX) for the group of samples (that is, Carita, Labuan, Tambelan, Pekalongan, and Kinabalu) selected for testing Hardenberg's hypothesis of two or three stocks in the periphery of the Java Sea (see Introduction).

There were some discrepancies among loci regarding the geographic patterns of genetic differentiation, as indicated by population-pairwise  $\hat{\theta}$ -estimates at each locus (Table 2). At the *Cytochrome b* locus, all significant pairwise  $\hat{\theta}$ -estimates involved either population Makassar or Toli-Toli *vs.* any other population in the Indo-Malay archipelago. At the *Aldolase B-1* locus, the Arafura Sea population distinguished itself from all the other populations. At the *GnRH 3-2* locus, significant genetic differences were found between the Arafura Sea population and a homogeneous group constituted by populations Sandakan, Makassar and Toli-Toli, while each of these two entities was not significantly distinct from the other populations in the Indo-Malay archipelago. Slight, albeit non-significant, genetic differences were also observed at locus *GnRH3-2* between the Sandakan / Makassar / Toli-Toli group and the western populations.

Considering each locus separately, the patterns of genetic differentiation were meaningful from a geographical perspective (Fig. 1). However, the geographic barrier uncovered by genetic analysis at the *Cytochrome b* locus was not evident when considering the *Aldolase B-1* locus, and vice-versa, while at locus *GnRH 3-2* a geographic barrier was evident only between those populations that were singled out at either of the two other loci. The synthetic picture emerging from the present multiple-locus analysis (see Fig. 4 presenting the results of CA) was that of three genetically differentiated geographic populations, being the Arafura Sea population, the Sulawesi Sea / Makassar Strait population, and the rest of the Indo-Malay archipelago.

The distribution of size classes in the Toli-Toli sample was bimodal (data not shown), designating two separate size-cohorts. No allele-frequency differences were evident between the two size-cohorts at loci *Cytochrome b* ( $\chi^2_{[1]}=1.167$ ;  $P=0.28$ ) and *Aldolase B-1* ( $\chi^2_{[2]}=0.359$ ;  $P=0.836$ ) while significant heterogeneity between cohorts was detected at locus *GnRH3-2* ( $\chi^2_{[4]}=10.413$ ;  $P=0.034$ ).

No pairwise genotypic or nucleo-cytoplasmic disequilibria between loci was evident, as indicated by the generally low  $R_{ij}$  values per sample, none of which was significant under the null hypothesis of linkage equilibrium (Table 3).

## Discussion

### *A single population or two sympatric species?*

Three differentiated mitochondrial lineages were present in Indian scad mackerel in the Indo-Malay archipelago. Former haplogroup *M* of Perrin and Borsa (2001) is now recognized to consist of two separate lineages. One lineage comprises haplotypes *b1*, *b2* and *i* formerly identified by Perrin and Borsa (2001), whereas haplotype *M* should now be singled out as a phylogenetically separate lineage. The geographic distribution of the three lineages was heterogeneous, with the western populations (those represented by samples Kelang, Carita, Labuan, Tambelan, Pekalongan, Kinabalu, and Sandakan) harbouring both the lineage represented by haplogroup *A* and that represented by haplotypes *b1*, *b2* and *i*, and the easternmost population (Arafura) possessing haplotype *A* only. Only the geographically intermediate population sampled in Makassar Strait / Sulawesi Sea harboured haplotype *M*, at high frequency, together with haplotype *A*. Following these observations we hypothesized that the admixture in the latter population of haplotype-*A* and haplotype-*M* individuals may reflect the occurrence of two cryptic species. Mitochondrial phylogenies have in some instances been used for elucidating systematic relationships among closely related species in fishes (e.g. Miya and Nishida, 1997; Rocha-Olivares *et al*, 1999; Agnès and Teugels, 2001; Colborn *et al*, 2001). The co-occurrence in some samples of separate mitochondrial lineages may indeed be indicative of reproductive isolation in marine fishes (e.g. Hoarau and Borsa, 2000; Borsa and Quignard, 2001; Pfeiler *et al*, 2002). Here, the alternative, null hypothesis that Toli-Toli and Makassar samples were drawn from a single panmictic population could not be rejected because genotypic frequencies at two nuclear loci did not depart from those expected under panmixia. Also, there was no evidence of nucleo-cytoplasmic disequilibrium in these two samples. In other terms, there was no indication that *A* and *M* individuals corresponded to different species. Therefore, the co-occurrence in the Sulawesi Sea / Makassar Strait region of these two distinct haplotypes indicates past geographic isolation that has been followed by secondary contact.

The present results emphasize the utility to systematics of recombinant (nuclear) markers in complement to mtDNA sequences. .

### *Population structure*

Surface currents in the western part of the Indo-Malay archipelago seasonally vary according to typical monsoon cycles (Fig. 1; Wyrki, 1961; Anonymous, 1978). During the wet monsoon (November to March), currents flow towards the East with speeds of ca. 1 to 2 knots. During the dry monsoon (May to September), the circulation is completely reversed with currents flowing towards the West (0.5 to 1 knots). During the intermonsoon (April to October), the winds and the currents are weak and variable (Anonymous, 1978). The monsoon cycle induces shifts in water circulation and changes in salinity, which coincides with the seasonal migration of adults and has an influence on the dispersion of the larvae of pelagic fishes (Hardenberg, 1937). According to Hardenberg (1937), two or perhaps three stocks of “layang” (a term used by this author to designate both *Decapterus russelli* and round scad mackerel, *D. macrosoma*) are present in the periphery of the Java Sea (see Introduction), which follow different migration routes and timings in relation with monsoons. However, the high connectivity of the seas in the Indo-Malay archipelago may favour high levels of gene flow between populations. Genetic data so far have failed to confirm Hardenberg's (1937) recognition of several stocks in the periphery of the Java Sea in both *D. russelli* (present results) and *D. macrosoma* (Borsa, 2003). At the scale of the entire Indo-Malay archipelago,

however, at least three distinct populations of *D. russelli* were here identified: (i) Arafura Sea, (ii) Sulawesi Sea and Makassar Strait, (iii) the rest of the Indo-Malay archipelago. Sound fishery management commands that these three geographically delineated, genetically differentiated populations be considered as separate management units.

That no correlation was observed between size class and either mitochondrial haplogroup or genotype at locus *Aldolase B-1* in Sample Toli-Toli, but a significant correlation was observed with genotype at locus *GnRH 3-2* suggests, perhaps, age-related selection at the latter locus or at a closely linked locus. Alternatively, this result could indicate subtle, random genetic differences between cohorts (see e.g. Ruzzante *et al*, 1996), warranting more insight into the fine-scale genetic and demographic structures of Indian scad mackerel. Another hypothesis is that the two cohorts may be drawn from genetically differentiated populations that mix outside the reproductive season ('admixture hypothesis' referred to hereafter). In this case, larger sample sizes for each cohort would be necessary to provide evidence of genetic differences at locus *GnRH 3-2* and perhaps at the two other loci. However, the fact that Samples Toli-Toli and Makassar did not exhibit any genetic differences at all three loci implies that under the admixture hypothesis the very same proportions of individuals from different cohorts were present in the two samples, which is unlikely.

#### *Concordant evolutionary patterns point to Pleistocene vicariance*

Assuming that the average rate of nucleotide change of the cytochrome *b* gene in marine fishes lies between the extremes of mammals and sharks, that is, respectively, 2.5% and 1.0% per million year (Irwin *et al*, 1991; Martin *et al*, 1992), the ca. 2.2% *Cytochrome b* divergence between lineages *A* and *M* in *Decapterus russelli* would correspond to between approximately 1.1 and 0.4 million year of allopatric isolation. This indicates that the split between haplogroups *A* and *M* dates to the mid-Pleistocene. The geographic distribution of haplotypes suggests the differences may result from vicariance. Through the Pleistocene, the Sunda and Sahul Shelves were repeatedly exposed above sea-level, constituting land barriers between the northeastern Indian Ocean and the western Pacific Ocean. The Java Sea and the Arafura Sea were completely above sea-level, the South China Sea was completely isolated from the Indian Ocean in its South and from the Sulu Sea in its East, and the Makassar Strait was much narrower than Present (Voris, 2000), presumably restricting the oceanic flow between the Sulawesi Sea and the rest of the Indo-Malay archipelago. The pattern of Makassar Strait / Sulawesi Sea populations being isolated by Pleistocene sea levels has been suggested for a mantis shrimp (Barber *et al*, 2000; 2002). Nucleotidic distances between Indian and Pacific populations in each of two monophyletic groups of nominal species of butterflyfishes (McMillan and Palumbi, 1995) were of the same order as between *D. russelli* haplogroups *A* and *M*. Two haplogroups separated by ca. 2.3% nucleotide divergence at the *Cytochrome b* locus were also detected in round scad mackerel (*D. macrosoma*) (Borsa, 2003). A similarly shallow phylogeographic break was reported in false clownfish on either side of the Malay Peninsula (Nelson *et al*, 2000) and between barramundi populations on either side of Torres Strait (Chenoweth *et al*, 1998). What thus appears to be a recurrent feature of the mitochondrial phylogenies of coastal fishes in the Indo-Malay region lends support to the hypothesis that changes in sea level associated with Pleistocene glaciations in higher latitudes provoked the geographic isolation of populations in the Indo-Malay region. Genetic divergence between geographically isolated population might have reached a level associated with possible reproductive isolation in coastal, sedentary species such as butterflyfishes (McMillan and Palumbi, 1995) but migratory, pelagic fishes such as *D. russelli* might have experienced shorter durations of geographic isolation. This would have prevented total reproductive isolation, hence left the possibility of secondary contact when the sea-level rose again.

#### *Population bottlenecks*

Bottlenecks followed by demographic expansions leave a genetic signature in extant populations in the form of a departure from the equilibrium between mutation and genetic drift, whereby rare alleles derive by a few mutations from a single or a few prevalent alleles (Grant and Bowen, 1998). The general aspect of the mitochondrial haplotype network in *Decapterus russelli* was consistent with the above hypothesis. Tajima's test was applicable for

all samples except Arafura, which exhibit sample monomorphism. Strong, negative  $D$ -values revealed bottlenecks in all populations except Makassar and Toli-Toli. The highly significant, positive  $D$ -values inferred for the two latter conversely points to a disequilibrium between mutation and drift. This was presumably caused by the occurrence in these samples of two differentiated mitochondrial lineages, which stems from recent secondary contact between populations that were reproductively isolated for a long period. The apparent discrepancy here observed between nuclear and mitochondrial markers is precisely what is expected from the scenario of a vicariance event separating two populations for a long time, followed by secondary contact. This is because of their different modes of transmission and evolutionary dynamics. Since mtDNA is uniparentally inherited, its effective population size is four-fold smaller than that of nuclear genes (Palumbi and Baker 1994; FitzSimmons *et al*, 1997). MtDNA is therefore more sensitive to demographic events such as reduction in population size and geographic isolation. Mitochondrial genetic differences between the Makassar Strait / Sulawesi Sea population and the Arafura Sea population were two to ten times higher than those at nuclear loci, roughly consistent with the expected four-fold ratio derived from their respective effective population sizes. Additional explanations are necessary to account for the higher discrepancy among genetic markers observed between the Makassar Strait / Sulawesi Sea population and the western populations, such as sex-biased gene flow (e.g. male dispersal / female philopatry; Mossman and Waser 1999) or selection at nuclear loci linked to the marker loci.

Low haplotype and nucleotide diversities point to low effective population sizes, due to either variance in reproductive success, cycles of demographic expansion and regression, or recent bottlenecks or foundation events (Grant and Bowen, 1998). Although no significant genetic differences were apparent at the *Cytochrome b* locus between the Arafura Sea population and the other predominantly haplotype-*A* populations, this sample was remarkably monomorphic. To explain the presence of the single haplotype *A* in Arafura Sea *Decapterus russelli*, a recent bottleneck or founding event should have involved no more than a few females, and the Arafura Sea population should be considered as geographically isolated from the rest of the Indo-Malay archipelago. The data from loci *Aldolase B-1* and *GnRH 3-2* indeed concur with the founder population hypothesis. As expected under this hypothesis, the effective number of alleles at locus *GnRH 3-2* was lowest in the Arafura Sea sample. At *Aldolase B-1* the alleles sampled in the Arafura Sea were the three most frequent ones in the rest of the Indo-Malay archipelago, even though the effective number of alleles was higher (Table 1), as a presumed consequence of genetic drift. As expected from a scenario of geographic isolation, allele frequencies at *Aldolase B-1* in the Arafura Sea were radically different from those in the rest of the Indo-Malay archipelago.

At the apogee of Pleistocene glaciations, when most continental seas were above sea-level, the habitat of coastal pelagic fishes was considerably reduced, implying much lower effective population sizes than Present. The rise in sea-level and the concomitant expansion of potentially favourable habitat may have allowed demographic expansion and the re-claiming of shallow seas such as the Java Sea and the Arafura Sea.

#### *Present barriers to gene flow*

Both nuclear and mitochondrial markers showed sharp geographic structure in *Decapterus russelli* in the Indo-Malay archipelago. How might allele-frequency differences between populations of this pelagic species have been maintained to the Present? *D. russelli* has a high dispersal potential, both by juveniles and adults which are known to undertake massive seasonal migrations (Hardenberg, 1937), and by larvae (Delsman, 1926). Eggs and larvae are transported passively by the strong marine currents that flow through the archipelago (Fig. 1). The putative secondary contact between different populations following the increase of the sea level should have induced a re-homogenization of allele frequencies, which did not happen. We therefore have to admit that *D. russelli* populations still exchange very few effective migrants despite the present, apparent continuity of the pelagic habitat. This suggests either reproductive homing, or the choice by adults of spawning areas that also constitute retention zones for eggs and larvae, or both.

It would seem sensible to assume that the extreme migrating abilities of some pelagic fishes favour wide-scale geographic homogeneity in allele frequencies. The observation of sharp geographic differences in Indian scad mackerel (present results) and some other highly

mobile, pelagic species (e.g. Kotoulas *et al*, 1995; Ward *et al*, 1997) shows that migrating abilities may also be associated with increased potential for homing, hence for reproductive isolation.

## Acknowledgements

We are grateful to R. Andamari, A.S. Cabanban, G. Lesage, M. Potier, Rumini, B. Sadhotomo, V. Sarpedonti, Suwarso, and Suprpto for helping in the collection of samples; to A.S. Cabanban and J.-M. Ecoutin for hospitality in Malaysia and in Indonesia; to F. Bonhomme, B. Crouau-Roy, and J.-D. Durand for comments on earlier versions of this paper; to R. Andamari, A.S. Cabanban, J.-R. Durand, J.-M. Ecoutin, M. Fatuchri Sukadi, P. Levang, J. Panfili, S. Prakoso, J. Roch and J. Widodo for facilitating research and contacts in Indonesia, Malaysia, South Maluku and West Papua; to M. Tortelier for her help in drawing Figure 1; to two anonymous referees for helpful comments and corrections. Finally, we would like to express our gratitude to F. Bonhomme for invaluable support at Station Méditerranéenne de l'Environnement Littoral and Laboratoire Génome et Populations. Part of the present work was undertaken as research operations of Pelfish and Genetikan, two French-Indonesian (IRD-CRIFI) cooperative projects on the assessment of pelagic fish resources in the Indonesian seas. Funded by Réseau Diversité Marine, IRD / DRV, IFREMER URM 16 and CNRS UMR 5000.

## References

- Agnès J-F, Teugels GG (2001) The *Bathyclarias-Clarias* species flock. A new model to understand rapid speciation in African Great Lakes. *C R Acad Sci Paris, Sér III* **324**: 683-688.
- Alvarado Bremer JR, Stequert B, Robertson NW, Ely B (1998) Genetic evidence for inter-oceanic subdivision of bigeye tuna (*Thunnus obesus*) populations. *Mar Biol* **132**: 547-557.
- Anonymous (1978) *Grand archipel d'Asie (partie méridionale): Indonésie*. Instructions Nautiques K6, Service Hydrographique et Océanographique de la Marine, Paris.
- Arnaud S, Bonhomme F, Borsa P (1999) Mitochondrial DNA analysis of the genetic relationships among populations of scad mackerel (*Decapterus macarellus*, *D. macrosoma* and *D. russelli*) in South-East Asia. *Mar Biol* **13**: 699-707.
- Barber PH, Palumbi SR, Erdmann MV, Moosa MK (2000) Biogeography. A marine Wallace's line? *Nature*, **406**: 692-693.
- Barber PH, Palumbi SR, Erdmann MV, Moosa MK (2002) Sharp genetic breaks among populations of *Haptosquilla pulchella* (Stomatopoda) indicate limits to larval transport: patterns, causes, and consequences. *Mol Ecol* **11**: 659-674.
- Belkhir K, Borsa P, Raufaste N, Chikhi L, Bonhomme F (2000) *GENETIX version 4.02, logiciel sous Windows™ pour la génétique des populations*. Université Montpellier 2, Montpellier, France.
- Benzécri J-P (1982) *L'analyse des données. 2. L'analyse des correspondances*. Dunod, Paris.
- Benzie JAH, Ballment E, Forbes AT, Demetriades NT, Sugama K, Haryanti, Moria S (2002) Mitochondrial DNA variation in Indo-Pacific populations of the giant tiger prawn, *Penaeus monodon*. *Mol Ecol* **11**: 2553-2569.
- Borsa P (2003) Genetic structure of round scad mackerel *Decapterus macrosoma* (Carangidae) in the Indo-Malay archipelago. *Mar Biol* **142**: 575-581.
- Borsa P, Quignard J-P (2001) Systematics of the Atlantic-Mediterranean soles *Pegusa impar*, *P. lascaris*, *Solea aegyptiaca*, *S. senegalensis*, and *S. solea* (Pleuronectiformes: Soleidae). *Can J Zool* **79**: 2297-2302.
- Chenoweth SF, Hughes JM, Keenan CP, Lavery S (1998) When oceans meet: a teleost shows secondary intergradation at an Indian-Pacific interface. *Proc Roy Soc Lond B* **265**: 415-420
- Chow S, Okamoto H, Miyabe N, Hiramatsu K, Barut N (2000) Genetic divergence between Atlantic and Indo-Pacific stocks of bigeye tuna (*Thunnus obesus*) and admixture around South Africa. *Mol Ecol* **9**: 221-227.

- Clement M., Posada D. Crandall K.A., 2000. TCS: a computer program to estimate gene genealogies. *Mol Ecol* **9**: 1657-1659.
- Colborn J, Crabtree RE, Shaklee JB, Pfeiler E, Bowen BW (2001) The evolutionary enigma of bonefishes (*Albula* spp.): cryptic species and ancient separations in a globally distributed shorefish. *Evolution* **55**: 807-820.
- Crow JF, Kimura M (1970) *An introduction to population genetics theory*. Alpha, Edina.
- De Jong JK (1940) A preliminary investigation of the spawning habits of some fishes of the Java Sea. *Treubia* **17**: 307-330.
- Delsman HC (1926) Fish eggs and larvae from the Java sea.5. *Caranx kurra*, *macrosoma*, and *crumenophthalmus*. *Treubia* **8**: 199-211.
- Felsenstein J (1985) Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* **39**: 783-791.
- Fieux M, Andrié C, Delécluse P, Ilahude AG, Kartavtseff A, Mantsi F, Molcard R, Swallow JC (1994) Measurements within the Pacific-Indian oceans throughflow region. *Deep-Sea Res* **41**: 1091-1130.
- FitzSimmons NN, Moritz C, Limpus CJ, Pope L, Prince R (1997) Geographic structure of mitochondrial and nuclear gene polymorphisms in Australian green turtle populations and male-biased gene flow. *Genetics* **147**: 1843--1854
- Garnier-Géré P, Dillmann C (1992) A computer program for testing pairwise linkage disequilibria in subdivided populations. *J Hered* **83**: 239.
- Gopurenko D, Hughes JM (2002) Regional patterns of genetic structure among Australian populations of the mud crab, *Scylla serrata* (Crustacea:Decapoda):evidence from mitochondrial DNA. *Mar Freshw Res* **53**: 849 –857.
- Grant WS, Bowen BW (1998) Shallow population histories in deep evolutionary lineages of marine fishes: insights from sardines and anchovies and lessons for conservation. *J Hered* **89**: 415-426.
- Graves JE, McDowell JR (1995). Inter-ocean genetic divergence of istiophorid billfishes. *Mar Biol* **122**: 193-203.
- Guinand B (1996) Use of a multivariate model using allele frequency distributions to analyse patterns of genetic differentiation among populations. *Biol J Linn Soc* **58**: 173-195.
- Hardenberg JDF (1937) Preliminary report on a migration of fish in the Java sea. *Treubia* **16**: 295-300.
- Hassan M, Lemaire C, Fauvelot C, Bonhomme F (2002) Seventeen new exon-primed intron-crossing polymerase chain reaction amplifiable introns in fish *Mol Ecol Notes* **2**: 334-340.
- Hayashi K (1991) PCR-SSCP: a simple and sensitive method for detection of mutations in the genomic DNA. *PCR Meth Appl* **1**: 34-38.
- Hewitt G (2000) The genetic legacy of the Quaternary ice ages. *Nature* **405**: 907-913.
- Hoarau G, Borsa P (2000) Extensive gene flow within sibling species in the deep-sea fish *Beryx splendens*. *C R Acad Sci Sér III* **323**: 315-325.
- Hoarau G, Holla S, Lescasse R, Stam WT, Olsen JL (2002) Heteroplasmy and evidence for recombination in the mitochondrial control region of the flatfish *Platichthys flesus*. *Mol Biol Evol* **19**: 2261-2264.
- Irwin DM, Kocher TD, Wilson AC (1991) Evolution of the cytochrome *b* gene of mammals. *J Mol Evol* **32**: 128-144.
- Kotoulas G, Magoulas A, Tsimenides N, Zouros E (1995) Marked mitochondrial DNA differences between Mediterranean and Atlantic populations of the swordfish, *Xiphias gladius*. *Mol Ecol* **4**: 473-481.
- Kumar S, Tamura K, Jakobsen IB, Nei M (2001) *MEGA2: molecular evolutionary genetics analysis software*. Arizona State University, Tempe.
- Lebreton J-D, Roux M, Banco G, Bacou A-M (1990) *BIOMEKO (biométrie-écologie). Logiciel d'écologie statistique pour PC et compatibles*. Centre National de la Recherche Scientifique., Montpellier.
- Lemaire C (2001) Flux génique et adaptations locales: étude de la divergence génétique en milieu marin chez *Dicentrarchus labrax* (Linné, 1758). Thèse de Doctorat, Institut National Agronomique Paris-Grignon, Paris.
- Lessa EP, Applebaum G (1993) Screening techniques for detecting allelic variation in DNA sequences. *Mol Ecol* **2**: 119-129.

- Mansor MI, Abdullah S (1995) Growth and mortality of Indian mackerel (*Rastrelliger kanagurta*) and slender scad (*Decapterus russelli*) off the East Coast of Peninsular Malaysia. *Sci Mar* **59**: 533-547.
- Martin AP, Naylor GJP, Palumbi SR (1992) Rates of mitochondrial evolution in sharks are slow compared with mammals. *Nature* **357**: 153-155.
- Mc Millan WO, Palumbi SR (1995) Concordant evolutionary patterns among Indo-West Pacific butterflyfishes. *Proc Roy Soc Lond B* **260**, 229-236.
- Miya M, Nishida M (1997) Speciation in the open ocean. *Nature* **389**, 803-804.
- Mora C, Chittaro PM, Sale PF, Kritzer JP, Ludsin SA (2003) Patterns and processes in reef fish diversity. *Nature* **421**: 933-936.
- Mossman CA, Waser PM (1999) Genetic detection of sex-biased dispersal. *Mol Ecol* **8**: 1063-1067.
- Nelson JS, Hoddell RJ, Chou LM, Chan WK, Phang VPE (2000) Phylogeographic structure of false clownfish, *Amphiprion ocellaris*, explained by sea level changes on the Sunda shelf. *Mar Biol* **137**: 727-736.
- Ovenden JR, Salini J, O'Connor S, Street R (2004) Pronounced genetic population structure in a potentially vagile fish species (*Pristipomoides multidens*, Teleostei, Perciformes, Lutjanidae) from the East Indies triangle. *Mol Ecol* **13**: 1991-1999.
- Palumbi SR, Baker CS (1994) Contrasting population structure from nuclear intron sequences and mtDNA of humpback whales. *Mol Biol Evol* **11**: 426-435
- Palumbi S, Martin A, Romano S, McMillan WO, Stice L, Grabowski G (1991) *The simple fool's guide to PCR, version 2.0*. Department of Zoology and Kewalo Marine Laboratory, University of Hawaii, Honolulu
- Perrin C, Borsa P (2001) Mitochondrial DNA analysis of the geographic structure of Indian scad mackerel, *Decapterus russelli* (Carangidae) in the Indo-Malay archipelago. *J Fish Biol* **59**: 1421-1426.
- Pfeiler E, Colborn J, Douglas MR, *et al.* (2002) Systematic status of bonefishes (*Albula* spp.) from the eastern Pacific Ocean inferred from analyses of allozymes and mitochondrial DNA. *Env Biol Fishes* **63**: 151-159.
- Potier M, Boely T (1990) Influence de paramètres de l'environnement sur la pêche à la senne tournante et coulissante en mer de Java. *Aquat Living Resour* **3**: 193-205.
- Rocha-Olivares A, Rosenblatt RH, Vetter RD (1999) Cryptic species of rockfishes (*Sebastes*: Scorpaenidae) in the Southern Hemisphere inferred from mitochondrial lineages. *J Hered* **90**: 409-411.
- Ruzzante DE, Taggart CT, Cook D (1996) Spatial and temporal variation in the genetic composition of a larval cod (*Gadus morhua*) aggregation: cohort contribution and genetic stability. *Can J Fish Aquat Sci* **53**: 2695-2705.
- Sadhotomo B (1998) Bioécologie des principales espèces pélagiques exploitées en mer de Java. Thèse de Doctorat, Université Montpellier 2, Montpellier.
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular cloning: a laboratory manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor.
- Schneider S, Roessli D, Excoffier L (2000) *ARLEQUIN: a software for population genetic data analysis*. Genetics and Biometry Laboratory, University of Geneva, Geneva.
- Smith-Vaniz WF (1999). Carangidae. In: *The living resources of the western central Pacific. Volume 4. Bony fishes part 2 (Mugilidae to Carangidae)* (eds Carpenter KE, Niem VH), pp. 2659-2756. FAO Species Identification Guide for Fishery Purposes, Food and Agriculture Organization of the United Nations, Rome.
- Sokal RR, Rohlf FJ (1995) *Biometry. The principles and practice of statistics in biological research, 3rd edn*. Freeman and Co, New York.
- Stepien CA, Randall JE, Rosenblatt RH (1994) Genetic and morphological divergence of a circumtropical complex of goatfishes: *Mulloidichthys vanicolensis*, *M. dentatus*, and *M. martinicus*. *Pac Sci* **48**: 44-56.
- Tajima F (1989) The effect of change in population size on DNA polymorphisms. *Genetics* **123**: 437-460.
- Templeton AR, Crandall KA, Sing CF (1992) A cladistic analysis of phenotypic associations with haplotypes inferred from restriction endonuclease mapping and DNA sequence data. III. Cladogram estimation. *Genetics* **132**: 619-633.

- Voris HK (2000) Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time durations. *J Biogeogr* **27**: 1153-1167.
- Ward RD, Elliott NG, Innes BH, Smolenski AJ, Grewe PM (1997). Global population structure of yellowfin tuna, *Thunnus albacares*, inferred from allozyme and mitochondrial DNA variation. *Fishery Bull US* **95**: 566-575.
- Weir BS, Cockerham CC (1984) Estimating *F*-statistics for the analysis of population structure. *Evolution* **38**: 1358-1370.
- Williams ST, Benzie JAH (1998) evidence of a biogeographic break between populations of a high dispersal starfish: congruent regions within the Indo-West pacific defined by color morphs, mtDNA, and allozyme data. *Evolution* **52**: 87-99.
- Wray W, Boulikas T, Wray VP, Hancock R (1981) Silver staining of proteins in polyacrylamide gels. *Anal Biochem* **118**: 197-203.
- Wyrski K (1961). Physical oceanography of the southeast Asian waters. *Naga Rep, Scripps Inst Oceanogr, Univ California* **2**: 1-95.

**Table 1** *Decapterus russelli*. Allele frequencies and summary statistics at Loci *Cytochrome b*, *Aldolase B-1* and *GnRH3-2* for ten samples from the Indo-Malay archipelago.  $\hat{f}$ , Weir et Cockerham's (1984) estimator of Wright's fixation index and  $P$ , associated probability under the null hypothesis of panmixia. Gene ( $\hat{h}$ ) and nucleotide ( $\hat{\pi}$ ) diversities  $\pm$  standard deviations ( $SD$ ) estimated using ARLEQUIN (Schneider *et al.*, 2000). Effective number of alleles per sample estimated as  $n_e = 1/\sum x_i^2$ , where  $x_i$  is frequency of the  $i^{\text{th}}$  allele (Crow and Kimura, 1970). Tajima's test of selective neutrality:  $D$  and  $P(D)$ , associated probability (of  $D$  random <  $D$  observed; 1000 random simulations), estimated using ARLEQUIN.  $N$ , sample size.

[Footnote to Table 1]:

\*Samples Carita, Labuan, Tambelan, Pekalongan, Kota Kinabalu and Sandakan were genotyped by Perrin and Borsa (2001) at the Cytochrome b locus.

† Compound alleles at locus GnRH 3-2 were "30", including alleles 25, 30 and 35 which were distinct on some gels only (see Fig. 2C); "40", similarly including alleles 40 and 45 (itself possibly including two slightly different size-alleles) (see Fig. 2C); "90", similarly including alleles 80 and 90 on some gels; "110", including alleles 100, 110 and 120 (see Fig. 2C)

Locus, Allele	Sample									
	Kelang	Carita	Labuan	Tambelan	Pekalongan	Kinabalu	Sandakan	Toli-Toli	Makassar	Arafura
<i>Cytochrome b</i> *										
39	–	–	–	0.02	–	–	–	–	–	–
A	0.83	0.90	0.91	0.92	0.95	0.95	0.92	0.27	0.36	1.00
a	0.08	0.02	–	–	–	–	0.02	–	–	–
b	–	–	–	–	–	–	0.02	–	–	–
b'	–	0.02	–	–	–	–	–	–	–	–
b''	–	–	–	–	–	–	0.02	–	–	–
bb	–	–	0.03	–	–	–	–	–	–	–
bb'	–	–	–	–	–	–	0.02	–	–	–
d	–	–	–	–	–	–	0.02	–	–	–
dD	–	–	–	–	0.01	–	–	–	–	–
dF	–	–	–	–	0.01	–	–	–	–	–
e	–	–	–	0.02	–	–	–	–	–	–
e'	–	0.02	0.01	0.02	–	0.03	–	–	–	–
ee	–	–	–	0.02	–	0.03	–	–	–	–
ii	0.08	–	–	–	–	–	–	–	–	–
m	–	–	0.01	–	–	–	–	–	–	–
b1	–	–	0.01	–	–	–	–	–	–	–
b2	–	0.02	0.03	–	–	–	–	–	–	–
i	–	–	–	–	0.01	–	–	–	–	–
M	–	–	–	–	–	–	–	0.73	0.64	–
(N)	(12)	(42)	(79)	(52)	(67)	(37)	(66)	(30)	(22)	(51)
$\hat{b}$	0.318	0.184	0.170	0.149	0.088	0.107	0.147	0.405	0.485	0.000
$\pm SD$	$\pm 0.016$	$\pm 0.080$	$\pm 0.057$	$\pm 0.067$	$\pm 0.048$	$\pm 0.068$	$\pm 0.059$	$\pm 0.078$	$\pm 0.064$	–
$n_e$	1.425	1.232	1.205	1.179	1.108	1.106	1.179	1.651	1.855	1.000
$\hat{\pi}$	0.0011	0.0016	0.0019	0.0010	0.0010	0.0009	0.0007	0.0081	0.0097	0.0000
$\pm SD$	$\pm 0.0007$	$\pm 0.0003$	$\pm 0.0004$	$\pm 0.0003$	$\pm 0.0002$	$\pm 0.0003$	$\pm 0.0002$	$\pm 0.0018$	$\pm 0.0024$	–
D	-1.451	-2.111	-1.781	-1.946	-2.331	-1.754	-1.836	1.720	2.378	–
P(D)	0.033	0.000	0.014	0.003	0.001	0.018	0.006	0.963	0.997	–
<i>Aldolase B-1</i>										
$r_2$	–	–	–	0.01	–	–	–	–	–	–
$r_1$	0.19	0.07	0.08	0.08	0.13	0.11	0.13	0.10	0.14	0.67
$r_{0.5}$	–	0.04	0.02	0.01	0.02	0.04	–	–	0.06	0.02
$r_0$	0.75	0.86	0.88	0.86	0.84	0.78	0.86	0.87	0.81	0.31
$r_{0.5}$	0.06	–	–	0.01	–	–	–	0.03	–	–
$r_{.1}$	–	0.03	0.01	0.01	0.01	0.04	0.02	–	–	–
$r_{.3}$	–	–	0.01	0.01	–	0.02	–	–	–	–
(N)	(8)	(37)	(66)	(39)	(51)	(23)	(59)	(15)	(18)	(32)
$\hat{f}$	0.429	-0.089	0.182	-0.091	0.146	0.202	-0.005	-0.087	-0.155	-0.024
P	0.222	0.503	0.071	0.455	0.179	0.129	0.674	0.800	0.489	0.557
$\hat{b}$	0.425	0.248	0.222	0.259	0.275	0.380	0.253	0.246	0.338	0.458
$\pm SD$	$\pm 0.047$	$\pm 0.065$	$\pm 0.047$	$\pm 0.064$	$\pm 0.053$	$\pm 0.087$	$\pm 0.048$	$\pm 0.098$	$\pm 0.092$	$\pm 0.046$
$n_e$	1.661	1.339	1.280	1.340	1.383	1.602	1.321	1.302	1.472	1.834
<i>GnRH 3-2</i> †										
30	–	0.03	0.05	0.01	0.06	0.04	0.07	0.07	0.03	0.05
40	0.40	0.34	0.39	0.38	0.30	0.41	0.21	0.17	0.18	0.53
50	0.30	0.36	0.28	0.28	0.39	0.32	0.44	0.52	0.42	0.32
60	0.15	0.02	0.05	0.06	0.05	0.01	0.03	0.05	0.03	–
70	–	0.03	0.01	0.01	0.01	–	–	–	–	–
90	0.05	–	0.01	0.01	0.04	0.04	0.02	–	0.03	–
110	0.10	0.22	0.21	0.24	0.15	0.16	0.22	0.19	0.32	0.11
(N)	(10)	(32)	(57)	(34)	(58)	(34)	(63)	(21)	(19)	(33)
$\hat{f}$	0.074	-0.007	-0.036	0.111	0.015	0.000	0.147	0.223	-0.046	0.162
P	0.451	0.540	0.407	0.177	0.468	0.567	0.041	0.077	0.513	0.124
$\hat{b}$	0.935	0.874	0.877	0.894	0.904	0.919	0.860	0.935	0.923	0.895
$\pm SD$	$\pm 0.025$	$\pm 0.019$	$\pm 0.013$	$\pm 0.013$	$\pm 0.013$	$\pm 0.014$	$\pm 0.021$	$\pm 0.015$	$\pm 0.018$	$\pm 0.012$
$n_e$	3.509	3.381	3.574	3.517	3.671	3.340	3.421	2.917	3.186	2.513

**Table 2** *Decapterus russelli*. Estimates of Weir and Cockerham's (1984)  $\hat{\theta}$  between pairs of samples at one mitochondrial (*Cytochrome b*) and two nuclear (*Aldolase B-1*, *GnRH3-2*) loci.

Locus, Sample	Sample								
	Carita	Labuan	Tambelan	Pekalongan	Kinabalu	Sandakan	Makassar	Toli-Toli	Arafura
<i>Cytochrome b</i>									
Kelang	-0.013	0.010	0.014	0.067	0.033	0.011	<b>0.406**</b>	<b>0.509***</b>	0.238*
Carita		-0.012	-0.012	0.003	-0.011	-0.010	<b>0.543***</b>	<b>0.621***</b>	0.045*
Labuan			-0.007	0.003	-0.007	-0.004	<b>0.594***</b>	<b>0.662***</b>	0.029*
Tambelan				-0.003	-0.018	-0.008	<b>0.587***</b>	<b>0.657***</b>	0.029
Pekalongan					-0.009	-0.003	<b>0.669***</b>	<b>0.725***</b>	0.009
Kinabalu						-0.010	<b>0.597***</b>	<b>0.666***</b>	0.025
Sandakan							<b>0.606***</b>	<b>0.673***</b>	0.024
Makassar								-0.023	<b>0.736***</b>
Toli-Toli									<b>0.778***</b>
<i>Aldolase B-1</i>									
Kelang	0.021	0.022	0.009	-0.015	-0.035	-0.003	-0.031	-0.020	<b>0.303***</b>
Carita		-0.009	-0.009	-0.003	-0.003	0.000	-0.001	-0.012	<b>0.490***</b>
Labuan			-0.009	-0.003	0.005	-0.002	0.000	-0.019	<b>0.525***</b>
Tambelan				-0.006	-0.003	-0.004	-0.002	-0.018	<b>0.481***</b>
Pekalongan					-0.008	-0.009	-0.015	-0.019	<b>0.451***</b>
Kinabalu						-0.001	-0.021	-0.012	<b>0.381***</b>
Sandakan							-0.006	-0.016	<b>0.474***</b>
Makassar								-0.011	<b>0.382***</b>
Toli-Toli									<b>0.441***</b>
<i>GnRH3-2</i>									
Kelang	-0.003	-0.011	-0.016	-0.008	-0.016	0.030	0.046	0.063	-0.003
Carita		-0.005	-0.009	-0.004	-0.007	0.006	0.008	0.023	0.021*
Labuan			-0.011	0.008	-0.006	0.030**	0.033*	0.056**	0.014
Tambelan				0.008	-0.007	0.028	0.024	0.054*	0.016
Pekalongan					0.001	0.003	0.014	0.011	0.032*
Kinabalu						0.029*	0.039*	0.052*	-0.001
Sandakan							-0.010	-0.011	<b>0.077***</b>
Makassar								-0.001	<b>0.102**</b>
Toli-Toli									<b>0.127***</b>

\*  $P < 0.05$  ; \*\*  $P < 0.010$  ; \*\*\*  $P < 0.001$  [1000 permutations under GENETIX (Belkhir *et al*, 2000)]. In bold : values remaining significant after sequential Bonferroni correction (Sokal et Rohlf, 1995). For this correction, nine series of nine pairwise comparisons were considered independently at each locus.

**Table 3** *Decapterus russelli*. Test of linkage disequilibrium at three locus pairs (1000 permutations using GENETIX).  $R_{ij}$ , average correlation coefficient estimate between alleles at two different loci;  $P$ , probability of a pseudo-value superior or equal to observed  $R_{ij}$  under null hypothesis  $R_{ij}=0$ .

Sample	<i>Cytochrome b</i> */ <i>Aldolase B-1</i>		<i>Cytochrome b</i> */ <i>GnRH 3-2</i>		<i>Aldolase B-1</i> / <i>GnRH 3-2</i>	
	$R_{ij}$	$P$	$R_{ij}$	$P$	$R_{ij}$	$P$
Kelang	--	--	--	--	0.727	0.890
Carita	0.066	0.894	0.136	0.361	0.170	0.199
Labuan	0.059	0.602	0.070	0.666	0.089	0.616
Tambelan	--	--	--	--	0.133	0.532
Pekalongan	0.111	0.207	0.085	0.487	0.098	0.762
Kinabalu	--	--	--	--	0.277	0.124
Sandakan	--	--	--	--	0.093	0.661
Makassar	0.192	0.524	0.202	0.885	0.205	0.592
Toli-Toli	0.246	0.552	0.117	0.574	0.304	0.389
Arafura	--	--	--	--	0.155	0.619
Makassar + Toli-Toli	0.115	0.716	0.075	0.932	0.134	0.697
Total sample	0.044	0.327	0.063	0.189	0.039	0.858

\* *Cytochrome b* haplotypes were grouped into two haplogroups, *A* and *M*, after Fig. 3


## Captions to Figures

**Figure 1** *Decapterus russelli*. Sampling locations (open circles) in the Indo-Malay archipelago. Abbreviations for samples, sampling dates and sample sizes as following. *K*: Kelang, November 1998,  $N=12$ ; *C*: Carita, October 1997,  $N=42$ ; *L*: Labuan, May 1995 and October 1997,  $N=79$ ; *T*: Tambelan, April 1995,  $N=52$ ; *P*: Pekalongan, April 1995,  $N=67$ ; *Kk*: Kota Kinabalu, October 1997,  $N=37$ ; *S*: Sandakan, October 1997,  $N=66$ ; *M*: Makassar, September 1998,  $N=22$ ; *Tt*: Toli-Toli, May 1995,  $N=30$ ; *A*: Arafura Sea, April 1998,  $N=51$ . *MS*, Makassar Strait; *SS*, Sunda Strait. **Bold arrows**: dominant surface currents; **dotted arrows**: seasonal surface currents (after Wyrтки, 1961; Anonymous, 1978). *Inset*: land areas (shaded) in the Indo-Malay region at the apogee of Pleistocene glaciations, inferred from Present bathymetric charts and using as coastline -120 m below Present sea-level (see Voris, 2000)


**Figure 2** *Decapterus russelli*. **A** Single-stranded DNA conformation polymorphism at the *Cytochrome b* locus. Composite drawing of all 19 SSCP phenotypes found in the total sample. **B** Length polymorphism at locus *Aldolase B-Intron 1*, revealed by electrophoresis on denaturing polyacrylamide gel. Six of the seven alleles thus detected in the total sample are shown. Lane 1: inferred genotype was  $r_2/r_2$ ; lanes 2, 8:  $r_0/r_0$ ; lanes 3, 4, 5:  $r_0/r_{.3}$ ; lanes 6, 7:  $r_0/r_{.1}$ ; lane 9:  $r_{0.5}/r_0$ ; lane 10:  $r_1/r_1$ . **C** Intron-length polymorphism at a Gonadotropin Releasing Hormone locus (*GnRH 3-2*), revealed by electrophoretic differences on denaturing polyacrylamide gel. Nine out of the 10 alleles detected in the total sample by using this technique are shown. Lane 1: inferred genotype was  $45/50$ ; lane 2:  $50/50$ ; lane 3:  $40/50$ ; lane 4:  $45/100$ ; lane 5:  $60/90$ ; lane 6:  $35/50$ ; lane 7:  $45/50$ ; lane 8:  $45/110$ ; lane 9:  $30/50$

**Figure 3** *Decapterus russelli*. Statistical parsimony network of mtDNA haplotypes. Dot circles designate haplotypes whose frequency was  $<0.01$  in the total sample. Small squares designate putative mutationally intermediate haplotypes. Nucleotide sequences deposited in GENBANK under accession nos AF307494-AF307510 and AF515757-AF515759

**Figure 4** Correspondence analysis (CA) of *Decapterus russelli* samples, where each sample was defined by its allele frequencies at loci *Cytochrome b*, *Aldolase B-1*, and *GnRH 3-2*. Projection of samples on CA's principal plane (Axis 1 and Axis 2). Inertia of each axis (that is, the percentage of total genetic variation that is explained by it; see Benzecri, 1982) is given in brackets. Ellipses delineate groups of samples that did not exhibit significant pairwise  $\hat{\theta}$  between them at any one locus (see Table 1). *Abbreviations for samples* as in legend to Fig. 1.




**A** *A a 39 b b' b'' bb bb' d dF dD e e' ee m h i ii M*



**B** 1 2 3 4 5 6 7 8 9 10

$r_2$  —  
 $r_1$  —  
 $r_{0.5}$  —  
 $r_0$  —  
 $r_{-1}$  —  
  
 $r_{-3}$  —



**C** 1 2 3 4 5 6 7 8 9

30 —  
 35 —  
 40 —  
 45 —  
 50 —  
 60 —  
  
 90 —  
 100 —  
 110 —

