

Asymptotic behavior of Structures made of Plates

G. Griso

Laboratoire J.-L. Lions–CNRS, Boîte courrier 187,
Université Pierre et Marie Curie, 4 place Jussieu, 75005 Paris, France,
Email: griso@ann.jussieu.fr

Abstract. The aim of this work is to study the asymptotic behavior of a structure made of plates of thickness 2δ when $\delta \rightarrow 0$. This study is carried on within the frame of linear elasticity by using the unfolding method. It is based on several decompositions of the structure displacements and on the passing to the limit in fixed domains.

We begin with studying the displacements of a plate. We show that any displacement is the sum of an elementary displacement concerning the normal lines on the middle surface of the plate and a residual displacement linked to these normal lines deformations. An elementary displacement is linear with respect to the variable x_3 . It is written $\mathcal{U}(\hat{x}) + \mathcal{R}(\hat{x}) \wedge x_3 \vec{e}_3$ where \mathcal{U} is a displacement of the mid-surface of the plate. We show a priori estimates and convergence results when $\delta \rightarrow 0$. We characterize the limits of the unfolded displacements of a plate as well as the limits of the unfolded of the strained tensor.

Then we extend these results to the structures made of plates. We show that any displacement of a structure is the sum of an elementary displacement of each plate and of a residual displacement. The elementary displacements of the structure (e.d.p.s.) coincide with elementary rods displacements in the junctions. Any e.d.p.s. is given by two functions belonging to $H^1(\mathbf{S}; \mathbb{R}^3)$ where \mathbf{S} is the skeleton of the structure (the plates mid-surfaces set). One of these functions : \mathcal{U} is the skeleton displacement. We show that \mathcal{U} is the sum of an extensional displacement and of an inextensional one. The first one characterizes the membrane displacements and the second one is a rigid displacement in the direction of the plates and it characterizes the plates flexion.

Eventually we pass to the limit as $\delta \rightarrow 0$ in the linearized elasticity system, on the one hand we obtain a variational problem that is satisfied by the limit extensional displacement, and on the other hand, a variational problem satisfied by the limit of inextensional displacements.

Résumé. Le but de ce travail est d'étudier le comportement asymptotique d'une structure formée de plaques d'épaisseur 2δ lorsque $\delta \rightarrow 0$. Cette étude est menée dans le cadre de l'élasticité linéaire en utilisant la méthode de l'éclatement. Elle est basée sur plusieurs décompositions des déplacements de la structure, et sur le passage à la limite dans des domaines fixes.

On commence par une étude des déplacements d'une plaque. On montre que tout déplacement d'une plaque est la somme d'un déplacement élémentaire concernant les normales à la surface moyenne de la plaque et d'un déplacement résiduel lié aux déformations de ces normales. Un déplacement élémentaire est affine par rapport à la variable x_3 , il s'écrit $\mathcal{U}(\hat{x}) + \mathcal{R}(\hat{x}) \wedge x_3 \vec{e}_3$ où \mathcal{U} est un déplacement de la surface moyenne de la plaque. On établit des estimations a priori et des résultats de convergence lorsque $\delta \rightarrow 0$. On caractérise les limites des éclatés des déplacements d'une plaque, ainsi que les limites des éclatés du tenseur des déformations.

On étend ensuite ces résultats aux structures formées de plaques. On montre que tout déplacement d'une structure est la somme d'un déplacement élémentaire de chaque plaque et d'un déplacement résiduel. Les déplacements élémentaires de la structure (d.e.s.p.) coïncident avec des déplacements élémentaires de poutres dans les jonctions. Tout d.e.s.p. est donné par deux fonctions appartenant à $H^1(\mathbf{S}; \mathbb{R}^3)$ où \mathbf{S} est le squelette de la structure (l'ensemble des surfaces moyennes des plaques). L'une de ces fonctions : \mathcal{U} est le déplacement du squelette. On montre que \mathcal{U} est la somme d'un déplacement extensionnel et d'un déplacement inextensionnel. Le premier caractérise les déplacements membranaires des surfaces moyennes, le second est un déplacement rigide dans la direction des plaques; il caractérise la flexion des plaques.

Pour finir on passe à la limite pour $\delta \rightarrow 0$ dans le système de l'élasticité linéaire, on obtient d'une part un problème variationnel vérifié par la limite des déplacements extensionnels, et d'autre part un problème variationnel vérifié par la limite des déplacements inextensionnels.

1. Introduction

Many articles and books have been dedicated to the mathematical justification of plates models (see for example [1,2]). A first study concerning the asymptotic behavior of a structure made of two thin plates of thickness ε , is due to Le Dret [7]. The obtained asymptotic model derives from the three-dimensional system of elasticity thanks to a thin domain standard technique (the plates are transformed into a fixed domain). At the limit, Le Dret obtains a two-dimensional system coupling the flexion displacements of the two mid-surfaces of the plates.

Our study continues [4] and [5]. In this paper we use again the notions of elementary displacements and of extensional and inextensional displacements and we extend them to the plates displacements and to the displacements of structures made of plates. Our paper is organised into three parts. In the first one we study the displacements of a plate, the second one is devoted to the displacements of a structure made of plates from which we deduce the asymptotic behavior of a structure made of thin plates. And in the third part we prove the technical lemmas used in the two first parts of our paper.

In Section 2 we consider a plate of thickness 2δ . We first introduce the elementary displacements of a plate (Definition 2.1). These are the displacements of the normal lines of the mid-surface of the plate. An elementary displacement is linear with respect to the variable x_3 . It is written $\mathcal{U}(\hat{x}) + \mathcal{R}(\hat{x}) \wedge x_3 \vec{e}_3$ where \mathcal{U} is a displacement of the mid-surface. By such a displacement the normal line is transformed into a line which is generally no longer perpendicular to the mid-surface. With each displacement u of the plate we associate an elementary displacement U_e (Definition 2.2). Theorem 2.3 gives estimates of appropriate norms of U_e and of the displacement $u - U_e$ in terms of δ . Using the elementary displacement U_e we show (formula (2.3)) that the displacement u is the sum of a Kirchhoff-Love displacement and of a residual one \tilde{u} , which satisfies estimate (2.4). We are now equipped to obtain the asymptotic behavior of a displacements sequence $(u_\delta)_{\delta>0}$ with strain energy of order δ . This is the main result of this section and it is given in Theorem 2.6. The previous decomposition allows us to give a simple interpretation (see Theorem 2.6) of the limits of the unfolding $\mathcal{T}_\delta(\gamma_{ij}(u_\delta))$ of the strain tensor $\gamma_{ij}(u_\delta)$ (where the unfolding operator \mathcal{T}_δ is given in Definition 2.5) in terms of the derivatives limits of the Kirchhoff-Love displacements and of the residual displacements. There is not a unique associated elementary displacement that satisfies estimates (2.2). In Definition 2.2 we give the simplest one. But the one we give in Definition 2.9 is more suitable for the study of a structure made of plates.

The structure \mathcal{S}_δ made of plates of thickness 2δ is introduced in Section 3. Our hypotheses about the skeleton of the structure \mathbf{S} (i.e. the plates mid-surfaces set) allow us to consider a wide range of structures. We extend to them the notions and decompositions of Section 2. Definition 3.1 gives us the elementary displacements of plates-structure (e.d.p.s.). These displacements coincide with elementary plate displacements in each plate and there are rods elementary displacements in the junctions (see [5]). Any e.d.p.s. is known by two functions belonging to $H^1(\mathbf{S}; \mathbb{R}^3)$. The first one \mathcal{U} is the skeleton displacement, the second one gives the rotations of the normal lines of the mid-surfaces. We show that \mathcal{U} is the sum of an extensional displacement and of an inextensional one (Definitions 3.6 and 3.5). The first one characterizes the membrane displacements and the second one is a rigid displacement in the direction of the plates and it characterizes the plates flexion. Corollary of Lemma 3.7 gives estimates for them with an appropriate norm. In subsection 3.4. we consider an e.d.p.s. sequence $(u_\delta)_{\delta>0}$ with strain energy of order δ . Thanks to all these decompositions we give the limits of the unfolding $\mathcal{T}_\delta(\gamma_{ij}(u_\delta))$ of the strain tensor as in the case of a plate. We also characterize the space of the inextensional limits displacements. In the last subsection, we give the limit for $\delta \rightarrow 0$ of the linearized elasticity system (3.9), written in \mathcal{S}_δ , where the applied forces F_δ satisfy assumptions (3.11). The main results are Theorem 3.8 and Theorem 3.10. In the first one we show

that the extensional displacement limit is the solution of a second-order system, and in the second one we show that the limit of the inextensional displacement is the solution of a fourth-order system.

In this work we use the Einstein convention of summation over repeated indices. As a rule, the Greek indices α and β take values in $\{1, 2\}$ and the Latin indices i, i', j and j' take values in $\{1, 2, 3\}$.

2. The plate displacements

2.1. The elementary plate displacements

The Euclidian space \mathbb{R}^3 is related to the frame $(O; \vec{e}_1, \vec{e}_2, \vec{e}_3)$. Let ω be a bounded domain in \mathbb{R}^2 with a lipschitzian boundary. The plate $\Omega_\delta = \omega \times]-\delta, \delta[$, $\delta > 0$, is the open set having as middle surface ω and as thickness 2δ . The direction of the normal lines of ω is given by \vec{e}_3 . The reference plate is the open set $\Omega = \omega \times]-1, 1[$.

The running point of Ω_δ (respectively Ω) is denoted $x = (x_1, x_2, x_3) = (\hat{x}, x_3)$, (resp. (\hat{x}, t_3)) where $\hat{x} \in \omega$ and $t_3 \in]-1, 1[$.

For any open set ω' of \mathbb{R}^n , $n \in \{2, 3\}$, and any displacement u belonging to $H^1(\omega', \mathbb{R}^n)$, we put

$$\mathcal{E}(u, \omega') = \int_{\omega'} \gamma_{ij}(u) \gamma_{ij}(u), \quad \gamma_{ij}(u) = \frac{1}{2} \left\{ \frac{\partial u_i}{\partial x_j} + \frac{\partial u_j}{\partial x_i} \right\}, \quad \mathcal{D}(u, \omega') = \int_{\omega'} \frac{\partial u_i}{\partial x_j} \frac{\partial u_i}{\partial x_j}.$$

Definition 2.1 : An elementary plate displacement (e.p.d.) is an element Φ belonging to $H^1(\Omega_\delta, \mathbb{R}^3)$, such that

$$\Phi(x) = \mathcal{A}(\hat{x}) + \mathcal{B}(\hat{x}) \wedge x_3 \vec{e}_3, \quad \text{a.e. } x = (\hat{x}, x_3) \in \omega \times]-\delta, \delta[= \Omega_\delta,$$

where \mathcal{A} and \mathcal{B} belong to $H^1(\omega, \mathbb{R}^3)$; \mathcal{A} is the first component and \mathcal{B} the second component of the e.p.d. Φ .

Elementary plate displacement associated with a displacement of $H^1(\Omega_\delta, \mathbb{R}^3)$.

Definition 2.2 : With any displacement $u \in H^1(\Omega_\delta, \mathbb{R}^3)$, we associate the elementary plate displacement U_e defined as

$$(2.1) \quad \begin{cases} U_e(x) = \mathcal{U}(\hat{x}) + \mathcal{R}(\hat{x}) \wedge x_3 \vec{e}_3, & x = (\hat{x}, x_3) \in \Omega_\delta, \\ \mathcal{U}(\hat{x}) = \frac{1}{2\delta} \int_{-\delta}^{\delta} u(\hat{x}, x_3) dx_3, & \mathcal{R}(\hat{x}) = \frac{3}{2\delta^3} \int_{-\delta}^{\delta} x_3 \vec{e}_3 \wedge u(\hat{x}, x_3) dx_3. \end{cases}$$

The component \mathcal{R}_3 of \mathcal{R} is equal to 0.

Theorem 2.3 : The elementary plate displacement U_e verifies

$$(2.2) \quad \mathcal{E}(U_e, \Omega_\delta) + \mathcal{D}(u - U_e, \Omega_\delta) + \frac{1}{\delta^2} \|u - U_e\|_{L^2(\Omega_\delta, \mathbb{R}^3)}^2 \leq C \mathcal{E}(u, \Omega_\delta).$$

The constants depend only on ω .

Proof : See Annex A. □

Proposition 2.4 : Any displacement u belonging to $H^1(\Omega_\delta, \mathbb{R}^3)$ is the sum of a Kirchhoff-Love displacement and a residual one \tilde{u}

$$(2.3) \quad u(x) = \left(\mathcal{U}_1(\hat{x}) - x_3 \frac{\partial \mathcal{U}_3}{\partial x_1}(\hat{x}) \right) \vec{e}_1 + \left(\mathcal{U}_2(\hat{x}) - x_3 \frac{\partial \mathcal{U}_3}{\partial x_2}(\hat{x}) \right) \vec{e}_2 + \mathcal{U}_3(\hat{x}) \vec{e}_3 + \tilde{u}(x), \quad x \in \Omega_\delta,$$

where \mathcal{U} is the first component of the e.p.d. U_e . The residual displacement \tilde{u} belongs to $L^2(\omega, H^1(]-\delta, \delta[, \mathbb{R}^3))$ and verifies

$$(2.4) \quad \frac{1}{\delta^2} \|\tilde{u}\|_{L^2(\Omega_\delta, \mathbb{R}^3)}^2 + \left\| \frac{\partial \tilde{u}}{\partial x_3} \right\|_{L^2(\Omega_\delta, \mathbb{R}^3)}^2 \leq C \mathcal{E}(u, \Omega_\delta).$$

Proof : We define \tilde{u} by

$$\tilde{u}(x) = u(x) - \left(\mathcal{U}_1(\hat{x}) - x_3 \frac{\partial \mathcal{U}_3}{\partial x_1}(\hat{x}) \right) \vec{e}_1 - \left(\mathcal{U}_2(\hat{x}) - x_3 \frac{\partial \mathcal{U}_3}{\partial x_2}(\hat{x}) \right) \vec{e}_2 - \mathcal{U}_3(\hat{x}) \vec{e}_3.$$

This displacement belongs to $L^2(\omega, H^1(\cdot - \delta, \delta[, \mathbb{R}^3))$. We obtain (2.4) using the estimates of Theorem 2.3. \square

2.2. Limit of a plate displacements sequence.

Definition 2.5 : The unfolding operator \mathcal{T}_δ from $L^2(\Omega_\delta, \mathbb{R}^n)$ into $L^2(\Omega, \mathbb{R}^n)$ is defined by

$$(2.5) \quad \mathcal{T}_\delta(\phi)(\hat{x}, t_3) = \phi(\hat{x}, \delta t_3), \quad \text{a.e. in } \Omega.$$

For any element $\phi \in H^1(\Omega_\delta)$, we have $\mathcal{T}_\delta(\phi) \in H^1(\Omega)$ and

$$(2.6) \quad \frac{\partial \mathcal{T}_\delta(\phi)}{\partial x_\alpha} = \mathcal{T}_\delta\left(\frac{\partial \phi}{\partial x_\alpha}\right), \quad \frac{\partial \mathcal{T}_\delta(\phi)}{\partial t_3} = \delta \mathcal{T}_\delta\left(\frac{\partial \phi}{\partial x_3}\right).$$

Theorem 2.6 : Let $(u_\delta)_{\delta > 0}$ be a sequence of displacements of $H^1(\Omega_\delta, \mathbb{R}^3)$ verifying

$$(2.7) \quad \mathcal{E}(u_\delta, \Omega_\delta) \leq C\delta.$$

There exist $(a_\delta, b_\delta) \in \mathbb{R}^3 \times \mathbb{R}^3$ and extracted sequences (still denoted in the same way), such that

$$(2.8) \quad \begin{cases} \mathcal{U}_{1,\delta} - a_{1,\delta} + x_2 b_{3,\delta} \rightharpoonup U_1, & \mathcal{U}_{2,\delta} - a_{2,\delta} - x_1 b_{3,\delta} \rightharpoonup U_2 & \text{weakly in } H^1(\omega) \\ \delta \{ \mathcal{U}_{3,\delta} - a_{3,\delta} + x_1 b_{2,\delta} - x_2 b_{1,\delta} \} \rightharpoonup U_3 & \text{weakly in } H^1(\omega). \end{cases}$$

Moreover U_3 belongs to $H^2(\omega)$. We have the following weak convergences of the unfolded of u_δ , \tilde{u}_δ and of the components of the strain tensor :

$$(2.9) \quad \begin{cases} \mathcal{T}_\delta(u_{1,\delta} - a_{1,\delta} + x_2 b_{3,\delta}) \rightharpoonup U_1 - t_3 \frac{\partial U_3}{\partial x_1}, & \mathcal{T}_\delta(u_{2,\delta} - a_{2,\delta} - x_1 b_{3,\delta}) \rightharpoonup U_2 - t_3 \frac{\partial U_3}{\partial x_2}, \\ \delta \mathcal{T}_\delta(u_{3,\delta} - a_{3,\delta} + x_1 b_{2,\delta} - x_2 b_{1,\delta}) \rightharpoonup U_3, & \text{weakly in } H^1(\Omega), \\ \frac{1}{\delta} \mathcal{T}_\delta(\tilde{u}_\delta) \rightharpoonup \tilde{u} & \text{weakly in } L^2(\omega, H^1(\cdot - 1, 1[, \mathbb{R}^3)), \\ \mathcal{T}_\delta(\gamma_{\alpha\beta}(u_\delta)) \rightharpoonup \frac{1}{2} \left\{ \frac{\partial U_\alpha}{\partial x_\beta} + \frac{\partial U_\beta}{\partial x_\alpha} \right\} - t_3 \frac{\partial^2 U_3}{\partial x_\alpha \partial x_\beta}, & \mathcal{T}_\delta(\gamma_{\alpha 3}(u_\delta)) \rightharpoonup \frac{1}{2} \frac{\partial \tilde{u}_\alpha}{\partial t_3}, \\ \mathcal{T}_\delta(\gamma_{33}(u_\delta)) \rightharpoonup \frac{\partial \tilde{u}_3}{\partial t_3} & \text{weakly in } L^2(\Omega). \end{cases}$$

Proof : With each u_δ we associate the e.p.d. $U_{e,\delta}$ with components \mathcal{U}_δ and \mathcal{R}_δ . From (2.4) the displacement $\mathcal{U}_{M,\delta} = \mathcal{U}_{1,\delta} \vec{e}_1 + \mathcal{U}_{2,\delta} \vec{e}_2$ has a strain energy $\mathcal{E}(\mathcal{U}_{M,\delta}, \omega) \leq \frac{C}{\delta} \mathcal{E}(U_{e,\delta}, \Omega_\delta) \leq \frac{C}{\delta} \mathcal{E}(u_\delta, \Omega_\delta) \leq C$. The classical Korn inequality applied to $\mathcal{U}_{M,\delta}$ affirms the existence of a rigid displacement $r_{M,\delta}(\hat{x}) = \begin{pmatrix} a_{1,\delta} - b_{3,\delta} x_2 \\ a_{2,\delta} + b_{3,\delta} x_1 \end{pmatrix}$, such that

$$\|\mathcal{U}_{M,\delta} - r_{M,\delta}\|_{L^2(\omega, \mathbb{R}^2)}^2 + \mathcal{D}(\mathcal{U}_{M,\delta} - r_{M,\delta}, \omega) \leq C \mathcal{E}(\mathcal{U}_{M,\delta}, \omega) \leq \frac{C}{\delta} \mathcal{E}(u_\delta, \Omega_\delta) \leq C.$$

If $b_{\alpha,\delta}$ is the mean of $\mathcal{R}_{\alpha,\delta}$ on ω , we obtain from the Poincaré-Wirtinger inequality

$$\|\mathcal{R}_{\alpha,\delta} - b_{\alpha,\delta}\|_{L^2(\omega)}^2 \leq C \|\nabla \mathcal{R}_{\alpha,\delta}\|_{[L^2(\omega)]^2}^2 \leq \frac{C}{\delta^3} \mathcal{E}(u_\delta, \Omega_\delta) \leq \frac{C}{\delta^2}$$

The estimate of $\mathcal{E}(U_{e,\delta}, \Omega_\delta)$ obtained in Theorem 2.3, gives

$$(2.10) \quad \left\| \frac{\partial \mathcal{U}_{3,\delta}}{\partial x_1} + \mathcal{R}_{2,\delta} \right\|_{L^2(\omega)}^2 + \left\| \frac{\partial \mathcal{U}_{3,\delta}}{\partial x_2} - \mathcal{R}_{1,\delta} \right\|_{L^2(\omega)}^2 \leq \frac{C}{\delta} \mathcal{E}(u_\delta, \Omega_\delta) \leq C,$$

hence

$$\left\| \frac{\partial \mathcal{U}_{3,\delta}}{\partial x_1} + b_{2,\delta} \right\|_{L^2(\omega)}^2 + \left\| \frac{\partial \mathcal{U}_{3,\delta}}{\partial x_2} - b_{1,\delta} \right\|_{L^2(\omega)}^2 \leq \frac{C}{\delta^3} \mathcal{E}(u_\delta, \Omega_\delta) \leq \frac{C}{\delta^2}.$$

Now we apply the Poincaré-Wirtinger inequality to the function $\mathcal{U}_{3,\delta} + b_{2,\delta}x_1 - b_{1,\delta}x_2$. There exists $a_{3,\delta}$ such that

$$\|\mathcal{U}_{3,\delta} - a_{3,\delta} + b_{2,\delta}x_1 - b_{1,\delta}x_2\|_{L^2(\omega)}^2 \leq \frac{C}{\delta^3} \mathcal{E}(u_\delta, \Omega_\delta) \leq \frac{C}{\delta^2}.$$

The sequences $\mathcal{U}_{M,\delta} - r_{M,\delta}$, $\delta\{\mathcal{U}_{3,\delta} - a_{3,\delta} + b_{2,\delta}x_1 - b_{1,\delta}x_2\}$, $\delta(\mathcal{R}_{\alpha,\delta} - b_{\alpha,\delta})$ and $\frac{1}{\delta}\mathcal{T}_\delta(\tilde{u}_\delta)$ are bounded in $H^1(\omega, \mathbb{R}^2)$ (respectively $H^1(\omega)$ and $L^2(\omega, H^1(\cdot - 1, 1], \mathbb{R}^3)$). We extract from these sequences some subsequences, still denoted in the same way, such that

$$(2.11) \quad \begin{cases} \mathcal{U}_{1,\delta} - a_{1,\delta} + x_2 b_{3,\delta} \rightharpoonup U_1, & \mathcal{U}_{2,\delta} - a_{2,\delta} - x_1 b_{3,\delta} \rightharpoonup U_2 \quad \text{weakly in } H^1(\omega), \\ \delta\{\mathcal{U}_{3,\delta} - a_{3,\delta} + x_1 b_{2,\delta} - x_2 b_{1,\delta}\} \rightharpoonup U_3 \quad \text{weakly in } H^1(\omega), \\ \delta(\mathcal{R}_{\alpha,\delta} - b_{\alpha,\delta}) \rightharpoonup \mathcal{R}_\alpha \quad \text{weakly in } H^1(\omega), \\ \frac{1}{\delta}\mathcal{T}_\delta(\tilde{u}_\delta) \rightharpoonup \tilde{u} \quad \text{weakly in } L^2(\omega, H^1(\cdot - 1, 1], \mathbb{R}^3). \end{cases}$$

The limits of the sequences $\delta\left\{\frac{\partial \mathcal{U}_{3,\delta}}{\partial x_1} + \mathcal{R}_{2,\delta}\right\}$ and $\delta\left\{\frac{\partial \mathcal{U}_{3,\delta}}{\partial x_2} - \mathcal{R}_{1,\delta}\right\}$ are equal to zero by (2.10), hence the equalities

$$(2.12) \quad \frac{\partial U_3}{\partial x_1} = -\mathcal{R}_2, \quad \frac{\partial U_3}{\partial x_2} = \mathcal{R}_1,$$

and the belonging of U_3 to $H^2(\omega)$. From the limits (2.11) and from the equalities (2.12) we immediately deduce the limits of the unfolded $\mathcal{T}_\delta(u_{1,\delta} - a_{1,\delta} + x_2 b_{3,\delta})$, $\mathcal{T}_\delta(u_{2,\delta} - a_{2,\delta} - x_1 b_{3,\delta})$ and $\delta\mathcal{T}_\delta(u_{3,\delta} - a_{3,\delta} + x_1 b_{2,\delta} - x_2 b_{1,\delta})$ in $H^1(\Omega)$.

To calculate the components of the strain tensor we use the equality (2.3)

$$\begin{aligned} \gamma_{\alpha\beta}(u_\delta) &= \frac{1}{2} \left\{ \frac{\partial \mathcal{U}_{\alpha,\delta}}{\partial x_\beta} + \frac{\partial \mathcal{U}_{\beta,\delta}}{\partial x_\alpha} \right\} - x_3 \frac{\partial^2 \mathcal{U}_{3,\delta}}{\partial x_\alpha \partial x_\beta} + \frac{1}{2} \left\{ \frac{\partial \tilde{u}_{\alpha,\delta}}{\partial x_\beta} + \frac{\partial \tilde{u}_{\beta,\delta}}{\partial x_\alpha} \right\}, \\ \gamma_{\alpha 3}(u_\delta) &= \frac{1}{2} \left\{ \frac{\partial \tilde{u}_{\alpha,\delta}}{\partial x_3} + \frac{\partial \tilde{u}_{3,\delta}}{\partial x_\alpha} \right\}, \quad \gamma_{33}(u_\delta) = \frac{\partial \tilde{u}_{3,\delta}}{\partial x_3}. \end{aligned}$$

These equalities are transformed through unfolding. All the sequences $\mathcal{T}_\delta(\gamma_{ij}(u_\delta))$ are bounded in $L^2(\Omega)$ and they have a limit in $H^{-1}(\Omega)$, which can be explained thanks to the convergences (2.11) and the equalities (2.12). Hence the last limits of (2.9). \square

Remark 2.7 : We consider again the sequence of displacements $(u_\delta)_{\delta>0}$ of Theorem 2.7. We put $(U'_{e,\delta})_{\delta>0}$ another sequence of e.p.d. verifying

$$(2.13) \quad \mathcal{E}(U'_{e,\delta}, \Omega_\delta) + \mathcal{D}(u_\delta - U'_{e,\delta}, \Omega_\delta) + \frac{1}{\delta^2} \|u_\delta - U'_{e,\delta}\|_{L^2(\Omega_\delta, \mathbb{R}^3)}^2 \leq C\mathcal{E}(u_\delta, \Omega_\delta) \leq C\delta,$$

where the constant is independent on δ . Then we obtain

$$\|U_\delta - U'_\delta\|_{L^2(\omega, \mathbb{R}^3)} \leq C\delta \quad \|\mathcal{R}_{\alpha,\delta} - \mathcal{R}'_{\alpha,\delta}\|_{L^2(\omega)} \leq C.$$

The displacement u_δ is decomposed now into the sum of a new Kirchhoff-Love displacement and a new residual one \tilde{u}'_δ

$$u_\delta = \left(\mathcal{U}'_{1,\delta} - x_3 \frac{\partial \mathcal{U}'_{3,\delta}}{\partial x_1} \right) \vec{e}_1 + \left(\mathcal{U}'_{2,\delta} - x_3 \frac{\partial \mathcal{U}'_{3,\delta}}{\partial x_2} \right) \vec{e}_2 + \mathcal{U}'_{3,\delta} \vec{e}_3 + \tilde{u}'_\delta$$

The displacement \tilde{u}'_δ verifies the inequality $\|\tilde{u}'_\delta\|_{L^2(\Omega_\delta, \mathbb{R}^3)}^2 + \delta^2 \left\| \frac{\partial \tilde{u}'_\delta}{\partial x_3} \right\|_{L^2(\Omega_\delta, \mathbb{R}^3)}^2 \leq C\delta^3$. After extraction of subsequences expressed by the same notation, we obtain the convergences

$$\begin{cases} \mathcal{U}_\delta - \mathcal{U}'_\delta \longrightarrow 0 & \delta(\mathcal{R}_\delta - \mathcal{R}'_\delta) \longrightarrow 0 & \text{strongly in } L^2(\omega, \mathbb{R}^3) \\ \frac{1}{\delta} \mathcal{T}_\delta(\tilde{u}'_\delta) \rightharpoonup \tilde{u}' & \text{weakly in } L^2(\omega, H^1([-1, 1], \mathbb{R}^3)). \end{cases}$$

The limits of the unfolded $\mathcal{T}_\delta(\gamma_{i3}(u_\delta))$ give $\frac{\partial \tilde{u}}{\partial t_3} = \frac{\partial \tilde{u}'}{\partial t_3}$. Except for the limit of the sequence of the unfolded $1/\delta \mathcal{T}_\delta(\tilde{u}'_\delta)$, the limits (2.8) and (2.9) do not depend on the decomposition of the displacement u_δ into the sum of an e.p.d. and a residual displacement. What matters is to be able to approximate u_δ with the help of an e.p.d. that verifies the estimates (2.13). It is to be noticed that the mere knowing of the limits of the unfolded of the stain tensor components of the sequence $(u_\delta)_{\delta>0}$ is not enough to determine completely the residual displacement \tilde{u} . It is obtained but for a function of $L^2(\omega, \mathbb{R}^3)$.

2.3. A second decomposition of a plate displacement

We consider now a round-rimmed plate Ω'_δ with a middle surface ω_δ . We denote

$$\begin{aligned} \Omega'_\delta &= \{x \in \mathbb{R}^3 \mid \text{dist}(x, \omega) < \delta\}, & \Gamma_\delta &= \{x \in \mathbb{R}^3 \mid \text{dist}(x, \partial\omega) < \delta\} \\ \tilde{\Omega}_\delta &= \{x \in \mathbb{R}^2 \mid \text{dist}(x, \omega) < 2\delta\} \times]-\delta, \delta[= \omega_{2\delta} \times]-\delta, \delta[. \end{aligned}$$

Lemma 2.8 : *For any $\delta \in]0, \delta_0]$, there exists an extension operator \mathcal{P}_δ , linear and continuous from $H^1(\Omega'_\delta, \mathbb{R}^3)$ into $H^1(\tilde{\Omega}_\delta, \mathbb{R}^3)$ such that*

$$(2.14) \quad \mathcal{P}_\delta(u)|_{\Omega'_\delta} = u, \quad \begin{cases} \mathcal{E}(\mathcal{P}_\delta(u), \tilde{\Omega}_\delta) \leq C\mathcal{E}(u, \Omega'_\delta) \\ \mathcal{E}(\mathcal{P}_\delta(u), \tilde{\Omega}_\delta \setminus \Omega'_\delta) \leq C\mathcal{E}(u, \Gamma_\delta) \end{cases}$$

The constants do not depend on δ .

Proof : See Annex B. □

The extension of u to $\tilde{\Omega}_\delta$ is still denoted u .

A second elementary plate displacement associated with a displacement of $H^1(\Omega'_\delta, \mathbb{R}^3)$.

Definition 2.9 : With any $u \in H^1(\Omega'_\delta, \mathbb{R}^3)$ we associate the e.p.d. U'_e defined by

$$(2.15) \quad \begin{cases} U'_e(x) = \mathcal{U}'(\hat{x}) + \mathcal{R}'(\hat{x}) \wedge x_3 \vec{e}_3, & x \in \Omega'_\delta, \\ \mathcal{U}'(\hat{x}) = \frac{6}{\pi\delta^3} \int_{B(\hat{x}; \delta/2)} u(M) dM, & \mathcal{R}'(\hat{x}) = \frac{60}{\pi\delta^5} \int_{B(\hat{x}; \delta/2)} \widehat{xM} \wedge u(M) dM, & \hat{x} \in \omega_\delta. \end{cases}$$

Theorem 2.10 : *We have the following inequalities:*

$$(2.16) \quad \begin{cases} \delta^3 \|\nabla \mathcal{R}'\|_{L^2(\omega_\delta, \mathbb{R}^6)}^2 + \delta \left\| \frac{\partial \mathcal{U}'}{\partial x_\alpha} - \mathcal{R}' \wedge \vec{e}_\alpha \right\|_{L^2(\omega_\delta, \mathbb{R}^3)}^2 + \mathcal{E}(U'_e, \Omega'_\delta) \leq C\mathcal{E}(u, \Omega'_\delta), \\ \mathcal{D}(u - U'_e, \Omega'_\delta) \leq C\mathcal{E}(u, \Omega'_\delta), \quad \|u - U'_e\|_{L^2(\Omega'_\delta, \mathbb{R}^3)}^2 \leq C\delta^2 \mathcal{E}(u, \Omega'_\delta). \end{cases}$$

The constants depend only on ω .

Proof : We now consider the covering $\{\omega'_{\delta,n}\}_{n \in N_\delta}$ (see Lemma 4.1 in Annex A). We put $\omega''_{\delta,n} = \{\hat{x} \in \omega'_{\delta,n} \mid \text{dist}(\hat{x}, \partial\omega'_{\delta,n}) > \delta/2\}$ and $\mathcal{O}'_{\delta,n} = \omega'_{\delta,n} \times]-\delta, \delta[$, $\mathcal{O}''_{\delta,n} = \omega''_{\delta,n} \times]-\delta, \delta[$, $n \in N_\delta$. The family $\{\omega''_{\delta,n}\}_{1 \leq n \leq N_\delta}$ verifies

$$\text{measure}\left(\bigcup_{n \in N_\delta} \omega''_{\delta,n} \setminus \omega_\delta\right) = 0$$

From Lemma 2.3 in [4] there exists a rigid displacement r_n such that

$$(2.17) \quad \mathcal{D}(u - r_n, \mathcal{O}'_{\delta,n}) + \frac{1}{\delta^2} \|u - r_n\|_{L^2(\mathcal{O}'_{\delta,n}, \mathbb{R}^3)}^2 \leq C\mathcal{E}(u, \mathcal{O}'_{\delta,n}), \quad r_n(x) = a_n + b_n \wedge \overrightarrow{A_n x}, \quad (a_n, b_n) \in \mathbb{R}^3.$$

The constants do not depend on n nor δ . We calculate the mean of $(u - r_n)(M)$ and of $\widehat{xM} \wedge (u - r_n)(M)$ in the ball $B(\hat{x}, \delta/2)$, $\hat{x} \in \omega''_{\delta,n}$. Due to (2.17) we obtain

$$(2.18) \quad \begin{cases} \|\mathcal{U}'(\hat{x}) - a_n - b_n \wedge \overrightarrow{A_n \hat{x}}\|_2^2 \leq \frac{C}{\delta} \mathcal{E}(u, \mathcal{O}'_{\delta,n}) \implies \|\mathcal{U}' - a_n - b_n \wedge \overrightarrow{A_n \cdot}\|_{L^2(\omega''_{\delta,n}, \mathbb{R}^3)}^2 \leq C\delta \mathcal{E}(u, \mathcal{O}'_{\delta,n}), \\ \text{and } \|\mathcal{R}'(\hat{x}) - b_n\|_2^2 \leq \frac{C}{\delta^3} \mathcal{E}(u, \mathcal{O}'_{\delta,n}) \implies \|\mathcal{R}' - b_n\|_{L^2(\omega''_{\delta,n}, \mathbb{R}^3)}^2 \leq \frac{C}{\delta} \mathcal{E}(u, \mathcal{O}'_{\delta,n}), \end{cases}$$

where $\|\cdot\|_2$ refers to the euclidian norm of \mathbb{R}^3 . From the inequalities (2.17), (2.18) and after elimination of the rigid displacement r_n , we obtain $\|u - U'_e\|_{L^2(\mathcal{O}''_{n,\delta}, \mathbb{R}^3)}^2 \leq C\delta^2 \mathcal{E}(u, \mathcal{O}'_{n,\delta})$. We add all these inequalities to obtain $\|u - U'_e\|_{L^2(\Omega'_\delta, \mathbb{R}^3)}^2 \leq C\delta^2 \mathcal{E}(u, \Omega'_\delta)$.

The components \mathcal{U}' and \mathcal{R}' of U'_e belong to $H^1(\omega_\delta, \mathbb{R}^3)$. The partial derivatives of these functions are

$$\frac{\partial \mathcal{U}'}{\partial x_\alpha}(\hat{x}) = \frac{6}{\pi\delta^3} \int_{B(\hat{x}; \delta/2)} \frac{\partial u}{\partial x_\alpha}(M) dM, \quad \frac{\partial \mathcal{R}'}{\partial x_\alpha}(\hat{x}) = \frac{24}{\pi\delta^5} \int_{B(\hat{x}; \delta/2)} \widehat{xM} \wedge \frac{\partial u}{\partial x_\alpha}(M) dM.$$

Let us calculate the means of $\frac{\partial}{\partial x_\alpha}(u - r_n)(M)$ and of $\widehat{xM} \wedge \frac{\partial}{\partial x_\alpha}(u - r_n)(M)$ in the ball $B(\hat{x}, \delta/2)$, $\hat{x} \in \omega''_{\delta,n}$. Thanks to (2.17) we obtain

$$(2.19) \quad \begin{cases} \left\| \frac{\partial \mathcal{U}'}{\partial x_\alpha}(\hat{x}) - b_n \wedge \vec{e}_\alpha \right\|_2^2 + \delta^2 \left\| \frac{\partial \mathcal{R}'}{\partial x_\alpha}(\hat{x}) \right\|_2^2 \leq \frac{C}{\delta^3} \mathcal{E}(u, \mathcal{O}'_{\delta,n}) \\ \implies \left\| \frac{\partial \mathcal{U}'}{\partial x_\alpha} - b_n \wedge \vec{e}_\alpha \right\|_{L^2(\omega''_{\delta,n}, \mathbb{R}^3)}^2 + \delta^2 \left\| \frac{\partial \mathcal{R}'}{\partial x_\alpha} \right\|_{L^2(\omega''_{\delta,n}, \mathbb{R}^3)}^2 \leq \frac{C}{\delta} \mathcal{E}(u, \mathcal{O}'_{\delta,n}), \end{cases}$$

hence, on the one hand $\delta \left\| \frac{\partial \mathcal{U}'}{\partial x_\alpha} - \mathcal{R}' \wedge \vec{e}_\alpha \right\|_{L^2(\omega''_{n,\delta}, \mathbb{R}^3)}^2 \leq C\mathcal{E}(u, \mathcal{O}'_{n,\delta})$ using (2.18) and on the other hand $\delta^3 \|\nabla \mathcal{R}\|_{L^2(\omega''_{n,\delta}, \mathbb{R}^6)}^2 \leq C\mathcal{E}(u, \mathcal{O}'_{n,\delta})$. We add all these inequalities

$$\delta \left\| \frac{\partial \mathcal{U}'}{\partial x_\alpha} - \mathcal{R}' \wedge \vec{e}_\alpha \right\|_{L^2(\omega_\delta, \mathbb{R}^3)}^2 \leq C\mathcal{E}(u, \Omega'_\delta), \quad \delta^3 \|\nabla \mathcal{R}\|_{L^2(\omega_\delta, \mathbb{R}^6)}^2 \leq C\mathcal{E}(u, \Omega'_\delta).$$

From (2.18) and (2.19) once more we obtain

$$\mathcal{D}(u - U'_e, \mathcal{O}''_{n,\delta}) \leq C\mathcal{E}(u, \mathcal{O}'_{n,\delta}) \implies \mathcal{D}(u - U'_e, \Omega'_\delta) \leq C\mathcal{E}(u, \Omega'_\delta).$$

Theorem 2.10 is proved. \square

3. The displacements of a structure and the asymptotic behavior of a structure made of plates

3.1. The structure made of plates

We work on a set of N plane bounded domains with polygonal boundary, included in \mathbb{R}^3 , $(\omega_l)_{1 \leq l \leq N}$. The skeleton \mathbf{S} is the union of $(\bar{\omega}_l)_{1 \leq l \leq N}$. A *face* of \mathbf{S} is a closed set $\bar{\omega}_l$. An *edge* of \mathbf{S} is a maximal segment shared by a set of faces or a maximal segment belonging to the boundary of a face. A *vertex* of \mathbf{S} is an extremity of an edge.

Hypotheses : We suppose

H1 • for any pair of faces $(\bar{\omega}_l, \bar{\omega}_p)$, there exists a sequence of faces $\bar{\omega}_l = \bar{\omega}_{l_0}, \bar{\omega}_{l_1}, \dots, \bar{\omega}_{l_k} = \bar{\omega}_p$ such that $\bar{\omega}_{l_r}$ and $\bar{\omega}_{l_{r+1}}$ have an edge in common, $0 \leq r \leq k-1$,

H2 • for any vertex A and any pair of faces $(\bar{\omega}_l, \bar{\omega}_p)$ containing A , there exists a sequence of faces $\bar{\omega}_l = \bar{\omega}_{l_0}, \bar{\omega}_{l_1}, \dots, \bar{\omega}_{l_k} = \bar{\omega}_p$ such that $\bar{\omega}_{l_r}$ and $\bar{\omega}_{l_{r+1}}$ have an edge in common containing A , $0 \leq r \leq k-1$,

H3 • the skeleton \mathbf{S} is fixed all along some edges.




Figure 1. The skeleton \mathbf{S} .

We denote

- Γ_0 the fixed part of the skeleton,
- \mathcal{J} the set of edges common to several faces,
- \mathcal{N} the set of vertexes common to several faces.

The structure made of plates is the domain $\mathcal{S}_\delta = \{x \in \mathbb{R}^3 \mid \text{dist}(x, \mathbf{S}) < \delta\}$. This structure is made of the gathering of the plates $\Omega'_{l,\delta}$ with thickness 2δ , with middle surface $\omega_{l,\delta}$ and with rounded rim. Each domain $\Omega'_{l,\delta}$ is equipped with a local frame $(O^{(l)}; \bar{e}_1^{(l)}, \bar{e}_2^{(l)}, \bar{e}_3^{(l)})$, $O^{(l)} \in \omega_l$, $\bar{e}_3^{(l)}$ is the normal direction to the face $\bar{\omega}_l$. The plate $\Omega'_{l,\delta}$ contains the plate $\Omega_{l,\delta} = \omega_l \times]-\delta, \delta[$ (\mathbb{R}^3 being equipped with the above local frame). The reference plate is the open set $\Omega_l = \omega_l \times]-1, 1[$ obtained through the transformation of $\Omega_{l,\delta}$ by the orthogonal affinity of ratio $1/\delta$.

The structure \mathcal{S}_δ is fixed to part $\Gamma_{0,\delta} = \{x \in \partial\mathcal{S}_\delta \mid \text{dist}(x, \Gamma_0) = \delta\}$ of its boundary. For each edge $J \in \mathcal{J}$ we choose a unit vector \bar{e}_J in the direction of the edge.

We consider $\mathbf{S}_\delta = \bigcup_{l=1}^N \bar{\omega}_{l,\delta}$, the set of the middle surfaces of the plates.

There exist two constants $\delta_0 > 0$ and $\mu_0 > 0$ depending only on the skeleton \mathbf{S} such that for any $\delta \in]0, \delta_0[$ the parts common to several plates are in the union of the junctions

$$\bigcup_{J \in \mathcal{J}} \{x \in \mathbb{R}^3 \mid \text{dist}(x, J) < \eta_0 \delta\}.$$

The restriction of a function ϕ , defined on \mathbf{S} , (resp. $\mathbf{S}_\delta, \mathcal{S}_\delta$), to ω_l , (resp. $\omega_{l,\delta}, \Omega'_{l,\delta}$) is denoted $\phi^{(l)}$. In the same way, we denote $x_\alpha^{(l)} = x^{(l)} \cdot \bar{e}_\alpha^{(l)}$ and $x_3^{(l)} = x^{(l)} \cdot \bar{e}_3^{(l)}$ the local variables ($x^{(l)} = (\hat{x}^{(l)}, x_3^{(l)}) = (x_1^{(l)}, x_2^{(l)}, x_3^{(l)})$). The space $H^1(\mathbf{S}, \mathbb{R}^n)$ (resp. $H^1(\mathbf{S}_\delta, \mathbb{R}^n)$) is the set of the functions ϕ defined a.e. in \mathbf{S} (resp. \mathbf{S}_δ), with values in \mathbb{R}^n , such that the restriction $\phi^{(l)}$ belongs to $H^1(\omega_l, \mathbb{R}^n)$ (resp. $H^1(\omega_{l,\delta}, \mathbb{R}^n)$) and such that for any edge $J \in \mathcal{J}$ and any pair of faces $(\bar{\omega}_l, \bar{\omega}_k)$ containing J , we have the equality of the restrictions to J , $(\phi^{(l)})|_J = (\phi^{(k)})|_J$ in $H^{1/2}(J, \mathbb{R}^n)$.

The space $H_{\Gamma_0}^1(\mathbf{S}, \mathbb{R}^n)$ is the subspace of $H^1(\mathbf{S}, \mathbb{R}^n)$ the elements of which are a.e. equal to zero on Γ_0 . We equip $H^1(\mathbf{S}_\delta, \mathbb{R}^3)$ with the inner product

$$[\mathcal{U}, \mathcal{V}] = \sum_{l=1}^N \int_{\omega_{l,\delta}} \left\{ \mathcal{U}^{(l)} \cdot \mathcal{V}^{(l)} + \frac{\partial \mathcal{U}^{(l)}}{\partial x_\alpha^{(l)}} \cdot \frac{\partial \mathcal{V}^{(l)}}{\partial x_\alpha^{(l)}} \right\}.$$

The associated norm is denoted $\|\cdot\|$.

3.2. The elementary displacements of plates structure

Definition 3.1 : An elementary displacement of a plates structure (e.d.p.s.) is a displacement $\Phi \in H^1(\mathcal{S}_\delta, \mathbb{R}^3)$, such that there exist two elements \mathcal{A} and \mathcal{B} belonging to $H^1(\mathbf{S}_\delta, \mathbb{R}^3)$, such that for any $l \in \{1, \dots, N\}$,

$$\Phi^{(l)}(x) = \mathcal{A}^{(l)}(\hat{x}^{(l)}) + \mathcal{B}^{(l)}(\hat{x}^{(l)}) \wedge x_3^{(l)} \bar{e}_3^{(l)}$$

is an e.p.d. of the plate $\Omega'_{l,\delta}$.

The functions \mathcal{A} and \mathcal{B} are respectively the first component and the second component of the e.d.p.s. Φ . The function \mathcal{A} accounts for the displacement of the skeleton faces while \mathcal{B} accounts for the rotation of the normal directions to the plates and of the rotation of the faces around the edges.

Theorem 3.2 : For any displacement $u \in H_{\Gamma_0}^1(\mathcal{S}_\delta, \mathbb{R}^3)$ there exists an e.d.p.s. U_e of components $(\mathcal{U}, \mathcal{R}) \in H_{\Gamma_0}^1(\mathbf{S}_\delta, \mathbb{R}^3) \times H_{\Gamma_0}^1(\mathbf{S}_\delta, \mathbb{R}^3)$ such that

$$(3.1) \quad \begin{cases} \sum_{l=1}^N \left\{ \delta^3 \|\nabla \mathcal{R}^{(l)}\|_{L^2(\omega_{l,\delta}, \mathbb{R}^6)}^2 + \delta \left\| \frac{\partial \mathcal{U}^{(l)}}{\partial x_\alpha^{(l)}} - \mathcal{R}^{(l)} \wedge \bar{e}_\alpha^{(l)} \right\|_{L^2(\omega_{l,\delta}, \mathbb{R}^3)}^2 \right\} \leq C \mathcal{E}(u, \mathcal{S}_\delta), \\ \mathcal{E}(U_e, \mathcal{S}_\delta) + \mathcal{D}(u - U_e, \mathcal{S}_\delta) \leq C \mathcal{E}(u, \mathcal{S}_\delta), \quad \|u - U_e\|_{L^2(\mathcal{S}_\delta, \mathbb{R}^3)}^2 \leq C \delta^2 \mathcal{E}(u, \mathcal{S}_\delta). \end{cases}$$

Proof : Let u be in $H_{\Gamma_0}^1(\mathcal{S}_\delta, \mathbb{R}^3)$. Thanks to Lemma 4.1, for any l belonging to $\{1, \dots, N\}$, we extend the restriction $u^{(l)}$ to the plate $\Omega'_{l,\delta}$, into a displacement of $\tilde{\Omega}_{l,\delta}$ (the plate of thickness 2δ and of middle surface $\omega_{l,2\delta}$). Therefore, using the formulas (2.15), we can define an e.p.d. $U_e'^{(l)}$ of the plate $\Omega'_{l,\delta}$ verifying (2.16). Both components $\mathcal{U}'^{(l)}$ and $\mathcal{R}'^{(l)}$ of $U_e'^{(l)}$ are the restrictions of elements belonging to $H_{\Gamma_0}^1(\mathbf{S}_\delta, \mathbb{R}^3)$ (corollary of Lemma 2.8 in Annex B).

Then we build a new e.d.p.s. U_e equal to U_e' in the open set $\bigcup_{J \in \mathcal{J}} \{x \in \mathcal{S}_\delta \mid \text{dist}(x, J) > 2\eta_0 \delta\}$ and equal to an elementary displacement of rods structures in the junctions (see Annex B). We then can deduce (3.1). \square

Proposition 3.3 (Korn inequality) : For any displacement $u \in H_{\Gamma_0}^1(\mathcal{S}_\delta, \mathbb{R}^3)$, we have

$$(3.2) \quad \delta \|\mathcal{R}\|^2 + \delta \|\mathcal{U}\|^2 + \mathcal{D}(u, \mathcal{S}_\delta) + \|u\|_{L^2(\mathcal{S}_\delta, \mathbb{R}^3)}^2 \leq \frac{C}{\delta^2} \mathcal{E}(u, \mathcal{S}_\delta).$$

The constant does not depend on δ .

Proof : The estimates (3.1) of the gradients of the $\mathcal{R}^{(l)}$ functions, the nullity of \mathcal{R} on Γ_0 and the hypothesis **H1** allow us to obtain step by step $\|\mathcal{R}\|_{L^2(\mathbf{S}_\delta, \mathbb{R}^3)}^2 \leq C/\delta^3 \mathcal{E}(u, \mathcal{S}_\delta)$. This inequality and (3.1) give then an upperbound of the L^2 norms of the functions gradients $\mathcal{U}^{(l)}$. The nullity of \mathcal{U} on Γ_0 and the hypothesis **H1** imply then that $\|\mathcal{U}\|_{L^2(\mathbf{S}_\delta, \mathbb{R}^3)}^2 \leq C/\delta^3 \mathcal{E}(u, \mathcal{S}_\delta)$. From these estimates of \mathcal{U} and \mathcal{R} follow $\mathcal{D}(U_e, \mathcal{S}_\delta) \leq C/\delta^2 \mathcal{E}(u, \mathcal{S}_\delta)$ and $\|U_e\|_{L^2(\mathbf{S}_\delta, \mathbb{R}^3)}^2 \leq C/\delta^2 \mathcal{E}(u, \mathcal{S}_\delta)$. Then again, thanks to (3.1), we obtain the estimates of the L^2 norm of u and of its gradient. \square

3.3. Inextensional displacements, extensional displacements

The space $H_{\rho, \Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$ is the set of the functions ϕ defined a.e. in \mathbf{S} , with values in \mathbb{R}^3 , such that:

- for any $l \in \{1, \dots, N\}$, the restrictions $\phi_1^{(l)}$ and $\phi_2^{(l)}$ belong to $H^1(\omega_l)$ and $\phi_3^{(l)}$ belongs to

$$H_{\rho}^1(\omega_l) = \left\{ \psi \in L^2(\omega_l) \mid \sqrt{\rho} \nabla \psi \in [L^2(\omega_l)]^2 \right\}$$

where $\rho(\hat{x}) = \text{dist}(\hat{x}, \mathcal{N})$ (distance from the point $\hat{x} \in \mathbf{S}_\delta$ to the vertexes belonging to several faces),

- for any edge $J \in \mathcal{J}$ and any pair of faces $(\bar{\omega}_l, \bar{\omega}_k)$ containing J , we have $(\phi^{(l)})|_J = (\phi^{(k)})|_J$ in $H^{1/2}(J, \mathbb{R}^3)$,
- the function ϕ is equal to zero on Γ_0 .

We equip $H_{\rho, \Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$ with the inner product

$$\langle \mathcal{U}, \mathcal{V} \rangle_{\rho} = \sum_{l=1}^N \int_{\omega_l} \left\{ \gamma_{\alpha\beta}(\mathcal{U}^{(l)}) \gamma_{\alpha\beta}(\mathcal{V}^{(l)}) + \rho \nabla \mathcal{U}_3^{(l)} \cdot \nabla \mathcal{V}_3^{(l)} \right\},$$

and with the norm $\|\mathcal{U}\|_{\rho} = \sqrt{\langle \mathcal{U}, \mathcal{U} \rangle_{\rho}}$. The usual norm on $H_{\rho, \Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$ is

$$\|\mathcal{U}\|_{\rho} = \sqrt{\sum_{l=1}^N \int_{\omega_l} \left\{ |\nabla \mathcal{U}_1^{(l)}|^2 + |\nabla \mathcal{U}_2^{(l)}|^2 + \rho |\nabla \mathcal{U}_3^{(l)}|^2 \right\}}$$

Lemma 3.4 : *The norms $\|\cdot\|_{\rho}$ and $|\cdot|_{\rho}$ are equivalent in $H_{\rho, \Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$. Moreover the space $H_{\Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$ is dense in $H_{\rho, \Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$.*

Proof : See Annex C. \square

Definition 3.5 : An **inextensional displacement of the skeleton** is an element U belonging to $H_{\rho, \Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$ such that

$$\forall l \in \{1, \dots, N\}, \quad \gamma_{\alpha\beta}(U^{(l)}) = 0, \quad \text{in } \omega_l.$$

The membrane component $U_M^{(l)} = U_1^{(l)} \bar{e}_1^{(l)} + U_2^{(l)} \bar{e}_2^{(l)}$ of an inextensional displacement is a rigid displacement of the face $\bar{\omega}_l$. The inextensional displacements space of the skeleton is denoted $D_I(\mathbf{S})$.

Definition 3.6 : An **extensional displacement of the skeleton** is an element of the orthogonal $D_E(\mathbf{S})$ of $D_I(\mathbf{S})$ in $H_{\rho, \Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$.

The set of extensional displacements is equipped with the semi-norm

$$\|U\|_E = \sqrt{\sum_{l=1}^N \int_{\omega_l} \gamma_{\alpha\beta}(U^{(l)}) \gamma_{\alpha\beta}(U^{(l)})}, \quad U \in D_E(\mathbf{S}).$$

The semi-norm $\|\cdot\|_E$ is a norm, because if $U \in D_E(\mathbf{S})$ is such that $\|U\|_E = 0$ then, $\gamma_{\alpha\beta}(U^{(l)}) = 0$ for any l . The displacement U is then of inextensional type and is equal to zero.

Lemma 3.7 : *The norms $\|\cdot\|_E$ and $|\cdot|_\rho$ are equivalent in $D_E(\mathbf{S})$.*

Proof : See Annex C. □

Corollary of Lemma 3.7 : Let u be a displacement belonging to $H_{\Gamma_0}^1(\mathcal{S}_\delta, \mathbb{R}^3)$ and U_e the e.d.p.s. given by Theorem 3.2. The restriction to \mathbf{S} of the first component \mathcal{U} of U_e can be written as the sum of an extensional displacement and an inextensional displacement,

$$(3.3) \quad \mathcal{U} = U_E + U_I, \quad U_E \in D_E(\mathbf{S}), \quad U_I \in D_I(\mathbf{S}).$$

According to the inequalities (3.1) and Lemma 3.7 we have

$$(3.4) \quad |U_E|_\rho^2 \leq C \|U_E\|_E^2 \leq \frac{C}{\delta} \mathcal{E}(u, \mathcal{S}_\delta), \quad |U_I|_\rho^2 \leq \frac{C}{\delta^3} \mathcal{E}(u, \mathcal{S}_\delta).$$

The constants are independent of δ .

3.4. The limit displacements

Let $(u_\delta)_{\delta>0}$ be a sequence of displacements belonging to $H_{\Gamma_0}^1(\mathcal{S}_\delta, \mathbb{R}^3)$ and verifying

$$(3.5) \quad \mathcal{E}(u_\delta, \mathcal{S}_\delta) \leq C\delta,$$

where the constant is independent of δ . Thanks to the estimates (3.2), (3.4) and (2.4), from the sequences $\delta\mathcal{U}_\delta, \delta\mathcal{R}_\delta, \delta U_{I,\delta}, U_{E,\delta}$ and $\frac{1}{\delta}\mathcal{T}_\delta(\tilde{u}_\delta^{(l)})$ we extract some sub-sequences, still denoted in the same way and which weakly converge,

$$(3.6) \quad \left\{ \begin{array}{ll} \delta\mathcal{U}_\delta \rightharpoonup U_I, & \delta\mathcal{R}_\delta \rightharpoonup \mathcal{R} \quad \text{weakly in } H_{\Gamma_0}^1(\mathbf{S}, \mathbb{R}^3), \\ \delta U_{I,\delta} \rightharpoonup U_I & \text{weakly in } D_I(\mathbf{S}), \\ U_{E,\delta} \rightharpoonup U_E & \text{weakly in } D_E(\mathbf{S}), \\ \frac{1}{\delta}\mathcal{T}_\delta(\tilde{u}_\delta^{(l)}) \rightharpoonup \tilde{u}^{(l)} & \text{weakly in } L^2(\omega_l, H^1(-1, 1], \mathbb{R}^3)). \end{array} \right.$$

The sequences $\delta\mathcal{U}_\delta$ and $\delta U_{I,\delta}$ have the same limit in $H_{\rho,\Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$ because the sequence $\delta U_{E,\delta}$ converges to 0 in $H_\rho^1(\mathbf{S}, \mathbb{R}^3)$. After passing to the limit and from (3.1) comes

$$(3.7) \quad \forall l \in \{1, \dots, N\}, \quad \forall \alpha \in \{1, 2\}, \quad \frac{\partial U_I^{(l)}}{\partial x_\alpha^{(l)}} = \mathcal{R}^{(l)} \wedge \tilde{e}_\alpha^{(l)}.$$

Now we define the space of the inextensional displacements limits. We put

$$\mathcal{D}_I(\mathbf{S}) = \left\{ \mathcal{A} \in D_I(\mathbf{S}) \cap H_{\Gamma_0}^1(\mathbf{S}, \mathbb{R}^3) \mid \exists \mathcal{B} \in H_{\Gamma_0}^1(\mathbf{S}, \mathbb{R}^3), \frac{\partial \mathcal{A}^{(l)}}{\partial x_\alpha^{(l)}} = \mathcal{B}^{(l)} \wedge \tilde{e}_\alpha^{(l)}, \quad \forall l \in \{1, \dots, N\} \right\}$$

For any $\mathcal{A} \in \mathcal{D}_I(\mathbf{S})$, there is only one \mathcal{B} which we denote $\widehat{\nabla}\mathcal{A}$. Then we have

$$\forall l \in \{1, \dots, N\}, \quad \forall \alpha \in \{1, 2\}, \quad \frac{\partial \mathcal{A}^{(l)}}{\partial x_\alpha^{(l)}} = \widehat{\nabla}\mathcal{A}^{(l)} \wedge \tilde{e}_\alpha^{(l)}.$$

We equip $\mathcal{D}_I(\mathbf{S})$ with the norm $\|\mathcal{A}\|_I = \|\widehat{\nabla}\mathcal{A}\|_{H^1(\mathbf{S}, \mathbb{R}^3)}$. The inextensional limit displacement U_I belongs to $\mathcal{D}_I(\mathbf{S})$.

3.5. Limit of the unfolded displacements and limit of the unfolded strain tensor components

Using (2.8) and after transformation by unfolding we have the following limits in the referring plates :

$$(3.8) \quad \begin{cases} \delta \mathcal{T}_\delta(u_\delta^{(l)}) \rightharpoonup U_I^{(l)} & \text{weakly in } H^1(\Omega_l, \mathbb{R}^3) \\ \mathcal{T}_\delta(u_{\alpha,\delta}^{(l)} - U_{I,\alpha,\delta}^{(l)}) \rightharpoonup U_{E,\alpha}^{(l)} - t_3^{(l)} \frac{\partial U_{I,3}^{(l)}}{\partial x_\alpha^{(l)}} & \text{weakly in } H^1(\Omega_l), \\ \mathcal{T}_\delta(\gamma_{\alpha\beta}(u_\delta^{(l)})) \rightharpoonup \frac{1}{2} \left\{ \frac{\partial U_{E,\alpha}^{(l)}}{\partial x_\beta^{(l)}} + \frac{\partial U_{E,\beta}^{(l)}}{\partial x_\alpha^{(l)}} \right\} - t_3^{(l)} \frac{\partial^2 U_{I,3}^{(l)}}{\partial x_\alpha^{(l)} \partial x_\beta^{(l)}} & \text{weakly in } L^2(\Omega_l), \\ \mathcal{T}_\delta(\gamma_{\alpha 3}(u_\delta^{(l)})) \rightharpoonup \frac{1}{2} \frac{\partial \tilde{u}_\alpha^{(l)}}{\partial t_3^{(l)}}, \quad \mathcal{T}_\delta(\gamma_{33}(u_\delta^{(l)})) \rightharpoonup \frac{\partial \tilde{u}_3^{(l)}}{\partial t_3^{(l)}} & \text{weakly in } L^2(\Omega_l). \end{cases}$$

3.6. Elasticity problem

The plates are made of an homogeneous and isotropic material. Our equations are given within the framework of linearised elasticity. In \mathcal{S}_δ let the elasticity system be

$$(3.9) \quad \begin{cases} -\frac{\partial}{\partial x_j} \left\{ a_{ij'i'j'} \frac{\partial u_{i',\delta}}{\partial x_{j'}} \right\} = F_{i,\delta} & \text{in } \mathcal{S}_\delta, \\ u_\delta = 0 & \text{on } \Gamma_{0,\delta}, \\ a_{ij'i'j'} \frac{\partial u_{i',\delta}}{\partial x_{j'}} n_j = 0 & \text{in } \Gamma_\delta, \quad \Gamma_\delta = \partial \mathcal{S}_\delta \setminus \Gamma_{0,\delta}. \end{cases}$$

The variational formulation of the problem (3.9) is

$$(3.10) \quad \begin{cases} u_\delta \in H_{\Gamma_0}^1(\mathcal{S}_\delta; \mathbb{R}^3) \\ \int_{\mathcal{S}_\delta} a_{ij'i'j'} \gamma_{ij}(u_\delta) \gamma_{i'j'}(v) = \int_{\mathcal{S}_\delta} F_\delta \cdot v & \forall v \in H_{\Gamma_0}^1(\mathcal{S}_\delta; \mathbb{R}^3) \end{cases}$$

where $a_{ij'i'j'} = \lambda \delta_{ij} \delta_{i'j'} + \mu (\delta_{ii'} \delta_{jj'} + \delta_{ij'} \delta_{ji'})$. The constants λ and μ are the Lamé constants of the material. The plates $\Omega_{l,\delta}$ are submitted to volume applied forces . Among these forces we make a distinction between those concerning the extensional displacements and those concerning the inextensional displacements.

$$(3.11) \quad F_\delta(x) = \sum_{l=1}^N \{ \delta f_I(\hat{x}^{(l)}) + f_E(\hat{x}^{(l)}) \} \mathbf{1}_{\Omega_{l,\delta}}(x), \quad f_I, f_E \in L^2(\mathbf{S}, \mathbb{R}^3),$$

where $\Omega_{l,\delta} = \omega_l \times]-\delta, \delta[$ (in the local frame) and where $\mathbf{1}_{\Omega_{l,\delta}}$ is the characteristic function of the open set $\Omega_{l,\delta}$. Hence several volume forces are stacked up in the junctions.

The function f_E verifies the condition of orthogonality

$$(3.12) \quad \forall V \in D_I(\mathbf{S}), \quad \int_{\mathbf{S}} f_E \cdot V = 0.$$

Let $(\mathcal{U}_\delta, \mathcal{R}_\delta)$ be the two components of the e.d.p.s. associated to the solution u_δ of the problem (3.10). In the plate $\Omega_{l,\delta}$, the displacement $u_\delta^{(l)}$ is the sum of the e.p.d. $\mathcal{U}_\delta^{(l)}(\hat{x}^{(l)}) + \mathcal{R}_\delta^{(l)}(\hat{x}^{(l)}) \wedge x_3^{(l)} \bar{e}_3^{(l)}$ and a residual displacement. The displacement \mathcal{U}_δ is the sum of an extensional displacement $U_{E,\delta}$ and of an inextensional displacement $U_{I,\delta}$. Then, thanks to (3.1) and (3.4), we have

$$(3.13) \quad \left| \frac{1}{2\delta} \int_{\mathcal{S}_\delta} F_\delta \cdot u_\delta - \int_{\mathbf{S}} f_E \cdot U_{E,\delta} - \delta \int_{\mathbf{S}} f_I \cdot U_{I,\delta} \right| \leq C \{ \|f_E\|_{L^2(\mathbf{S}, \mathbb{R}^3)} + \|f_I\|_{L^2(\mathbf{S}, \mathbb{R}^3)} \} \sqrt{\mathcal{E}(u_\delta, \mathcal{S}_\delta)}$$

hence

$$\left| \int_{\mathcal{S}_\delta} F_\delta \cdot u_\delta \right| \leq C \{ \|f_E\|_{L^2(\mathbf{S}, \mathbb{R}^3)} + \|f_I\|_{L^2(\mathbf{S}, \mathbb{R}^3)} \} \sqrt{\delta} \sqrt{\mathcal{E}(u_\delta, \mathcal{S}_\delta)}.$$

We deduce that the solution of the variational problem (3.10) verifies the estimation

$$(3.14) \quad \mathcal{E}(u_\delta, \mathcal{S}_\delta) \leq C\delta \{ \|f_E\|_{L^2(\mathbf{S}, \mathbb{R}^3)}^2 + \|f_I\|_{L^2(\mathbf{S}, \mathbb{R}^3)}^2 \}.$$

3.7. Asymptotic behavior of the stress tensor

We begin with determining the partial derivatives of the residual displacements $\tilde{u}^{(l)}$ in the normal directions to the plates.

Let ϕ be a displacement of $H^1(\Omega_l, \mathbb{R}^3)$, equal to zero in the neighborhood of all the sets $J \times]-1, 1[$ where J is an edge of the face $\bar{\omega}_l$. For δ small, the displacement $\phi_\delta(x) = \delta\phi(\hat{x}^{(l)}, \frac{x_3^{(l)}}{\delta})$ is an acceptable displacement of the full structure \mathcal{S}_δ . We have the following strong convergences of the unfolded of the strained tensor components of ϕ_δ :

$$(3.15) \quad \begin{cases} \mathcal{T}_\delta(\gamma_{\alpha\beta}(\phi_\delta)) \longrightarrow 0 & \text{strongly in } L^2(\Omega_l), \\ \mathcal{T}_\delta(\gamma_{\alpha 3}(\phi_\delta)) \longrightarrow \frac{1}{2} \frac{\partial \phi_\alpha}{\partial t_3^{(l)}} & \text{strongly in } L^2(\Omega_l), \\ \mathcal{T}_\delta(\gamma_{33}(\phi_\delta)) \longrightarrow \frac{\partial \phi_3}{\partial t_3^{(l)}} & \text{strongly in } L^2(\Omega_l). \end{cases}$$

We now take ϕ_δ as a test-displacement in (3.10), we transform by unfolding the integral on $\Omega_{l,\delta}$ into an integral on Ω_l and after dividing by the thickness of the plate we pass to the limit. We obtain

$$(3.16) \quad \int_{\Omega_l} \left[\lambda \left\{ \frac{\partial U_{E,1}^{(l)}}{\partial x_1^{(l)}} - t_3^{(l)} \frac{\partial^2 U_{I,3}^{(l)}}{\partial x_1^{(l),2}} + \frac{\partial U_{E,2}^{(l)}}{\partial x_2^{(l)}} - t_3^{(l)} \frac{\partial^2 U_{I,3}^{(l)}}{\partial x_2^{(l),2}} \right\} + (\lambda + 2\mu) \frac{\partial \tilde{u}_3^{(l)}}{\partial t_3^{(l)}} \right] \frac{\partial \phi_3}{\partial t_3^{(l)}} + \mu \left[\frac{\partial \tilde{u}_1^{(l)}}{\partial t_3^{(l)}} \frac{\partial \phi_1}{\partial t_3^{(l)}} + \frac{\partial \tilde{u}_2^{(l)}}{\partial t_3^{(l)}} \frac{\partial \phi_2}{\partial t_3^{(l)}} \right] = 0$$

because the right member of (3.10) tends to 0 ($\frac{1}{2\delta} \left| \int_{\Omega_{l,\delta}} F_\delta \cdot \phi_\delta \right| \leq C\delta \|\phi\|_{L^2(\Omega_l, \mathbb{R}^3)}$).

The set of these test-displacements is a dense subset in $L^2(\omega_l, H^1(]-1, 1[, \mathbb{R}^3))$. Hence the equality (3.16) is verified for any element of $L^2(\omega_l, H^1(]-1, 1[, \mathbb{R}^3))$. We deduce the partial derivatives $\frac{\partial \tilde{u}^{(l)}}{\partial t_3^{(l)}}$ in terms of the first partial derivatives of $U_E^{(l)}$ and of the second partial derivatives of $U_I^{(l)}$,

$$(3.17) \quad \frac{\partial \tilde{u}_1^{(l)}}{\partial t_3^{(l)}} = \frac{\partial \tilde{u}_2^{(l)}}{\partial t_3^{(l)}} = 0, \quad \frac{\partial \tilde{u}_3^{(l)}}{\partial t_3^{(l)}} = \frac{\lambda}{\lambda + 2\mu} \left\{ -\frac{\partial U_{E,1}^{(l)}}{\partial x_1^{(l)}} - \frac{\partial U_{E,2}^{(l)}}{\partial x_2^{(l)}} + t_3^{(l)} \Delta U_{I,3}^{(l)} \right\}.$$

We give now the weak limit in $L^2(\Omega_l)$ of the unfolded of the stress tensor components

$$(3.18) \quad \begin{cases} \mathcal{T}_\delta(\sigma_{11}(u_\delta^{(l)})) \rightharpoonup \frac{E}{1-\nu^2} \left[\frac{\partial U_{E,1}^{(l)}}{\partial x_1^{(l)}} - t_3^{(l)} \frac{\partial^2 U_{I,3}^{(l)}}{\partial x_1^{(l),2}} + \nu \left\{ \frac{\partial U_{E,2}^{(l)}}{\partial x_2^{(l)}} - t_3^{(l)} \frac{\partial^2 U_{I,3}^{(l)}}{\partial x_2^{(l),2}} \right\} \right], \\ \mathcal{T}_\delta(\sigma_{12}(u_\delta^{(l)})) \rightharpoonup \mu \left\{ \frac{\partial U_{E,1}^{(l)}}{\partial x_2^{(l)}} + \frac{\partial U_{E,2}^{(l)}}{\partial x_1^{(l)}} - 2t_3^{(l)} \frac{\partial^2 U_{I,3}^{(l)}}{\partial x_1^{(l)} \partial x_2^{(l)}} \right\}, \\ \mathcal{T}_\delta(\sigma_{22}(u_\delta^{(l)})) \rightharpoonup \frac{E}{1-\nu^2} \left[\frac{\partial U_{E,2}^{(l)}}{\partial x_2^{(l)}} - t_3^{(l)} \frac{\partial^2 U_{I,3}^{(l)}}{\partial x_2^{(l),2}} + \nu \left\{ \frac{\partial U_{E,1}^{(l)}}{\partial x_1^{(l)}} - t_3^{(l)} \frac{\partial^2 U_{I,3}^{(l)}}{\partial x_1^{(l),2}} \right\} \right], \\ \mathcal{T}_\delta(\sigma_{i3}(u_\delta^{(l)})) \rightharpoonup 0. \end{cases}$$

3.8. The extensional displacement U_E or the problem of coupled membrane plates

Theorem 3.8 : *The extensional displacement U_E is the solution of the variational problem*

$$(3.19) \quad \frac{E}{1-\nu^2} \sum_{l=1}^N \int_{\omega_l} [(1-\nu)\gamma_{\alpha\beta}(U_E^{(l)})\gamma_{\alpha\beta}(V^{(l)}) + \nu\gamma_{\alpha\alpha}(U_E^{(l)})\gamma_{\beta\beta}(V^{(l)})] = \int_{\mathbf{S}} f_E \cdot V, \quad \forall V \in D_E(\mathbf{S}).$$

where E is the Young modulus and ν the Poisson constant . □

The proof of Theorem 3.8 requires the next lemma.

Lemma 3.9 : *For any element $V \in H_{\Gamma_0}^1(\mathbf{S}_{\delta_0}, \mathbb{R}^3)$, there exists a sequence of displacements $(V_\delta)_{0 < \delta \leq \delta_0}$ belonging to $H_{\Gamma_0}^1(\mathbf{S}_{\delta_0}, \mathbb{R}^3) \cap H_{\Gamma_0}^1(\mathcal{S}_\delta, \mathbb{R}^3)$ such that*

$$(3.20) \quad V_\delta \longrightarrow V \quad \text{strongly in } H_{\Gamma_0}^1(\mathbf{S}_{\delta_0}, \mathbb{R}^3).$$

Proof : See Annex E. □

Proof of Theorem 3.8 : Let V be an element of $H_{\Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$, we extend V into an element, still denoted V , of the space $H_{\Gamma_0}^1(\mathbf{S}_{\delta_0}, \mathbb{R}^3)$. We take V_δ as a test-displacement in (3.10), we transform, by unfolding, the integrals on the plates into integrals on the reference plates then we divide by 2δ . Thus we are led to take into account again and again the neighborhoods of the edges belonging to \mathcal{J} .

Let J be an edge common to several faces. For any face $\bar{\omega}_l$ containing J , we have

$$(3.21) \quad \mathbf{1}_{\{\hat{x}^{(l)} \in \omega_{l,\delta} \mid \text{dist}(\hat{x}^{(l)}, J) < \eta_0 \delta\}} \mathcal{T}_\delta(\gamma_{ij}(V_\delta^{(l)})) \longrightarrow 0 \quad \text{strongly in } L^2(\Omega_l).$$

The part of \mathcal{S}_δ neighbour of the edge J and common to several plates is contained into the cylinder $\{x \in \mathbb{R}^3 \mid \text{dist}(x, J) < \eta_0 \delta\}$. Thanks to the convergences (3.21) its contribution in the limit problem is equal to zero. Then we can make δ tends to 0 in order to obtain (3.19) with the displacement V . The limit of the right handside term of (3.10) is given by (3.13).

The set $H_{\Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$ is dense in $H_{\rho, \Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$ (Lemma B.3), which gives (3.19) with any displacement of $D_E(\mathbf{S})$. □

3.9. The inextensional displacement U_I or the problem of coupled bending plates

Theorem 3.10 : *The inextensional displacement U_I is the solution of the variational problem*

$$(3.22) \quad \frac{E}{3(1-\nu^2)} \sum_{l=1}^N \int_{\omega_l} \left[(1-\nu) \frac{\partial^2 U_{I,3}^{(l)}}{\partial x_\alpha^{(l)} \partial x_\beta^{(l)}} \frac{\partial^2 V_3^{(l)}}{\partial x_\alpha^{(l)} \partial x_\beta^{(l)}} + \nu \Delta U_{I,3}^{(l)} \Delta V_3^{(l)} \right] = \int_{\mathbf{S}} f_I \cdot V, \quad \forall V \in \mathcal{D}_I(\mathbf{S}).$$

The proof of Theorem 3.10 requires the next lemma.

Lemma 3.11 : *For any element $V \in \mathcal{D}_I(\mathbf{S})$, there exists a sequence of displacements $(W_\delta)_{0 < \delta \leq \delta_0}$ such that*

$$(3.23) \quad W_\delta \in H_{\Gamma_0}^1(\mathcal{S}_\delta, \mathbb{R}^3), \quad \text{and} \quad \begin{cases} \mathcal{T}_\delta(W_\delta^{(l)}) \longrightarrow V^{(l)} \quad \text{strongly in } L^2(\Omega_l, \mathbb{R}^3), \\ \mathcal{T}_\delta(\gamma_{\alpha\beta}(W_\delta^{(l)})) \longrightarrow -t_3^{(l)} \frac{\partial^2 V_3^{(l)}}{\partial x_\alpha^{(l)} \partial x_\beta^{(l)}} \quad \text{strongly in } L^2(\Omega_l), \\ \mathcal{T}_\delta(\gamma_{k3}(W_\delta^{(l)})) \longrightarrow 0 \quad \text{strongly in } L^2(\Omega_l). \end{cases}$$

Proof : See Annex E. □

Proof of Theorem 3.10 : Let W be an element of $\mathcal{D}_I(\mathbf{S})$. For any edge J and any face $\bar{\omega}_l$ containing J we have

$$(3.24) \quad \mathbf{1}_{\{\hat{x}^{(l)} \in \omega_{l,\delta} \mid \text{dist}(\hat{x}^{(l)}, J) < \eta_0 \delta\}} \mathcal{T}_\delta(\gamma_{ij}(W_\delta^{(l)})) \longrightarrow 0 \quad \text{strongly in } L^2(\Omega_l).$$

We take W_δ as a test-displacement in (3.10). We transform, by unfolding, the integrals on the plates into integrals on the reference plates, then we divide by 2δ . We pass to the limit (thanks to (3.24) the contribution of the immediate junction neighborhoods tends to 0). We obtain (3.22) with the test-displacement V . \square

Remark 3.12 : The problems (3.19) and (3.22) are coercive. It results that the whole encountered sequences converges to their limit. We are going to show now that these convergences are strong. We consider the formal displacement U of the structure \mathcal{S}_δ defined in each plate by

$$U^{(l)}(x) = U_E^{(l)}(\hat{x}^{(l)}) + \frac{1}{\delta} U_I^{(l)}(\hat{x}^{(l)}) + \widehat{\nabla} U_I^{(l)}(\hat{x}^{(l)}) \wedge x_3^{(l)} \bar{e}_3^{(l)}, \quad x \in \Omega_{l,\delta}.$$

Let $\mathbf{1}_{\mathcal{J}_\delta^c}$ be the characteristic function of the complement in \mathcal{S}_δ of the union of the edges neighborhoods $\bigcup_{J \in \mathcal{J}} \{x \in \mathcal{S}_\delta \mid \text{dist}(x, J) < \eta_0 \delta\}$. In the reference plate Ω_l , we have the convergences

$$\mathcal{T}_\delta(\gamma_{ij}(u_\delta^{(l)})) \mathcal{T}_\delta(\mathbf{1}_{\mathcal{J}_\delta^c}) \rightharpoonup 0 \quad \text{weakly in } L^2(\Omega_l).$$

Hence

$$\left\{ \begin{aligned} & \sum_{l=1}^N \int_{\Omega_l} \sigma_{ij}(U^{(l)}) \gamma_{ij}(U^{(l)}) \leq \liminf_{\delta \rightarrow 0} \sum_{l=1}^N \int_{\Omega_l} \mathcal{T}_\delta(\sigma_{ij}(u_\delta^{(l)})) \mathcal{T}_\delta(\gamma_{ij}(u_\delta^{(l)})) \mathcal{T}_\delta(\mathbf{1}_{\mathcal{J}_\delta^c}) \\ & = \liminf_{\delta \rightarrow 0} \frac{1}{2\delta} \int_{\mathcal{S}_\delta} \sigma_{ij}(u_\delta) \gamma_{ij}(u_\delta) \mathbf{1}_{\mathcal{J}_\delta^c} \leq \overline{\lim}_{\delta \rightarrow 0} \frac{1}{2\delta} \int_{\mathcal{S}_\delta} \sigma_{ij}(u_\delta) \gamma_{ij}(u_\delta) \\ & = \overline{\lim}_{\delta \rightarrow 0} \frac{1}{2\delta} \int_{\mathcal{S}_\delta} F_\delta \cdot u_\delta = \lim_{\delta \rightarrow 0} \frac{1}{2\delta} \int_{\mathcal{S}_\delta} F_\delta \cdot u_\delta = \int_{\mathbf{S}} f_E \cdot U_E + \int_{\mathbf{S}} f_I \cdot U_I \end{aligned} \right.$$

The first term of (3.24) is the sum of the left handside members of (3.19) and (3.22). Hence the above inequalities are equalities. Besides

$$\begin{aligned} & \liminf_{\delta \rightarrow 0} \frac{1}{2\delta} \int_{\mathcal{S}_\delta} \sigma_{ij}(u_\delta) \gamma_{ij}(u_\delta) \mathbf{1}_{\mathcal{J}_\delta^c} \leq \liminf_{\delta \rightarrow 0} \frac{1}{2\delta} \int_{\mathcal{S}_\delta} \sigma_{ij}(u_\delta) \gamma_{ij}(u_\delta) \leq \overline{\lim}_{\delta \rightarrow 0} \frac{1}{2\delta} \int_{\mathcal{S}_\delta} \sigma_{ij}(u_\delta) \gamma_{ij}(u_\delta) \\ \text{and } & \liminf_{\delta \rightarrow 0} \frac{1}{2\delta} \int_{\mathcal{S}_\delta} \sigma_{ij}(u_\delta) \gamma_{ij}(u_\delta) \mathbf{1}_{\mathcal{J}_\delta^c} \leq \overline{\lim}_{\delta \rightarrow 0} \frac{1}{2\delta} \int_{\mathcal{S}_\delta} \sigma_{ij}(u_\delta) \gamma_{ij}(u_\delta) \mathbf{1}_{\mathcal{J}_\delta^c} \leq \overline{\lim}_{\delta \rightarrow 0} \frac{1}{2\delta} \int_{\mathcal{S}_\delta} \sigma_{ij}(u_\delta) \gamma_{ij}(u_\delta) \end{aligned}$$

We deduce that $\lim_{\delta \rightarrow 0} \frac{1}{2\delta} \int_{\mathcal{S}_\delta} \sigma_{ij}(u_\delta) \gamma_{ij}(u_\delta) (1 - \mathbf{1}_{\mathcal{J}_\delta^c}) = 0$. All the sequences of the unfolded of the strained tensor components strongly converge in $L^2(\Omega_l)$. We have also the strong convergences

$$\left\{ \begin{aligned} & \delta \mathcal{U}_\delta \longrightarrow U_I, \quad \delta \mathcal{R}_\delta \longrightarrow \mathcal{R} \quad \text{strongly in } H^1(\mathbf{S}, \mathbb{R}^3), \\ & \delta U_{I,\delta} \longrightarrow U_I \quad \text{strongly in } D_I(\mathbf{S}), \\ & U_{E,\delta} \longrightarrow U_E \quad \text{strongly in } D_E(\mathbf{S}). \end{aligned} \right.$$

3.10. Complements

The orthogonal condition (3.12) requires an explanation . First, for any function $\phi \in H^1(\omega_l)$ equal to zero on the edges, the displacement Φ defined by

$$\Phi^{(l)} = \phi \bar{e}_3^{(l)}, \quad \text{in } \omega_l, \text{ and by 0 in the other faces,}$$

belongs to $D_I(\mathbf{S})$. We deduce that the function $f_{E,3}^{(l)}$ is orthogonal to ϕ and then, by density of these test-functions in $L^2(\omega_l)$, we get

$$(3.25) \quad \forall l \in \{1, \dots, N\}, \quad f_{E,3}^{(l)} = 0.$$

Let $D_{I,0}(\mathbf{S})$ be the space of the inextensional displacements equal to zero on the edges belonging to \mathcal{J} and let $(D_{I,0}(\mathbf{S}))^\perp$ be its orthogonal in $D_I(\mathbf{S})$ for the inner product $\langle \cdot, \cdot \rangle_\rho$. The subset $(D_{I,0}(\mathbf{S}))^\perp$ is of finite dimension. The condition (3.12) is then equivalent to

$$(3.26) \quad \forall V \in (D_{I,0}(\mathbf{S}))^\perp, \quad \int_{\mathbf{S}} f_E \cdot V = 0.$$

This last condition results in a finite number of equalities related to the means in the faces $\bar{\omega}_l$ of the components $f_{E,\alpha}^{(l)}$ of f_E .

4. Annexes

4.1 Annex A. Proof of Theorem 2.3

The proof of Theorem 2.3 is based on Lemma 2.3 in [4] and on Lemma 4.1.

We denote

$$\omega_\eta = \{\hat{x} \in \mathbb{R}^2 \mid \text{dist}(\hat{x}, \omega) < \eta\}, \quad \eta > 0.$$

Lemma 4.1 : *There exist $R > 0$ and $\delta_0 > 0$, depending only on ω , such that for any $\delta \in]0, \delta_0]$, $\omega_{2\delta}$ is covered by a family of open sets, of diameter less than $R\delta$, star-shaped with respect to a disc of radius $\delta/2$ and such that any point of $\omega_{2\delta}$ belongs to a finite number (independent of δ) of open sets of that family.*

Proof : The open set $\mathcal{A}_{pq} =](p - 1/2)\delta, (p + 3/2)\delta[\times](q - 1/2)\delta, (q + 3/2)\delta[$, $(p, q) \in \mathbb{Z}^2$, has a diameter of $2\sqrt{2}\delta$ and is star-shaped with respect to the disc of center $((p + 1/2)\delta, (q + 1/2)\delta)$ and of radius $\delta/2$. Let \mathcal{I}_δ be the set of the pairs (p, q) of \mathbb{Z}^2 such that $\mathcal{A}_{pq} \subset \omega$. The distance between the boundary of ω and $\bigcup_{(p,q) \in \mathcal{I}_\delta} \mathcal{A}_{pq}$ is less than 3δ .

Let us proceed now to the covering of the neighborhood of the boundary of ω .

The boundary of ω is lipschitzian. Hence there exist constants A, B, C, M strictly positive, a finite number N of local coordinate systems (x_{1r}, x_{2r}) in $(O_r; \vec{e}_{1r}, \vec{e}_{2r})$ and maps $f_r : [-A, A] \rightarrow \mathbb{R}$, Lipschitz continuous with ratio M , $1 \leq r \leq N$, such that

$$\left\{ \begin{array}{l} \partial\omega = \bigcup_{r=1}^N \left\{ (x_{1r}, x_{2r}) \mid x_{2r} = f_r(x_{1r}), \quad x_{1r} \in]-A, A[\right\}, \\ \left\{ x \in \omega \mid \text{dist}(x, \partial\omega) < C \right\} \subset \bigcup_{r=1}^N \left\{ (x_{1r}, x_{2r}) \mid f_r(x_{1r}) < x_{2r} < f_r(x_{1r}) + B, \quad |x_{1r}| \leq A \right\} \subset \omega, \\ \omega_C \setminus \omega \subset \bigcup_{r=1}^N \left\{ (x_{1r}, x_{2r}) \mid f_r(x_{1r}) - B < x_{2r} < f_r(x_{1r}), \quad |x_{1r}| \leq A \right\} \subset \mathbb{R}^2 \setminus \omega. \end{array} \right.$$

Through the use of easy geometrical arguments we show that if $4\delta \leq \inf\{C, B/\sqrt{1+M^2}\}$, we have

$$\begin{aligned} \omega_{2\delta} \setminus \omega &\subset \bigcup_{r=1}^N \left\{ (x_{1r}, x_{2r}) \mid f_r(x_{1r}) - 2\delta\sqrt{1+M^2} < x_{2r} < f_r(x_{1r}), \quad |x_{1r}| \leq A \right\}, \\ \left\{ x \in \omega \mid \text{dist}(x, \partial\omega) < 4\delta \right\} &\subset \bigcup_{r=1}^N \left\{ (x_{1r}, x_{2r}) \mid f_r(x_{1r}) < x_{2r} < f_r(x_{1r}) + 4\delta\sqrt{1+M^2}, \quad |x_{1r}| \leq A \right\}. \end{aligned}$$

For any $a \in]-A, A - 2\delta[$, the domains

$$\mathcal{B}_{\delta,a,r} = \{(x_{1r}, x_{2r}) \mid f_r(x_{1r}) - (6M + 2)\delta < x_{2r} < f_r(x_{1r}) + (6M + 2)\delta, x_{1r} \in]a, a + 2\delta[\}$$

and $\mathcal{B}_{\delta,a,r} \cap \omega$ are star-shaped with respect to the disc of center $(a + \delta, f_r(a) + (3M + 1)\delta)$ and of radius $\delta/2$. These open sets have a diameter less than $6(3M + 1)\delta = R\delta$.

For $0 < \delta \leq \delta_0 = \inf\{B/(6M + 2), A/2, C/4\}$ the open sets \mathcal{A}_{pq} ($(p, q) \in \mathcal{I}_\delta$), $\mathcal{B}_{\delta,a_p,r}$, where $a_p = p\delta \in]-A, A[$ ($p \in \mathbb{Z}$), $\mathcal{B}_{\delta,-A,r}$ and $\mathcal{B}_{\delta,A-2\delta,r}$ ($r \in \{1, \dots, N\}$) cover $\omega_{2\delta}$; their diameter is less than $R\delta$ and they are star-shaped with respect to a disc of radius $\delta/2$. Any point of $\omega_{2\delta}$ belongs to a finite number (depending only on ω) of open sets of that family. \square

We denote $\{\omega'_{\delta,n}\}_{n \in N_\delta}$ the covering of $\omega_{2\delta}$ obtained in Lemma A.1 and $\{\omega_{\delta,n}\}_{n \in N_\delta}$ the covering of ω defined by $\omega_{\delta,n} = \omega'_{\delta,n} \cap \omega$, $n \in N_\delta$.




Figure 2. The domain $\mathcal{B}_{\delta,a,r}$

Proof of Theorem 2.3 : The open set $\omega_{\delta,n}$ is star-shaped with respect to a disc of center A_n and of radius $\delta/2$. We put $\mathcal{O}_{\delta,n} = \omega_{\delta,n} \times]-\delta, \delta[\subset \Omega_\delta$, $n \in N_\delta$. The domain $\mathcal{O}_{\delta,n}$ has a diameter less than $(R + 2)\delta$, and is star-shaped with respect to a ball of center A_n and of radius $\delta/2$. From Lemma 2.3 of [4], there exists a rigid displacement r_n such that

$$(4.1) \quad \mathcal{D}(u - r_n, \mathcal{O}_{\delta,n}) + \frac{1}{\delta^2} \|u - r_n\|_{L^2(\mathcal{O}_{\delta,n}, \mathbb{R}^3)}^2 \leq C\mathcal{E}(u, \mathcal{O}_{\delta,n}), \quad r_n(x) = a_n + b_n \wedge \overrightarrow{A_n x}, \quad (a_n, b_n) \in \mathbb{R}^3.$$

The constant depends only on R .

We calculate the mean of $(u - r_n)(x)$ and of $x_3 \vec{e}_3 \wedge (u - r_n)(x)$ on the intervals $\{\hat{x}\} \times]-\delta, \delta[$, $\hat{x} \in \omega_{\delta,n}$, then we integrate on $\omega_{n,\delta}$ the inequalities we have obtained. Thanks to (4.1), we have

$$(4.2) \quad \int_{\omega_{\delta,n}} |\mathcal{U}(\hat{x}) - a_n - b_n \wedge \overrightarrow{A_n \hat{x}}|^2 d\hat{x} \leq C\delta\mathcal{E}(u, \mathcal{O}_{\delta,n}), \quad \int_{\omega_{\delta,n}} |\mathcal{R}_\alpha(\hat{x}) - b_{\alpha,n}|^2 d\hat{x} \leq \frac{C}{\delta}\mathcal{E}(u, \mathcal{O}_{\delta,n}).$$

In (4.1) we eliminate the displacement r_n thanks to the estimations (4.2). Hence we have $\|u - U_e\|_{L^2(\mathcal{O}_{\delta,n}, \mathbb{R}^3)}^2 \leq C\delta^2\mathcal{E}(u, \mathcal{O}_{\delta,n})$, then we add these inequalities and we obtain

$$\|u - U_e\|_{L^2(\Omega_\delta, \mathbb{R}^3)}^2 \leq C\delta^2\mathcal{E}(u, \Omega_\delta).$$

Both components of e.p.d. U_e belong to $H^1(\omega, \mathbb{R}^3)$. We calculate the mean of the gradient of $(u - r_n)(x)$, then the mean of $x_3 \vec{e}_3 \wedge \nabla(u - r_n)(x)$ on the intervals $\{\hat{x}\} \times]-\delta, \delta[$, $\hat{x} \in \omega_{\delta, n}$. Using (4.1) we obtain

$$(4.3) \quad \left\| \frac{\partial \mathcal{U}}{\partial x_\alpha} - b_n \wedge \vec{e}_\alpha \right\|_{L^2(\omega_{\delta, n}, \mathbb{R}^3)}^2 + \delta^2 \left\| \frac{\partial \mathcal{R}}{\partial x_\alpha} \right\|_{L^2(\omega_{\delta, n}, \mathbb{R}^2)}^2 \leq \frac{C}{\delta} \mathcal{E}(u, \mathcal{O}_{\delta, n}),$$

hence, after elimination of b_n in the first inequality,

$$(4.4) \quad \begin{cases} \left\| \frac{\partial \mathcal{U}_1}{\partial x_1} \right\|_{L^2(\omega_{\delta, n})}^2 + \left\| \frac{\partial \mathcal{U}_2}{\partial x_2} \right\|_{L^2(\omega_{\delta, n})}^2 + \left\| \frac{\partial \mathcal{U}_1}{\partial x_2} + \frac{\partial \mathcal{U}_2}{\partial x_1} \right\|_{L^2(\omega_{\delta, n})}^2 \\ + \left\| \frac{\partial \mathcal{U}_3}{\partial x_1} + \mathcal{R}_2 \right\|_{L^2(\omega_{\delta, n})}^2 + \left\| \frac{\partial \mathcal{U}_3}{\partial x_2} - \mathcal{R}_1 \right\|_{L^2(\omega_{\delta, n})}^2 \leq \frac{C}{\delta} \mathcal{E}(u, \mathcal{O}_{\delta, n}) \end{cases}$$

From (4.3) and (4.4) we deduce the estimate of $\mathcal{E}(U_e, \mathcal{O}_{\delta, n})$

$$\mathcal{E}(U_e, \mathcal{O}_{\delta, n}) \leq C \mathcal{E}(u, \mathcal{O}_{\delta, n}) \quad \text{hence} \quad \mathcal{E}(U_e, \Omega_\delta) \leq C \mathcal{E}(u, \Omega_\delta).$$

From (4.3), (4.4), (4.1) and after elimination of the gradient of r_n we also deduce

$$\mathcal{D}(u - U_e, \mathcal{O}_{\delta, n}) \leq C \mathcal{E}(u, \mathcal{O}_{\delta, n}), \quad \text{hence} \quad \mathcal{D}(u - U_e, \Omega_\delta) \leq C \mathcal{E}(u, \Omega_\delta).$$

Theorem 2.3 is proved. □

4.2 Annex B. About the second decomposition of a plate displacement

4.2.a Extension of a plate displacement

Let ω be a polygonal bounded domain in \mathbb{R}^2 . The boundary of ω is made of a finite number of segments. Let \mathcal{C} be a connected component of $\partial\omega$. There exists $\delta'_0 > 0$ such that for any $\delta \in]0, \delta'_0]$ the domains

$$\mathcal{C}_\delta = \left\{ x \in \mathbb{R}^3 \mid \text{dist}(x, \mathcal{C}) < \delta \right\}, \quad \text{and} \quad \mathcal{C}_{3\delta} = \left\{ x \in \mathbb{R}^3 \mid \text{dist}(x, \mathcal{C}) < 3\delta \right\}$$

are rods structures. Then there exists $\mu'_0 > 0$ such that all the balls centered in a vertex of \mathcal{C} , and of radius $3\eta'_0 \delta$ contain the junctions of the rods belonging to $\mathcal{C}_{3\delta}$.

We recall that for any $\delta \in]0, \delta'_0]$, there exists an extension operator, linear and continuous, P_δ from $H^1(\mathcal{C}_\delta)$ into $H^1(\mathcal{C}_{3\delta})$ such that for any $\phi \in H^1(\mathcal{C}_\delta)$,

$$(4.5) \quad P_\delta(\phi)|_{\mathcal{C}_\delta} = \phi, \quad \|P_\delta(\phi)\|_{L^2(\mathcal{C}_{3\delta})} + \delta \|\nabla P_\delta(\phi)\|_{[L^2(\mathcal{C}_{3\delta})]^3} \leq C \left\{ \|\phi\|_{L^2(\mathcal{C}_\delta)} + \delta \|\nabla \phi\|_{[L^2(\mathcal{C}_\delta)]^3} \right\}.$$

The constant does not depend on δ .

Proof of Lemma 2.8 : We begin with extending u in the neighborhood of a connected component of $\partial\omega$. Let \mathcal{C} be a connected component of $\partial\omega$. The restriction of u to \mathcal{C}_δ is a displacement belonging to $H^1(\mathcal{C}_\delta, \mathbb{R}^3)$. Hence there exists an elementary displacement of a rods structure (e.d.r.s.) $U_{e,R}$ (see [5]) which coincides with a rigid displacement in each set $B(A, 3\eta'_0 \delta) \cap \mathcal{C}_\delta$ where A is a vertex of \mathcal{C} and which verifies

$$(4.6) \quad \mathcal{E}(U_{e,R}, \mathcal{C}_\delta) + \mathcal{D}(u - U_{e,R}, \mathcal{C}_\delta) + \frac{1}{\delta^2} \|u - U_{e,R}\|_{L^2(\mathcal{C}_\delta, \mathbb{R}^3)}^2 \leq C \mathcal{E}(u, \mathcal{C}_\delta).$$

The displacement $U_{e,R}$ is also an e.d.r.s. of $\mathcal{C}_{3\delta}$ and $\mathcal{E}(U_{e,R}, \mathcal{C}_{3\delta}) \leq C \mathcal{E}(u, \mathcal{C}_\delta)$. The displacement

$$u'_{\mathcal{C}_{3\delta}} = U_{e,R} + P_\delta(u - U_{e,R})$$

is an extension of u to the set $\mathcal{C}_{3\delta}$. From (4.6) we have the following inequalities:

$$(4.7) \quad \mathcal{E}(u_{\mathcal{C}_{3\delta}}, \mathcal{C}_{3\delta}) \leq 2\{\mathcal{E}(U_{e,R}, \mathcal{C}_{3\delta}) + \mathcal{E}(P_\delta(u - U_{e,R}), \mathcal{C}_{3\delta})\} \leq C\mathcal{E}(u, \mathcal{C}_\delta)$$

In the same way we build an extension of u in the neighborhood of the other connected components of $\partial\omega$. The extension $\mathcal{P}_\delta(u)$ is then the displacement which coincides with u in Ω'_δ and which is equal to one of the previous extensions in $\tilde{\Omega}_\delta \setminus \Omega'_\delta$. The estimates (2.14) are the immediate consequences of the inequalities (4.7) obtained in the neighborhood of each connected components of ω . \square

Remark 4.2 : If one of the edges of the boundary of ω is fixed we can take an e.d.r.s. with its two components equal to zero on this edge without modifying the estimates (4.6) and then extend u beyond this edge by 0. \square

Remark 4.3 : We also can construct an extension operator P_δ when ω is of lipschitzian boundary with the help of a few changes. \square

4.2.b Modification of an e.p.d. in the neighborhood of an edge.

Let J be an edge contained in the face $\bar{\omega}$, J_δ the rod

$$J_\delta = \{x \in \mathbb{R}^3 \mid \text{dist}(x, J) < \delta\} \subset \Omega'_\delta,$$

and u a displacement of the plate $\tilde{\Omega}_\delta$. Without being detrimental to the general case we can suppose that the edge's direction is \vec{e}_1 and that one of these extremities is the chosen origin on the face, so that J is identified with the segment $[0, L] \times \{0\}$ where L is the edge's length.

The restriction of the displacement u to the rod J_δ can be decomposed into the sum of the elementary rod displacement (e.r.d.) $U_{e,R}$ of components \mathcal{U}_R and \mathcal{R}_R and of a residual displacement. We choose an e.r.d. $U_{e,R}$ coinciding with a rigid displacement in the balls centered in the extremities of J and of radius $\eta_0\delta$ (see [5]). We have

$$U_{e,R}(x) = \mathcal{U}_R(x_1) + \mathcal{R}_R(x_1) \wedge (x_2\vec{e}_2 + x_3\vec{e}_3).$$

We know (see [4] and [5]) that the components \mathcal{U}_R and \mathcal{R}_R of $U_{e,R}$ belong to $H^1(J, \mathbb{R}^3)$, and verify

$$(4.8) \quad \begin{cases} \delta^2 \left\| \frac{d\mathcal{R}_R}{dx_1} \right\|_{L^2([0, L], \mathbb{R}^3)}^2 + \left\| \frac{d\mathcal{U}_R}{dx_1} - \mathcal{R}_R \wedge \vec{e}_1 \right\|_{L^2([0, L], \mathbb{R}^3)}^2 \leq \frac{C}{\delta^2} \mathcal{E}(u, J_\delta) \\ \mathcal{E}(U_{e,R}, J_\delta) + \mathcal{D}(u - U_{e,R}, J_\delta) + \frac{1}{\delta^2} \|u - U_{e,R}\|_{L^2(J_\delta, \mathbb{R}^3)}^2 \leq C\mathcal{E}(u, J_\delta) \end{cases}$$

The functions \mathcal{U}_R and \mathcal{R}_R are extended into functions belonging to $H_{loc}^1(\mathbb{R}, \mathbb{R}^3)$ (by construction \mathcal{R}_R is constant and \mathcal{U}_R is linear in a neighborhood of the extremities of J). These extensions are then identified with elements belonging to $H_{loc}^1(\mathbb{R}^2, \mathbb{R})$ depending only on the variable x_1 .

Let

$$J'_\delta = \left\{ x \in \Omega'_\delta \mid \text{dist}(\hat{x}, J) < (2\eta_0 + 1)\delta \right\}$$

be the rod and \hat{J}_δ the neighborhood of J in ω_δ ,

$$\hat{J}_\delta = \left\{ x \in \omega_\delta \mid \text{dist}(\hat{x}, J) < 2\eta_0\delta \right\}$$

From the estimates (4.8) of the restriction of $u - U_{e,R}$ to J_δ we deduce the following estimates of the restriction of $u - U_{e,R}$ to J'_δ :

$$(4.9) \quad \mathcal{D}(u - U_{e,R}, J'_\delta) + \frac{1}{\delta^2} \|u - U_{e,R}\|_{L^2(J'_\delta, \mathbb{R}^3)}^2 \leq C\mathcal{E}(u, J'_\delta).$$

The constant depends on L and η_0 .

The displacement u of the plate Ω'_δ is decomposed into the sum of an elementary plate displacement $U_{e,P}$, given by (2.15), and of a residual displacement,

$$U_{e,P}(x) = \mathcal{U}_P(\hat{x}) + \mathcal{R}_P(\hat{x}) \wedge x_3 \vec{e}_3, \quad x \in \Omega'_\delta.$$

Besides the inequalities (2.16), we also have

$$(4.10) \quad \begin{cases} \delta^3 \|\nabla \mathcal{R}_P\|_{L^2(\hat{J}_\delta, \mathbb{R}^6)}^2 + \delta \left\| \frac{\partial \mathcal{U}_P}{\partial x_\alpha} - \mathcal{R}_P \wedge \vec{e}_\alpha \right\|_{L^2(\hat{J}_\delta, \mathbb{R}^3)}^2 \leq C \mathcal{E}(u, J'_\delta), \\ \mathcal{D}(u - U_{e,P}, J'_\delta) + \frac{1}{\delta^2} \|u - U_{e,P}\|_{L^2(J'_\delta, \mathbb{R}^3)}^2 \leq C \mathcal{E}(u, J'_\delta), \end{cases}$$

This allows us to compare the different elementary displacements. We obtain

$$(4.11) \quad \|U_{e,R} - \mathcal{U}_P\|_{L^2(\hat{J}_\delta, \mathbb{R}^3)}^2 + \delta^2 \|\mathcal{R}_R - \mathcal{R}_P\|_{L^2(\hat{J}_\delta, \mathbb{R}^3)}^2 \leq C \delta \mathcal{E}(u, J'_\delta).$$

The estimate of $\|\mathcal{R}_R - \mathcal{R}_P\|_{L^2(\hat{J}_\delta, \mathbb{R}^3)}^2$ follows from the nullity of $\mathcal{R}_R - \mathcal{R}_P$ on J .

We are now going to modify the e.r.d. $U_{e,P}$ in the neighborhood of J .

We consider a function m belonging to $\mathcal{C}^\infty(\mathbb{R}^+, [0, 1])$ such that

$$(4.12) \quad m(t) = 1 \quad \forall t \geq 2, \quad m(t) = 0 \quad \forall t \leq 1, \quad |m'(t)| \leq 2 \quad \forall t \in \mathbb{R}.$$

We define the components, \mathcal{U}' and \mathcal{R}' , of a new e.r.d. U'_e by

$$(4.13) \quad \begin{cases} \mathcal{U}'(\hat{x}) = U_{e,R}(\hat{x}) \left(1 - m\left(\frac{\text{dist}(\hat{x}, J)}{\eta_0 \delta}\right)\right) + \mathcal{U}_P(\hat{x}) m\left(\frac{\text{dist}(\hat{x}, J)}{\eta_0 \delta}\right), \\ \mathcal{R}'(\hat{x}) = \mathcal{R}_R(\hat{x}) \left(1 - m\left(\frac{\text{dist}(\hat{x}, J)}{\eta_0 \delta}\right)\right) + \mathcal{R}_P(\hat{x}) m\left(\frac{\text{dist}(\hat{x}, J)}{\eta_0 \delta}\right), & \hat{x} \in \omega_\delta, \\ U'_e(x) = \mathcal{U}'(\hat{x}) + \mathcal{R}'(\hat{x}) \wedge x_3 \vec{e}_3, & x \in \Omega'_\delta. \end{cases}$$

Hence we have by construction of U'_e ,

$$\begin{aligned} & \text{if } x \in \Omega'_\delta \text{ and } \text{dist}(\hat{x}, J) < \eta_0 \delta \text{ then } U'_e(x) = U_{e,R}(x) \\ & \text{if } x \in \Omega'_\delta \text{ and } \text{dist}(\hat{x}, J) > 2\eta_0 \delta \text{ then } U'_e(x) = U_{e,P}(x). \end{aligned}$$

Thanks to (4.8), (4.10), (4.12) and (2.16) the e.p.d. U'_e verifies

$$\begin{aligned} & \delta^3 \|\nabla \mathcal{R}'\|_{L^2(\omega_\delta, \mathbb{R}^6)}^2 + \left\| \frac{\partial \mathcal{U}'}{\partial x_\alpha} - \mathcal{R}' \wedge \vec{e}_\alpha \right\|_{L^2(\omega_\delta, \mathbb{R}^3)}^2 + \mathcal{E}(U'_e, \Omega'_\delta) \leq C \mathcal{E}(u, \Omega'_\delta), \\ & \mathcal{D}(u - U'_e, \Omega'_\delta) \leq C \mathcal{E}(u, \Omega'_\delta), \quad \|u - U'_e\|_{L^2(\Omega'_\delta, \mathbb{R}^3)}^2 \leq C \delta^2 \mathcal{E}(u, \Omega'_\delta). \end{aligned}$$

The constants depend only on ω , J and η_0 .

4.3 Annex C. About the spaces $H_{\rho, \Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$ and $D_E(\mathbf{S})$

For any $\theta_0 \in]0, \pi[$ and any $r > 0$, we denote $C_{r, \theta_0} = \{(x, y) \in B(O; r) \mid 0 < \theta < \theta_0\}$, J_0 the segment of origin O and of extremity $A = (0, 1)$ and we denote J_{θ_0} the segment of origin O and extremity $B = (\cos(\theta_0), \sin(\theta_0))$.

We denote $r = \text{dist}(x, O)$, $x \in \mathbb{R}^2$.

Lemma 4.3 : Let ϕ belong to $H^1(C_{1,\theta_0})$, for any $\alpha \in]0, 1]$, we have

$$(4.14) \quad \int_{C_{1,\theta_0}} |\phi|^2 r^{\alpha-2} \leq \frac{4}{\alpha} \|\phi\|_{L^2(C_{1,\theta_0})}^2 + \frac{2}{\alpha^2} \|\nabla \phi\|_{[L^2(C_{1,\theta_0})]^2}$$

Proof : We recall that for any $u \in H^1(0, L)$ and for any $\alpha \in]0, 1]$, we have

$$(4.15) \quad \int_0^L |u(t)|^2 t^{\alpha-1} dt \leq \frac{4}{\alpha L^{2-\alpha}} \int_0^L |u(t)|^2 t dt + \frac{2}{\alpha^2 L^\alpha} \int_0^L |u'(t)|^2 t dt$$

Let us take $\phi \in C^\infty(\overline{C_{1,\theta_0}})$. We apply the inequality 4.4 to the restriction of ϕ to a radius coming from the origin and contained in C_{1,θ_0} . This gives

$$\int_0^1 \frac{|\phi(r \cos(\theta), r \sin(\theta))|^2}{r^{1-\alpha}} dr \leq \frac{4}{\alpha} \int_0^1 |\phi(r \cos(\theta), r \sin(\theta))|^2 r dr + \frac{2}{\alpha^2} \int_0^1 \left| \frac{\partial \phi}{\partial r}(r \cos(\theta), r \sin(\theta)) \right|^2 r dr.$$

We then integrate with respect to θ between 0 and θ_0 and we obtain (4.14). The density of $C^\infty(\overline{C_{1,\theta_0}})$ into $H^1(C_{1,\theta_0})$ gives the inequality for any function of the space $H^1(C_{1,\theta_0})$. \square

Lemma 4.4 : Let u be in $H^{1/2}(J_0)$, v be in $H^{1/2}(J_{\theta_0})$ and $\alpha \in]0, 2]$. There exists a function w belonging to $H_{loc}^1(C_{1,\theta_0}) \cap L^2(C_{1,\theta_0})$ such that

$$\begin{aligned} w|_{J_0} &= u, & w|_{J_{\theta_0}} &= v, & \text{and} \\ \|w\|_{L^2(C_{1,\theta_0})}^2 + \int_{C_{1,\theta_0}} |\nabla w|^2 r^\alpha &\leq \frac{C}{\alpha^2} \{ \|u\|_{H^{1/2}(J_0)}^2 + \|v\|_{H^{1/2}(J_{\theta_0})}^2 \}. \end{aligned}$$

The constant depends only on θ_0 . Moreover w belongs to $W^{1,p}(C_{1,\theta_0})$ for any p , such that $1 \leq p < \frac{4}{2+\alpha}$.

Proof : We denote J'_0 (resp. J'_{θ_0}), the segment of same direction as J_0 (resp. J_{θ_0}) and of length $\tan(\theta_0/2)$. The function u (resp. v) extends by reflexion into an element still denoted u (resp. v) belonging to $H^{1/2}(J'_0)$ (resp. $H^{1/2}(J'_{\theta_0})$).

Let \tilde{u} and \tilde{v} be the functions belonging to $H^{1/2}(J'_0 \cup J'_{\theta_0})$ defined by

$$\begin{aligned} \tilde{u}|_{J'_0} &= u, & \tilde{u}|_{J'_{\theta_0}}(t\vec{e}_{\theta_0}) &= u(t\vec{e}_0), \\ \tilde{v}|_{J'_0}(t\vec{e}_0) &= v(t\vec{e}_{\theta_0}), & \tilde{v}|_{J'_{\theta_0}} &= v, \end{aligned} \quad t \in]0, \tan(\theta_0/2)[$$

where $\vec{e}_\theta = \cos(\theta)\vec{e}_1 + \sin(\theta)\vec{e}_2$, $\theta \in [0, \theta_0]$. There exists a continuous lifting operator from $H^{1/2}(J'_0 \cup J'_{\theta_0})$ into $H^1(C_{\tan(\theta_0/2), \theta_0})$. Let U (resp. V) be the lifting of \tilde{u} (resp. \tilde{v}). In the triangle T_{θ_0} of vertexes $(0, 0)$, $(0, \tan(\theta_0/2))$ and $\tan(\theta_0/2)(\cos(\theta_0), \sin(\theta_0))$, containing C_{1,θ_0} and contained in $C_{\tan(\theta_0/2), \theta_0}$, we define w by

$$w(x_1, x_2) = U(x_1, x_2) \frac{x_1 \sin(\theta_0) - x_2 \cos(\theta_0)}{x_2(1 - \cos(\theta_0)) + x_1 \sin(\theta_0)} + V(x_1, x_2) \frac{x_2}{x_2(1 - \cos(\theta_0)) + x_1 \sin(\theta_0)}.$$

In the above expression the coefficients of $U(x_1, x_2)$ and $V(x_1, x_2)$ are barycentric coordinates of point (x_1, x_2) belonging to T_{θ_0} . By construction we have $w|_{J_0} = u$ and $w|_{J_{\theta_0}} = v$. The function w belongs to $L^2(T_{\theta_0})$ and

$$\|w\|_{L^2(T_{\theta_0})} \leq \|U\|_{L^2(C_{\tan(\theta_0/2), \theta_0})} + \|V\|_{L^2(C_{\tan(\theta_0/2), \theta_0})} \leq C \{ \|u\|_{H^{1/2}(J_0)}^2 + \|v\|_{H^{1/2}(J_{\theta_0})}^2 \}.$$

We then calculate the partial derivatives of w and we conclude that w belongs to $H_{loc}^1(T_{\theta_0})$. Moreover we have

$$|\nabla w(x_1, x_2)| \leq C \{ |\nabla U(x_1, x_2)| + |\nabla V(x_1, x_2)| + |(U - V)(x_1, x_2)| r^{-1} \} \quad (x_1, x_2) \in C_{\tan(\theta_0/2), \theta_0}$$

The constant depends on θ_0 . Thanks to the inequality of Lemma 4.3, we have

$$\begin{aligned} \int_{C_{1, \theta_0}} |U - V|^2 r^{\alpha-2} &\leq \frac{4}{\alpha} \|U - V\|_{L^2(C_{1, \theta_0})}^2 + \frac{2}{\alpha^2} \|\nabla(U - V)\|_{[L^2(C_{1, \theta_0})]^2}^2 \\ \implies \int_{C_{1, \theta_0}} |\nabla w|^2 r^\alpha &\leq \frac{C}{\alpha^2} \{ \|U\|_{H^1(C_{\tan(\theta_0/2), \theta_0})}^2 + \|V\|_{H^1(C_{\tan(\theta_0/2), \theta_0})}^2 \}. \end{aligned}$$

Eventually we obtain the estimate of Lemma 4.4. Moreover we have

$$\int_{C_{1, \theta_0}} |\nabla w|^p \leq \left\{ \int_{C_{1, \theta_0}} |\nabla w|^2 r^\alpha \right\}^{\frac{p}{2}} \left\{ \int_{C_{1, \theta_0}} r^{-\frac{\alpha p}{2-p}} \right\}^{\frac{2-p}{2}}$$

Hence w belongs to $W^{1,p}(C_{1, \theta_0})$ if $\frac{\alpha p}{2-p} < 2$. That is to say for $1 \leq p < \frac{4}{2+\alpha}$. \square

Corollary : If $\alpha = 1$, the function w belongs to $W^{1,p}(C_{1, \theta_0})$ for any $1 \leq p < 4/3$. \square

Proof of Lemma 3.4 :

Step 1 The norms are equivalent.

Let be V in $H_{\rho, \Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$. We applied the classical Korn inequality to the membrane displacements $V_M^{(l)} = V_1^{(l)} \vec{e}_1^{(l)} + V_2^{(l)} \vec{e}_2^{(l)}$ and then we add all the inequalities to obtain

$$\sum_{l=1}^N \{ \|\nabla V_1^{(l)}\|_{[L^2(\omega_l)]^2}^2 + \|\nabla V_2^{(l)}\|_{[L^2(\omega_l)]^2}^2 \} \leq C (|V|_\rho^2 + \|V\|_{L^2(\mathbf{S}, \mathbb{R}^3)}^2)$$

hence

$$\|V\|_\rho \leq C \{ |V|_\rho + \|V\|_{L^2(\mathbf{S}, \mathbb{R}^3)} \}.$$

The space $H_{\rho, \Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$ is embedded in $L^2(\mathbf{S}, \mathbb{R}^3)$ (see Lemma 4.4). Then we prove by contradiction that there exists a constant C_0 such that $\|V\|_\rho \leq C_0 |V|_\rho$. Moreover we can immediately see that there exists C_1 such that $|V|_\rho \leq C_1 \|V\|_\rho$. The norms $|\cdot|_\rho$ and $\|\cdot\|_\rho$ are therefore equivalent.

Step 2 The space $H_{\Gamma_0}^1(\mathbf{S}, \mathbb{R}^3) \cap L^\infty(\mathbf{S}, \mathbb{R}^3)$ is dense in $H_{\rho, \Gamma_0}^1(\mathbf{S}, \mathbb{R}^3) \cap L^\infty(\mathbf{S}, \mathbb{R}^3)$.

Let be $V \in H_{\rho, \Gamma_0}^1(\mathbf{S}, \mathbb{R}^3) \cap L^\infty(\mathbf{S}, \mathbb{R}^3)$, we consider the sequence $(V_\delta)_{\delta>0}$ defined by

$$V_\delta(\hat{x}) = V(\hat{x}) \sum_{A \in \mathcal{N}} m \left(\frac{\text{dist}(\hat{x}, A)}{\delta} \right), \quad \hat{x} \in \mathbf{S}$$

where m is given by (4.12). The displacement V_δ is equal to zero in the neighborhood of each vertex belonging to \mathcal{N} . This displacement belongs to $H_{\Gamma_0}^1(\mathbf{S}, \mathbb{R}^3) \cap L^\infty(\mathbf{S}, \mathbb{R}^3)$ and we have

$$\|V_\delta - V\|_{L^2(\mathbf{S}, \mathbb{R}^3)} \leq C\delta \|V\|_{L^\infty(\mathbf{S}, \mathbb{R}^3)}, \quad \|V_\delta\|_{L^\infty(\mathbf{S}, \mathbb{R}^3)} \leq C \|V\|_{L^\infty(\mathbf{S}, \mathbb{R}^3)}.$$

We calculate the gradient of the restriction of $V_\delta^{(l)}$ to each face $\bar{\omega}_l$. Using the L^2 estimate of $V_\delta - V$, we obtain $|V_\delta|_\rho \leq C \{ |V|_\rho + \|V\|_{L^\infty(\mathbf{S}, \mathbb{R}^3)} \}$. The constant does not depend on δ . The sequence $(V_\delta)_{\delta>0}$ weakly converges to V in $H_{\rho, \Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$, which gives the density of $H_{\Gamma_0}^1(\mathbf{S}, \mathbb{R}^3) \cap L^\infty(\mathbf{S}, \mathbb{R}^3)$ into $H_{\rho, \Gamma_0}^1(\mathbf{S}, \mathbb{R}^3) \cap L^\infty(\mathbf{S}, \mathbb{R}^3)$.

Step 3 The space $H_{\rho,\Gamma_0}^1(\mathbf{S}, \mathbb{R}^3) \cap L^\infty(\mathbf{S}, \mathbb{R}^3)$ is dense in $H_{\rho,\Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$.

We consider the truncature function T_M from \mathbb{R}^3 into \mathbb{R}^3 defined by

$$T_M(x) = \begin{cases} x & \text{if } \|x\|_2 < M \\ \frac{x}{\|x\|_2} M & \text{if } \|x\|_2 \geq M, \end{cases}$$

where M belongs to \mathbb{R}_+^* and where $\|\cdot\|_2$ is the euclidian norm of \mathbb{R}^3 . The map T_M is piecewise \mathcal{C}^1 verifying $T_M(0) = 0$ and $\|\nabla T_M\|_{[L^\infty(\mathbb{R}^3, \mathbb{R}^3)]^3} \leq C$ (the constant does not depend on M).

Let $V \in H_{\rho,\Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$, $V_M = T_M(V)$ belongs to $H_{\rho,\Gamma_0}^1(\mathbf{S}, \mathbb{R}^3) \cap L^\infty(\mathbf{S}, \mathbb{R}^3)$ and verifies

$$\|V_M\|_{L^2(\mathbf{S}, \mathbb{R}^3)} \leq \|V\|_{L^2(\mathbf{S}, \mathbb{R}^3)}, \quad \|V_M\|_{L^\infty(\mathbf{S}, \mathbb{R}^3)} \leq M, \quad |V_M|_\rho \leq C|V|_\rho.$$

The constant does not depend on M . When M tends to infinity, V_M tends strongly to V in $H_{\rho,\Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$. Hence the density of $H_{\rho,\Gamma_0}^1(\mathbf{S}, \mathbb{R}^3) \cap L^\infty(\mathbf{S}, \mathbb{R}^3)$ in $H_{\rho,\Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$. \square

Proof of Lemma 3.7 : We put $V \in D_E(\mathbf{S})$. As in the proof of Lemma 3.4 we get

$$\sum_{l=1}^N \{ \|\nabla V_1^{(l)}\|_{[L^2(\omega_l)]^2}^2 + \|\nabla V_2^{(l)}\|_{[L^2(\omega_l)]^2}^2 \} \leq C(\|V\|_E^2 + \|V\|_{L^2(\mathbf{S}, \mathbb{R}^3)}^2)$$

We put $J \in \mathcal{J}$ a common edge to the faces $\bar{\omega}_l$ and $\bar{\omega}_k$. The restrictions to J of the membrane displacements $V_M^{(l)}$ and $V_M^{(k)}$ completely define the restriction $V|_J$. Hence we get

$$\sum_{J \in \mathcal{J}} \|V|_J\|_{H^{1/2}(J, \mathbb{R}^3)}^2 \leq C(\|V\|_E^2 + \|V\|_{L^2(\mathbf{S}, \mathbb{R}^3)}^2)$$

With the help of Lemma 4.4 we build a displacement $W \in H_{\rho,\Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$ such that

$$W_\alpha^{(l)} = V_\alpha^{(l)}, \quad \forall l \in \{1, \dots, N\}, \quad W|_J = V|_J, \quad \forall J \in \mathcal{J},$$

and verifying

$$|W|_\rho^2 \leq C \sum_{l=1}^N \{ \|\nabla V_1^{(l)}\|_{[L^2(\omega_l)]^2}^2 + \|\nabla V_2^{(l)}\|_{[L^2(\omega_l)]^2}^2 \} + C \sum_{J \in \mathcal{J}} \|V|_J\|_{H^{1/2}(J, \mathbb{R}^3)}^2 \leq C(\|V\|_E^2 + \|V\|_{L^2(\mathbf{S}, \mathbb{R}^3)}^2).$$

The displacement $V - W$ is of inextensional type and hence orthogonal to V , hence

$$(4.16) \quad |V|_\rho \leq |W|_\rho \leq C(\|V\|_E + \|V\|_{L^2(\mathbf{S}, \mathbb{R}^3)}).$$

Now we show that the norm $\|\cdot\|_E$ is equivalent to the norm $|\cdot|_\rho$ in $D_E(\mathbf{S})$. We already have $\|V\|_E \leq |V|_\rho$ for any $V \in D_E(\mathbf{S})$. We suppose that the norms are not equivalent. For any $n \in \mathbb{N}^*$, we can find $V_n \in D_E(\mathbf{S})$ such that $\|V_n\|_E \leq 1/n$ and $|V_n|_\rho = 1$. The sequence $(V_n)_{n \in \mathbb{N}^*}$ being bounded in $H_\rho^1(\mathbf{S}, \mathbb{R}^3)$, we can then extract a sub-sequence, still denoted in the same way, such that

$$V_n \rightharpoonup V \quad \text{weakly in } H_{\rho,\Gamma_0}^1(\mathbf{S}, \mathbb{R}^3).$$

The limit V belongs also to $D_E(\mathbf{S})$. Let us make n tend to infinity in the inequality $\|V_n\|_E \leq 1/n$, we obtain $\gamma_{\alpha\beta}(V^{(l)}) = 0$. The displacement V is of inextensional type, and hence is equal to zero.

If $p \in [1, 4/3[$ the space $H_\rho^1(\mathbf{S}, \mathbb{R}^3)$ is continuously imbedded in $W^{1,p}(\mathbf{S}, \mathbb{R}^3)$ (see Lemma 4.4) and if $p \in]1, 4/3[$ the space $W^{1,p}(\mathbf{S}, \mathbb{R}^3)$ is compactly imbedded in $L^2(\mathbf{S}, \mathbb{R}^3)$. Hence the sequence $(V_n)_{n \in \mathbb{N}^*}$ converges strongly to 0 in $L^2(\mathbf{S}, \mathbb{R}^3)$, hence $\|V_n\|_{L^2(\mathbf{S}, \mathbb{R}^3)} \rightarrow 0$. From (4.16) follows then that the sequence $(V_n)_{n \in \mathbb{N}^*}$ converges strongly to 0 in $H_{\rho, \Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$. This stands in contradiction with $|V_n|_\rho = 1$. \square

4.4 Annex D. The inextensional displacements

4.4.a. The inextensional displacements of $D_I(\mathbf{S})$

Let U be an inextensional displacement. From the definition of the inextensional displacements we have $\gamma_{\alpha\beta}(U^{(l)}) = 0$ in $\bar{\omega}_l$. Hence, in each face the membrane displacement $U_M^{(l)} = U_1^{(l)} \bar{e}_1^{(l)} + U_2^{(l)} \bar{e}_2^{(l)}$ is a rigid displacement. The restriction of U to an edge $J \in \mathcal{J}$ is then

$$(4.17) \quad U_{|J}(M) = \vec{A}_J + \vec{B}_J \wedge \overrightarrow{AM} \quad \forall M \in J$$

where A_J is a vertex of the edge. The vectors \vec{A}_J and \vec{B}_J depend only on the edge. We choose \vec{B}_J orthogonal to \bar{e}_J to have the unicity of this vector.

4.4.b. The inextensional displacements of $\mathcal{D}_I(\mathbf{S})$

A displacement $\mathcal{A} \in \mathcal{D}_I(\mathbf{S})$ verifies $\frac{\partial \mathcal{A}^{(l)}}{\partial x_i^{(l)}} = \widehat{\nabla} \mathcal{A}^{(l)} \wedge \bar{e}_i^{(l)}$. This displacement belongs also to $D_I(\mathbf{S})$, hence $\widehat{\nabla} \mathcal{A}_3^{(l)}$ is constant in each face. Let A be a vertex belonging to \mathcal{N} and let J and L be two edges sharing the vertex A . The functions \mathcal{A} and $\widehat{\nabla} \mathcal{A}$ belong to $H^1(\mathbf{S}, \mathbb{R}^3)$. The hypothesis **H2** implies that

$$(4.18) \quad \|\mathcal{A}_{|J \cup L}\|_{H^{1/2}(J \cup L, \mathbb{R}^3)} \leq C \|\mathcal{A}\|_I, \quad \|\widehat{\nabla} \mathcal{A}_{|J \cup L}\|_{H^{1/2}(J \cup L, \mathbb{R}^3)} \leq C \|\mathcal{A}\|_I.$$

From (4.17) we have $\mathcal{A}_{|J}(M) = \vec{A}_J + \vec{B}_J \wedge \overrightarrow{AM}$ and $\mathcal{A}_{|L}(M) = \vec{A}_L + \vec{B}_L \wedge \overrightarrow{AM}$, hence $\vec{A}_J = \vec{A}_L$. We denote $\mathcal{A}(A)$ this value which is common to all the edges containing the vertex A .

We also have $\vec{B}_J \wedge \bar{e}_J = \widehat{\nabla} \mathcal{A}_{|J} \wedge \bar{e}_J$. The vector $\widehat{\nabla} \mathcal{A}_{|J} \wedge \bar{e}_J$ is constant along the edge J , hence

$$\vec{B}_J \cdot (\bar{e}_J \wedge \bar{e}_L) = \vec{B}_L \cdot (\bar{e}_J \wedge \bar{e}_L).$$

There exists a vector $\vec{B}_{JL} \in \mathbb{R}^3$ (depending on \vec{B}_J and \vec{B}_L) such that

$$\vec{B}_J \wedge \bar{e}_J = \vec{B}_{JL} \wedge \bar{e}_J, \quad \vec{B}_L \wedge \bar{e}_L = \vec{B}_{JL} \wedge \bar{e}_L.$$

Since $\widehat{\nabla} \mathcal{A}_{|L}(M) \wedge \bar{e}_L = \vec{B}_{JL} \wedge \bar{e}_L = \vec{B}_{JL} \wedge \bar{e}_L$ for any $M \in L$. From (4.18) we deduce that

$$\int_0^{L_J} \frac{|\widehat{\nabla} \mathcal{A}_{|J}(A + t\bar{e}_J) \wedge \bar{e}_L - \vec{B}_{JL} \wedge \bar{e}_L|^2}{t} dt \leq C \|\mathcal{A}\|_I,$$

hence

$$\int_0^{L_J} \frac{|\widehat{\nabla} \mathcal{A}_{|J}(A + t\bar{e}_J) - \vec{B}_{JL}|^2}{t} dt \leq C \|\mathcal{A}\|_I,$$

because we have $\widehat{\nabla} \mathcal{A}_{|J}(M) \wedge \bar{e}_J = \vec{B}_{JL} \wedge \bar{e}_J$ for any $M \in J$. The vector \vec{B}_{JL} does not depend on the edge L . Hence this vector is independant from the edges that go via A , and is denoted $\widehat{\nabla} \mathcal{A}(A)$.

The restriction of the displacement \mathcal{A} to any edge that goes via A is the restriction to that edge of a rigid displacement depending only on the vertex A ,

$$\forall J \in \mathcal{J}, \quad \forall M \in J, \quad \mathcal{A}_{|J}(M) = \mathcal{A}(A) + \widehat{\nabla} \mathcal{A}(A) \wedge \overrightarrow{AM}.$$

4.5 Annex E. The test functions

Proof of Lemma 3.9 : Let U be an element belonging to $H_{\Gamma_0}^1(\mathbf{S}_{\delta_0}, \mathbb{R}^3)$.

We suppose that the real δ_0 is such that the two balls centered in the extremities of the edges and of radius $8\eta_0\delta_0$ do not share any common point.

Step 1 For any δ in the interval $]0, \delta_0]$, we build a displacement $U_{\delta,1}$ constant in the neighborhood of each vertex belonging to \mathcal{N} and approaching U .

We begin with modifying U in the neighborhood of an vertex. Let A be an vertex common to the faces $\bar{\omega}_{l_1}, \dots, \bar{\omega}_{l_p}$; $\bar{A}^{(l_i)}$ the mean value of $U^{(l_i)}$ in the disc $B(A; \delta) \cap \omega_{l_i, \delta_0}$ and \bar{A} the mean value of the vectors $\bar{A}^{(l_i)}$. If the vertex A belongs to Γ_0 we replace \bar{A} by $\vec{0}$.

We define the displacement $U_{\delta,A}$ in $\bar{\omega}_{l, \delta_0}$ by

$$U_{\delta,A}^{(l)}(\hat{x}^{(l)}) = U(\hat{x}^{(l)})m\left(\frac{r^{(l)}}{2\eta_0\delta}\right) + \left\{1 - m\left(\frac{r^{(l)}}{2\eta_0\delta}\right)\right\}\bar{A}, \quad l \in \{l_1, \dots, l_p\}.$$

where m has been introduced by (4.12) and where $r^{(l)} = \text{dist}(\hat{x}^{(l)}, A)$. In $B(A; 2\eta_0\delta) \cap \bar{\omega}_{l, \delta_0}$ the displacement $U_{\delta,A}^{(l)}$ is by construction constant and equal to \bar{A} . We have

$$\frac{\partial U_{\delta,A}^{(l)}}{\partial x_{\alpha}^{(l)}} - \frac{\partial U^{(l)}}{\partial x_{\alpha}^{(l)}} = \left\{U^{(l)} - \bar{A}\right\}m'\left(\frac{r^{(l)}}{2\eta_0\delta}\right)\frac{x_{\alpha}^{(l)}}{2\eta_0\delta r^{(l)}} - \left\{1 - m\left(\frac{r^{(l)}}{2\eta_0\delta}\right)\right\}\frac{\partial U^{(l)}}{\partial x_{\alpha}^{(l)}}.$$

Let us estimate the L^2 norm of the gradient of $U_{\delta,A}^{(l)} - U^{(l)}$, $l \in \{l_1, \dots, l_p\}$,

$$\|\nabla U_{\delta,A}^{(l)} - \nabla U^{(l)}\|_{[L^2(\omega_{l, \delta_0}, \mathbb{R}^3)]^2}^2 \leq C\|\nabla U^{(l)}\|_{[L^2(B(A; 4\eta_0\delta) \cap \bar{\omega}_{l, \delta_0}, \mathbb{R}^3)]^2}^2 + \frac{C}{\delta^2}\|U^{(l)} - \bar{A}\|_{L^2(B(A; 4\eta_0\delta) \cap \bar{\omega}_{l, \delta_0}, \mathbb{R}^3)}^2$$

The Poincaré-Wirtinger inequality allows us to estimate the L^2 norm of $U^{(l)} - \bar{A}^{(l)}$ in the disc $B(A; \delta) \cap \omega_{l, \delta_0}$,

$$\|U^{(l)} - \bar{A}^{(l)}\|_{L^2(B(A; \delta) \cap \bar{\omega}_{l, \delta_0}, \mathbb{R}^3)}^2 \leq C\delta^2\|\nabla U^{(l)}\|_{[L^2(B(A; \delta) \cap \bar{\omega}_{l, \delta_0}, \mathbb{R}^3)]^2}^2.$$

If J is an edge of vertex A contained in $\bar{\omega}_l$, then we have

$$\|U|_J - \bar{A}^{(l)}\|_{L^2(B(A; \delta) \cap J, \mathbb{R}^3)}^2 \leq C\delta\|\nabla U^{(l)}\|_{[L^2(B(A; \delta) \cap \bar{\omega}_{l, \delta_0}, \mathbb{R}^3)]^2}^2$$

For any other face $\bar{\omega}_{k, \delta_0}$ containing $J \cap B(A; \delta)$, we also have the above estimate, hence

$$\|\bar{A}^{(k)} - \bar{A}^{(l)}\|_2^2 \leq C\left\{\|\nabla U^{(l)}\|_{[L^2(B(A; \delta) \cap \bar{\omega}_{l, \delta_0}, \mathbb{R}^3)]^2}^2 + \|\nabla U^{(k)}\|_{[L^2(B(A; \delta) \cap \bar{\omega}_{k, \delta_0}, \mathbb{R}^3)]^2}^2\right\}.$$

We deduce that

$$\begin{aligned} \|U^{(l)} - \bar{A}\|_{L^2(B(A; \delta) \cap \omega_{l, \delta_0}, \mathbb{R}^3)}^2 &\leq C\delta^2 \sum_{i=1}^p \|\nabla U^{(l_i)}\|_{[L^2(B(A; \delta) \cap \omega_{l_i, \delta_0}, \mathbb{R}^3)]^2}^2 \\ \implies \|U^{(l)} - \bar{A}\|_{L^2(B(A; 4\eta_0\delta) \cap \omega_{l, \delta_0}, \mathbb{R}^3)}^2 &\leq C\delta^2 \sum_{k=1}^p \|\nabla U^{(l_k)}\|_{[L^2(B(A; 4\eta_0\delta) \cap \omega_{l_k, \delta_0}, \mathbb{R}^3)]^2}^2 \end{aligned}$$

And eventually

$$\|U_{\delta,A} - U\|_{H^1(\mathbf{S}_{\delta_0}, \mathbb{R}^3)} \leq C \sum_{k=1}^p \|\nabla U^{(l_k)}\|_{[L^2(B(A; 4\eta_0\delta) \cap \omega_{l_k, \delta_0}, \mathbb{R}^3)]^2}.$$

We can do the same with all the structure vertexes. We denote $U_{\delta,1}$ the displacement obtained after having modified U in a neighborhood of each vertex. Hence we have

$$\|U_{\delta,1} - U\|_{H^1(\mathbf{S}_{\delta_0, \mathbb{R}^3})} \leq C \sum_{A \in \mathcal{N}} \|U\|_{H^1(B(A; 4\eta_0\delta) \cap \mathbf{S}_{\delta_0, \mathbb{R}^3})}.$$

Step 2 Let J be an edge belonging to \mathcal{J} . This edge is common to the faces $\bar{\omega}_{l_1}, \dots, \bar{\omega}_{l_p}$. We denote $V_{\delta, J}^{(l_k)}$ the element of $H^1(J, \mathbb{R}^3)$ defined by

$$V_{\delta, J}^{(l_k)}(\hat{x}_J) = \frac{1}{2\delta} \int_{-\delta}^{\delta} U_{\delta, 1}^{(l_k)}(\hat{x}_J + se_{J, \perp}^{(l_k)}) ds$$

and we denote $V_{\delta, J}$ the mean values of $V_{\delta, J}^{(l_k)}$ ($\{e_J^{(l_k)}, e_{J, \perp}^{(l_k)}\}$ is an orthonormal basis of the direction of the face $\bar{\omega}_{l_k}$). We have $V_{\delta, J}^{(l_k)} \in H^1(J, \mathbb{R}^3)$ and

$$(4.19) \quad \begin{cases} \|U - V_{\delta, J}\|_{L^2(\hat{J}_{\delta, \mathbb{R}^3})} \leq C\delta \|U_{\delta, 1}\|_{H^1(\hat{J}_{\delta, \mathbb{R}^3})} \leq C\delta \|U\|_{H^1(\hat{J}_{\delta, \mathbb{R}^3})} \\ \|U - V_{\delta, J}\|_{H^1(\hat{J}_{\delta, \mathbb{R}^3})} \leq C\|U_{\delta, 1}\|_{H^1(\hat{J}_{\delta, \mathbb{R}^3})} \leq C\|U\|_{H^1(\hat{J}_{\delta, \mathbb{R}^3})} \end{cases}$$

where $\hat{J}_{\delta} = \{\hat{x} \in \mathbf{S}_{\delta_0} \mid \text{dist}(\hat{x}, J) < \eta_0\delta\}$, \hat{J}_{δ} is the union of two-dimensional sets of breadth $2\eta_0\delta$ and of length $L_J + 2\eta_0\delta$. If the edge J is contained in Γ_0 , we take $V_{\delta, J} = 0$, in this case we have again the estimate (4.19). The displacement

$$U_{\delta} = \sum_{J \in \mathcal{J}} U_{\delta, 1} \left\{ 1 - m\left(\frac{\text{dist}(\cdot, J)}{\eta_0\delta}\right) \right\} + \sum_{J \in \mathcal{J}} V_{\delta, J} m\left(\frac{\text{dist}(\cdot, J)}{\eta_0\delta}\right)$$

belongs to $H_{\Gamma_0}^1(\mathbf{S}_{\delta}, \mathbb{R}^3)$ and verifies

$$\|U_{\delta} - U\|_{H^1(\mathbf{S}_{\delta_0, \mathbb{R}^3})} \leq C \sum_{A \in \mathcal{N}} \|U\|_{H^1(B(A; 4\eta_0\delta) \cap \mathbf{S}_{\delta_0, \mathbb{R}^3})} + C \sum_{J \in \mathcal{J}} \|U\|_{H^1(\hat{J}_{\delta, \mathbb{R}^3})}$$

The constant is independant of δ . Lemma 3.9 is proved. \square

Proof of Lemma 3.11 : Let be $U \in \mathcal{D}_I(\mathbf{S})$. We recall that there exists only one function in $H_{\Gamma_0}^1(\mathbf{S}, \mathbb{R}^3)$

denoted $\widehat{\nabla}U$ such that $\frac{\partial U^{(l)}}{\partial x_{\alpha}^{(l)}} = \widehat{\nabla}U^{(l)} \wedge \bar{e}_{\alpha}^{(l)}$, $l \in \{1, \dots, N\}$, $\alpha \in \{1, 2\}$.

Step 1 Extension of $U^{(l)}$ and of $\widehat{\nabla}U^{(l)}$ to $\omega_{l, \delta}$. The displacement

$$u^{(l)}(x) = \frac{1}{\delta} \left\{ U^{(l)}(\hat{x}^{(l)}) + \widehat{\nabla}U^{(l)}(\hat{x}^{(l)}) \wedge x_3^{(l)} \bar{e}_3^{(l)} \right\}, \quad x \in \Omega_{l, \delta} = \omega_l \times]-\delta, \delta[$$

of the plate $\Omega_{l, \delta}$ extends into a displacement still denoted $u^{(l)}$ of the plate $\Omega_{l, \delta}'' = \omega_{l, \delta} \times]-\delta, \delta[$. The extension $u^{(l)}$ is by construction equal to zero on $\Gamma_{0, \delta} \cap \Omega_{l, \delta}''$. We have

$$\mathcal{E}(u^{(l)}, \Omega_{l, \delta}'') \leq C\mathcal{E}(u^{(l)}, \Omega_{l, \delta}) \leq C\delta \|\widehat{\nabla}U^{(l)}\|_{H^1(\omega_{l, \mathbb{R}^3})}^2 \quad \mathcal{E}(u^{(l)}, \Omega_{l, \delta}'' \setminus \Omega_{l, \delta}) \leq C\delta \|\widehat{\nabla}U^{(l)}\|_{H^1(\Gamma_{l, \delta, \mathbb{R}^3})}^2$$

where $\Gamma_{l, \delta} = \{\hat{x}^{(l)} \in \omega_l \mid \text{dist}(\hat{x}^{(l)}, \partial\omega_l) < 2\eta_0\delta\}$. Let $U_e^{(l)}$ be the e.p.d. associated to $u^{(l)}$ by the formulas (2.1), its components $\frac{1}{\delta}\mathcal{U}^{(l)}$ and $\frac{1}{\delta}\mathcal{R}^{(l)}$ are the restrictions to $\omega_{l, 2\delta}$ of elements (denoted $\frac{1}{\delta}\mathcal{U}$, $\frac{1}{\delta}\mathcal{R}$) belonging to $H^1(\mathbf{S}_{\delta}, \mathbb{R}^3)$. They verify $\mathcal{U}_{|\omega_l}^{(l)} = U^{(l)}$, $\mathcal{R}_{|\omega_l}^{(l)} = \widehat{\nabla}U^{(l)}$ and

$$(4.20) \quad \begin{cases} \left\| \frac{\partial \mathcal{U}_3^{(l)}}{\partial x_1} + \mathcal{R}_2^{(l)} \right\|_{L^2(\omega_{l, \delta})} + \left\| \frac{\partial \mathcal{U}_3^{(l)}}{\partial x_2} - \mathcal{R}_1^{(l)} \right\|_{L^2(\omega_{l, \delta})} \leq C\delta \|\widehat{\nabla}U^{(l)}\|_{H^1(\Gamma_{l, \delta, \mathbb{R}^3})} \\ \left\| \nabla \mathcal{R}_1^{(l)} \right\|_{L^2(\omega_{l, \delta} \setminus \omega_{l, \mathbb{R}^2})} + \left\| \nabla \mathcal{R}_2^{(l)} \right\|_{L^2(\omega_{l, \delta} \setminus \omega_{l, \mathbb{R}^2})} \leq C\|\widehat{\nabla}U^{(l)}\|_{H^1(\Gamma_{l, \delta, \mathbb{R}^3})} \end{cases}$$

Step 2 We denote $V_\delta^{(l)}$ the displacement

$$V_\delta^{(l)}(x) = \frac{1}{\delta} \left\{ \mathcal{U}^{(l)}(\widehat{x}^{(l)}) + \mathcal{R}^{(l)}(\widehat{x}^{(l)}) \wedge x_3^{(l)} \vec{e}_3^{(l)} \right\}, \quad x \in \Omega'_{l,\delta}.$$

We modify $V_\delta^{(l)}$ in the neighborhood of the vertexes belonging to \mathcal{N} .

Let A be an vertex belonging to \mathcal{N} , for any edge J containing A , we have

$$U_{|J}(x) = U(A) + \widehat{\nabla}U(A) \wedge \overrightarrow{Ax} = r_A(x), \quad x \in J.$$

For any face $\overline{\omega}_l$ containing the vertex A , we define the displacement $U_{\delta,A}^{(l)}$ by

$$U_{\delta,A}^{(l)}(x) = V_\delta^{(l)}(x) m \left(\frac{\text{dist}(x,A)}{2\eta_0\delta} \right) + \frac{1}{\delta} \left\{ 1 - m \left(\frac{\text{dist}(x,A)}{2\eta_0\delta} \right) \right\} r_A(x), \quad x \in \Omega'_{l,\delta},$$

where m has been introduced by (4.7). We have $U_{\delta,A}^{(l)}(x) = 1/\delta r_A(x)$ in $B(A, 2\eta_0\delta) \cap \Omega'_{l,\delta}$. Let us remind that for any edge J containing the vertex A , we have $\mathcal{R}_{|J} = \widehat{\nabla}U_{|J}$ and $\mathcal{R}_{|J} \wedge \vec{e}_J = \widehat{\nabla}U_{|J} \wedge \vec{e}_J = \widehat{\nabla}U(A) \wedge \vec{e}_J$. Hence, thanks to (4.20), we have the following inequality :

$$\begin{aligned} & \|\mathcal{R} - \widehat{\nabla}U(A)\|_{L^2(B(A,4\eta_0\delta) \cap \mathbf{S}_{\delta,\mathbb{R}^3})} + \left\| \frac{\partial \mathcal{U}_3^{(l)}}{\partial x_\alpha} - \widehat{\nabla}U(A) \wedge \vec{e}_\alpha^{(l)} \right\|_{L^2(B(A,4\eta_0\delta) \cap \mathbf{S}_{\delta,\mathbb{R}^3})} \\ & \leq C\delta \left\{ \|\widehat{\nabla}U\|_{H^1(B(A,4\eta_0\delta) \cap \mathbf{S}_{\delta,\mathbb{R}^3})} + \|\widehat{\nabla}U^{(l)}\|_{H^1(\Gamma_{l,\delta,\mathbb{R}^3})} \right\}. \end{aligned}$$

This inequality implies that

$$\begin{aligned} & \|U_{\delta,A}^{(l)} - V_\delta^{(l)}\|_{L^2(B(A,4\eta_0\delta) \cap \Omega'_{l,\delta,\mathbb{R}^3})}^2 \leq C\delta^3 \left\{ \|\widehat{\nabla}U\|_{H^1(B(A,4\eta_0\delta) \cap \mathbf{S}_{\delta,\mathbb{R}^3})} + \|\widehat{\nabla}U^{(l)}\|_{H^1(\Gamma_{l,\delta,\mathbb{R}^3})} \right\}, \\ & \text{and} \quad \mathcal{E}(U_{\delta,A}^{(l)} - V_\delta^{(l)}, \Omega'_{l,\delta}) \leq C\delta \left\{ \|\widehat{\nabla}U\|_{H^1(B(A,4\eta_0\delta) \cap \mathbf{S}_{\delta,\mathbb{R}^3})} + \|\widehat{\nabla}U^{(l)}\|_{H^1(\Gamma_{l,\delta,\mathbb{R}^3})} \right\}. \end{aligned}$$

The displacement $U_{\delta,1}^{(l)} = \sum_{A \in \overline{\omega}_l \cap \mathcal{N}} U_{\delta,A}^{(l)}$ coincides with a rigid displacement independent of l in the neighborhood of each vertex contained in the face $\overline{\omega}_l$ and verifies

$$\mathcal{E}(U_{\delta,1}^{(l)} - V_\delta^{(l)}, \Omega'_{l,\delta}) \leq C\delta \left\{ \sum_{A \in \overline{\omega}_l \cap \mathcal{N}} \|\widehat{\nabla}U\|_{H^1(B(A,4\eta_0\delta) \cap \mathbf{S}_{\delta,\mathbb{R}^3})} + \|\widehat{\nabla}U^{(l)}\|_{H^1(\Gamma_{l,\delta,\mathbb{R}^3})} \right\}.$$

Step 3 We modify $U_{\delta,1}^{(l)}$ in the neighborhood of each edge belonging to \mathcal{J} .

Let J be an edge belonging to several faces and $\overline{\omega}_l$ a face of \mathbf{S} containing J . We take the orthornormal frame $(O_J; \vec{e}_J, \vec{e}_{J,\perp}^{(l)}, \vec{e}_3^{(l)})$ linked to the edge J and to the plate $\Omega_{l,\delta}$ containing this edge, O_J is an extremity of J and \vec{e}_J the direction of the edge ($x_{1J}^{(l)} = \widehat{x}^{(l)} \cdot \vec{e}_J \in [0, L_J]$, L_J the length of the edge). In this frame we consider the neighborhood of J

$$J_{\eta\delta}^{(l)} =]0, L_J[\times] - \eta\delta, \eta\delta[\times] - \delta, \delta[, \quad \eta \geq 1.$$

The restriction of $U_{\delta,1}^{(l)}$ to $J_\delta^{(l)}$ ($\eta = 1$) is decomposed into the sum of an elementary rod displacement $U_{e,J}^{(l)}$ and of a residual displacement,

$$U_{e,J}^{(l)}(x_{1J}, x_{2J}, x_3^{(l)}) = \mathcal{U}_J^{(l)}(x_{1J}) + \mathcal{R}_J^{(l)}(x_{1J}) \wedge (x_{2J} \vec{e}_{J,\perp}^{(l)} + x_3^{(l)} \vec{e}_3^{(l)}).$$

Let us remind that (see [4]) the components $\mathcal{U}_J^{(l)}$ and $\mathcal{R}_J^{(l)}$ of $U_{e,J}^{(l)}$ belong to $H^1(J, \mathbb{R}^3)$, and verify

$$\begin{cases} \delta^2 \left\| \frac{d\mathcal{R}_J^{(l)}}{dx_{1J}} \right\|_{L^2(J, \mathbb{R}^3)}^2 + \left\| \frac{d\mathcal{U}_J^{(l)}}{dx_{1J}} - \mathcal{R}_J^{(l)} \wedge \vec{e}_J \right\|_{L^2(J, \mathbb{R}^3)}^2 \leq \frac{C}{\delta^2} \mathcal{E}(U_{\delta,1}^{(l)}, J_\delta^{(l)}) \leq \frac{C}{\delta^2} \mathcal{E}(u^{(l)}, J_\delta^{(l)}) \\ \mathcal{D}(U_{\delta,1}^{(l)} - U_{e,J}^{(l)}, J_\delta^{(l)}) + \frac{1}{\delta^2} \|U_{\delta,1}^{(l)} - U_{e,J}^{(l)}\|_{L^2(J_\delta^{(l)}, \mathbb{R}^3)}^2 \leq C \mathcal{E}(U_{\delta,1}^{(l)}, J_\delta^{(l)}) \leq C \mathcal{E}(u^{(l)}, J_\delta^{(l)}) \end{cases}$$

By construction, the displacement $U_{e,J}^{(l)}$ coincides with a rigid displacement in the neighborhood of the edge extremities. We deduce that

$$\|U_{|J} - \mathcal{U}_J^{(l)}\|_{L^2(J, \mathbb{R}^3)}^2 \leq C \mathcal{E}(u^{(l)}, J_\delta^{(l)}), \quad \|\widehat{\nabla} U_{|J} - \mathcal{R}_J^{(l)}\|_{L^2(J, \mathbb{R}^3)}^2 \leq \frac{C}{\delta^2} \mathcal{E}(u^{(l)}, J_\delta^{(l)})$$

The edge J belongs to the faces $\bar{\omega}_{l_1}, \dots, \bar{\omega}_{l_p}$. Let $U_{e,J}$ be the elementary rod displacement equal to the mean value of the displacements $U_{e,J}^{(l_1)}, \dots, U_{e,J}^{(l_p)}$. The components of $U_{e,J}$ being \mathcal{U}_J and \mathcal{R}_J , we have

$$\begin{cases} \delta^2 \left\| \frac{d\mathcal{R}_J}{dx_{1J}} \right\|_{L^2(J, \mathbb{R}^3)}^2 + \left\| \frac{d\mathcal{U}_J}{dx_{1J}} - \mathcal{R}_J \wedge \vec{e}_J \right\|_{L^2(J, \mathbb{R}^3)}^2 \leq \frac{C}{\delta^2} \sum_{i=1}^p \mathcal{E}(u^{(l_i)}, J_\delta^{(l_i)}) \\ \sum_{i=1}^p \left\{ \mathcal{D}(U_{\delta,1}^{(l_i)} - U_{e,J}, J_\delta^{(l_i)}) + \frac{1}{\delta^2} \|U_{\delta,1}^{(l_i)} - U_{e,J}\|_{L^2(J_\delta^{(l_i)}, \mathbb{R}^3)}^2 \right\} \leq C \sum_{i=1}^p \mathcal{E}(u^{(l_i)}, J_\delta^{(l_i)}) \end{cases}$$

We deduce (see [4]) that

$$\begin{cases} \sum_{i=1}^p \mathcal{E}(U_{e,J}, J_{2\eta_0\delta}^{(l_i)}) \leq C \sum_{i=1}^p \mathcal{E}(u^{(l_i)}, J_\delta^{(l_i)}) \\ \sum_{i=1}^p \left\{ \mathcal{E}(U_{\delta,1}^{(l_i)} - U_{e,J}, J_{2\eta_0\delta}^{(l_i)}) + \frac{1}{\delta^2} \|U_{\delta,1}^{(l_i)} - U_{e,J}\|_{L^2(J_{2\eta_0\delta}^{(l_i)}, \mathbb{R}^3)}^2 \right\} \leq C \sum_{i=1}^p \mathcal{E}(u^{(l_i)}, J_{2\eta_0\delta}^{(l_i)}) \end{cases}$$

Now we modify the displacement $U_{\delta,1}^{(l)}$ in the neighborhood of the edge J ,

$$U_{\delta,J}^{(l)}(x) = U_{\delta,1}^{(l)}(x) \left\{ 1 - m \left(\frac{\text{dist}(\hat{x}^{(l)}, J)}{\eta_0\delta} \right) \right\} + U_{e,J}(x) m \left(\frac{\text{dist}(\hat{x}^{(l)}, J)}{\eta_0\delta} \right), \quad x \in \Omega'_{l,\delta}.$$

Then we have

$$\sum_{i=1}^p \mathcal{E}(U_{\delta,J}^{(l_i)} - U_{\delta,1}^{(l_i)}, \Omega'_{l_i,\delta}) \leq C \sum_{i=1}^p \mathcal{E}(u^{(l_i)}, J_{2\eta_0\delta}^{(l_i)})$$

Eventually the displacement U_δ obtained by modifying $U_{\delta,1}^{(l)}$ in the neighborhood of each edge of \mathcal{J} belongs to $H_{\Gamma_0}^1(\mathcal{S}_\delta, \mathbb{R}^3)$ and for any $l \in \{1, \dots, N\}$ verifies (3.23). \square

REFERENCES

- [1] P.G. Ciarlet, Ph. Destuynder. A justification of two-dimensional linear plate model, J. Mécanique, Vol. 18 (2.2), 1979, 315–344.
- [2] D. Cioranescu, A. Damlamian and G. Griso, Periodic Unfolding and Homogenization. CRAS, Ser. I 335 (2002) 99-104.
- [3] Ph. Destuynder. Une théorie asymptotique des plaques minces en élasticité linéaire. RMA 2, Masson (1986).
- [4] G. Griso. Asymptotic behavior of curved rods by the unfolding method. (To appear).
- [5] G. Griso. Asymptotic behavior of structures made of curved rods. (To appear).
- [6] G. Griso. Asymptotic behavior of structures made of plates. CRAS, Ser. I 336 (2003) 101-106.
- [7] H. Le Dret. Modeling of a folded plate, Comput. Mech., 5 (1990), 401–416.