

Evolution as a confounding parameter in scaling relations for galaxies

D. Fraix-Burnet

Abstract Early-type galaxies are characterized by many scaling relations. Evolutionary classifications find that some of these correlations are indeed generated by diversification. With a simple mathematical formalism, we show that even the so-called fundamental plane, a relatively tight correlation between three variables, can be easily explained as the artifact of the effect of another parameter influencing all, without any physical hypothesis. In other words, the fundamental plane is probably a confounding correlation, i.e. not physically causal. The complexity of the physics of galaxies and of their evolution suggests that the confounding parameter must be related to the level of diversification reached by the galaxies. Consequently, many scaling relations for galaxies are probably evolutionary correlations, explained by the statistical general evolution of most properties of galaxies.

1 The fundamental plane as a confounding correlation

The fundamental plane for early-type galaxies is a correlation between effective radius, the central velocity dispersion and the surface brightness within the effective radius [1, 2, 3]. Let us consider that the effective radius r_e , the central velocity dispersion σ and the luminosity L are all power-law functions of a same generic parameter \tilde{X} :

$$\begin{cases} r_e = A_1 \tilde{X}^p \\ \sigma = A_2 \tilde{X}^s \\ L = A_3 \tilde{X}^t \end{cases} \quad (1)$$

Université Joseph Fourier - Grenoble 1 / CNRS, Institut de Planétologie et d'Astrophysique de Grenoble, BP 53, F-38041 Grenoble cedex 9, France, e-mail: fraix@obs.ujf-grenoble.fr

The surface brightness μ_e can be expressed as $\mu_e = -2.5 \log(L/4\pi r_e^2) + m = (-2.5t + 5p) \log \tilde{X} + 2.5 \log(4\pi) + m$ where m is a constant of normalisation. Any linear correlation of the form

$$\log r_e = a \log \sigma + b \mu_e + c \quad (2)$$

translates to

$$\begin{cases} p = sa + (-2.5t + 5p)b \\ \log A_1 = a \log A_2 + b (2.5 \log(4\pi A_1^2/A_3) + m) + c. \end{cases} \quad (3)$$

If a solution can be found for a and b from Eq. 3, then the equation of the fundamental plane Eq. 2 is obtained. Conversely, the observations provide a , b and c , so that it is possible to derive p , s and t . There is no need of any further assumption to explain the fundamental plane.

2 Evolutionary correlations

In the course of diversification, many properties of galaxies change, and they tend to statistically change in a more or less monotonous way. It seems difficult to avoid the evolution to act as a confounding factor. It is a well-known problem of comparative methods in phylogeny [4].

We thus propose that the main confounding parameter is $\tilde{X} = T$ with T an indicator of the level of diversification, being something like an evolutionary clock not necessarily easily related to time or redshift. Indeed, the evolutionary clock, i.e. the factor $\tilde{X} = T$, can be hidden, not understandable analytically and not directly observable. It is related to an evolutionary classification that gathers objects according to their history. This work is published in [5].

References

- [1] Djorgovski, S., Davis, M.: Fundamental properties of elliptical galaxies, *ApJ*, **313**, 59-68 (1987)
- [2] Dressler, A., Lynden-Bell, D., Burstein, D., Davies, R. L., Faber, S. M., Terlevich, R., Wegner, G.: Spectroscopy and photometry of elliptical galaxies. I - A new distance estimator. *ApJ*, **313**, 42-58 (1987)
- [4] Felsenstein, J.: Phylogenies and the Comparative Method, *The American Naturalist*, **125**, 1-15 (1985)
- [5] Fraix-Burnet, D.: The Fundamental Plane of early-type galaxies as a confounding correlation, *MNRAS*, **416**, L36-L40, (2011) (arxiv:1106.3154)
- [3] Fraix-Burnet, D., Dugué M., Chattopadhyay T., Chattopadhyay A. K., Davoust E.: Structures in the fundamental plane of early-type galaxies, *MNRAS*, **407**, 2207-2222 (2010) (arXiv:1005.5645)