

ON A CATEGORY OF CLUSTER ALGEBRAS

IBRAHIM ASSEM, GRÉGOIRE DUPONT AND RALF SCHIFFLER

ABSTRACT. We introduce a category of cluster algebras with fixed initial seeds. This category has countable coproducts, which can be constructed combinatorially, but no products. We characterise isomorphisms and monomorphisms in this category and provide combinatorial methods for constructing special classes of monomorphisms and epimorphisms. In the case of cluster algebras from surfaces, we describe interactions between this category and the geometry of the surfaces.

CONTENTS

Introduction	1
Notations	2
1. Rooted cluster algebras	3
2. The category of rooted cluster algebras	6
3. Rooted cluster isomorphisms	9
4. Rooted cluster monomorphisms	13
5. Coproducts, products and amalgamated sums	19
6. Surjective rooted cluster morphisms	23
7. Surgery	33
Acknowledgements	36
References	36

INTRODUCTION

Cluster algebras are particular commutative algebras which were introduced by Fomin and Zelevinsky in [FZ02] in order to provide a combinatorial framework for studying total positivity and canonical bases in algebraic groups. Since then, a fast-growing literature focused on the numerous interactions of these algebras with various areas of mathematics like Lie theory, Poisson geometry, representation theory of algebras or mathematical physics. The study of the cluster algebras as algebraic structures in themselves can essentially be found in the seminal series of articles [FZ02, FZ03, BFZ05, FZ07] and their ring-theoretic properties were recently studied in [GLS11].

For a long time, an obstacle to the good understanding of cluster algebras was that they are defined *recursively* by applying a combinatorial process called *mutation*. However, the interactions of cluster algebras with either the (combinatorial) Teichmüller theory or the representation theory of algebras led to various closed formulae which enlightened the structure of cluster algebras, see [MSW11, DWZ10, Pla11b, HL10].

In order to get a better comprehension of cluster algebras, the next step is thus to define a categorical framework for their study. Therefore, one has to find what are the “right” morphisms between cluster algebras. The most natural idea is to look at ring homomorphisms which commute

Date: January 27, 2012.

with mutations. For bijective morphisms from a coefficient-free skew-symmetric cluster algebra to itself, this gave rise to the notion of *cluster automorphisms*, introduced in [ASS11]. However, in more general settings this idea turns out to be slightly too restrictive in order to get enough morphisms between non-isomorphic cluster algebras.

In this article, we slightly relax this first idea and propose a similar definition of cluster morphism between arbitrary skew-symmetrisable cluster algebras of geometric type (with non-invertible coefficients). The definition is still based on the idea that morphisms between cluster algebras should commute with mutations but, broadly speaking, we also allow the morphisms to specialise certain cluster variables to integral values or to send frozen variables to exchangeable ones (see Definition 2.2).

With this notion of morphisms, we obtain a category **Clus** with countable coproducts (Lemma 5.1) but generally without products (Proposition 5.4). We prove that in **Clus**, the isomorphisms are the bijective morphisms (Corollary 3.10), the monomorphisms are the injective monomorphisms (Proposition 4.3) while the epimorphisms are not necessarily surjective (Remark 6.1).

Inspired by the interactions between geometry and the combinatorics of cluster algebras associated with surfaces in the sense of [FST08], we define for arbitrary cluster algebras concepts of *gluings* and *cuttings* which provide natural classes of monomorphisms and epimorphisms in **Clus** (see Sections 5 and 7).

We also study specialisations of cluster variables in this categorical context. We prove that the usual specialisations of *frozen* variables to 1 yield epimorphisms in **Clus**. More surprisingly, for cluster algebras from surfaces or for acyclic cluster algebras, we prove that specialisations of exchangeable cluster variables also give rise to epimorphisms in **Clus** (Theorems 6.11 and 6.13).

The article is organised as follows. After a brief section recalling our conventions, Section 1 recalls the principal definitions on cluster algebras which we will use along the article. In Section 2, we introduce the notion of *rooted cluster morphisms* and the category **Clus** and we establish some basic properties. Section 3 is devoted to the study of isomorphisms in **Clus**, generalising previous results of [ASS11] on cluster automorphisms. Section 4 is devoted to the study of monomorphisms in **Clus**. Section 5 concerns the study of products and coproducts in **Clus** and their interactions with the geometry of surfaces. Section 6 is devoted to the study of epimorphisms in **Clus** and gives rise to a combinatorial process called *surgery*, whose interactions with the geometry of surfaces are studied in Section 7.

NOTATIONS

In this article, every ring A has an identity 1_A and every ring homomorphism $A \rightarrow B$ is assumed to send the identity 1_A to the identity 1_B . We denote by **Ring** the category of rings with ring homomorphisms.

If I is a countable set, we denote by $M_I(\mathbb{Z})$ the ring of locally finite matrices with integral entries indexed by $I \times I$ (we recall that a matrix $B = (b_{ij})_{i,j \in I}$ is *locally finite* if for every $i \in I$, the families $(b_{ij})_{j \in I}$ and $(b_{ji})_{j \in I}$ have finite support). We say that B is *skew-symmetrisable* if there exists a family of non-negative integers $(d_i)_{i \in I}$ such that $d_i b_{ij} = -d_j b_{ji}$ for any $i, j \in I$. If $J \subset I$, we denote by $B[J] = (b_{ij})_{i,j \in J}$ the submatrix of B formed by the entries labelled by $J \times J$.

If I and J are sets, we use the notation $I \setminus J = \{i \in I \mid i \notin J\}$ independently of whether J is contained in I or not. By a *countable* set we mean a set of cardinality at most \aleph_0 .

If R is a subring of a ring S , and if $\mathcal{X} \subset S$ is a set, we denote by $R[\mathcal{X}]$ the ring of all polynomials with coefficients in R evaluated in the elements of S . Note that this is not necessarily isomorphic to a ring of polynomials.

We recall that a *concrete* category is a category whose objects have underlying sets and whose morphisms between objects induce maps between the corresponding sets.

In order to make some statements clearer, it might be convenient for the reader to use a combinatorial representation of the skew-symmetrisable locally finite matrices as *valued quivers*. If $B \in M_I(\mathbb{Z})$ is a locally finite skew-symmetric matrix, we associate with B a valued quiver Q_B whose points are indexed by I and such that for any $i, j \in I$, if $b_{ij} > 0$ (so that $b_{ji} < 0$), then we draw a valued arrow

$$i \xrightarrow{(b_{ij}, -b_{ji})} j$$

in Q_B . As B is skew-symmetrisable the valued quiver Q_B has no oriented cycles of length at most two. In the case where B is skew-symmetric, if $i, j \in I$ are such that $b_{ij} > 0$, we usually draw b_{ij} arrows from i to j in Q_B instead of a unique arrow with valuation $(b_{ij}, -b_{ji})$.

1. ROOTED CLUSTER ALGEBRAS

In this section we recall the definition of a cluster algebra of geometric type. As opposed to the initial definition formulated in [FZ02], we consider *non-invertible* coefficients.

1.1. Seeds and mutations.

Definition 1.1 (Seeds). A *seed* is a triple $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ such that :

- (1) \mathbf{x} is a countable set of undeterminates over \mathbb{Z} , called the *cluster* of Σ ;
- (2) $\mathbf{ex} \subset \mathbf{x}$ is a subset of \mathbf{x} whose elements are the *exchangeable variables* of Σ ;
- (3) $B = (b_{xy})_{x,y \in \mathbf{x}} \in M_{\mathbf{x}}(\mathbb{Z})$ is a (locally finite) skew-symmetrisable matrix, called the *exchange matrix* of Σ .

The variables in $\mathbf{x} \setminus \mathbf{ex}$ are the *frozen variables* of Σ . A seed $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ is *coefficient-free* if $\mathbf{ex} = \mathbf{x}$ and in this case, we simply write $\Sigma = (\mathbf{x}, B)$. A seed is *finite* if \mathbf{x} is a finite set.

Given a seed Σ , the field $\mathcal{F}_{\Sigma} = \mathbb{Q}(x \mid x \in \mathbf{x})$ is called the associated *ambient field*.

Definition 1.2 (Mutation). Given a seed $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ and an exchangeable variable $x \in \mathbf{ex}$, the image of the *mutation* of Σ in the direction x is the seed

$$\mu_x(\Sigma) = (\mathbf{x}', \mathbf{ex}', B')$$

given by

- (1) $\mathbf{x}' = (\mathbf{x} \setminus \{x\}) \sqcup \{x'\}$ where

$$xx' = \prod_{\substack{y \in \mathbf{x} ; \\ b_{xy} > 0}} y^{b_{xy}} + \prod_{\substack{y \in \mathbf{x} ; \\ b_{xy} < 0}} y^{-b_{xy}}.$$

- (2) $\mathbf{ex}' = (\mathbf{ex} \setminus \{x\}) \sqcup \{x'\}$.
- (3) $B' = (b'_{yz}) \in M_{\mathbf{x}}(\mathbb{Z})$ is given by

$$b'_{yz} = \begin{cases} -b_{yz} & \text{if } x = y \text{ or } x = z ; \\ b_{yz} + \frac{1}{2}(|b_{yx}|b_{xz} + b_{yx}|b_{xz}|) & \text{otherwise.} \end{cases}$$

For any $y \in \mathbf{x}$ we denote by $\mu_{x,\Sigma}(y)$ the variable corresponding to y in the cluster of the seed $\mu_x(\Sigma)$, that is, $\mu_{x,\Sigma}(y) = y$ if $y \neq x$ and $\mu_{x,\Sigma}(x) = x'$ where x' is defined as above. If there is no risk of confusion, we simply write $\mu_x(y)$ instead of $\mu_{x,\Sigma}(y)$.

The set \mathbf{x}' is again a free generating set of \mathcal{F}_{Σ} and the mutation is involutive in the sense that $\mu_{x'} \circ \mu_x(\Sigma) = \Sigma$, for each $x \in \mathbf{ex}$.

Definition 1.3 (Admissible sequence of variables). Let $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ be a seed. We say that (x_1, \dots, x_l) is Σ -*admissible* if x_1 is exchangeable in Σ and if, for every $i \geq 2$, the variable x_i is exchangeable in $\mu_{x_{i-1}} \circ \dots \circ \mu_{x_1}(\Sigma)$.

Given a seed $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$, its *mutation class* is the set $\text{Mut}(\Sigma)$ of all seeds which can be obtained from Σ by applying successive mutations along finite admissible sequences of variables. In other words,

$$\text{Mut}(\Sigma) = \{\mu_{x_n} \circ \cdots \circ \mu_{x_1}(\Sigma) \mid n > 0 \text{ and } (x_1, \dots, x_n) \text{ is } \Sigma\text{-admissible}\}.$$

Two seeds in the same mutation class are *mutation-equivalent*.

1.2. Rooted cluster algebras.

Definition 1.4 (Rooted cluster algebra). Let Σ be a seed. The *rooted cluster algebra with initial seed* Σ is the pair (Σ, \mathcal{A}) where \mathcal{A} is the \mathbb{Z} -subalgebra of \mathcal{F}_Σ given by :

$$\mathcal{A} = \mathcal{A}(\Sigma) = \mathbb{Z} \left[x \mid x \in \bigcup_{(\mathbf{x}, \mathbf{ex}, B) \in \text{Mut}(\Sigma)} \mathbf{x} \right] \subset \mathcal{F}_\Sigma.$$

The variables (exchangeable variables and frozen variables respectively) arising in the clusters of seeds which are mutation-equivalent to Σ are the *cluster variables* (or the *exchangeable variables* and *frozen variables* respectively) of the rooted cluster algebra (Σ, \mathcal{A}) . We denote by \mathcal{X}_Σ the set of cluster variables in (Σ, \mathcal{A}) .

In order to simplify notations, a rooted cluster algebra (Σ, \mathcal{A}) is in general simply denoted by $\mathcal{A}(\Sigma)$ but one should keep in mind that a rooted cluster algebra is always viewed together *with* its initial seed.

Remark 1.5. This definition authorises seeds whose clusters are empty. Such seeds are called *empty seeds* and by convention the rooted cluster algebra corresponding to an empty seed is \mathbb{Z} .

Example 1.6. If $\Sigma = (\mathbf{x}, \emptyset, B)$ has no exchangeable variables, then

$$\mathcal{A}(\Sigma) = \mathbb{Z}[x \mid x \in \mathbf{x}]$$

is a polynomial ring in countably many variables.

Remark 1.7. In the original definition of cluster algebras given in [FZ02], the frozen variables are supposed to be invertible in the cluster algebra. However, it is known that cluster variables in a cluster algebra of geometric type are Laurent polynomials in the exchangeable variables with polynomial coefficients in the frozen ones (see for instance [FZ03, Proposition 11.2]). Therefore, the cluster algebra structure can essentially be considered without inverting the coefficients. Also, several “natural” examples of cluster algebras arise with non-invertible coefficients, as for instance cluster algebras arising in Lie theory as polynomial rings (see [GLS11, §6.4]) or cluster structures on rings of homogeneous coordinates on Grassmannians (see [GSV10, §2.1] or Section 6.6).

Of course, if one wants to recover the initial definition from ours, it is enough to localise the cluster algebra at the frozen variables. Nevertheless, some of our results (in particular the crucial Lemma 4.1) require that frozen variables be non-invertible.

1.3. Rooted cluster algebras from surfaces. In this article, we are often interested in the particular class of rooted cluster algebras associated with marked surfaces in the sense of [FST08]. We recall that a *marked surface* is a pair (S, M) where S is an oriented 2-dimensional Riemann surface and M is a finite set of marked points in the closure of S such that each connected component of the boundary ∂S of the surface S contains at least one marked point in M . We also assume that none of the connected components of (S, M) is a *degenerate marked surface*, that is, a surface which is homeomorphic to one of the following surfaces :

- a sphere with one, two or three punctures,
- an unpunctured or a once-punctured monogon,
- an unpunctured digon or an unpunctured triangle.

All curves in (S, M) are considered up to isotopy with respect to the set M of marked points. Therefore, two curves γ and γ' are called *distinct* if they are not isotopic. Two curves γ and γ' are called *compatible* if there exist representatives of their respective isotopy classes which do not intersect in $S \setminus M$.

An *arc* is a curve joining two marked points in (S, M) which is compatible with itself. An arc is a *boundary arc* if it is isotopic to a connected component of $\partial S \setminus M$, otherwise it is *internal*.

A *triangulation* of (S, M) is a maximal collection of arcs which are pairwise distinct and compatible. The arcs of the triangulation cut the surface into triangles (which may be *self-folded*).

With any triangulation T of (S, M) we can associate a skew-symmetric matrix B^T as in [FST08]. For the convenience of the reader, we recall this construction in the case where T has no self-folded triangles (for the general case we refer the reader to [FST08, §4]). For any triangle Δ in T , we define a matrix B^Δ , indexed by the arcs in T and given by

$$B_{\gamma, \gamma'}^\Delta = \begin{cases} 1 & \text{if } \gamma \text{ and } \gamma' \text{ are sides of } \Delta \text{ and } \gamma' \text{ follows } \gamma \text{ in the positive direction ;} \\ -1 & \text{if } \gamma \text{ and } \gamma' \text{ are sides of } \Delta \text{ and } \gamma' \text{ follows } \gamma \text{ in the negative direction ;} \\ 0 & \text{otherwise.} \end{cases}$$

The matrix B^T is then given by

$$B^T = \sum_{\Delta} B^\Delta$$

where Δ runs over all the triangles in T .

In terms of quivers, the quiver Q_T corresponding to B^T is the quiver such that :

- the points in Q_T are the arcs of T ,
- the frozen points in Q_T are the boundary arcs of T ,
- there is an arrow $\gamma \rightarrow \gamma'$ if and only if γ and γ' are sides of a same triangle and γ' follows γ in the positive direction,
- a maximal collection of 2-cycles is removed.

Example 1.8. Figure 1 shows an example of quiver obtained from a triangulation of a hexagon. Points in white correspond to frozen variables and points in black correspond to exchangeable variables. The dashed arrows, joining frozen points, are precisely those which we remove in the *simplification of the seed* (see Definition 3.5).




FIGURE 1. The quiver of a triangulation of a hexagon.

Then, we can associate with T the *seed* $\Sigma_T = (\mathbf{x}_T, \mathbf{ex}_T, B_T)$ where :

- \mathbf{x}_T is indexed by the arcs in T ;
- \mathbf{ex}_T is indexed by the internal arcs in T ;
- B^T is the matrix defined above.

The *rooted cluster algebra associated with the triangulation* T is therefore $\mathcal{A}(\Sigma_T)$.

It is proved in [FST08] that if T and T' are two triangulations of (S, M) , then Σ_T and $\Sigma_{T'}$ can be joined by a sequence of mutations. Therefore, up to a ring isomorphism, $\mathcal{A}(\Sigma_T)$ does not depend on the choice of the triangulation T and is called *the cluster algebra* $\mathcal{A}(S, M)$ associated with (S, M) , see [FST08] for further details.

More generally, it is possible to associate to a marked surface a cluster algebra with an arbitrary choice of coefficients and not only the coefficients arising from the boundary arcs considered above. However, for geometric statements (see for instance Sections 4.1, 6.3 or 7.3), it is usually more natural to consider coefficients associated with the boundary arcs.

Finally, in order to avoid technicalities, we only consider *untagged* triangulations but all the results we present can easily be extended to the case of tagged triangulations.

2. THE CATEGORY OF ROOTED CLUSTER ALGEBRAS

2.1. Rooted cluster morphisms.

Definition 2.1 (Biadmissible sequences). Let $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ and $\Sigma' = (\mathbf{x}', \mathbf{ex}', B')$ be two seeds and let $f : \mathcal{F}_\Sigma \rightarrow \mathcal{F}_{\Sigma'}$ be a map. A sequence $(x_1, \dots, x_n) \subset \mathcal{A}(\Sigma)$ is called (f, Σ, Σ') -*biadmissible* (or simply *biadmissible* if there is no risk of confusion) if it is Σ -admissible and if $(f(x_1), \dots, f(x_n))$ is Σ' -admissible.

We fix two seeds $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ and $\Sigma' = (\mathbf{x}', \mathbf{ex}', B')$.

Definition 2.2 (Rooted cluster morphisms). A *rooted cluster morphism* from $\mathcal{A}(\Sigma)$ to $\mathcal{A}(\Sigma')$ is a ring homomorphism from $\mathcal{A}(\Sigma)$ to $\mathcal{A}(\Sigma')$ such that :

(CM1) $f(\mathbf{x}) \subset \mathbf{x}' \sqcup \mathbb{Z}$;

(CM2) $f(\mathbf{ex}) \subset \mathbf{ex}' \sqcup \mathbb{Z}$;

(CM3) For every (f, Σ, Σ') -biadmissible sequence (x_1, \dots, x_n) , we have

$$f(\mu_{x_n} \circ \dots \circ \mu_{x_1, \Sigma}(y)) = \mu_{f(x_n)} \circ \dots \circ \mu_{f(x_1), \Sigma'}(f(y))$$

for any y in \mathbf{x} .

We sometimes say that f *commutes with biadmissible mutations* if it satisfies (CM3).

Remark 2.3. A rooted cluster morphism may send a frozen cluster variable to an exchangeable cluster variable whereas (CM2) prevents the opposite from happening.

Remark 2.4. Given an explicit ring homomorphism f between two rooted cluster algebras, (CM3) is difficult to check since it requires to test every biadmissible sequence, and there are in general infinitely many. However, we shall see that for instance for isomorphisms, it is sometimes possible to simplify this situation (see Lemma 3.8).

Proposition 2.5. *The composition of rooted cluster morphisms is a rooted cluster morphism.*

Proof. We fix three rooted cluster algebras $\mathcal{A}_1, \mathcal{A}_2$ and \mathcal{A}_3 with respective initial seeds Σ_1, Σ_2 and Σ_3 where $\Sigma_i = (\mathbf{x}_i, \mathbf{ex}_i, B^i)$ and consider rooted cluster morphisms $f : \mathcal{A}_1 \rightarrow \mathcal{A}_2$ and $g : \mathcal{A}_2 \rightarrow \mathcal{A}_3$. The composition $g \circ f$ is a ring homomorphism from \mathcal{A}_1 to \mathcal{A}_3 . Moreover, we have :

(CM1) $(g \circ f)(\mathbf{x}_1) = g(f(\mathbf{x}_1)) \subset g(\mathbf{x}_2 \sqcup \mathbb{Z}) \subset \mathbf{x}_3 \sqcup \mathbb{Z}$;

(CM2) $(g \circ f)(\mathbf{ex}_1) = g(f(\mathbf{ex}_1)) \subset g(\mathbf{ex}_2 \sqcup \mathbb{Z}) \subset \mathbf{ex}_3 \sqcup \mathbb{Z}$;

(CM3) Let (x_1, \dots, x_n) be a $((g \circ f), \Sigma_1, \Sigma_3)$ -biadmissible sequence. We claim that (x_1, \dots, x_n) is (f, Σ_1, Σ_2) -biadmissible and that $(f(x_1), \dots, f(x_n))$ is (g, Σ_2, Σ_3) -biadmissible. Indeed, since f satisfies (CM2), an exchangeable variable $x \in \mathbf{ex}_1$ is sent either to an exchangeable variable in \mathbf{ex}_2 or to an integer. If $f(x) \in \mathbb{Z}$, because g is a ring homomorphism, then $g(f(x)) \in \mathbb{Z}$ and therefore $(g \circ f)(x)$ is not exchangeable, a contradiction. Thus, (x_1, \dots, x_n) is (f, Σ_1, Σ_2) -biadmissible. It follows that $(f(x_1), \dots, f(x_n))$ is Σ_2 -admissible

and therefore, as $(g(f(x_1)), \dots, g(f(x_n)))$ is Σ_3 -admissible, the sequence $(f(x_1), \dots, f(x_n))$ is (g, Σ_2, Σ_3) -biadmissible.

Now, since f satisfies **(CM3)**, for every i such that $1 \leq i \leq n$ and any $x \in \mathbf{x}_1$, we have

$$f(\mu_{x_i} \circ \dots \circ \mu_{x_1}(x)) = \mu_{f(x_i)} \circ \dots \circ \mu_{f(x_1)}(f(x)).$$

and, since g satisfies **(CM3)**, for any (g, Σ_2, Σ_3) -biadmissible sequence (y_1, \dots, y_n) and any i such that $1 \leq i \leq n$ and $y \in \mathbf{x}_2$, we have

$$g(\mu_{y_i} \circ \dots \circ \mu_{y_1}(y)) = \mu_{g(y_i)} \circ \dots \circ \mu_{g(y_1)}(g(y)).$$

Then, as $f(\mathbf{x}_1) \subset \mathbf{x}_2 \sqcup \mathbb{Z}$, we get

$$(g \circ f)(\mu_{x_i} \circ \dots \circ \mu_{x_1}(x)) = \mu_{(g \circ f)(x_i)} \circ \dots \circ \mu_{(g \circ f)(x_1)}((g \circ f)(x))$$

for any $x \in \mathbf{x}$ so that $(g \circ f)$ satisfies **(CM3)**.

Thus, $g \circ f : \mathcal{A}_1 \rightarrow \mathcal{A}_3$ is a rooted cluster morphism. □

Therefore, we can set the following definition :

Definition 2.6. The *category of rooted cluster algebras* is the category **Clus** defined by :

- The objects in **Clus** are the rooted cluster algebras ;
- The morphisms between two rooted cluster algebras are the rooted cluster morphisms.

Remark 2.7. One should observe the importance of the condition **(CM2)** in the proof of Proposition 2.5 in order to obtain well-defined compositions of rooted cluster morphisms. If this condition is removed from the definition of a rooted cluster morphism, one can easily construct examples of rooted cluster morphisms whose composition is not a rooted cluster morphism.

For instance, consider $\Sigma_1 = (x, x, [0])$, $\Sigma_2 = (x, \emptyset, [0])$ and $\Sigma_3 = \left((x, y), (x, y), \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \right)$.

Let f denote the identity in $\mathbb{Q}(x)$ and let g denote the canonical inclusion of $\mathbb{Q}(x)$ in $\mathbb{Q}(x, y)$. By construction f and g satisfy **(CM1)** and not **(CM2)**. Since there are neither non-empty (f, Σ_1, Σ_2) -biadmissible sequences nor non-empty (g, Σ_2, Σ_3) -biadmissible sequences, f and g satisfy **(CM3)**. The composition $g \circ f$ also satisfies **(CM1)** and (x) is $(g \circ f, \Sigma_1, \Sigma_3)$ -biadmissible. However,

$$(g \circ f)(\mu_x(x)) = \frac{2}{x} \neq \frac{1+y}{x} = \mu_{(g \circ f)(x)}((g \circ f)(x)).$$

Therefore, $g \circ f$ does not satisfy **(CM3)**.

2.2. Ideal rooted cluster morphisms.

Definition 2.8 (Image seed). Let $f : \mathcal{A}(\Sigma) \rightarrow \mathcal{A}(\Sigma')$ be a rooted cluster morphism. The *image seed* of Σ under f is

$$f(\Sigma) = (\mathbf{x}' \cap f(\mathbf{x}), \mathbf{e}\mathbf{x}' \cap f(\mathbf{e}\mathbf{x}), B'[f(\mathbf{x})]).$$

Lemma 2.9. Let $f : \mathcal{A}(\Sigma) \rightarrow \mathcal{A}(\Sigma')$ be a rooted cluster morphism and let (y_1, \dots, y_l) be an $f(\Sigma)$ -admissible sequence. Then (y_1, \dots, y_l) is Σ' -admissible and

$$\mu_{y_1} \circ \dots \circ \mu_{y_l, f(\Sigma)}(y_1) = \mu_{y_1} \circ \dots \circ \mu_{y_l, \Sigma'}(y_1).$$

Proof. Let $\Sigma = (\mathbf{x}, \mathbf{e}\mathbf{x}, B)$ and $\Sigma' = (\mathbf{x}', \mathbf{e}\mathbf{x}', B')$. Because $\mu_{y_1}(B'[f(\mathbf{x})]) = B'[\mu_{y_1}(f(\mathbf{x}))]$, it is enough to prove the statement for $l = 1$ and to proceed by induction. By definition, exchangeable variables in $f(\Sigma)$ are exchangeable in Σ' . Now, if y_1 is $f(\Sigma)$ -admissible, we have

$$\mu_{y_1, f(\Sigma)}(y_1) = \frac{1}{y_1} \left(\prod_{\substack{b'_{zy_1} > 0 \\ z \in \mathbf{x}' \cap f(\mathbf{x})}} z^{b'_{zy_1}} + \prod_{\substack{b'_{zy_1} < 0 \\ z \in \mathbf{x}' \cap f(\mathbf{x})}} z^{-b'_{zy_1}} \right).$$

Because y_1 is exchangeable in $f(\Sigma)$, there exists some $x_1 \in \mathbf{ex}$ such that $y_1 = f(x_1)$ and we have

$$\mu_{x_1, \Sigma}(x_1) = \frac{1}{x_1} \left(\prod_{\substack{b_{ux_1} > 0 \\ u \in \mathbf{x}}} u^{b_{ux_1}} + \prod_{\substack{b_{ux_1} < 0 \\ u \in \mathbf{x}}} u^{-b_{ux_1}} \right).$$

Therefore, since f satisfies **(CM3)**, we get

$$\mu_{y_1, \Sigma'}(y_1) = f(\mu_{x_1, \Sigma}(x_1)) = \frac{1}{y_1} \left(\prod_{\substack{b_{ux_1} > 0 \\ u \in \mathbf{x}}} f(u)^{b_{ux_1}} + \prod_{\substack{b_{ux_1} < 0 \\ u \in \mathbf{x}}} f(u)^{-b_{ux_1}} \right).$$

And by definition

$$\mu_{y_1, \Sigma'}(y_1) = \frac{1}{y_1} \left(\prod_{\substack{b'_{zy_1} > 0 \\ z \in \mathbf{x}'}} z^{b'_{zy_1}} + \prod_{\substack{b'_{zy_1} < 0 \\ z \in \mathbf{x}'}} z^{-b'_{zy_1}} \right).$$

Therefore, $b'_{zy_1} = 0$ for any $z \in \mathbf{x}' \setminus f(\mathbf{x})$ so that $\mu_{y_1, \Sigma'}(y_1) = \mu_{y_1, f(\Sigma)}(y_1)$. \square

Lemma 2.10. *Let $f : \mathcal{A}(\Sigma) \rightarrow \mathcal{A}(\Sigma')$ be a rooted cluster morphism. Then $\mathcal{A}(f(\Sigma)) \subset f(\mathcal{A}(\Sigma))$.*

Proof. Let $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ and $\Sigma' = (\mathbf{x}', \mathbf{ex}', B')$. We have to prove that any cluster variable in $\mathcal{A}(f(\Sigma))$ belongs to $f(\mathcal{A}(\Sigma))$. Fix an arbitrary cluster variable $y \in \mathcal{A}(f(\Sigma))$, then there exists an $f(\Sigma)$ -admissible sequence (y_1, \dots, y_l) such that $y = \mu_{y_l} \circ \dots \circ \mu_{y_1}(y_1)$.

We claim that any $f(\Sigma)$ -admissible sequence (y_1, \dots, y_l) lifts to an $(f, \Sigma, f(\Sigma))$ -biadmissible sequence (x_1, \dots, x_l) . If $l = 1$, then $y_1 = f(x_1)$ for some $x_1 \in \mathbf{ex}$ and thus (x_1) is biadmissible. Assume now that for $k < l$ the sequence (y_1, \dots, y_k) lifts to a biadmissible sequence (x_1, \dots, x_k) . Then y_{k+1} is exchangeable in $\mu_{y_k} \circ \dots \circ \mu_{y_1}(f(\Sigma))$ and thus there exists some $x \in \mathbf{ex}$ such that $y_{k+1} = \mu_{y_k} \circ \dots \circ \mu_{y_1}(f(x))$. Because f satisfies **(CM3)**, we get

$$y_{k+1} = f(\mu_{x_k} \circ \dots \circ \mu_{x_1}(x)).$$

Therefore, $x_{k+1} = \mu_{x_k} \circ \dots \circ \mu_{x_1}(x)$ is exchangeable in $\mu_{x_k} \circ \dots \circ \mu_{x_1}(\Sigma)$ and (y_1, \dots, y_{k+1}) lifts to (x_1, \dots, x_{k+1}) . The claim follows by induction.

If $l = 0$, then by definition of $f(\Sigma)$, the elements in the cluster of $f(\Sigma)$ belong to $f(\mathcal{A}(\Sigma))$. If $l \geq 0$, then it follows from the claim that (y_1, \dots, y_l) lifts to an $(f, \Sigma, f(\Sigma))$ -biadmissible sequence (x_1, \dots, x_l) in $\mathcal{A}(\Sigma)$. Moreover, we have

$$\begin{aligned} y &= \mu_{y_l} \circ \dots \circ \mu_{y_1, f(\Sigma)}(y_1) \\ &= \mu_{f(x_l)} \circ \dots \circ \mu_{f(x_1), f(\Sigma)}(f(x_1)) \\ &= \mu_{f(x_l)} \circ \dots \circ \mu_{f(x_1), \Sigma'}(f(x_1)) \\ &= f(\mu_{x_l} \circ \dots \circ \mu_{x_1, \Sigma}(x_1)) \in f(\mathcal{A}(\Sigma)). \end{aligned}$$

where the third equality follows from Lemma 2.9 and the last one from the fact that f satisfies **(CM3)**. \square

Definition 2.11 (Ideal rooted cluster morphism). A rooted cluster morphism $f : \mathcal{A}(\Sigma) \rightarrow \mathcal{A}(\Sigma')$ is called *ideal* if $\mathcal{A}(f(\Sigma)) = f(\mathcal{A}(\Sigma))$.

We shall meet along the article several natural classes of rooted cluster morphisms which are ideal (see for instance Corollary 4.5, Proposition 6.4 or Proposition 6.10). However, we do not know whether or not every rooted cluster morphism is ideal. We may thus state the following problem :

Problem 2.12. *Characterise the rooted cluster morphisms which are ideal.*

Example 2.13. Consider the seeds

$$\Sigma = \left((x_1, x_2, x_3), (x_2), \begin{bmatrix} 0 & 1 & 0 \\ -1 & 0 & 1 \\ 0 & -1 & 0 \end{bmatrix} \right) \text{ and } \Sigma' = \left((y_1, y_2, y_3), (y_1, y_2, y_3), \begin{bmatrix} 0 & 1 & 0 \\ -1 & 0 & 1 \\ 0 & -1 & 0 \end{bmatrix} \right)$$

and the ring homomorphism

$$f : \begin{cases} \mathcal{F}_\Sigma & \longrightarrow & \mathcal{F}_{\Sigma'} \\ x_1 & \mapsto & 1 \\ x_2 & \mapsto & y_1 \\ x_3 & \mapsto & y_2 \end{cases}$$

The only (f, Σ, Σ') -biadmissible sequence to consider is (x_2) and

$$f(\mu_{x_2}(x_2)) = f\left(\frac{x_1 + x_3}{x_2}\right) = \frac{1 + y_2}{y_1} = \mu_{y_1}(y_1) = \mu_{f(x_2)}(f(x_2))$$

so that f is a rooted cluster morphism $\mathcal{A}(\Sigma) \rightarrow \mathcal{A}(\Sigma')$. Moreover we have

$$f(\Sigma) = \left((y_1, y_2), (y_1) \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \right).$$

Therefore

$$\mathcal{A}(\Sigma) = \mathbb{Z}[x_1, x_2, x_3, \frac{x_1 + x_3}{x_2}] \text{ and } \mathcal{A}(f(\Sigma)) = \mathbb{Z}[y_1, y_2, \frac{1 + y_2}{y_1}]$$

so that $\mathcal{A}(f(\Sigma)) = f(\mathcal{A}(\Sigma))$ and therefore f is ideal.

Definition 2.14 (Rooted cluster ideal). A (ring theoretic) ideal I in a rooted cluster algebra $\mathcal{A}(\Sigma)$ is called a *rooted cluster ideal* if the quotient $\mathcal{A}(\Sigma)/I$ can be endowed with a structure of rooted cluster algebra such that the canonical projection is a rooted cluster morphism.

Proposition 2.15. *Let $f : \mathcal{A}(\Sigma) \rightarrow \mathcal{A}(\Sigma')$ be an ideal rooted cluster morphism. Then $\text{Ker}(f)$ is a rooted cluster ideal.*

Proof. Let $f : \mathcal{A}(\Sigma) \rightarrow \mathcal{A}(\Sigma')$ be an ideal rooted cluster morphism. Then f induces a ring isomorphism

$$\mathcal{A}(\Sigma)/\text{Ker}(f) \simeq f(\mathcal{A}(\Sigma)) = \mathcal{A}(f(\Sigma))$$

endowing $\mathcal{A}(\Sigma)/\text{Ker}(f)$ with a structure of rooted cluster algebra with initial seed $f(\Sigma)$.

Moreover, since f is a rooted cluster morphism, the morphism $\bar{f} : \mathcal{A}(\Sigma) \rightarrow \mathcal{A}(f(\Sigma))$ induced by f is also a rooted cluster morphism and therefore $\mathcal{A}(\Sigma) \rightarrow \mathcal{A}(\Sigma)/\text{Ker}(f)$ is a rooted cluster morphism. \square

3. ROOTED CLUSTER ISOMORPHISMS

In this section we characterise isomorphisms in **Clus** which we call *rooted cluster isomorphisms*. These results are generalisations of those obtained in [ASS11] for coefficient-free skew-symmetric cluster algebras. We recall that an *isomorphism* in **Clus** is an invertible morphism.

We start with a general lemma on surjective morphisms :

Lemma 3.1. *Let $\Sigma = (\mathbf{x}_1, \mathbf{ex}_1, B^1)$ and $\Sigma_2 = (\mathbf{x}_2, \mathbf{ex}_2, B^2)$ be two seeds and $f : \mathcal{A}(\Sigma_1) \rightarrow \mathcal{A}(\Sigma_2)$ be a surjective ring homomorphism satisfying (CM1). Then $\mathbf{x}_2 \subset f(\mathbf{x}_1)$ and $\mathbf{ex}_2 \subset f(\mathbf{ex}_1)$.*

Proof. Let $z \in \mathbf{x}_2$. Since f is surjective, there exists $y \in \mathcal{A}(\Sigma_1)$ such that $f(y) = z$. According to the Laurent phenomenon, there exists a Laurent polynomial L such that $y = L(x|x \in \mathbf{x}_1)$. Therefore, $z = f(y) = L(f(x)|x \in \mathbf{x}_1)$. Since f satisfies **(CM1)**, we know that $f(\mathbf{x}_1) \subset \mathbf{x}_2 \sqcup \mathbb{Z}$. If $z \notin f(\mathbf{x}_1)$, since \mathbf{x}_2 is a transcendence basis of \mathcal{F}_{Σ_2} , we get a contradiction. Therefore, $z \in f(\mathbf{x}_1)$ and thus $\mathbf{x}_2 \subset f(\mathbf{x}_1)$.

Fix now $z \in \mathbf{ex}_2$. According to the above discussion, we know that $f^{-1}(z) \cap \mathbf{x}_1 \neq \emptyset$. Since f is surjective, there exists $X \in \mathcal{A}(\Sigma_1)$ such that $f(X) = \mu_{z, \Sigma_2}(z)$. Now we know that $X \in \mathbb{Z}[\mathbf{x}_1 \setminus \mathbf{ex}_1][\mathbf{ex}_1^{\pm 1}]$. Therefore, if $f^{-1}(z) \cap \mathbf{ex}_1 = \emptyset$, then $f^{-1}(z) \cap \mathbf{x}_1 \subset (\mathbf{x}_1 \setminus \mathbf{ex}_1)$ and therefore, X is a sum of Laurent monomials with non-negative partial degree with respect to any $t \in f^{-1}(z) \cap \mathbf{x}_1$. Therefore, $f(X)$ is a sum of Laurent monomials with non-negative partial degree with respect to z , a contradiction since $\mu_{z, \Sigma_2}(z) = f(X)$ is the sum of two Laurent monomials with partial degree -1 with respect to z . Therefore $f^{-1}(z) \cap \mathbf{ex}_1 \neq \emptyset$ so that $\mathbf{ex}_2 \subset f(\mathbf{ex}_1)$. \square

Corollary 3.2. *Let $\Sigma_i = (\mathbf{x}_i, \mathbf{ex}_i, B^i)$ be a seed for $i \in \{1, 2\}$ and let $f : \mathcal{A}(\Sigma_1) \rightarrow \mathcal{A}(\Sigma_2)$ be a bijective ring homomorphism satisfying **(CM1)**. Then f induces a bijection from \mathbf{x}_1 to \mathbf{x}_2 . If moreover f satisfies **(CM2)**, then f induces a bijection from \mathbf{ex}_1 to \mathbf{ex}_2 .*

Proof. Since f is injective and satisfies **(CM1)**, it induces an injection $\mathbf{x}_1 \rightarrow \mathbf{x}_2$. Since f is also surjective, it follows from Lemma 3.1 that $\mathbf{x}_2 \subset f(\mathbf{x}_1)$ and $\mathbf{ex}_2 \subset f(\mathbf{ex}_1)$. Therefore, f induces a bijection from \mathbf{x}_1 to \mathbf{x}_2 . If moreover f satisfies **(CM2)**, then f induces an injection from \mathbf{ex}_1 to \mathbf{ex}_2 and thus it induces a bijection from \mathbf{ex}_1 to \mathbf{ex}_2 . \square

Definition 3.3 (Isomorphic seeds). Two seeds $\Sigma = (\mathbf{x}_1, \mathbf{ex}_1, B^1)$ and $\Sigma_2 = (\mathbf{x}_2, \mathbf{ex}_2, B^2)$ are called *isomorphic* if there exists a bijection $\phi : \mathbf{x}_1 \rightarrow \mathbf{x}_2$ inducing a bijection $\phi : \mathbf{ex}_1 \rightarrow \mathbf{ex}_2$ and such that $b_{\phi(x), \phi(y)}^2 = b_{xy}^1$ for every $x, y \in \mathbf{x}_1$. We then write $\Sigma_1 \simeq \Sigma_2$ and $B^1 \simeq B^2$.

Definition 3.4 (Opposite seed). Given a seed $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$, the *opposite seed* is

$$\Sigma^{\text{op}} = (\mathbf{x}, \mathbf{ex}, -B).$$

Definition 3.5 (Simplification of a seed). Given a seed $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$, we set $\bar{B} = (\bar{b}_{xy})_{x, y \in \mathbf{x}} \in M_{\mathbf{x}}(\mathbb{Z})$ where

$$\bar{b}_{xy} = \begin{cases} 0 & \text{if } x, y \in \mathbf{x} \setminus \mathbf{ex} \\ b_{xy} & \text{otherwise.} \end{cases}$$

The *simplification of the seed* Σ is defined as $\bar{\Sigma} = (\mathbf{x}, \mathbf{ex}, \bar{B})$.

Remark 3.6. In terms of valued quivers, simplifying the seed simply corresponds to removing all the arrows between the frozen points. An example is shown in Figure 1 where the arrows between frozen points are shown dashed.

Definition 3.7 (Locally rooted cluster morphism). Let $\Sigma_1 = (\mathbf{x}_1, \mathbf{ex}_1, B^1)$ and $\Sigma_2 = (\mathbf{x}_2, \mathbf{ex}_2, B^2)$ be two seeds. A ring homomorphism f from $\mathcal{A}(\Sigma_1)$ to $\mathcal{A}(\Sigma_2)$ is called a *locally rooted cluster morphism* if it satisfies **(CM1)**, **(CM2)** and **(CM3loc)**: for any $x \in \mathbf{ex}_1$ and any $y \in \mathbf{x}_1$, we have, $f(\mu_{x, \Sigma_1}(y)) = \mu_{f(x), \Sigma_2}(f(y))$.

As we now prove, for bijective ring homomorphisms, it is possible to simplify considerably the condition **(CM3)** (compare [ASS11, Proposition 2.4]).

Lemma 3.8. *Let Σ_1 and Σ_2 be two seeds and let $f : \mathcal{A}(\Sigma_1) \rightarrow \mathcal{A}(\Sigma_2)$ be a bijective locally rooted cluster morphism. Then :*

- (1) f is a rooted cluster morphism ;
- (2) $\bar{\Sigma}_1 \simeq \bar{\Sigma}_2$ or $\bar{\Sigma}_1 \simeq (\bar{\Sigma}_2)^{\text{op}}$.

Proof. For $i \in \{1, 2\}$, we set $\Sigma_i = (\mathbf{x}_i, \mathbf{ex}_i, B^i)$. Let f be as in the hypothesis. It follows from Corollary 3.2 that f induces bijections from \mathbf{x}_1 to \mathbf{x}_2 and from \mathbf{ex}_1 to \mathbf{ex}_2 .

For every $x \in \mathbf{ex}_1$, we have

$$\begin{aligned} f(\mu_x(x)) &= f\left(\frac{1}{x} \left(\prod_{\substack{z \in \mathbf{x}_1; \\ b_{xz}^1 > 0}} z^{b_{xz}^1} + \prod_{\substack{z \in \mathbf{x}_1; \\ b_{xz}^1 < 0}} z^{-b_{xz}^1} \right)\right) \\ &= \frac{1}{f(x)} \left(\prod_{\substack{z \in \mathbf{x}_1; \\ b_{xz}^1 > 0}} f(z)^{b_{xz}^1} + \prod_{\substack{z \in \mathbf{x}_1; \\ b_{xz}^1 < 0}} f(z)^{-b_{xz}^1} \right). \end{aligned}$$

Because f satisfies **(CM3loc)**, we have

$$\begin{aligned} f(\mu_x(x)) &= \mu_{f(x)}(f(x)) \\ &= \frac{1}{f(x)} \left(\prod_{\substack{f(z) \in \mathbf{ex}_2; \\ b_{f(x)f(z)}^2 > 0}} f(z)^{b_{f(x)f(z)}^2} + \prod_{\substack{f(z) \in \mathbf{ex}_2; \\ b_{f(x)f(z)}^2 < 0}} f(z)^{-b_{f(x)f(z)}^2} \right) \end{aligned}$$

and because \mathbf{x}_2 is algebraically independent, we get $b_{xz}^1 = b_{f(x)f(z)}^2$ for any $x \in \mathbf{ex}_1$ and any $z \in \mathbf{x}_1$ or $b_{xz}^1 = -b_{f(x)f(z)}^2$ for any $x \in \mathbf{ex}_1$ and any $z \in \mathbf{x}_1$, that is, $\overline{B^1} \simeq \overline{B^2}$ or $\overline{B^1} \simeq (-\overline{B^2})$. Therefore, $\overline{\Sigma_1} \simeq \overline{\Sigma_2}$ or $\overline{\Sigma_1} \simeq (\overline{\Sigma_2})^{\text{op}}$.

Since the mutations in $\mathcal{A}(\Sigma_1)$ and $\mathcal{A}(\Sigma_2)$ are entirely encoded in the simplifications $\overline{\Sigma_1}$ and $\overline{\Sigma_2}$ of the exchange matrices, it follows easily that f is a rooted cluster morphism. \square

Theorem 3.9. *Let $\mathcal{A}(\Sigma_1)$ and $\mathcal{A}(\Sigma_2)$ be two rooted cluster algebras. Then $\mathcal{A}(\Sigma_1)$ and $\mathcal{A}(\Sigma_2)$ are isomorphic in **Clus** if and only if $\overline{\Sigma_1} \simeq \overline{\Sigma_2}$ or $\overline{\Sigma_1} \simeq \overline{\Sigma_2}^{\text{op}}$.*

Proof. As **Clus** is a concrete category, an isomorphism $f : \mathcal{A}(\Sigma_1) \rightarrow \mathcal{A}(\Sigma_2)$ is necessarily bijective. Therefore, Corollary 3.2 and Lemma 3.8 imply that f induces a bijection $\mathbf{x}_1 \xrightarrow{\sim} \mathbf{x}_2$ such that $\overline{B^1} \simeq \overline{B^2}$ or $\overline{B^1} \simeq (\overline{B^2})^{\text{op}}$. Moreover, it also follows from Corollary 3.2 that f is a bijection $\mathbf{ex}_1 \xrightarrow{\sim} \mathbf{ex}_2$ so that it induces an isomorphism of seeds $\overline{\Sigma_1} \simeq \overline{\Sigma_2}$ or $\overline{\Sigma_1} \simeq \overline{\Sigma_2}^{\text{op}}$.

Conversely, if $\overline{\Sigma_1} \simeq \overline{\Sigma_2}$ or $\overline{\Sigma_1} \simeq \overline{\Sigma_2}^{\text{op}}$, we consider the bijection $\sigma : \mathbf{x}_1 \rightarrow \mathbf{x}_2$ inducing the isomorphism of seeds. It thus induces naturally a ring isomorphism $f_\sigma : \mathcal{F}_{\Sigma_1} \rightarrow \mathcal{F}_{\Sigma_2}$ and it is easily seen that f_σ is a rooted cluster isomorphism. \square

Corollary 3.10. *The isomorphisms in **Clus** coincide with the bijective rooted cluster morphisms.*

Proof. As **Clus** is a concrete category, isomorphisms are bijective. Conversely, if we consider a bijective rooted cluster morphism $\mathcal{A}(\Sigma_1) \rightarrow \mathcal{A}(\Sigma_2)$, then it follows from Lemma 3.8 that f induces an isomorphism of seeds $\overline{\Sigma_1} \simeq \overline{\Sigma_2}$ or $\overline{\Sigma_1} \simeq \overline{\Sigma_2}^{\text{op}}$ so that f is an isomorphism in **Clus**. \square

Corollary 3.11. *Let $\mathcal{A}(\Sigma_1)$ and $\mathcal{A}(\Sigma_2)$ be two rooted cluster algebras and let $f : \mathcal{A}(\Sigma_1) \rightarrow \mathcal{A}(\Sigma_2)$ be a rooted cluster isomorphism. Then the following hold :*

- (1) any Σ_1 -admissible sequence is (f, Σ_1, Σ_2) -biadmissible ;
- (2) any Σ_2 -admissible sequence lifts to a unique (f, Σ_1, Σ_2) -biadmissible sequence ;
- (3) $f(\mathcal{X}_{\Sigma_1}) = \mathcal{X}_{\Sigma_2}$. \square

Remark 3.12. For non-bijective morphisms, one can find locally rooted cluster morphisms which are not rooted cluster morphisms. For instance, let $\Sigma_1 = (\mathbf{x}_1, \mathbf{ex}_1, B^1)$ and $\Sigma_2 = (\mathbf{x}_2, \mathbf{ex}_2, B^2)$ where

$$\mathbf{ex}_1 = \mathbf{x}_1 = \{x_1, x_2, x_3\}, \quad \mathbf{ex}_2 = \mathbf{x}_2 = \{u_1, u_2\},$$

$$B^1 = \begin{bmatrix} 0 & 1 & 0 \\ -1 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix} \text{ and } B^2 = \begin{bmatrix} 0 & 1 \\ -2 & 0 \end{bmatrix}$$

and consider the ring homomorphism

$$\pi : \begin{cases} \mathbb{Q}(x_1, x_2, x_3) & \longrightarrow & \mathbb{Q}(u_1, u_2) \\ x_1, x_3 & \mapsto & u_1 \\ x_2 & \mapsto & u_2. \end{cases}$$

Then

$$\pi(\mu_{x_1}(x_1)) = \pi(\mu_{x_3}(x_3)) = \pi\left(\frac{1+x_2}{x_1}\right) = \frac{1+u_2}{u_1} = \mu_{u_1}(u_1)$$

and

$$\pi(\mu_{x_2}(x_2)) = \pi\left(\frac{1+x_1x_3}{x_2}\right) = \frac{1+u_1^2}{u_2} = \mu_{u_2}(u_2)$$

so that π commutes with biadmissible sequences of length one.

However

$$\mu_{x_2}(\mathbf{x}_1) = \left\{x_1, x'_2 = \frac{1+x_1x_3}{x_2}, x_3\right\}$$

$$(\mu_{x_1} \circ \mu_{x_2})(\mathbf{x}_1) = \left\{\frac{1+x_2+x_1x_3}{x_1x_2}, x'_2, x_3\right\}$$

and

$$(\mu_{x'_2} \circ \mu_{x_1} \circ \mu_{x_2})(\mathbf{x}_1) = \left\{\frac{1+x_2+x_1x_3}{x_1x_2}, \frac{1+x_2}{x_1}, x_3\right\}$$

but

$$\mu_{u_2}(\mathbf{x}_2) = \left\{u_1, u'_2 = \frac{1+u_1^2}{u_2}\right\}$$

$$(\mu_{u_1} \circ \mu_{u_2})(\mathbf{x}_2) = \left\{\frac{1+u_2+u_1^2}{u_1u_2}, u'_2\right\}$$

and

$$(\mu_{u'_2} \circ \mu_{u_1} \circ \mu_{u_2})(\mathbf{x}_2) = \left\{\frac{1+u_2+u_1^2}{u_1u_2}, \frac{1+2u_2+u_2^2+u_1^2}{u_1^2u_2}\right\}.$$

so that

$$\pi\left(\frac{1+x_2}{x_1}\right) = \frac{1+u_2}{u_1} \neq \frac{1+2u_2+u_2^2+u_1^2}{u_1^2u_2}$$

and thus π is not a rooted cluster morphism between the cluster algebras $\mathcal{A}(\Sigma_1)$ and $\mathcal{A}(\Sigma_2)$.

Note that the morphism π is induced by the folding of the quiver $Q_{B^1} : 1 \longleftarrow 2 \longrightarrow 3$ with respect to the automorphism group exchanging 1 and 3. For general results concerning the interactions of foldings with cluster algebras, we refer the reader for instance to [Dem11].

Remark 3.13. a) Two rooted cluster algebras associated with mutation-equivalent seeds are *not* necessarily isomorphic in the category **Clus** since mutation-equivalent seeds are in general neither isomorphic nor opposite.

b) The *cluster automorphisms* considered in [ASS11] correspond in our context to rooted cluster isomorphisms from $\mathcal{A}(\Sigma)$ to itself when Σ is finite, skew-symmetric and coefficient-free. The groups of cluster automorphisms have been computed for seeds associated with Dynkin or affine quivers, see [ASS11, §3.3].

c) A strong isomorphism $\mathcal{A}(\Sigma_1) \rightarrow \mathcal{A}(\Sigma_2)$ in the sense of [FZ03] is a rooted cluster isomorphism such that $\overline{\Sigma_1} \simeq \overline{\Sigma_2}$.

4. ROOTED CLUSTER MONOMORPHISMS

We recall that a *monomorphism* in a category is a morphism f such that if there exist morphisms g and h such that $fg = fh$, then $g = h$.

Lemma 4.1. *Let $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ be a seed, let $\mathbf{y} \subset \mathbf{x}$ and let $\Theta = (\mathbf{y}, \emptyset, C)$ be another seed. Then the canonical ring homomorphism $\mathcal{F}_\Theta \rightarrow \mathcal{F}_\Sigma$ induces an injective rooted cluster morphism $\mathcal{A}(\Theta) \rightarrow \mathcal{A}(\Sigma)$.*

Proof. The canonical ring homomorphism $\mathcal{F}_\Theta \rightarrow \mathcal{F}_\Sigma$ sends x to x for any $x \in \mathbf{y}$ therefore, it satisfies **(CM1)** and **(CM2)**. Moreover, since there are no exchangeable variables in Θ , it automatically satisfies **(CM3)**. We thus only have to prove that it induces a ring homomorphism $\mathcal{A}(\Theta) \rightarrow \mathcal{A}(\Sigma)$, and this is clear because $\mathcal{A}(\Theta) = \mathbb{Z}[x \mid x \in \mathbf{y}] \subset \mathbb{Z}[x \mid x \in \mathbf{x}] \subset \mathcal{A}(\Sigma)$. \square

Remark 4.2. In order for Lemma 4.1 to hold, it is necessary to consider non-invertible coefficients because if the image of a frozen variable is exchangeable then the image of its inverse would have to be the inverse of the exchangeable variable, which is not in the cluster algebra.

Proposition 4.3. *Monomorphisms in **Clus** coincide with injective rooted cluster morphisms.*

Proof. Let $\Sigma_i = (\mathbf{x}_i, \mathbf{ex}_i, B^i)$ be seeds for $1 \leq i \leq 3$ and consider rooted cluster morphisms

$$\mathcal{A}(\Sigma_1) \begin{array}{c} \xrightarrow{g} \\ \xrightarrow{h} \end{array} \mathcal{A}(\Sigma_2) \xrightarrow{f} \mathcal{A}(\Sigma_3).$$

Since **Clus** is a concrete category, every injective morphism in **Clus** is a monomorphism. We thus only need to prove the converse.

Let f be a non-injective rooted cluster morphism. Because it satisfies **(CM1)**, we get $f(\mathbf{x}_2) \subset \mathbf{x}_3 \sqcup \mathbb{Z}$. If $f(\mathbf{x}_2) \subset \mathbf{x}_3$ and if the restriction of f to \mathbf{x}_2 is injective, f sends a transcendence basis of \mathcal{F}_{Σ_2} to an algebraically independent family in \mathcal{F}_{Σ_3} and therefore, it induces an injective ring homomorphism $\mathcal{F}_{\Sigma_2} \rightarrow \mathcal{F}_{\Sigma_3}$ so that it is itself injective, a contradiction. Thus, there are two cases to consider :

- there exists $x \in \mathbf{x}_2$ such that $f(x) \in \mathbb{Z}$,
- there exist $x, y \in \mathbf{x}_2$ such that $x \neq y$ and $f(x) = f(y)$.

In the first case, it follows from Lemma 4.1 that we can consider the rooted cluster morphisms $h, g : \mathbb{Z}[x] \rightarrow \mathcal{A}(\Sigma_2)$ given by $g(x) = x$ and $h(x) = f(x) \in \mathbb{Z}$. Then $fh(x) = fg(x) = f(x)$ so that $fh = fg$ but $f \neq g$. Thus f is not a monomorphism in **Clus**.

In the second case, it also follows from Lemma 4.1 that we can consider the rooted cluster morphisms $h, g : \mathbb{Z}[x, y] \rightarrow \mathcal{A}(\Sigma_2)$ given by $g(x) = h(y) = x$ and $g(y) = h(x) = y$. Thus, $fg = fh$ but $g \neq h$ and therefore f is not a monomorphism in **Clus**. \square

As a consequence, the study of monomorphisms in **Clus** restricts to the study of injective rooted cluster morphisms.

Lemma 4.4. *Let $\Sigma_1 = (\mathbf{x}_1, \mathbf{ex}_1, B_1)$ and $\Sigma_2 = (\mathbf{x}_2, \mathbf{ex}_2, B^2)$ be two seeds and let $f : \mathcal{A}(\Sigma_1) \rightarrow \mathcal{A}(\Sigma_2)$ be an injective rooted cluster morphism. Then f induces an isomorphism of seeds $\overline{\Sigma_1} \simeq \overline{f(\Sigma_1)}$ or $\overline{\Sigma_1} \simeq (\overline{f(\Sigma_1)})^{\text{op}}$.*

Proof. As f is injective and satisfies **(CM1)**, we have $f(\mathbf{x}_1) \subset \mathbf{x}_2$ and since it satisfies **(CM2)**, we have $f(\mathbf{ex}_1) = \mathbf{ex}_2 \cap f(\mathbf{x}_1)$. Let $x \in \mathbf{ex}_1$. Since f satisfies **(CM3)**, we have $f(\mu_{x, \Sigma_1}(x)) =$

$\mu_{f(x), \Sigma_2}(f(x))$ so that

$$\prod_{\substack{y \in \mathbf{x}_1, \\ b_{xy}^1 > 0}} f(y)^{b_{xy}^1} + \prod_{\substack{y \in \mathbf{x}_1, \\ b_{xy}^1 < 0}} f(y)^{-b_{xy}^1} = \prod_{\substack{z \in \mathbf{x}_2, \\ b_{f(x)z}^2 > 0}} z^{b_{f(x)z}^2} + \prod_{\substack{z \in \mathbf{x}_2, \\ b_{f(x)z}^2 < 0}} z^{-b_{f(x)z}^2}.$$

Therefore, $b_{f(x)z}^2 = 0$ for any $z \notin f(\mathbf{x}_1)$ and either $b_{f(x)z}^2 = b_{xy}^1$ for any $y \in \mathbf{x}_1$ such that $f(y) = z$ or $b_{f(x)z}^2 = -b_{xy}^1$ for any $y \in \mathbf{x}_1$ such that $f(y) = z$, which proves the lemma. \square

Corollary 4.5. *Any injective rooted cluster morphism is ideal.*

Proof. Let $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ and $\Sigma' = (\mathbf{x}', \mathbf{ex}', B')$ be two seeds and let $f : \mathcal{A}(\Sigma) \rightarrow \mathcal{A}(\Sigma')$ be an injective rooted cluster morphism. As a cluster algebra does only depend on the simplification of the seed, we can assume that both $\Sigma = \overline{\Sigma}$ and $\Sigma' = \overline{\Sigma}'$. According to Lemma 4.4, f induces an isomorphism of seeds $\Sigma \simeq f(\Sigma)$ or $\Sigma \simeq \overline{f(\Sigma)}$. It follows that every Σ -admissible sequence is (f, Σ, Σ') -biadmissible.

In order to prove that f is ideal, it is enough to prove that for any cluster variable x in $\mathcal{A}(\Sigma)$, the variable $f(x)$ is an element in $\mathcal{A}(f(\Sigma))$. Let thus z be a cluster variable in $\mathcal{A}(\Sigma)$. Then either z is a frozen variable in Σ and thus $f(z)$ is a frozen variable in $f(\Sigma)$ and we are done, or there exists a Σ -admissible sequence (x_1, \dots, x_l) such that $z = \mu_{x_l} \circ \dots \circ \mu_{x_1}(x)$ for some $x \in \mathbf{ex}$. Then (x_1, \dots, x_l) is (f, Σ, Σ') -biadmissible and because f satisfies **(CM3)**, we get

$$f(z) = f(\mu_{x_l} \circ \dots \circ \mu_{x_1, \Sigma}(x)) = \mu_{f(x_l)} \circ \dots \circ \mu_{f(x_1), \Sigma'}(f(x))$$

but it follows from Lemma 2.9 that

$$\mu_{f(x_l)} \circ \dots \circ \mu_{f(x_1), \Sigma'}(f(x)) = \mu_{f(x_l)} \circ \dots \circ \mu_{f(x_1), f(\Sigma)}(f(x)).$$

Therefore, $f(z)$ is a cluster variable in $\mathcal{A}(f(\Sigma))$ and thus $f(\mathcal{A}(\Sigma)) \subset \mathcal{A}(f(\Sigma))$. The reverse inclusion follows from Lemma 2.10 and therefore f is ideal. \square

As a byproduct of the proof of Corollary 4.5, we obtain :

Corollary 4.6. *Let Σ_1, Σ_2 be two seeds and $f : \mathcal{A}(\Sigma_1) \rightarrow \mathcal{A}(\Sigma_2)$ be an injective rooted cluster morphism. Then :*

- (1) *any Σ_1 -admissible sequence is (f, Σ_1, Σ_2) -biadmissible,*
- (2) *$f(\mathcal{X}_{\Sigma_1}) \subset \mathcal{X}_{\Sigma_2}$.* \square

We recall that a seed $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ is called :

- of *finite cluster type* if \mathcal{X}_{Σ} is finite,
- *acyclic* if the valued quiver obtained by deleting the arrows between frozen vertices in Q_B has no oriented cycles,
- *mutation-finite* if the mutation class of the exchange matrix of B is finite.


Corollary 4.7. *Let Σ_1, Σ_2 be two seeds and $f : \mathcal{A}(\Sigma_1) \rightarrow \mathcal{A}(\Sigma_2)$ be an injective rooted cluster morphism. Then :*

- (1) *If Σ_2 is of finite cluster type, then so is Σ_1 ;*
- (2) *If Σ_2 is acyclic, then so is Σ_1 ;*
- (3) *If Σ_2 is mutation-finite, then so is Σ_1 .*

Proof. The first assertion is a consequence of Corollary 4.6. The second assertion follows from Lemma 4.4 and from the fact that a subquiver of an acyclic (valued) quiver is acyclic. The third assertion follows from Lemma 4.4 and from the fact that a full subquiver of a mutation-finite (valued) quiver is mutation-finite. \square

4.1. Monomorphisms arising from triangulations of the n -gon. For any integer $m \geq 3$, we denote by Π_m the m -gon whose points are labelled cyclically from 1 to m . For $m \geq 4$, the cluster algebra $\mathcal{A}(\Pi_m)$ (with coefficients associated with boundary arcs) is a cluster algebra of type A_{m-3} .

We construct by induction a family $\{T_m\}_{m \geq 3}$ where each T_m is a *fan triangulation* of Π_m . We start with the triangle Π_3 whose points are denoted by 1,2 and 3. For any $m \geq 3$, the triangulation T_{m+1} of Π_{m+1} is obtained by gluing a triangle along the boundary arc joining m to 1 in T_m and the new marked point introduced by this triangle is labelled by $m+1$, as shown in the figure below.



For any $m \geq 4$, we denote by Σ_m the seed associated with T_m in $\mathcal{A}(\Pi_m)$. Cluster variables in $\mathcal{A}(\Pi_m)$ are identified with the arcs joining two marked points in Π_m and for any i and j such that $1 \leq i < j \leq m$, we denote by x_{ij} the variable corresponding to the arc joining i to j . The exchangeable variables are thus the variables corresponding to internal arcs. The exchange relations given by the mutations in $\mathcal{A}(\Pi_m)$ are the so-called *Plücker relations* :


$$x_{ij}x_{kl} = x_{ik}x_{jl} + x_{il}x_{kj} \text{ for } 1 \leq i < j < k < l \leq m.$$

For any $m' \geq m$, the inclusion of T_m in $T_{m'}$ defines a natural ring monomorphism $j_{m,m'} : \mathcal{F}_{\Sigma_m} \rightarrow \mathcal{F}_{\Sigma_{m'}}$. Since arcs (or internal arcs) in T_m are sent to arcs (or internal arcs, respectively) in $T_{m'}$, then $j_{m,m'}$ satisfies **(CM1)** and **(CM2)**. Moreover, since exchange relations in $\mathcal{A}(\Pi_m)$ and $\mathcal{A}(\Pi_{m'})$ correspond to Plücker relations in Π_m and $\Pi_{m'}$ respectively, it is easily seen that $j_{m,m'}$ satisfies **(CM3)**. Finally, since every admissible sequence of variables in $\mathcal{A}(\Pi_m)$ is $(j_{m,m'}, \Sigma_m, \Sigma_{m'})$ -biadmissible, it follows from the fact that $j_{m,m'}$ preserves the Plücker relations that it commutes with biadmissible mutations and that $j_{m,m'}(\mathcal{A}(\Pi_m)) \subset \mathcal{A}(\Pi_{m'})$.

Therefore, for $m < m'$, we have exhibited an injective rooted cluster morphism

$$j_{m,m'} : \mathcal{A}(\Pi_m) \rightarrow \mathcal{A}(\Pi_{m'}).$$


Example 4.8. Consider the cluster algebra $\mathcal{A}(\Pi_4)$ whose cluster variables are shown on the square below.



Then

$$\mathcal{A}(\Pi_4) = \mathbb{Z}[x_{ij} \mid 1 \leq i < j \leq 4] / (x_{13}x_{24} = x_{12}x_{34} + x_{14}x_{23}).$$

Now consider the cluster algebra $\mathcal{A}(\Pi_5)$. It has 4 additional cluster variables, two are exchangeable and two are frozen. We show these new variables on the picture below.



Then

$$\mathcal{A}(\Pi_5) = \mathbb{Z}[x_{ij} \mid 1 \leq i < j \leq 5] / (x_{ij}x_{kl} = x_{ik}x_{jl} + x_{il}x_{kj} \text{ for } 1 \leq i < j < k < l \leq 5).$$

In particular, the canonical morphism $j_{4,5} : \mathbb{Q}(x_{ij} \mid 1 \leq i < j \leq 4) \rightarrow \mathbb{Q}(x_{ij} \mid 1 \leq i < j \leq 5)$ induces an injective ring homomorphism $\mathcal{A}(\Pi_4) \rightarrow \mathcal{A}(\Pi_5)$ which is a rooted cluster monomorphism $\mathcal{A}(\Sigma_{T_4}) \rightarrow \mathcal{A}(\Sigma_{T_5})$.

Definition 4.9 (Full subseed of a seed). Given a seed $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ where $\mathbf{x} = (x_i, i \in I)$ and given a subset $J \subset I$, we set $\Sigma|_J$ the seed with cluster $\mathbf{x}|_J = (x_i, i \in J)$, with exchangeable variables $\mathbf{ex}|_J = \mathbf{ex} \cap \mathbf{x}|_J$ and exchange matrix $B[J]$. Such a seed is called a *full subseed* of Σ .

Remark 4.10. If Σ' is a full subseed of Σ , the canonical morphism $\mathcal{F}_{\Sigma'} \rightarrow \mathcal{F}_{\Sigma}$ does not necessarily induce an injective rooted cluster morphism $\mathcal{A}(\Sigma') \rightarrow \mathcal{A}(\Sigma)$. For instance, if Σ_m denotes the coefficient-free seed associated with the quiver

$$Q_m : 1 \rightarrow 2 \rightarrow \dots \rightarrow m$$

for any $m \geq 1$, then the canonical inclusion $\iota : \mathcal{F}_{\Sigma_m} \rightarrow \mathcal{F}_{\Sigma_{m+1}}$ does not induce a rooted cluster morphism $\mathcal{A}(\Sigma_m) \rightarrow \mathcal{A}(\Sigma_{m+1})$ because

$$\iota(\mu_{x_m, \Sigma_m}(x_m)) = \iota\left(\frac{1+x_{m-1}}{x_m}\right) \neq \frac{x_{m-1}+x_{m+1}}{x_m} = \mu_{x_m, \Sigma_{m+1}}(x_m).$$

However, as we shall see in Proposition 6.4, if Σ' is a full subseed of Σ , then there is a natural ideal surjective rooted cluster morphism $\mathcal{A}(\Sigma) \rightarrow \mathcal{A}(\Sigma')$.

We now describe a combinatorial operation on seeds which allows one to construct a class of injective rooted cluster morphisms.

4.2. Amalgamated sum of seeds. Let $\Sigma_1 = (\mathbf{x}_1, \mathbf{ex}_1, B^1)$ and $\Sigma_2 = (\mathbf{x}_2, \mathbf{ex}_2, B^2)$ be two seeds and $\mathcal{A}(\Sigma_1), \mathcal{A}(\Sigma_2)$ be the corresponding rooted cluster algebras.

Let $\Delta^1 \subset (\mathbf{x}_1 \setminus \mathbf{ex}_1)$ and $\Delta^2 \subset (\mathbf{x}_2 \setminus \mathbf{ex}_2)$ be two (possibly empty) subsets such that there is an isomorphism of seeds $\Sigma_1|_{\Delta^1} \simeq \Sigma_2|_{\Delta^2}$. In this case, we say that Σ_1 and Σ_2 are *glueable along Δ_1 and Δ_2* .

Let Δ be a family of undeterminates in bijection with Δ_1 and Δ_2 . We set

$$\mathbf{x}_1 \coprod_{\Delta_1, \Delta_2} \mathbf{x}_2 = (\mathbf{x}_1 \setminus \Delta_1) \sqcup (\mathbf{x}_2 \setminus \Delta_2) \sqcup \Delta.$$

As Δ_1 and Δ_2 consist of frozen variables, \mathbf{ex}_1 and \mathbf{ex}_2 are naturally identified with two disjoint subsets of $\mathbf{x}_1 \coprod_{\Delta_1, \Delta_2} \mathbf{x}_2$ and we set

$$\mathbf{ex}_1 \coprod_{\Delta_1, \Delta_2} \mathbf{ex}_2 = \mathbf{ex}_1 \sqcup \mathbf{ex}_2.$$

With respect to the partitions $\mathbf{x}_i = (\mathbf{x}_i \setminus \Delta_i) \sqcup \Delta$, for any $i \in \{1, 2\}$ the matrix B^i can be written :

$$B^i = \left[\begin{array}{c|c} B_{11}^i & B_{12}^i \\ \hline B_{21}^i & B_{\Delta} \end{array} \right]$$

where the matrices are possibly infinite.

We then set

$$B^1 \coprod_{\Delta_1, \Delta_2} B^2 = \left[\begin{array}{c|c|c} B_{11}^1 & 0 & B_{12}^1 \\ \hline 0 & B_{11}^2 & B_{12}^2 \\ \hline B_{21}^1 & B_{12}^2 & B_{\Delta} \end{array} \right]$$


Definition 4.11 (Amalgamated sum of seeds). With the above notations, the *amalgamated sum* of $\mathcal{A}(\Sigma_1)$ and $\mathcal{A}(\Sigma_2)$ along Δ_1, Δ_2 is the rooted cluster algebra $\mathcal{A}(\Sigma)$ where $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ with :

- (1) $\mathbf{x} = \mathbf{x}_1 \coprod_{\Delta_1, \Delta_2} \mathbf{x}_2$;
- (2) $\mathbf{ex} = \mathbf{ex}_1 \coprod_{\Delta_1, \Delta_2} \mathbf{ex}_2$;
- (3) $B = B^1 \coprod_{\Delta_1, \Delta_2} B^2$.

We use the notations

$$\Sigma = \Sigma_1 \coprod_{\Delta_1, \Delta_2} \Sigma_2 \text{ and } \mathcal{A}(\Sigma) = \mathcal{A}(\Sigma_1) \coprod_{\Delta_1, \Delta_2} \mathcal{A}(\Sigma_2).$$

Remark 4.12. In terms of valued quivers, the amalgamated sum of exchange matrices corresponds to the amalgamated sum of valued quivers. For instance, the following figure shows an example of amalgamated sum over the subquivers in the shaded area where points corresponding to exchangeable (or frozen) variables are black (or white, respectively).



Lemma 4.13. Let Σ_1 and Σ_2 be two seeds which are glueable along subsets Δ_1 and Δ_2 as above and let $\Sigma = \Sigma_1 \coprod_{\Delta_1, \Delta_2} \Sigma_2$. Then for any i such that $i \in \{1, 2\}$, the morphism $\mathcal{F}_{\Sigma_i} \rightarrow \mathcal{F}_{\Sigma}$ induced by the inclusion induces an injective rooted cluster morphism $\mathcal{A}(\Sigma_i) \rightarrow \mathcal{A}(\Sigma)$.

Proof. By construction of Σ , the canonical morphism $j_i : \mathcal{F}_{\Sigma_i} \rightarrow \mathcal{F}_{\Sigma}$ is injective and satisfies **(CM1)** and **(CM2)**. We now prove by induction on l that any Σ_1 -admissible sequence of length l is (j_1, Σ_1, Σ) -biadmissible and that j_1 commutes with mutations along biadmissible sequences of length l .

Let $x \in \mathbf{ex}_1$. Then $j_1(x) = x \in \mathbf{ex}$ so that (x) is (j_1, Σ_1, Σ) -biadmissible and therefore mutating in Σ_1 gives

$$\mu_x(x) = \prod_{\substack{y \in \mathbf{x}_1 ; \\ b_{xy}^1 > 0}} y^{b_{xy}^1} + \prod_{\substack{y \in \mathbf{x}_1 ; \\ b_{xy}^1 < 0}} y^{-b_{xy}^1}$$

and, since $b_{zy} = 0$ for any $z \in (\mathbf{x}_1 \setminus \Delta) \supset \mathbf{ex}_1$ and any $y \in \mathbf{x}_2 \setminus \Delta$, mutating in Σ gives

$$\begin{aligned} \mu_{x,\Sigma}(x) &= \prod_{\substack{y \in \mathbf{x} \\ b_{xy} > 0}} y^{b_{xy}} + \prod_{\substack{y \in \mathbf{x} \\ b_{xy} < 0}} y^{-b_{xy}} \\ &= \prod_{\substack{y \in (\mathbf{x}_1 \setminus \Delta_1) \sqcup \Delta \\ b_{xy} > 0}} y^{b_{xy}} + \prod_{\substack{y \in (\mathbf{x}_1 \setminus \Delta_1) \sqcup \Delta \\ b_{xy} < 0}} y^{-b_{xy}} \\ &= \prod_{\substack{y \in \mathbf{x}_1 \\ b_{xy}^1 > 0}} y^{b_{xy}^1} + \prod_{\substack{y \in \mathbf{x}_1 \\ b_{xy}^1 < 0}} y^{-b_{xy}^1} \\ &= j_1(\mu_{x,\Sigma_1}(x)) \end{aligned}$$

and thus j_1 is a locally rooted cluster morphism.

Assume now that we proved the claim for any $k < l$ and let $\Sigma^{(k)} = \mu_{x_k} \circ \dots \circ \mu_{x_1}(\Sigma)$ and $\Sigma_1^{(k)} = \mu_{x_k} \circ \dots \circ \mu_{x_1}(\Sigma_1)$. We denote by $B^{(k)} = (b_{xy}^{(k)})$ (or $B^{1,(k)} = (b_{xy}^{1,(k)})$) the exchange matrix of $\Sigma^{(k)}$ (or $\Sigma_1^{(k)}$, respectively). By the induction hypothesis, the cluster $\mathbf{x}^{(k)}$ of $\Sigma^{(k)}$ is $\mathbf{x}_1^{(k)} \sqcup (\mathbf{x}_2 \setminus \Delta) \sqcup \Delta$ where $\mathbf{x}_1^{(k)} \sqcup \Delta_1$ is the cluster of $\Sigma_1^{(k)}$. Then an easy induction proves that

$$b_{xy}^{(l)} = \begin{cases} 0 & \text{if } x \in \mathbf{x}_1^{(k)} \text{ and } y \in \mathbf{x}_2 \setminus \Delta \\ b_{xy}^{1,(l)} & \text{if } x, z \in \mathbf{x}_1^{(k)} \sqcup \Delta. \end{cases}$$

Because j_1 commutes with sequences of biadmissible mutations of length k , any variable x_{k+1} exchangeable in $\Sigma_1^{(k)}$ is also exchangeable in $\Sigma^{(k)}$ and a similar calculation proves that for any such exchangeable variable x_{k+1} , the morphism j_1 commutes with $\mu_{x_{k+1}} \circ \dots \circ \mu_{x_1}$. \square

Remark 4.14. If Δ_1 and Δ_2 do not consist of frozen variables, then the canonical morphism $\mathcal{F}_{\Sigma_i} \rightarrow \mathcal{F}_{\Sigma}$ may not satisfy **(CM3)**. Indeed, if one considers the seeds

$$\Sigma_1 = \left((x_1, x_2), (x_1, x_2), \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \right)$$

$$\Sigma_2 = \left((x_2, x_3), (x_3), \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \right)$$

so that

$$\Sigma = \Sigma_1 \coprod_{x_2, x_2} \Sigma_2 = \left((x_1, x_3, x_2), (x_1, x_3, x_2), \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & -1 \\ -1 & 1 & 0 \end{bmatrix} \right).$$

Then mutating along x_2 in Σ_1 gives $\frac{1+x_1}{x_2}$ whereas mutating along x_2 in Σ gives $\frac{x_1+x_3}{x_2}$. Therefore, the canonical morphism $\mathcal{F}_{\Sigma_1} \rightarrow \mathcal{F}_{\Sigma}$ does not satisfy **(CM3)**.

If $\Sigma_1 = (\mathbf{x}_1, \mathbf{ex}_1, B^1)$ and $\Sigma_2 = (\mathbf{x}_2, \mathbf{ex}_2, B^2)$ are glueable along Δ_1, Δ_2 and if we denote by Δ the common image of Δ_1 and Δ_2 in $\Sigma = \Sigma_1 \coprod_{\Delta_1, \Delta_2} \Sigma_2 = (\mathbf{x}, \mathbf{ex}, B)$, then the compositions of the canonical ring homomorphisms

$$\mathbb{Z}[\Delta] \simeq \mathbb{Z}[\Delta_1] \rightarrow \mathcal{A}(\Sigma_1) \text{ and } \mathbb{Z}[\Delta] \simeq \mathbb{Z}[\Delta_2] \rightarrow \mathcal{A}(\Sigma_2)$$

induce $\mathbb{Z}[\Delta]$ -algebra structures on $\mathcal{A}(\Sigma_1)$ and $\mathcal{A}(\Sigma_2)$. Also the inclusion $\mathbb{Z}[\Delta] \subset \mathcal{A}(\Sigma)$ induces a $\mathbb{Z}[\Delta]$ -algebra structure on $\mathcal{A}(\Sigma)$. Then we have the following proposition :

Proposition 4.15. *The canonical ring isomorphism $\mathbb{Q}(\mathbf{x}) \xrightarrow{\sim} \mathbb{Q}(\mathbf{x}_1) \otimes_{\mathbb{Q}(\Delta)} \mathbb{Q}(\mathbf{x}_2)$ induces an isomorphism of $\mathbb{Z}[\Delta]$ -algebras :*

$$\mathcal{A} \left(\Sigma_1 \coprod_{\Delta_1, \Delta_2} \Sigma_2 \right) \simeq \mathcal{A}(\Sigma_1) \otimes_{\mathbb{Z}[\Delta]} \mathcal{A}(\Sigma_2).$$

□

5. COPRODUCTS, PRODUCTS AND AMALGAMATED SUMS

5.1. Coproducts of rooted cluster algebras.

Lemma 5.1. *The category **Clus** admits countable coproducts.*

Proof. Let I be a countable set and let $\{\mathcal{A}(\Sigma_i)\}_{i \in I}$ be a countable family of rooted cluster algebras. For any $i \in I$, we set $\Sigma^i = (\mathbf{x}_i, \mathbf{ex}_i, B^i)$ and

$$\Sigma = \coprod_{i \in I} \Sigma_i = (\mathbf{x}, \mathbf{ex}, B)$$

where

$$\mathbf{x} = \bigsqcup_{i \in I} \mathbf{x}_i, \quad \mathbf{ex} = \bigsqcup_{i \in I} \mathbf{ex}_i,$$

and B is the block-diagonal matrix whose blocks are indexed by I and such that for any $i \in I$, the i -th diagonal block is B^i . Then B is locally finite and thus Σ is a well-defined seed. For any $i \in I$, we denote by j_i the canonical inclusion $\mathcal{F}_{\Sigma_i} \rightarrow \mathcal{F}_{\Sigma}$, which clearly induces a rooted cluster morphism $\mathcal{A}(\Sigma_i) \rightarrow \mathcal{A}(\Sigma)$.

Now assume that there exists a rooted cluster algebra $\mathcal{A}(\Theta)$ and for any $i \in I$ a rooted cluster morphism $g_i : \mathcal{A}(\Sigma_i) \rightarrow \mathcal{A}(\Theta)$. In order to prove that $\mathcal{A}(\Sigma)$ is the coproduct of the $\mathcal{A}(\Sigma_i)$ with $i \in I$, we need to prove that there exists a unique rooted cluster morphism $h : \mathcal{A}(\Sigma) \rightarrow \mathcal{A}(\Theta)$ such that $h \circ j_i = g_i$ for any $i \in I$. It is easily seen that such a rooted cluster morphism h exists if and only if there exists a ring homomorphism $h : \mathcal{F}_{\Sigma} \rightarrow \mathcal{F}_{\Theta}$ satisfying $h(x) = g_i(x)$ for any $i \in I$. This latter condition defines precisely one ring homomorphism $h : \mathcal{F}_{\Sigma} \rightarrow \mathcal{F}_{\Theta}$ and therefore, h exists and is unique. □

Corollary 5.2. *The full subcategory of **Clus** formed by rooted cluster algebras associated with finite seeds has finite coproducts.*

Remark 5.3. Topologically, this can be interpreted by saying that if (S, M) is a marked surface, with connected components S_1, \dots, S_n and with $M_i = M \cap S_i$ for any $1 \leq i \leq n$, then $\mathcal{A}(S, M)$ is the coproduct of the $\mathcal{A}(S_i, M_i)$ in **Clus**. Indeed, for any $1 \leq i \leq n$, let T_i be a triangulation of (S_i, M_i) . Then $T = \bigsqcup_i T_i$ is a triangulation of (S, M) and it follows immediately from the definitions that $\Sigma_T = \coprod_{i=1}^n \Sigma_{T_i}$ so that $\mathcal{A}(\Sigma_T) = \coprod_{i=1}^n \mathcal{A}(\Sigma_{T_i})$.

5.2. Products of rooted cluster algebras.

Proposition 5.4. ***Clus** does not generally admit products.*

Proof. The proof consists of the construction of an example of two rooted cluster algebras whose product is not defined in **Clus**. We consider the rooted cluster algebras associated with the seeds $\Sigma_1 = ((t_1), \emptyset, [0])$ and $\Sigma_2 = ((t_2), \emptyset, [0])$, so that $\mathcal{A}(\Sigma_1) = \mathbb{Z}[t_1]$ and $\mathcal{A}(\Sigma_2) = \mathbb{Z}[t_2]$. Assume that

there exists a product in **Clus**

$$\begin{array}{ccc} & \mathcal{A}(\Sigma) & \\ p_1 \swarrow & & \searrow p_2 \\ \mathbb{Z}[t_1] & & \mathbb{Z}[t_2]. \end{array}$$

with $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$. For any $i \in \{1, 2\}$, the morphism p_i is a rooted cluster morphism so that

$$p_i(\mathbf{x}) \subset \{t_i\} \sqcup \mathbb{Z} \text{ and } p_i(\mathbf{ex}) \subset \mathbb{Z}.$$

Consider the seed $\Sigma' = ((x), \emptyset, [0])$ and for any $i \in \{1, 2\}$, let $f_i : \mathbb{Z}[x] \rightarrow \mathbb{Z}[t_i]$ be the ring homomorphism sending x to t_i . Then f_i is a rooted cluster morphism $\mathcal{A}(\Sigma') = \mathbb{Z}[x] \rightarrow \mathcal{A}(\Sigma)$ for any $i \in \{1, 2\}$.

By definition of the product, there exists a unique rooted cluster morphism $h : \mathcal{A}(\Sigma') \rightarrow \mathcal{A}(\Sigma)$ such that the following diagram commutes:

$$\begin{array}{ccc} & \mathcal{A}(\Sigma') & \\ & \downarrow h & \\ f_1 \swarrow & \mathcal{A}(\Sigma) & \searrow f_2 \\ & \downarrow p_1 \quad \downarrow p_2 & \\ \mathbb{Z}[t_1] & & \mathbb{Z}[t_2]. \end{array}$$

In particular, for any $i \in \{1, 2\}$, there exists $x_i \in \mathbf{x}$ such that $p_i(x_i) = t_i$.

Now consider the seed $\Sigma' = ((v_1, v_2), \emptyset, [0])$ so that $\mathcal{A}(\Sigma') = \mathbb{Z}[v_1, v_2]$. For any $i \in \{1, 2\}$, let $f_i : \mathcal{A}(\Sigma') \rightarrow \mathcal{A}(\Sigma_i)$ be defined by $f_i(v_j) = \delta_{ij}t_j$ where δ_{ij} is the Kronecker symbol. Then each f_i is a rooted cluster morphism. Again by definition of product, there exists a unique rooted cluster morphism $h : \mathcal{A}(\Sigma') \rightarrow \mathcal{A}(\Sigma)$ such that the above diagram commutes. Since h satisfies **(CM1)**, for any $i \in \{1, 2\}$, we have $h(v_i) = x_i$ for some $x_i \in \mathbf{x}$ such that $p_i(x_i) = p_i(x'_i) = t_i$. If $x_i, x'_i \in \mathbf{x}$ are such that $p_i(x_i) = t_i$, then the morphism given by $h(v_i) = h(x'_i)$ induces a rooted cluster morphism $\mathcal{A}(\Sigma') \rightarrow \mathcal{A}(\Sigma)$ and by uniqueness, $h = h'$ and thus $x_i = x'_i$. Also, as $(p_1 \circ h)(v_1) = t_1$ and $(p_1 \circ h)(v_2) = 0$, we have $h(v_1) \neq h(v_2)$. Therefore, we obtained exactly two elements $h(v_1) = x_1$ and $h(v_2) = x_2$ in \mathbf{x} such that $p_1(x_1) = t_1, p_2(x_1) = 0, p_1(x_2) = 0$ and $p_2(x_2) = t_2$ and these elements are distinct.

Now consider again the seed $\Sigma' = ((x), \emptyset, [0])$ and $f_i : \mathbb{Z}[x] \rightarrow \mathbb{Z}[t_i]$ the morphism sending x to t_i for any $i \in \{1, 2\}$. Let h be the unique morphism such that the above diagram commutes. Then, by commutativity of the left triangle, we get $h(x) = x_1$ and by commutativity of the right triangle, we get $h(x) = x_2$, a contradiction. Therefore, $\mathcal{A}(\Sigma_1)$ and $\mathcal{A}(\Sigma_2)$ have no product in **Clus**. \square

5.3. Amalgamated sums. In this subsection, we prove that the amalgamated sums of seeds yield pushouts of injective morphisms in **Clus**.

Let $\Sigma_1 = (\mathbf{x}_1, \mathbf{ex}_1, B^1)$ and $\Sigma_2 = (\mathbf{x}_2, \mathbf{ex}_2, B^2)$ be two seeds and let Δ_1, Δ_2 be (possibly empty) subsets of \mathbf{x}_1 and \mathbf{x}_2 respectively such that Σ_1 and Σ_2 are glueable along Δ_1, Δ_2 . We recall that necessarily Δ_1 and Δ_2 consist of frozen variables.

For any $k = 1, 2$, it follows from Lemma 4.1 that we have a natural injective rooted cluster morphism :

$$i_k : \mathbb{Z}[\Delta] \rightarrow \mathcal{A}(\Sigma_k)$$

and from Lemma 4.13 that we have a natural injective rooted cluster morphism

$$j_k : \mathcal{A}(\Sigma_k) \rightarrow \mathcal{A}(\Sigma_1) \amalg_{\Delta_1, \Delta_2} \mathcal{A}(\Sigma_2).$$

Proposition 5.5. *With the previous notations, the diagram*

$$\begin{array}{ccc}
 \mathbb{Z}[\Delta] & \xrightarrow{i_1} & \mathcal{A}(\Sigma_1) \\
 i_2 \downarrow & & \downarrow j_1 \\
 \mathcal{A}(\Sigma_2) & \xrightarrow{j_2} & \mathcal{A}(\Sigma_1) \amalg_{\Delta_1, \Delta_2} \mathcal{A}(\Sigma_2)
 \end{array}$$

is the amalgamated sum of i_1 and i_2 in **Clus**.

Proof. One must first prove that the diagram commutes. Let x be a variable in the cluster Δ of $\Sigma_1 \cap_{\Delta_1, \Delta_2} \Sigma_2$. Then i_1 identifies canonically x with a variable x_1 in Δ_1 and i_2 identifies canonically x with a variable x_2 in Δ_2 . But j_1 and j_2 then identify canonically x_1 and x_2 with the variable x viewed as an element in $\mathbf{x}_1 \amalg_{\Delta_1, \Delta_2} \mathbf{x}_2$. Thus the diagram commutes.

Let now Σ be a seed such that there exists a commutative diagram in **Clus**

$$\begin{array}{ccc}
 \mathcal{A}(\Sigma_1 \cap_{\Delta_1, \Delta_2} \Sigma_2) & \xrightarrow{i_1} & \mathcal{A}(\Sigma_1) \\
 i_2 \downarrow & & \downarrow f_1 \\
 \mathcal{A}(\Sigma_2) & \xrightarrow{f_2} & \mathcal{A}(\Sigma)
 \end{array}$$

We show that there exists a unique rooted cluster morphism $h : \mathcal{A}(\Sigma_1) \amalg_{\Delta_1, \Delta_2} \mathcal{A}(\Sigma_2) \rightarrow \mathcal{A}(\Sigma)$ such that the following diagram commutes :

$$(1) \quad \begin{array}{ccc}
 & \mathcal{A}(\Sigma_1) \amalg_{\Delta_1, \Delta_2} \mathcal{A}(\Sigma_2) & \\
 j_1 \nearrow & \downarrow h & \nwarrow j_2 \\
 \mathcal{A}(\Sigma_1) & \xrightarrow{f_1} \mathcal{A}(\Sigma) \xleftarrow{f_2} & \mathcal{A}(\Sigma_2)
 \end{array}$$

For any $k = 1, 2$ we identify Δ_k with $\Delta \subset \mathbf{x}_1 \amalg_{\Delta_1, \Delta_2} \mathbf{x}_2$ via the morphism j_k . It follows from the commutativity of the first diagram that $f_1(x) = f_2(x)$ for any $x \in \Delta$. We thus set $h : \mathcal{F}_{\Sigma_1 \amalg_{\Delta_1, \Delta_2} \Sigma_2} \rightarrow \mathcal{F}_{\Sigma}$ via

$$h(x) = \begin{cases} f_1(x) = f_2(x) & \text{if } x \in \Delta ; \\ f_1(x) & \text{if } x \in \mathbf{x}_1 ; \\ f_2(x) & \text{if } x \in \mathbf{x}_2. \end{cases}$$

Because f_1 and f_2 are rooted cluster morphisms, h is necessarily also a rooted cluster morphism and therefore the diagram (1) commutes. Conversely, if h is a rooted cluster morphism such that the diagram (1) commutes, then h is entirely determined by its values on $\mathbf{x}_1 \amalg_{\Delta_1, \Delta_2} \mathbf{x}_2$ and it is easily seen that h must be the above morphism. \square

5.4. Topological interpretation of the amalgamated sums. In this subsection we prove that amalgamated sums of cluster algebras of surfaces correspond to cluster algebras associated with connected sums of surfaces.

Let (S_1, M_1) and (S_2, M_2) be marked surfaces in the sense of [FST08] and let ∂_1 and ∂_2 be boundary components respectively of S_1 and S_2 such that there exists a homeomorphism $h : \partial_1 \xrightarrow{\sim} \partial_2$ satisfying $h(\partial_1 \cap M_1) = \partial_2 \cap M_2$.

We denote by

$$(S, M) = (S_1, M_1) \coprod_{\partial_1, \partial_2} (S_2, M_2)$$

the *connected sum* of S_1 and S_2 along h , that is, S is the surface obtained by gluing S_1 and S_2 along the homeomorphism h and $M = (M_1 \setminus (M_1 \cap \partial_1)) \cup (M_2 \setminus (M_2 \cap \partial_2)) \cup M_\partial$ where M_∂ is the common image of $\partial_1 \cap M_1$ and $\partial_2 \cap M_2$ in the surface S (see for instance [Mas77, p.8]). We denote by ∂ the common image of ∂_1 and ∂_2 in the surface S .

Remark 5.6. Even if (S_1, M_1) and (S_2, M_2) are unpunctured surfaces, the surface (S, M) may have punctures, see for instance Figure 2.

Let T_1 and T_2 be triangulations of (S_1, M_1) and (S_2, M_2) respectively and let Σ_1 and Σ_2 be the corresponding seeds. Then ∂_1 is identified with a subset of the frozen variables in Σ_1 and ∂_2 is identified with a subset of the frozen variables in Σ_2 . In the connected sum (S, M) , the collection $T_1 \cup T_2$ defines a triangulation and we denote by Σ the associated seed. Then it follows from Proposition 5.5 that

$$\mathcal{A}(\Sigma^\partial) = \mathcal{A}(\Sigma_1) \coprod_{\partial_1, \partial_2} \mathcal{A}(\Sigma_2)$$

where the notation Σ^∂ means that we have frozen the variables corresponding to ∂ in Σ

Corollary 5.7. *In the category **Clus**, the rooted cluster algebra associated with the triangulation T of S where the arcs in ∂ are frozen is the amalgamated sum over the polynomial ring $\mathbb{Z}[\partial]$ of the rooted cluster algebras $\mathcal{A}(\Sigma_1)$ and $\mathcal{A}(\Sigma_2)$ associated with the triangulations T_1 and T_2 induced by T respectively on S_1 and S_2 .*

Example 5.8. Figure 2 shows such a gluing.




FIGURE 2. An example of connected sum of an annulus and a pair of pants along a boundary and the induced triangulations.

6. SURJECTIVE ROOTED CLUSTER MORPHISMS

We recall that an *epimorphism* in a category is a morphism f such that if there exist morphisms h and g such that $gf = hf$, then $g = h$. In this section, we focus on surjective rooted cluster morphisms, which are particular cases of epimorphisms since **Clus** is a concrete category.

Remark 6.1. As in the category **Ring**, epimorphisms in **Clus** are *not* necessarily surjective. Indeed, if one considers the seeds

$$\Sigma_1 = ((x_1), \emptyset, [0]), \text{ and } \Sigma_2 = ((x_1), (x_1), [0]),$$

then it follows from Lemma 4.1 that the identity morphism $\mathcal{F}_{\Sigma_1} = \mathbb{Q}(x_1) \rightarrow \mathbb{Q}(x_1) = \mathcal{F}_{\Sigma_2}$ induces a rooted cluster morphism $f : \mathbb{Z}[x_1] = \mathcal{A}(\Sigma_1) \rightarrow \mathcal{A}(\Sigma_2) = \mathbb{Z}[x_1, \frac{2}{x_1}]$. If Σ_3 is another seed and $g, h : \mathcal{A}(\Sigma_2) \rightarrow \mathcal{A}(\Sigma_3)$ are such that $hf = gf$, then $hf(x_1) = gf(x_1)$ so that $h(x_1) = g(x_1)$ and as g and h are ring homomorphisms, we also have $h(\frac{2}{x_1}) = g(\frac{2}{x_1})$ so that $h = g$. Therefore f is an epimorphism.

As f is injective, it follows from Proposition 4.3 that it is a monomorphism in **Clus** and thus, f is an example of a *bimorphism* (that is, both a monomorphism and an epimorphism) in **Clus** which is not an isomorphism.

Proposition 6.2. *Let Σ, Σ' be two seeds and $f : \mathcal{A}(\Sigma) \rightarrow \mathcal{A}(\Sigma')$ be a surjective rooted cluster morphism. Then :*

- (1) *any Σ' -admissible sequence lifts to an (f, Σ, Σ') -biadmissible sequence,*
- (2) $\mathcal{X}'_{\Sigma'} \subset f(\mathcal{X}_{\Sigma})$.

Proof. Let $y \in \mathbf{ex}'$. According to Lemma 3.1, there exists $x \in \mathbf{ex}$ such that $f(x) = y$ so that the Σ' -admissible sequence (y) lifts to the (f, Σ, Σ') -biadmissible sequence (x) . Now let (y_1, \dots, y_l) be a Σ' -admissible sequence. We prove by induction on l that (y_1, \dots, y_l) lifts to an (f, Σ, Σ') -biadmissible sequence. If $l = 1$, this follows from the above discussion. Otherwise, there exists $y \in \mathbf{x}'$ such that

$$y_l = \mu_{y_{l-1}} \circ \dots \circ \mu_{y_1}(y).$$

By the induction hypothesis, (y_1, \dots, y_{l-1}) lifts to an (f, Σ, Σ') -biadmissible sequence (x_1, \dots, x_{l-1}) and since f satisfies **(CM3)**, we get

$$y_l = \mu_{f(x_{l-1})} \circ \dots \circ \mu_{f(x_1)}(x) = f(\mu_{x_{l-1}} \circ \dots \circ \mu_{x_1}(x))$$

where x lifts y in \mathbf{ex} . Therefore, if $x_l = \mu_{x_{l-1}} \circ \dots \circ \mu_{x_1}(x)$, the sequence (x_1, \dots, x_l) lifts (y_1, \dots, y_l) . And moreover, $y_l \in f(\mathcal{X}_{\Sigma})$, which proves the corollary. \square

Corollary 6.3. *Let Σ_1 and Σ_2 be two seeds and $f : \mathcal{A}(\Sigma_1) \rightarrow \mathcal{A}(\Sigma_2)$ be a surjective rooted cluster morphism. Assume that Σ_1 is of finite cluster type. Then Σ_2 is of finite cluster type.*

6.1. Subseeds and surjective morphisms. Let $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ be a seed and let $\mathbf{x}' \subset \mathbf{x}$ be a subset. We set $\Sigma' = \Sigma|_{\mathbf{x}'}$ the corresponding full subseed (see Definition 4.9). We consider the surjective ring homomorphism

$$\text{Res}_{\Sigma, \Sigma'} : \begin{cases} \mathcal{F}_{\Sigma} & \longrightarrow & \mathcal{F}_{\Sigma'} \\ x & \mapsto & x \text{ if } x \in \mathbf{x}', \\ x & \mapsto & 1 \text{ if } x \in \mathbf{x} \setminus \mathbf{x}'. \end{cases}$$

Proposition 6.4. *$\text{Res}_{\Sigma, \Sigma'}$ induces an ideal, surjective, rooted cluster morphism $\mathcal{A}(\Sigma) \rightarrow \mathcal{A}(\Sigma')$.*

Proof. In order to simplify notations, we set $\text{Res} = \text{Res}_{\Sigma, \Sigma'}$ and $\Sigma' = (\mathbf{x}', \mathbf{ex}', B')$. Let $x \in \mathbf{ex}$. Then $\text{Res}(x) \in \mathbf{ex}'$ if and only if $x \in \mathbf{x}'$, that is, if and only if $x \in \mathbf{ex}'$. Moreover,

$$\text{Res}(\mu_{x, \Sigma}(x)) = \text{Res} \left(\frac{1}{x} \left(\prod_{\substack{b_{xy} > 0 \\ y \in \mathbf{x}}} y^{b_{xy}} + \prod_{\substack{b_{xy} < 0 \\ y \in \mathbf{x}}} y^{-b_{xy}} \right) \right) = \frac{1}{x} \left(\prod_{\substack{b_{xy} > 0 \\ y \in \mathbf{x}'}} y^{b_{xy}} + \prod_{\substack{b_{xy} < 0 \\ y \in \mathbf{x}'}} y^{-b_{xy}} \right) = \mu_{x, \Sigma'}(x).$$

Moreover, it is easily seen that $\mu_x(B)[\mathbf{x}'] = \mu_x(B[\mathbf{x}'])$. By induction, we see that every sequence (x_1, \dots, x_n) which is Σ -admissible is $(\text{Res}, \Sigma, \Sigma')$ -biadmissible and that Res commutes with biadmissible mutations. Therefore Res is a rooted cluster morphism.

Moreover, $\text{Res}(\mathcal{A}(\Sigma)) \subset \mathcal{A}(\Sigma')$ and $\Sigma' = \Sigma|_{\mathbf{x}'} = \text{Res}(\Sigma)$ so that Lemma 2.10 implies that Res is ideal. Finally, by definition of Σ' , every Σ' -admissible sequence (x_1, \dots, x_n) lifts to a biadmissible sequence and therefore, $\mathcal{A}(\Sigma') \subset \text{Res}(\mathcal{A}(\Sigma))$ so that Res is surjective. \square

6.2. Specialisations. It is well-known that specialising frozen variables to 1 allows one to realise coefficient-free cluster algebras from cluster algebras of geometric type, see for instance [FZ07]. In this subsection, we study the slightly more general case where an arbitrary cluster variable, frozen or not, is specialised to an integer (which can essentially be assumed to be 1). If the considered cluster variable is frozen, then one finds natural surjective rooted cluster morphisms. More surprisingly, as we prove in certain cases (and expect in general), specialising an *exchangeable* cluster variable to 1 also leads to rooted cluster morphisms.

We start with a technical lemma :

Lemma 6.5. *Let \mathcal{A} be a rooted cluster algebra and let \mathbf{x} be a cluster in \mathcal{A} . Let $m = \prod_{x \in \mathbf{x}} x^{d_x}$ be a Laurent monomial in the variables in \mathbf{x} , with $d_x \in \mathbb{Z}$ for any $x \in \mathbf{x}$. Then the following conditions are equivalent :*

- (1) *m is an element in \mathcal{A} ;*
- (2) *$d_x \geq 0$ for any $x \in \mathbf{x}$;*
- (3) *m is a monomial in \mathbf{x} .*

Proof. Let Σ be a seed containing the cluster \mathbf{x} . It is clear that the second and third assertions are equivalent and that the second implies the first one. Therefore, we only have to prove that the first one implies the second one. Assume that there exists some $x \in \mathbf{x}$ such that $d_x < 0$. Because the elements in \mathcal{A} are Laurent polynomials in the exchangeable variables of \mathbf{x} with polynomial coefficients in the frozen variables of \mathbf{x} (see for instance [FZ03, Proposition 11.2]), if $d_x < 0$, then the variable x is necessarily exchangeable. Let thus $\Sigma' = (\mathbf{x}', \mathbf{ex}', B') = \mu_x \Sigma$ with $\mathbf{x}' = (\mathbf{x} \setminus \{x\}) \sqcup \{x'\}$. Then the expansion of m in Σ' is

$$m = \prod_{y \in \mathbf{x} \setminus x} y^{d_y} \left(\frac{x'}{\prod_{b_{xz} > 0} z^{b_{xz}} + \prod_{b_{xz} < 0} z^{-b_{xz}}} \right)^{-d_x}.$$

In particular, m is not a Laurent polynomial in the cluster \mathbf{x}' and thus, according to the Laurent phenomenon (see [FZ02]), m does not belong to \mathcal{A} . \square

Let $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ and let $x \in \mathbf{x}$. We denote by $\Sigma \setminus \{x\}$ the seed $\Sigma|_{\mathbf{x} \setminus \{x\}} = (\mathbf{x}', \mathbf{ex}', B')$ where $\mathbf{x}' = \mathbf{x} \setminus \{x\}$, $\mathbf{ex}' = \mathbf{ex} \setminus \{x\}$ and $b'_{yz} = b_{yz}$ for any $y, z \neq x$ in \mathbf{x}' .

Definition 6.6 (Simple specialisation). Let $n \in \mathbb{Z}$. The *simple specialisation of x to n* is the ring homomorphism :

$$\sigma_{x,n} : \begin{cases} \mathcal{F}_{\Sigma} & \longrightarrow & \mathcal{F}_{\Sigma \setminus \{x\}} \\ x & \mapsto & n, \\ z & \mapsto & z \text{ if } z \in \mathbf{x} \setminus \{x\}. \end{cases}$$

The following lemma shows that, except degenerate cases, the only value to which we can specialise a (single) cluster variable is 1.

Lemma 6.7. *Let $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ be a seed, let $x \in \mathbf{x}$ and let $n \in \mathbb{Z}$. Assume that $\sigma_{x,n}$ induces a ring homomorphism $\mathcal{A}(\Sigma) \rightarrow \mathcal{A}(\Sigma \setminus \{x\})$. If there exists some $y \in \mathbf{ex}$ such that $b_{xy} \neq 0$, then $n \in \{-1, 1\}$. If there exists some $y \in \mathbf{ex}$ such that $b_{xy} \in 2\mathbb{Z} + 1$, then $n = 1$.*

Proof. Let $x \in \mathbf{x}$, $\mathbf{x}' = \mathbf{x} \setminus \{x\}$ and $\Sigma' = \Sigma \setminus \{x\}$. Assume that there exists $y \in \mathbf{ex}$ such that $b_{xy} \neq 0$. Without loss of generality, we assume that $b_{xy} > 0$. Then

$$\begin{aligned} \sigma(\mu_{y,\Sigma}(y)) &= \sigma\left(\frac{1}{y} \left(\prod_{b_{zy}>0} z^{b_{zy}} + \prod_{b_{yz}>0} z^{b_{yz}} \right)\right) \\ &= \frac{1}{y} \left(n^{b_{xy}} \prod_{\substack{b_{zy}>0 \\ z \neq x}} z^{b_{zy}} + \prod_{\substack{b_{yz}>0 \\ z \neq x}} z^{b_{yz}} \right) \\ &= \frac{1}{y} \left(n^{b_{xy}} \prod_{b'_{zy}>0} z^{b'_{zy}} + \prod_{b'_{yz}>0} z^{b'_{yz}} \right). \end{aligned}$$

But

$$\mu_{y,\Sigma'}(y) = \frac{1}{y} \left(\prod_{b'_{zy}>0} z^{b_{zy}} + \prod_{b'_{yz}>0} z^{b_{yz}} \right).$$

Thus, if σ induces a ring homomorphism between the rooted cluster algebras, we get $\sigma(\mu_{y,\Sigma}(y)) \in \mathcal{A}(\Sigma \setminus \{x\})$ and so the difference $\sigma(\mu_{y,\Sigma}(y)) - n^{b_{xy}} \mu_{y,\Sigma'}(y)$ also belongs to $\mathcal{A}(\Sigma \setminus \{x\})$.

But

$$\sigma(\mu_{y,\Sigma}(y)) - n^{b_{xy}} \mu_{y,\Sigma'}(y) = \frac{1}{y} (1 - n^{b_{xy}}) \prod_{b'_{yz}>0} z^{b'_{yz}}$$

is a Laurent monomial in the cluster \mathbf{x}' such that the exponent of y is $-1 < 0$, with $y \in \mathbf{ex}$. Therefore, it follows from 6.5 that necessarily $1 - n^{b_{xy}} = 0$, that is, $n^{b_{xy}} = 1$ and thus $n \in \{-1, 1\}$. If moreover b_{xy} is odd, we necessarily have $n = 1$. \square

Example 6.8. We exhibit an example where a simple specialisation to -1 does not induce a map at the level of the corresponding cluster algebras. Consider the cluster algebras of respective types A_3 and A_2 associated with the coefficient-free seeds

$$\Sigma = ((x_1, x_2, x_3), 1 \longrightarrow 2 \longleftarrow 3) \text{ and } \Sigma' = ((x_1, x_2), 1 \longrightarrow 2)$$

and consider the simple specialisation $\sigma = \sigma_{x_3,-1}$ of x_3 to -1 .

The image under σ of the cluster variable $\frac{1+x_1x_3}{x_2}$ is $\frac{1-x_1}{x_2}$. But if $\frac{1-x_1}{x_2}$ is in $\mathcal{A}(\Sigma')$, then, since $\frac{1+x_1}{x_2}$ is in $\mathcal{A}(\Sigma')$, we get $\frac{2}{x_2} \in \mathcal{A}(\Sigma')$. Then the expansion of $\frac{2}{x_2}$ in the cluster (x_1, x'_2) of the seed $\mu_2(\Sigma')$ is $2\frac{x'_2}{1+x_1}$ which is not a Laurent polynomial, a contradiction.

The following lemma shows that the study of simple specialisations in -1 can be reduced to the study of simple specialisations to 1 .

Lemma 6.9. *Let $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ be a seed and let $x \in \mathbf{ex}$ be such that $b_{xy} \in 2\mathbb{Z}$ for any $y \in \mathbf{ex}$. Then $\sigma_{x,-1}$ is a rooted cluster morphism if and only if $\sigma_{x,1}$ is. Moreover, in this case $\sigma_{x,-1}(a) = \pm \sigma_{x,1}(a)$ for any $a \in \mathcal{X}_\Sigma$.*

Proof. Both $\sigma_{x,-1}$ and $\sigma_{x,1}$ satisfy **(CM1)** and **(CM2)**. Now we observe that a sequence of variables is $(\sigma_{x,-1}, \Sigma, \Sigma \setminus \{x\})$ -biadmissible if and only if it is disjoint from x , which is also the condition for this sequence to be $(\sigma_{x,1}, \Sigma, \Sigma \setminus \{x\})$ -biadmissible. Let $a \neq x$ be an exchangeable variable in Σ . Then, since $b_{xy} \in 2\mathbb{Z}$ for any $y \in \mathbf{ex}$, it follows that $\sigma_{x,-1}(\mu_{a,\Sigma}(a)) = \sigma_{x,1}(\mu_{a,\Sigma}(a))$. Moreover, if $\mu_a \Sigma = \Sigma' = (\mathbf{x}', \mathbf{ex}', B')$, then it follows from the mutation rule for exchange matrices that $b'_{xy} \in 2\mathbb{Z}$ for any $y \in \mathbf{ex}'$. Therefore, by induction, $\sigma_{x,-1}$ satisfies **(CM3)** if and only if $\sigma_{x,1}$ does. Hence, it only remains to prove that $\sigma_{x,-1}$ induces a map $\mathcal{A}(\Sigma) \rightarrow \mathcal{A}(\Sigma \setminus x)$ if and only if $\sigma_{x,1}$ does. This follows from the fact that $\sigma_{x,-1}(a) \in \{-\sigma_{x,1}(a), \sigma_{x,1}(a)\}$ for any $a \in \mathcal{X}_\Sigma$, which is easily proved by induction. \square

Proposition 6.10. *Let $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ be a seed and let $x \in \mathbf{x}$. Then $\sigma_{x,1}$ induces an ideal surjective rooted cluster morphism $\mathcal{A}(\Sigma) \rightarrow \mathcal{A}(\Sigma \setminus \{x\})$ if and only if it induces a ring homomorphism $\mathcal{A}(\Sigma) \rightarrow \mathcal{A}(\Sigma \setminus \{x\})$.*

Proof. Let $\sigma = \sigma_{x,1}$. Then σ clearly satisfies **(CM1)** and **(CM2)**. In order to prove **(CM3)**, it is enough to notice that a Σ -admissible sequence (x_1, \dots, x_l) is $(\sigma, \Sigma, \Sigma \setminus \{x\})$ -biadmissible if and only if $x_k \neq x$ for any k such that $1 \leq k \leq l$ and to proceed by induction on l . It follows that σ is a rooted cluster morphism if and only if σ induces a ring homomorphism $\mathcal{A}(\Sigma) \rightarrow \mathcal{A}(\Sigma \setminus \{x\})$. Its surjectivity comes from the fact that any admissible sequence in $\Sigma \setminus \{x\}$ can naturally be lifted to a $(\sigma, \Sigma, \Sigma \setminus \{x\})$ -biadmissible sequence. In order to prove that σ is ideal, it is enough to observe that $\sigma(\Sigma) = \Sigma \setminus \{x\}$ so that $\sigma(\mathcal{A}(\Sigma)) = \mathcal{A}(\Sigma \setminus \{x\}) = \mathcal{A}(\sigma(\Sigma))$. \square

6.3. Specialisations for cluster algebras from surfaces. In this section, we prove that simple specialisations induce surjective rooted cluster morphisms for cluster algebras associated with surfaces.

Given a marked surface (S, M) and an internal arc γ in (S, M) , we denote by $d_\gamma(S, M)$ the (non-necessarily connected) marked surface obtained by cutting (S, M) along γ . The arc γ induces in $d_\gamma(S, M)$ two new boundary arcs which we denote by γ_1 and γ_2 . Now if T is a triangulation of (S, M) , then

$$d_\gamma T = (T \setminus \{\gamma\}) \sqcup \{\gamma_1, \gamma_2\}$$

is a triangulation of $d_\gamma(S, M)$.

Figures 3 and 4 present examples of cuttings of surfaces along an arc.




FIGURE 3. Cutting a disc with two punctures to get an annulus.

Note that the seed $\Sigma_{d_\gamma T}$ differs from the seed $\Sigma_T \setminus \{x_\gamma\}$. However, we have $\Sigma_{d_\gamma T}[T \setminus \{\gamma_1, \gamma_2\}] = \Sigma_T[T \setminus \{\gamma\}]$.




FIGURE 4. Cutting an annulus to get a disc without punctures.

Theorem 6.11. *Let (S, M) be a marked surface, T be a triangulation of (S, M) and Σ_T be the seed associated with T . Then for any $\gamma \in T$ which does not enclose a degenerate marked surface, the simple specialisation of x_γ to 1 induces an ideal surjective rooted cluster morphism in $\mathcal{A}(\Sigma_T) \rightarrow \mathcal{A}(\Sigma_T \setminus \{x_\gamma\})$.*

Proof. If γ is a boundary arc, then x_γ is a coefficient and the result is well-known, see [FZ07]. We may thus assume that γ is an internal arc.

According to Proposition 6.10, as $\sigma = \sigma_{x_\gamma, 1}$ is a ring homomorphism, in order to prove that σ is a rooted cluster morphism, it is enough to prove that the image of σ is contained in $\mathcal{A}(\Sigma_T \setminus \{x_\gamma\})$. For this we only need to prove that $\sigma(x_\eta) \in \mathcal{A}(\Sigma_T \setminus \{x_\gamma\})$ for any (possibly tagged) arc η in (S, M) . For the sake of simplicity we only prove it for an untagged arc η . The case of tagged arcs is a straightforward adaptation.


Let η be an arc in (S, M) . Resolving the intersections of η with γ (using for instance the resolutions described in [DP11] or more generally the skein relations described in [MW11]), we can write $x_\gamma x_\eta$ as a linear combination of products of x_θ where θ runs over a family of curves which do not intersect γ .

Every curve which does not cross γ induces a curve in the surface $d_\gamma(S, M)$. Let τ denote the specialisation of x_{γ_1} and x_{γ_2} to 1. Then, as γ_1 and γ_2 are boundary arcs, τ is a rooted cluster morphism from $\mathcal{A}(\Sigma_{d_\gamma T})$ to $\mathcal{A}(\Sigma_{d_\gamma T} \setminus \{x_{\gamma_1}, x_{\gamma_2}\}) = \mathcal{A}(\Sigma_T \setminus \{x_\gamma\})$. Moreover, for any arc θ in (S, M) which does not cross γ , we have $\sigma(x_\theta) = \tau(x_\theta)$.

Therefore, $\sigma(x_\eta) = \sigma(x_\gamma x_\eta)$ is a linear combination of $\tau(x_\theta)$ where θ runs over a family of curves which do not intersect γ . Now for each such curve, x_θ is an element of the cluster algebra $\mathcal{A}(d_\gamma T)$ and thus $\tau(x_\theta)$ is an element of the cluster algebra $\mathcal{A}(\Sigma_T \setminus \{x_\gamma\})$.

The fact that σ is surjective is clear since any admissible sequence for $\Sigma_T \setminus \{x_\gamma\}$ lifts to an admissible sequence for Σ_T . The fact that it is ideal comes from the fact that $\sigma(\Sigma_T) = \Sigma_T \setminus \{x_\gamma\}$. \square


Example 6.12. Consider the once-punctured torus \mathbb{T}_1 and fix a triangulation T of \mathbb{T}_1 . It has three arcs which we denote by 1, 2 and 3 and which we show as follows in the universal cover of \mathbb{T}_1 :



The seed corresponding to this triangulation is the coefficient-free seed

$$\Sigma_T = \left((x_1, x_2, x_3), \begin{bmatrix} 0 & 2 & -2 \\ -2 & 0 & 2 \\ 2 & -2 & 0 \end{bmatrix} \right).$$

Now, an arbitrary cluster variable in $\mathcal{A}(\mathbb{T}_1)$ corresponds to a certain arc. In this example we choose for instance the curve η shown below. Let us cut \mathbb{T}_1 along the arc 1, which we show dashed. Resolving intersections between η and 1, and applying skein relations we get $x_\eta x_1 = x_2 x_3 + x_2 x_\theta$ where θ does not intersect the arc 1.




We can easily compute

$$x_\theta = \frac{x_1^2 + x_2^2}{x_3}$$

so that $\sigma_{x_1,1}(x_\theta) = \frac{1+x_2^2}{x_3} \in \mathcal{A}(\Sigma \setminus \{x_1\})$. Let us also give a geometric argument.

Cutting \mathbb{T}_1 along the arc 1, we get the annulus $C_{1,1}$ with one marked point on each boundary component and $\{2, 3\}$ together with the two boundary arcs b and b' induce a triangulation of $C_{1,1}$. The arc θ in \mathbb{T}_1 induces an arc in $C_{1,1}$ and we denote by x'_θ the corresponding cluster variable in $\mathcal{A}(C_{1,1})$.



Then, it follows directly from the various expansion formulae for cluster variables associated with arcs (see for instance [ST09, MSW11] or [ADSS11]) that x_θ is given by x'_θ where the variables x_b and $x_{b'}$ corresponding to the two boundary components are identified with x_1 . Indeed, a direct computation gives

$$x'_\theta = \frac{x_b x_{b'} + x_2^2}{x_3}.$$

In particular, if we specialise $x_b, x_{b'}$ and x_1 to 1, we still get equality. But the cluster algebra associated with $C_{1,1}$ whose frozen variables are specialised to 1 is nothing but the cluster algebra associated with the seed $\mathcal{A}(\Sigma_T \setminus \{x_1\})$.

6.4. Specialisations in acyclic cluster algebras. We now prove an analogue result for acyclic cluster algebras using techniques coming from additive categorifications of cluster algebras. We refer the interested reader to [Kel10] and references therein.

Philosophically speaking, the proof of the following theorem follows the lines of the proof of Theorem 6.11 where, in the spirit of [BZ11], objects in categories should be thought as curves in a surface, extensions of objects as intersections of the corresponding curves and “Hall products” as skein relations.

Theorem 6.13. *Let $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ be a seed which is mutation-equivalent to a finite skew-symmetric acyclic seed. Then for any $x \in \mathbf{x}$, we have $\sigma_{x,1}(\mathcal{A}(\Sigma)) \subset \mathcal{A}(\Sigma \setminus \{x\}) \otimes_{\mathbb{Z}} \mathbb{Q}$.*

Proof. Without loss of generality we can assume that $\Sigma = \bar{\Sigma}$ is simplified. Let Q be an acyclic quiver corresponding to the exchange matrix of a seed which is mutation-equivalent to Σ .

Let \mathcal{C} be the cluster category of Q over an algebraically closed field \mathbb{k} introduced in [BMR⁺06]. Its suspension functor is denoted by [1]. Let $T = \bigoplus_{y \in \mathbf{x}} T_y$ be a cluster-tilting object in \mathcal{C} corresponding to the seed Σ , see [CK06, FK10].

Let T_x^\perp be the full subcategory of \mathcal{C} formed by the objects V such that $\text{Hom}_{\mathcal{C}}(T_x, V) = 0$. Then it follows from [IY08] that $\mathcal{C}' = T_x^\perp/T_x[1]$ is a Hom-finite 2-Calabi-Yau category and that $T' = \bigoplus_{y \in \mathbf{x} \setminus \{x\}} T_y$ is a cluster-tilting object in \mathcal{C}' . The quiver $Q_{T'}$ of the endomorphism ring of T' is a full subquiver of the quiver Q_T of the endomorphism ring of T . By assumption Q_T is mutation-acyclic. Thus, it follows from [BMR08] (see also [War11]) that $Q_{T'}$ is also mutation-acyclic and therefore \mathcal{C}' is triangle equivalent to a cluster category, see [KR08].

We denote by X_γ (or X'_γ , respectively) the cluster character associated to T on \mathcal{C} (or to T' on \mathcal{C}' , respectively), see [Pal08]. Since \mathcal{C} and \mathcal{C}' are cluster categories associated to acyclic quivers, it follows from [GLS10, Theorem 4] that the map X_γ (or X'_γ , respectively) has values in the cluster algebra $\mathcal{A}(\Sigma)$ (or $\mathcal{A}(\Sigma \setminus \{x\})$, respectively).

Let $\sigma = \sigma_{x,1}$ be the simple specialisation of x to 1. Let M be an object in \mathcal{C} . We prove by induction on the dimension of $\text{Hom}_{\mathcal{C}}(T_x, M)$ that $\sigma(X_M)$ is a finite \mathbb{Q} -linear combination of X'_Y where Y runs over the objects of \mathcal{C}' .

Assume first that $\text{Hom}_{\mathcal{C}}(T_x, M) = 0$, that is, M belongs to T_x^\perp . Then M can be decomposed as $\bar{M} \oplus T_x[1]^m$ for some $m \geq 1$ where \bar{M} has no direct summand isomorphic to $T_x[1]$. Therefore,

$$\sigma(X_M) = \sigma(X_{\bar{M} \oplus T_x[1]^m}) = \sigma(X_{\bar{M}} X_{T_x[1]}^m) = \sigma(X_{\bar{M}}) \sigma(X_{T_x[1]}^m) = \sigma(X_{\bar{M}}) \sigma(x^m) = \sigma(X_{\bar{M}}).$$

Since M belongs to T_x^\perp , the object \bar{M} belongs to $T_x^\perp/T_x[1]$ and is thus identified with an object in \mathcal{C}' . Now it follows easily from the definition of the cluster characters that $\sigma(X_{\bar{M}}) = X'_{\bar{M}}$. Therefore, $\sigma(X_M)$ belongs to $\mathcal{A}(\Sigma \setminus \{x\})$.

Assume now that $\text{Hom}_{\mathcal{C}}(T_x, M) \neq 0$. Therefore,

$$\text{Ext}_{\mathcal{C}}^1(T_x[1], M) = \text{Hom}_{\mathcal{C}}(T_x[1], M[1]) \simeq \text{Hom}_{\mathcal{C}}(T_x, M) \neq 0.$$

Then it follows from [Pal11] that the product $(\dim \text{Ext}_{\mathcal{C}}^1(T_x[1], M) X_{T_x[1]} X_M)$ is a \mathbb{Z} -linear combination of X_Y 's where Y runs over middle terms of non-split triangles of the form

$$(2) \quad M \longrightarrow Y \longrightarrow T_x[1] \longrightarrow M[1]$$

or of the form

$$(3) \quad T_x[1] \longrightarrow Y \xrightarrow{a} M \xrightarrow{b} T_x[2].$$

In other words, we have

$$(4) \quad xX_M = \sum_Y n_Y X_Y$$

where $(n_Y) \subset \mathbb{Q}$ is finitely supported on a set of isoclasses of objects Y in \mathcal{C} such that there exists triangles of the form (2) or (3).

We claim that $\dim \text{Hom}_{\mathcal{C}}(T_x, Y) < \dim \text{Hom}_{\mathcal{C}}(T_x, M)$ for any Y such that $n_Y \neq 0$. Indeed, assume first that Y is such that there exists a triangle of the form (2), that is,

$$T_x \xrightarrow{\alpha} M \xrightarrow{\beta} Y \longrightarrow T_x[1] \longrightarrow M[1].$$

Applying the homological functor $\text{Hom}_{\mathcal{C}}(T_x, -)$ to this triangle, we get the long exact sequence

$$\text{End}_{\mathcal{C}}(T_x) \xrightarrow{\alpha_*} \text{Hom}_{\mathcal{C}}(T_x, M) \xrightarrow{\beta_*} \text{Hom}_{\mathcal{C}}(T_x, Y) \longrightarrow 0,$$

so that the post-composition by β yields an epimorphism $\text{Hom}_{\mathcal{C}}(T_x, M) \xrightarrow{\beta_*} \text{Hom}_{\mathcal{C}}(T_x, Y)$. Since the sequence is exact, we have $\text{Ker}(\beta_*) = \text{Im}(\alpha_*)$. Moreover, $\text{End}_{\mathcal{C}}(T_x) \xrightarrow{\alpha_*} \text{Hom}_{\mathcal{C}}(T_x, M)$ is

non-zero since $\alpha_*(\mathbb{1}_{T_x}) = \alpha \neq 0$. Therefore, $\dim \operatorname{Hom}_{\mathcal{C}}(T_x, Y) < \dim \operatorname{Hom}_{\mathcal{C}}(T_x, M)$ in this case, as claimed.

Assume now that Y is such that there exists a triangle of the form (3), that is,

$$T_x[1] \longrightarrow Y \xrightarrow{a} M \xrightarrow{b} T_x[2].$$

Applying the homological functor $\operatorname{Hom}_{\mathcal{C}}(T_x, -)$ to this triangle, we get the long exact sequence

$$0 \longrightarrow \operatorname{Hom}_{\mathcal{C}}(T_x, Y) \xrightarrow{a_*} \operatorname{Hom}_{\mathcal{C}}(T_x, M) \xrightarrow{b_*} \operatorname{Hom}_{\mathcal{C}}(T_x, T_x[2]).$$

Therefore, the post-composition by a yields an injection $\operatorname{Hom}_{\mathcal{C}}(T_x, Y) \xrightarrow{a_*} \operatorname{Hom}_{\mathcal{C}}(T_x, M)$. In order to prove that the injection is proper, since the sequence is exact, we need to prove that the post-composition $\operatorname{Hom}_{\mathcal{C}}(T_x, M) \xrightarrow{b_*} \operatorname{Hom}_{\mathcal{C}}(T_x, T_x[2])$ is non-zero. Since \mathcal{C} is 2-Calabi-Yau, we have the commutative diagram

$$\begin{array}{ccc} \operatorname{Hom}_{\mathcal{C}}(T_x, M) & \xrightarrow{b_*} & \operatorname{Hom}_{\mathcal{C}}(T_x, T_x[2]) \\ \sim \downarrow & & \downarrow \sim \\ D\operatorname{Hom}_{\mathcal{C}}(M, T_x[2]) & \xrightarrow{Db^*} & D\operatorname{End}_{\mathcal{C}}(T_x[2]) \end{array}$$

where $D = \operatorname{Hom}_{\mathbb{k}}(-, \mathbb{k})$ is the standard duality and $\operatorname{End}_{\mathcal{C}}(T_x[2]) \xrightarrow{b^*} \operatorname{Hom}_{\mathcal{C}}(M, T_x[2])$ is the pre-composition by b . Since b is non-zero, $b^*(\mathbb{1}_{T_x[2]}) \neq 0$ and thus Db^* is non-zero so that b_* is non-zero. This proves the claim.

Therefore, equality (4) allows to write xX_M as a \mathbb{Q} -linear combination of elements X_Y where $\dim \operatorname{Hom}_{\mathcal{C}}(T_x, Y) < \dim \operatorname{Hom}_{\mathcal{C}}(T_x, M)$. If $\dim \operatorname{Hom}_{\mathcal{C}}(T_x, Y) = \dim \operatorname{Ext}_{\mathcal{C}}^1(T_x[1], Y) \neq 0$, we can again write xX_Y as a linear combination of cluster characters of objects for which the dimension is strictly smaller. Proceeding by induction, there exists some $n \geq 1$ such that $x^n X_M$ is a \mathbb{Q} -linear combination of elements of the form X_Y where $\dim \operatorname{Hom}_{\mathcal{C}}(T_x, Y) = 0$. Therefore $\sigma(X_M) = \sigma(x^n X_M)$ is a \mathbb{Q} -linear combination of $\sigma(X_Y)$ with $\dim \operatorname{Hom}_{\mathcal{C}}(T_x, Y) = 0$. Hence, it follows from the previous discussion that $\sigma(X_M)$ belongs to $\mathcal{A}(\Sigma \setminus \{x\})$. \square

Remark 6.14. In the proof of Theorem 6.13, the assumption that Σ is mutation-acyclic is only used in order to prove that the cluster characters take their values in the cluster algebras. In fact, the proof shows that for any cluster variable x in a seed Σ admitting an additive Hom-finite 2-Calabi-Yau categorification, the map $\sigma_{x,1}$ sends any cluster variable in $\mathcal{A}(\Sigma)$ to a \mathbb{Q} -linear combination of cluster characters associated to objects in a Hom-finite 2-Calabi-Yau categorification of $\Sigma \setminus \{x\}$. However, in general these latter cluster characters may not live in the cluster algebra but only in the upper cluster algebra, see for instance [Pla11a, Example 5.6.3]. Interactions of simple specialisations with lower and upper bounds of cluster algebras are studied in Subsection 6.5.

The following example illustrates the fact appearing in the proofs of Theorems 6.11 and 6.13 that the simple specialisation of an exchangeable variable to 1 does not send a cluster variable to a cluster variable in general, but rather sends it to a linear combination of elements in the cluster algebra.

Example 6.15. Consider the coefficient-free seed $\Sigma = ((x_1, x_2, x_3, x_4), B)$ where

$$B = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ -1 & -1 & -1 & 0 \end{bmatrix}$$

is the incidence matrix of the quiver Q of Dynkin type D_4 where 4 is a sink.

We consider the specialisation of x_4 to 1 so that we also consider the seed

$$\Sigma \setminus \{x_4\} = ((x_1, x_2, x_3), [0])$$

which is of type $A_1 \times A_1 \times A_1$. In particular,

$$\mathcal{A}(\Sigma \setminus \{x_4\}) = \mathbb{Z} \left[x_1, \frac{2}{x_1}, x_2, \frac{2}{x_2}, x_3, \frac{2}{x_3} \right] \subset \mathbb{Q}(x_1, x_2, x_3).$$

Consider the cluster variable in $\mathcal{A}(\Sigma)$

$$x = \frac{1 + x_1x_2x_3 + 3x_4 + 3x_4^2 + x_4^3}{x_1x_2x_3x_4}$$

which, in the context of [CC06, BMR⁺06], corresponds to the cluster character of the indecomposable representation of Q with dimension vector (1111).

Then

$$\sigma_{x_4,1}(x) = \frac{8}{x_1x_2x_3} + 1 = \frac{2}{x_1} \frac{2}{x_2} \frac{2}{x_3} + 1$$

is the sum of 1 and the cluster monomial in $\mathcal{A}(\Sigma \setminus \{x_4\})$ corresponding to the cluster character of the semisimple representation with dimension vector (111) of $Q \setminus \{4\}$.

Now consider the cluster variable in $\mathcal{A}(\Sigma)$

$$z = \frac{1 + 2x_1x_2x_3 + x_1^2x_2^2x_3^2 + 3x_4 + 3x_1x_2x_3x_4 + 3x_4^2 + x_4^3}{x_1x_2x_3x_4^2}$$

which corresponds to the cluster character of the indecomposable representation of Q with dimension vector (1112).

Then

$$\sigma_{x_4,1}(z) = \frac{2}{x_1} \frac{2}{x_2} \frac{2}{x_3} + 5 + x_1x_2x_3.$$

is a linear combination of three distinct cluster monomials in $\mathcal{A}(\Sigma \setminus \{x_4\})$.

6.5. Simple specialisations in general. In general, we expect that simple specialisations of cluster variables to 1 induce rooted cluster morphisms.

Problem 6.16. *Let $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ be a seed and let $x \in \mathbf{x}$. Then does $\sigma_{x,1}$ induce a surjective ideal rooted cluster morphism $\mathcal{A}(\Sigma) \rightarrow \mathcal{A}(\Sigma \setminus \{x\})$?*

We now prove that simple specialisations preserve upper and lower bounds of cluster algebras in general.

Given a seed $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$, and given an element $x \in \mathbf{ex}$, we denote by $\mu_x(\mathbf{ex})$ the set of exchangeable variables in $\mu_x(\Sigma)$ so that $\mathbb{Z}[\mathbf{x} \setminus \mathbf{ex}][\mu_x(\mathbf{ex})^{\pm 1}]$ is the set of Laurent polynomials in the exchangeable variables of $\mu_x(\Sigma)$ with polynomial coefficients in the frozen variables of $\mu_x(\Sigma)$. We also set

$$\mathbf{ex}' = \bigcup_{x \in \mathbf{ex}} \mu_x(\mathbf{ex})$$

to be the set of all the exchangeable variables in the seeds obtained from Σ by applying exactly one mutation.

Following [BFZ05], we set :

Definition 6.17. (1) The lower bound of $\mathcal{A}(\Sigma)$ is

$$\mathcal{L}(\Sigma) = \mathbb{Z}[\mathbf{x} \setminus \mathbf{ex}][\mathbf{ex} \cup \mathbf{ex}'].$$

(2) The upper bound of $\mathcal{A}(\Sigma)$ is

$$\mathcal{U}(\Sigma) = \mathbb{Z}[\mathbf{x} \setminus \mathbf{ex}][\mathbf{ex}^{\pm 1}] \cap \bigcap_{x \in \mathbf{ex}} \mathbb{Z}[\mathbf{x} \setminus \mathbf{ex}][\mu_x(\mathbf{ex})^{\pm 1}].$$

These are subalgebras of \mathcal{F}_Σ and we always have the inclusions $\mathcal{L}(\Sigma) \subset \mathcal{A}(\Sigma) \subset \mathcal{U}(\Sigma)$.

Proposition 6.18. *Let $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ be a seed and let $x \in \mathbf{x}$. Then :*

- (1) $\sigma_{x,1}(\mathcal{L}(\Sigma)) \subset \mathcal{L}(\Sigma \setminus \{x\})$;
- (2) $\sigma_{x,1}(\mathcal{U}(\Sigma)) \subset \mathcal{U}(\Sigma \setminus \{x\})$.

Proof. If x is frozen, the result is clear. Therefore, we fix some exchangeable variable x in \mathbf{ex} . In order to simplify the notations, we set $\sigma = \sigma_{x,1}$.

Let z be a cluster variable in Σ . If z is frozen, then $\sigma(z) = z$ so that $\sigma(z)$ is both in the upper and in the lower bounds of $\mathcal{A}(\Sigma \setminus \{x\})$. Assume now that z is exchangeable. If $z \neq x$, we have

$$\begin{aligned} \sigma(\mu_{z,\Sigma}(z)) &= \sigma \left(\frac{1}{z} \left(\prod_{\substack{b_{yz} > 0 \\ y \in \mathbf{x}}} y^{b_{yz}} + \prod_{\substack{b_{yz} < 0 \\ y \in \mathbf{x}}} y^{-b_{yz}} \right) \right) \\ &= \frac{1}{z} \left(\prod_{\substack{b_{yz} > 0 \\ y \in \mathbf{x} \setminus \{x\}}} y^{b_{yz}} + \prod_{\substack{b_{yz} < 0 \\ y \in \mathbf{x} \setminus \{x\}}} y^{-b_{yz}} \right) \\ &= \mu_{z,\Sigma \setminus \{x\}}(z). \end{aligned}$$

If $z = x$, we get

$$\sigma(\mu_{x,\Sigma}(x)) = \sigma \left(\frac{1}{x} \left(\prod_{\substack{b_{xy} > 0 \\ y \in \mathbf{x}}} y^{b_{xy}} + \prod_{\substack{b_{xy} < 0 \\ y \in \mathbf{x}}} y^{-b_{xy}} \right) \right) = \prod_{\substack{b_{xy} > 0 \\ y \in \mathbf{x} \setminus \{x\}}} y^{b_{xy}} + \prod_{\substack{b_{xy} < 0 \\ y \in \mathbf{x} \setminus \{x\}}} y^{-b_{xy}}.$$

It easily follows that $\sigma(\mathcal{L}(\Sigma)) \subset \mathcal{L}(\Sigma \setminus \{x\})$ and $\sigma(\mathcal{U}(\Sigma)) \subset \mathcal{U}(\Sigma \setminus \{x\})$. \square

6.6. An example of multiple specialisation in zero. For any $m \geq 4$, we denote by $\text{Gr}_2(m)$ the set of planes in \mathbb{C}^m and let $\mathbb{C}[\text{Gr}_2(m)]$ denote its ring of homogeneous coordinates. It is known that

$$\mathbb{C}[\text{Gr}_2(m)] \simeq \mathcal{A}(\Sigma_m) \otimes_{\mathbb{Z}} \mathbb{C}$$

where Σ_m is the seed constructed in §4.1, see for instance [Sco06] or [GSV10, Chapter 2]. The cluster variables in $\mathcal{A}(\Sigma_m)$ are identified with the Plücker coordinates $x_{k,l}$, with $1 \leq k < l \leq m$ in $\mathbb{C}[\text{Gr}_2(m)]$ in such a way that the Plücker coordinate $x_{k,l}$ corresponds to the arc joining k to l in Π_m , see [loc. cit.].

Let m, m' be integers such that $4 \leq m' \leq m$. The choice of an inclusion of $\mathbb{C}^{m'}$ into \mathbb{C}^m induces an embedding $\iota : \text{Gr}_2(m') \rightarrow \text{Gr}_2(m)$ and thus an epimorphism of \mathbb{C} -algebras :

$$\iota^* : \mathbb{C}[\text{Gr}_2(m)] \rightarrow \mathbb{C}[\text{Gr}_2(m')].$$

If $x'_{k,l}$, with $1 \leq k < l \leq m'$, are the Plücker coordinates on $\mathbb{C}[\text{Gr}_2(m')]$, then the morphism ι^* is given by

$$\iota^*(x_{k,l}) = \begin{cases} x'_{k,l} & \text{if } l \leq m' \\ 0 & \text{if } l > m' \end{cases}$$

for $1 \leq k < l \leq m$.

Proposition 6.19. *There exists a unique rooted cluster morphism $\pi_{m,m'} : \mathcal{A}(\Sigma_m) \rightarrow \mathcal{A}(\Sigma_{m'})$ such that $\iota^* = \pi \otimes_{\mathbb{Z}} \mathbb{1}_{\mathbb{C}}$.*

Proof. Consider the ring homomorphism $\pi_{m,m'} : \mathcal{F}_{\Sigma_m} \rightarrow \mathcal{F}_{\Sigma_{m'}}$ acting as ι^* on the Plücker coordinates. Then π defines a ring homomorphism from $\mathcal{A}(\Sigma_m)$ to $\mathcal{A}(\Sigma_{m'})$. Moreover, $\pi_{m,m'}$ satisfies **(CM1)** and **(CM2)** by construction.

Exchange relations in $\mathcal{A}(\Sigma_m)$ are given by

$$x_{ij}x_{kl} = x_{ik}x_{jl} + x_{il}x_{jk}$$

for any i, j, k, l such that $1 \leq i < k < j < l \leq m$ and similarly for $\mathcal{A}(\Sigma_{m'})$. Thus, π sends exactly the exchange relations involving only $x_{k,l}$ with $1 \leq i < k < j < l \leq m'$ in $\mathcal{A}(\Sigma_m)$ to the same exchange relations for $x'_{k,l}$ with $1 \leq i < k < j < l \leq m'$ in $\mathcal{A}(\Sigma_{m'})$. In other words, π commutes with mutations along biadmissible sequences and thus satisfies **(CM3)**.

For uniqueness, it is enough to observe that if such a morphism $\pi_{m,m'}$ exists then it necessarily acts as ι^* on the Plücker coordinates and thus it is unique. \square

7. SURGERY

In this section we introduce a combinatorial procedure, called *cutting*, which turns out to be the inverse process of the amalgamated sums considered in Section 4.2. More precisely, these cuttings provide epimorphisms in **Clus** which are retractions of the monomorphisms constructed from amalgamated sums of rooted cluster algebras.

7.1. Cutting along separating families of variables.

Definition 7.1 (Separating families). Let $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ be a seed. If there exist a subset $\Delta \subset (\mathbf{x} \setminus \mathbf{ex})$ and a partition $\mathbf{x} = \mathbf{x}_1 \sqcup \mathbf{x}_2 \sqcup \Delta$ such that, with respect to this partition, the exchange matrix B is of the form

$$B = \left[\begin{array}{c|c|c} B_{11}^1 & 0 & B_{12}^1 \\ \hline 0 & B_{11}^2 & B_{12}^2 \\ \hline B_{21}^1 & B_{21}^2 & B_{\Delta} \end{array} \right],$$

then we say that Δ separates \mathbf{x}_1 and \mathbf{x}_2 in Σ .

For $j \in \{1, 2\}$, we set $d_{\Delta}^j \Sigma = (\mathbf{x}_j \sqcup \Delta, \mathbf{ex} \cap \mathbf{x}_j, B^j)$ where

$$B^j = \begin{bmatrix} B_{11}^j & B_{12}^j \\ B_{21}^j & B_{22}^j \end{bmatrix}.$$


Definition 7.2 (Cutting). Let $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$ be a seed and let Δ be a separating family of variables as above. The *cutting of Σ along Δ* is the pair

$$d_{\Delta} \Sigma = (d_{\Delta}^1 \Sigma, d_{\Delta}^2 \Sigma).$$


Example 7.3. Consider for instance the matrix

$$B = \left[\begin{array}{cc|cc|c} 0 & 1 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 & 1 \\ \hline 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & -1 & 0 & 1 \\ \hline 1 & -1 & 1 & -1 & 0 \end{array} \right]$$

corresponding to the quiver



with point 5 frozen. Then cutting along 5 gives two oriented 3-cycles with one frozen point each :



The following lemmata prove that the cutting is the inverse operation to the amalgamated sum of seeds.

Lemma 7.4. *Let Σ_1 and Σ_2 be seeds which are glueable along Δ_1 and Δ_2 and let $\Sigma = \Sigma_1 \amalg_{\Delta_1, \Delta_2} \Sigma_2$. We denote by Δ the subset of the cluster of Σ corresponding to Δ_1 and Δ_2 . Then :*

- (1) Δ separates $\mathbf{x}_1 \setminus \Delta$ and $\mathbf{x}_2 \setminus \Delta$ in Σ ;
- (2) $d_\Delta^i \Sigma \simeq \Sigma_i$ for any $i \in \{1, 2\}$.

Proof. By definition, we have $\Sigma = \Sigma_1 \amalg_{\Delta_1, \Delta_2} \Sigma_2 = (\mathbf{x}, \mathbf{ex}, B)$ where $\mathbf{x} = (\mathbf{x}_1 \setminus \Delta_1) \sqcup (\mathbf{x}_2 \setminus \Delta_2) \sqcup \Delta$, $\mathbf{ex} = \mathbf{ex}_1 \sqcup \mathbf{ex}_2$ and

$$B = \left[\begin{array}{c|c|c} B_{11}^1 & 0 & B_{12}^1 \\ \hline 0 & B_{11}^2 & B_{12}^2 \\ \hline B_{21}^1 & B_{21}^2 & B_\Delta \end{array} \right].$$

Therefore, for any $i = 1, 2$, the cluster of $d_\Delta^i \Sigma$ is $\mathbf{x}_i \setminus \Delta_i \sqcup \Delta$, the exchangeable variables in this cluster are the exchangeable variables in Σ_i (none of them belongs to Δ by assumption) and the exchange matrix of this seed is

$$B^i = \left[\begin{array}{c|c} B_{11}^i & B_{12}^i \\ \hline B_{21}^i & B_\Delta \end{array} \right].$$

Therefore, $d_\Delta^i \Sigma \simeq \Sigma_i$. □

Conversely :

Lemma 7.5. *Let Σ be a seed and Δ a separating family of variables in Σ . Then $d_\Delta^1 \Sigma$ and $d_\Delta^2 \Sigma$ are glueable along the respective images Δ_1 and Δ_2 of Δ and*

$$d_\Delta^1 \Sigma \amalg_{\Delta_1, \Delta_2} d_\Delta^2 \Sigma \simeq \Sigma.$$

Proof. We write $\Sigma = (\mathbf{x}, \mathbf{ex}, B)$. Since Δ is separating in Σ , there exists a partition $\mathbf{x} = \mathbf{x}_1 \sqcup \mathbf{x}_2 \sqcup \Delta$ such that, adapted to this partition, B is given by

$$B = \left[\begin{array}{c|c|c} B_{11}^1 & 0 & B_{12}^1 \\ \hline 0 & B_{11}^2 & B_{12}^2 \\ \hline B_{21}^1 & B_{21}^2 & B_\Delta \end{array} \right]$$

and since Δ consists of frozen variables, $\mathbf{ex} = (\mathbf{ex} \cap \mathbf{x}_1) \sqcup (\mathbf{ex} \cap \mathbf{x}_2)$.

By definition, for $i = 1, 2$, we have $d_\Delta^i \Sigma = \Sigma_i$ where

$$\Sigma_i = (\mathbf{x}_i \sqcup \Delta, \mathbf{ex} \cap \mathbf{x}_i, B^i)$$

with

$$B^i = \left[\begin{array}{c|c} B_{11}^i & B_{12}^i \\ \hline B_{21}^i & B_\Delta \end{array} \right].$$

It follows that Σ_1 and Σ_2 are glueable along Δ , Δ and thus

$$\Sigma_1 \amalg_{\Delta, \Delta} \Sigma_2 \simeq (\mathbf{x}, \mathbf{ex}, B) = \Sigma.$$

□

7.2. Epimorphisms from cuttings. Let Σ be a seed and Δ a separating family of variables in Σ as above and set $\Sigma_i = d_\Delta^i \Sigma$ for any $i \in \{1, 2\}$.

Fix $i \in \{1, 2\}$. It follows from Lemmata 7.4, 7.5 and 4.13 that we have canonical monomorphisms in **Clus** :

$$j_i : \mathcal{A}(\Sigma_i) \longrightarrow \mathcal{A}(\Sigma).$$

Consider the ring homomorphism :

$$\text{pr}_i : \begin{cases} \mathcal{F}_\Sigma & \longrightarrow \mathcal{F}_{\Sigma_i} \\ x & \mapsto x \text{ if } x \in \mathbf{x}_i \sqcup \Delta, \\ x & \mapsto 0 \text{ if } x \in \mathbf{x}_j \text{ for } i \neq j. \end{cases}$$

Proposition 7.6. *For any $i \in \{1, 2\}$, the ring homomorphism pr_i induces a rooted cluster epimorphism $\mathcal{A}(\Sigma) \longrightarrow \mathcal{A}(\Sigma_i)$ which is a retraction for j_i .*

Proof. We first observe that

$$\text{pr}_i(\mathbf{x}) \subset \mathbf{x}_i \sqcup \Delta \sqcup \{0\} \text{ and } \text{pr}_i(\mathbf{ex}) \subset (\mathbf{ex} \cap \mathbf{x}_i) \sqcup \{0\}$$

so that pr_i satisfies **(CM1)** and **(CM2)**.

In order to prove that pr_i satisfies **(CM3)**, we prove as in Lemma 4.13 that the $(\text{pr}_i, \Sigma, \Sigma_i)$ -biadmissible sequences are precisely the Σ_i -admissible sequences and that Δ remains separating along biadmissible mutations. It follows as in the proof of Lemma 4.13 that pr_i commutes with biadmissible mutations and thus satisfies **(CM3)** and induces a surjective rooted cluster morphism $\mathcal{A}(\Sigma) \longrightarrow \mathcal{A}(\Sigma_i)$.

Finally, as $(\text{pr}_i \circ j_i)(x) = x$ for any $x \in \mathbf{x}_i \sqcup \Delta$, we get $\text{pr}_i \circ j_i = \mathbb{1}_{\mathcal{A}(\Sigma_i)}$. □

In general, we state the following problem :

Problem 7.7. *Determine which monomorphisms in **Clus** are sections.*


7.3. Topological interpretation of the surgery. Let (S, M) be a marked surface. Assume that there exists a collection Δ of (internal or boundary) arcs in (S, M) which can be concatenated in order to form a simple closed curve in (S, M) which delimits two non-degenerate marked subsurfaces (S_1, M_1) and (S_2, M_2) in (S, M) .

Consider a triangulation T of (S, M) containing Δ as a subset. Then T induces two triangulations T_1 and T_2 of (S_1, M_1) and (S_2, M_2) respectively in which Δ corresponds to a set of boundary arcs, that is, to frozen variables. We denote by Σ the seed corresponding to the triangulation T and by Σ_1, Σ_2 the seeds corresponding respectively to the triangulations T_1 and T_2 . Then Δ is a separating family of variables in Σ and


$$d_\Delta(\Sigma) = (\Sigma_1, \Sigma_2).$$

In other words, cutting Σ along Δ coincides with taking the seed associated with the triangulation of the surface obtained by cutting (S, M) along Δ .

Example 7.8. Consider the following triangulation of the disc with two marked points on the boundary and five punctures.



Let Δ denote the union of the four arcs which are dashed in the above figure. Cutting the surface along Δ gives two new marked surfaces, namely an unpunctured annulus with two marked points on a boundary and four marked points on the other and a disc with four marked points on the boundary and one puncture. The triangulation of the above disc containing Δ thus induces triangulations of the two cut surfaces.



Then we clearly see that the cluster algebra associated with the initial surface where the arcs in Δ are frozen is the amalgamated sum over the respective images of Δ of the two cut surfaces.

ACKNOWLEDGEMENTS

The first author gratefully acknowledges partial support from NSERC of Canada, FQRNT of Québec and the University of Sherbrooke.

This work was initiated while the second author was a CRM-ISM postdoctoral fellow at the University of Sherbrooke and was finished while he was a postdoctoral fellow at the Université Denis Diderot – Paris 7, funded by the ANR *Geométrie Tropicale et Algèbres Amassées*. The second author would like to thank the third author for his kind hospitality during his stay in University of Connecticut where this project started.

The third author has been supported by the NSF Grant 1001637 and by the University of Connecticut.

The authors wish to thank Bernhard Keller, Yann Palu, Pierre-Guy Plamondon, Jan Schröer and Vasilisa Shramchenko for fruitful discussions on the topic.

REFERENCES

- [ADSS11] Ibrahim Assem, Grégoire Dupont, Ralf Schiffler, and David Smith, *Friezes, strings and cluster variables*, Glasgow Mathematical Journal **54** (2011), no. 1, 27–60.
- [ASS11] Ibrahim Assem, Ralf Schiffler, and Vasilisa Shramchenko, *Cluster automorphisms*, Proc. LMS (2011), in press.
- [BFZ05] Arkady Berenstein, Sergey Fomin, and Andrei Zelevinsky, *Cluster algebras III: Upper bounds and double Bruhat cells*, Duke Mathematical Journal **126** (2005), no. 1, 1–52.
- [BMR⁺06] Aslak Bakke Buan, Robert Marsh, Markus Reineke, Idun Reiten, and Gordana Todorov, *Tilting theory and cluster combinatorics*, Adv. Math. **204** (2006), no. 2, 572–618.
- [BMR08] Aslak Bakke Buan, Robert Marsh, and Idun Reiten, *Cluster mutation via quiver representations*, Commentarii Mathematici Helvetici **83** (2008), no. 1, 143–177.
- [BZ11] Thomas Brüstle and Jie Zhang, *On the cluster category of a marked surface*, Algebra and Number Theory **to appear** (2011).
- [CC06] Philippe Caldero and Frédéric Chapoton, *Cluster algebras as Hall algebras of quiver representations*, Commentarii Mathematici Helvetici **81** (2006), 596–616.
- [CK06] Philippe Caldero and Bernhard Keller, *From triangulated categories to cluster algebras II*, Annales Scientifiques de l’Ecole Normale Supérieure **39** (2006), no. 4, 83–100.
- [Dem11] Laurent Demonet, *Categorification of skew-symmetrizable cluster algebras*, Algebras and Representation Theory **14** (2011), 1087–1162, 10.1007/s10468-010-9228-4.
- [DP11] Grégoire Dupont and Frédéric Palesi, *Quasi-cluster algebras associated to non-orientable surfaces*, arXiv:1105.1560v1 [math.RA] (2011).
- [DWZ10] Harm Derksen, Jerzy Weyman, and Andrei Zelevinsky, *Quivers with potentials and their representations II: applications to cluster algebras*, J. Amer. Math. Soc. **23** (2010), no. 3, 749–790.

- [FK10] Changjian Fu and Bernhard Keller, *On cluster algebras with coefficients and 2-Calabi-Yau categories*, Trans. Amer. Math. Soc. **362** (2010), 859–895.
- [FST08] Sergey Fomin, Michael Shapiro, and Dylan Thurston, *Cluster algebras and triangulated surfaces. I. Cluster complexes*, Acta Math. **201** (2008), no. 1, 83–146.
- [FZ02] Sergey Fomin and Andrei Zelevinsky, *Cluster algebras I: Foundations*, J. Amer. Math. Soc. **15** (2002), 497–529.
- [FZ03] ———, *Cluster algebras II: Finite type classification*, Inventiones Mathematicae **154** (2003), 63–121.
- [FZ07] ———, *Cluster algebras IV: Coefficients*, Compositio Mathematica **143** (2007), no. 1, 112–164.
- [GLS10] Christof Geiss, Bernard Leclerc, and Jan Schröer, *Generic bases for cluster algebras and the Chamber Ansatz*, arXiv:1004.2781v2 [math.RT] (2010).
- [GLS11] Christof Geiss, Bernard Leclerc, and Jan Schröer, *Factorial cluster algebras*, arXiv:1110.1199v2 [math.RA] (2011).
- [GSV10] Michael Gekhtman, Michael Shapiro, and Alek Vainshtein, *Cluster algebras and Poisson geometry*, Mathematical Surveys and Monographs, vol. 167, American Mathematical Society, Providence, RI, 2010.
- [HL10] David Hernandez and Bernard Leclerc, *Cluster algebras and quantum affine algebras*, Duke Math. J. **154** (2010), no. 2, 265–341.
- [IY08] Osamu Iyama and Yuji Yoshino, *Mutation in triangulated categories and rigid Cohen-Macaulay modules*, Invent. Math. **172** (2008), no. 1, 117–168.
- [Kel10] Bernhard Keller, *Cluster algebras, quiver representations and triangulated categories*, Triangulated categories, London Math. Soc. Lecture Note Ser., vol. 375, Cambridge Univ. Press, Cambridge, 2010, pp. 76–160.
- [KR08] Bernhard Keller and Idun Reiten, *Acyclic Calabi-Yau categories*, Compositio Mathematicae **144** (2008), no. 5, 1332–1348.
- [Mas77] William S. Massey, *Algebraic topology: an introduction*, Springer-Verlag, New York, 1977, Reprint of the 1967 edition, Graduate Texts in Mathematics, Vol. 56.
- [MSW11] Gregg Musiker, Ralf Schiffler, and Lauren Williams, *Positivity for cluster algebras from surfaces*, Adv. Math. **227** (2011), 2241–2308.
- [MW11] Gregg Musiker and Lauren Williams, *Matrix formulae and skein relations for cluster algebras from surfaces*, arXiv:1108.3382v1 [math.CO] (2011).
- [Pal08] Yann Palu, *Cluster characters for 2-Calabi-Yau triangulated categories*, Ann. Inst. Fourier (Grenoble) **58** (2008), no. 6, 2221–2248.
- [Pal11] ———, *Cluster characters II: A multiplication formula*, Proc. LMS to appear (2011).
- [Pla11a] Pierre-Guy Plamondon, *Catégories amassées aux espaces de morphismes de dimension infinie, applications*, Ph.D. thesis, Université Paris 7, 2011.
- [Pla11b] Pierre-Guy Plamondon, *Cluster algebras via cluster categories with infinite-dimensional morphism spaces*, Compos. Math. (2011), to appear.
- [Sco06] Joshua Scott, *Grassmannians and cluster algebras*, Proc. London Math. Soc. (3) **92** (2006), no. 2, 345–380.
- [ST09] Ralf Schiffler and Hugh Thomas, *On cluster algebras arising from unpunctured surfaces*, Int. Math. Res. Not. (2009), no. 17, 3160–3189.
- [War11] Matthias Warkentin, *Subquivers of mutation-acyclic quivers are mutation-acyclic*, arXiv:1102.3382v2 [math.RT] (2011).

UNIVERSITÉ DE SHERBROOKE, SHERBROOKE QC, CANADA
E-mail address: ibrahim.assem@usherbrooke.ca

INSTITUT DE MATHÉMATIQUES DE JUSSIEU-PARIS RIVE GAUCHE, PARIS, FRANCE
E-mail address: dupontg@math.jussieu.fr

UNIVERSITY OF CONNECTICUT, STORRS CT, USA
E-mail address: schiffler@math.uconn.edu