



HAL
open science

Structural definition of a novel CD4-induced epitope that is targeted by a single-headed immunoglobulin to effect broad and potent HIV neutralization

Priyamvada Acharya, Timothy S. Luongo, Julie Matz, Stephen D. Schmidt, Gwo-Yu Chuang, Ivelin Georgiev, Pascal Kessler, Yongping Yang, Patrick Chames, Loïc M Martin, et al.

► To cite this version:

Priyamvada Acharya, Timothy S. Luongo, Julie Matz, Stephen D. Schmidt, Gwo-Yu Chuang, et al.. Structural definition of a novel CD4-induced epitope that is targeted by a single-headed immunoglobulin to effect broad and potent HIV neutralization. *Retrovirology*, 2012, 9 (Suppl 2), pp.P346. inserm-00731776

HAL Id: inserm-00731776

<https://inserm.hal.science/inserm-00731776>

Submitted on 13 Sep 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



POSTER PRESENTATION

Open Access

Structural definition of a novel CD4-induced epitope that is targeted by a single-headed immunoglobulin to effect broad and potent HIV neutralization

P Acharya^{1*}, TS Luongo¹, J Matz², SD Schmidt¹, G Chuang¹, I Georgiev¹, P Kessler³, Y Yang¹, P Chames⁴, L Martin³, JR Mascola¹, PD Kwong¹

From AIDS Vaccine 2012
Boston, MA, USA. 9-12 September 2012

Background

HIV-1 enters cells by sequentially binding the CD4 receptor and a coreceptor, either CCR5 or CXCR4. Functional constraints result in a high degree of conservation of the receptor-binding sites making them potential targets for intervention. The coreceptor-binding site on HIV-1 envelope gp120 glycoprotein is protected from the humoral immune system by conformational masking and steric occlusion. The site becomes available after a conformational change in gp120 following CD4 engagement, but at that point in the entry process, the proximity of the viral and cellular membranes makes the site inaccessible to bulky antibody molecules. Thus, in spite of being highly conserved, this region has not been considered a viable vaccine target.

Methods

Single domain antibody vHH120.4 was isolated from llama immunized with gp120 covalently linked to a CD4-mimetic peptide. Full-length versions of this antibody (IgG2B and IgG3) were created and tested for neutralization. The structure of vHH120.4 bound to gp120 from the HIV-1 YU2 strain was determined at 2.1 Å resolution.

Results

Both IgG2B and IgG3 versions of the vHH120.4 potently neutralized over 95% of a panel of circulating Tier 2 HIV-1 isolates. Structural analyses of vHH120.4 bound to gp120 revealed a novel CD4i epitope that involves

antibody interactions with region on gp120 encompassing the bridging sheet and the base of the V3 loop, the β19 strand, the CD4-binding loop, and the glycan at Asn 386. This epitope overlaps the classically defined CD4-induced epitopes recognized by antibodies 17b and 48d, but is shifted towards the site of CD4 attachment.

Conclusion

The discovery of a neutralizing CD4-induced epitope indicates that not all CD4-induced sites are masked from neutralization. Whether human antibodies can also utilize the newly defined vHH120.4 epitope for effective neutralization remains to be determined; HIV-1 envelope probes designed to specifically select antibodies targeting this epitope are now being developed.

Author details

¹National Institutes of Health, Bethesda, MD, USA. ²Inserm, Marseille, France. ³CEA/iBiTecs, Gif sur Yvette, France. ⁴INSERM, Marseille, France.

Published: 13 September 2012

doi:10.1186/1742-4690-9-S2-P346

Cite this article as: Acharya et al.: Structural definition of a novel CD4-induced epitope that is targeted by a single-headed immunoglobulin to effect broad and potent HIV neutralization. *Retrovirology* 2012 **9**(Suppl 2):P346.

¹National Institutes of Health, Bethesda, MD, USA
Full list of author information is available at the end of the article