



HAL
open science

Boundary behaviour of harmonic functions on hyperbolic manifolds

Camille Petit

► **To cite this version:**

Camille Petit. Boundary behaviour of harmonic functions on hyperbolic manifolds. 2013. hal-00794012

HAL Id: hal-00794012

<https://hal.science/hal-00794012>

Preprint submitted on 24 Feb 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

BOUNDARY BEHAVIOUR OF HARMONIC FUNCTIONS ON HYPERBOLIC MANIFOLDS

CAMILLE PETIT

ABSTRACT. Let M be a complete simply connected manifold which is in addition Gromov hyperbolic, coercive and roughly starlike. For a given harmonic function on M , a local Fatou Theorem and a pointwise criteria of non-tangential convergence coming from the density of energy are shown: at almost all points of the boundary, the harmonic function converges non-tangentially if and only if the supremum of the density of energy is finite. As an application of these results, a Calderón-Stein Theorem is proved, that is, the non-tangential properties of convergence, boundedness and finiteness of energy are equivalent at almost every point of the boundary.

CONTENTS

5	1. Introduction	2
	2. Preliminaries	4
	2.1. Gromov hyperbolic spaces	5
	2.2. Roughly starlike manifolds	6
	2.3. Coercive manifolds	6
10	2.4. Comments	7
	3. Brownian motion and conditioning	7
	3.1. Brownian motion	7
	3.2. Conditioning	8
	3.3. Stochastic convergence	8
15	4. Harnack inequalities	8
	5. Non-tangential behaviour of Brownian motion	10
	5.1. A geometric Lemma	10
	5.2. Behaviour of Green functions	13
	5.3. Brownian motion and non-tangential sets	15
20	6. Local Fatou theorem	16
	7. Density of energy	17
	References	19

2010 *Mathematics Subject Classification.* Primary 31C05; 05C81; Secondary 60J45; 60D05; 60J50.

Key words and phrases. Harmonic functions, Gromov hyperbolic manifolds, Brownian motion, boundary at infinity, Fatou's theorem, non-tangential convergence.

1. INTRODUCTION

The interplay between the geometry of a complete Riemannian manifold M and the existence of non-constant harmonic functions on M has been studied by researcher in geometric analysis for decades. On one hand, S.T. Yau [Yau75] proved that, on a complete Riemannian manifold M of non-negative Ricci curvature, every positive harmonic function is constant. On the other hand, in the middle of the eighties, M.T. Anderson and R. Schoen [AS85] provided a complete description of the space of non-negative harmonic functions for manifolds of pinched negative curvature. They proved the identification between the sphere at infinity and the Martin boundary, that is, the boundary allowing an integral representation of non-negative harmonic functions by measures on the boundary. Major contributions to this latter issue were given by A. Ancona in a series of papers [Anc87, Anc88, Anc90] and will be discussed later in this article.

The study of non-tangential convergence of harmonic functions goes back to 1906 with P. Fatou's seminal paper [Fat06], where the following result is proved:

Any positive harmonic function on the unit disc admits non-tangential limits at almost every point θ of the boundary circle.

Recall that a function is said to *converge non-tangentially at θ* if it has a finite limit at θ on every non-tangential cone with vertex θ . Fatou type theorems have been proved in many different contexts since then and in particular on Gromov hyperbolic graphs and manifolds [Anc88, Anc90]. One of the issues is to replace the global positivity condition by local criteria of non-tangential convergence. Of special interest are two criteria which have been intensively studied: the criterion of non-tangential boundedness [Pri16, Cal50b] and the criterion of finiteness of non-tangential integral area [MZ38, Spe43, Cal50a, Ste61], also called the criteria of Lusin area. A function is *non-tangentially bounded at a boundary point θ* if it is bounded on every non-tangential cone with vertex θ . Similarly, a function is of *finite non-tangential integral area at θ* if on every non-tangential cone with vertex θ , it has a finite area integral. Calderón-Stein's Theorem ([Cal50b, Ste61]) asserts that for a harmonic function in the Euclidean half-space, notions of non-tangential convergence, non-tangential boundedness and finiteness of non-tangential area integral coincide at almost all points of the boundary. These results were reproved by J. Brossard ([Bro78]) using Brownian motion. J. Brossard also stated in [Bro88] the criterion of the density of the area integral, a notion first introduced by R.F. Gundy [Gun83].

As noticed by A. Korányi, hyperbolic spaces provide a convenient framework for studying Calderón-Stein like results. A first reason is that there should exist a lot of non-constant harmonic functions under negative curvature assumptions, whereas they are in some sense rare on manifolds satisfying non-negative curvature assumptions. Another reason is that several notions have simpler and more natural expressions in the setting of hyperbolic spaces. When we equip the Euclidean half-space with the hyperbolic Poincaré metric, Euclidean notions of non-tangential cone with vertex θ :

$$\Gamma_\alpha^\theta := \{(x, y) \in \mathbb{R}^\nu \times \mathbb{R}_+ \mid |x - \theta| < ay < a\},$$

of non-tangential area integral and of density of area integral:

$$\int_{\Gamma_\alpha^\theta} |\nabla u(x, y)|^2 y^{1-\nu} dx dy \quad \text{and} \quad \frac{1}{2} \int_{\Gamma_\alpha^\theta} y^{1-\nu} \Delta |u - r|(dx dy),$$

turn out to be respectively tubular neighborhoods of geodesic rays starting at a base point o :

$$\Gamma_c^\theta := \{z \mid \exists \gamma \text{ a geodesic ray from } o \text{ to } \theta \text{ such that } d(z, \gamma) < c\},$$

a true energy and a density of energy:

$$J_c^\theta := \int_{\Gamma_c^\theta} |\nabla u|^2 d\nu \quad \text{and} \quad D_c^r(\theta) := -\frac{1}{2} \int_{\Gamma_c^\theta} \Delta |u - r|(dx).$$

Following this philosophy, Calderón-Stein's result was extended to the framework of Riemannian manifolds of pinched negative curvature ([Mou95]) and trees ([Mou00, AP08, Mou10, Pic10]). The criterion introduced by J. Brossard of the density of area integral was also addressed in [Mou07] for Riemannian manifolds of pinched negative curvature. The present paper was motivated by the question whether these kinds of results also hold for Gromov hyperbolic spaces. In a recent paper [Pet12], we proved the criteria of non-tangential boundedness in the framework of Gromov hyperbolic graphs. The aim of this paper is to deal with the different criteria presented above in the case of Gromov hyperbolic manifolds.

Let us now describe the main results of this paper. We first introduce briefly the geometric setting. The geometric notions will be defined precisely in section 2. We say that a Riemannian manifold M is *roughly-starlike* if there exist a constant $K \geq 0$ and a base point $o \in M$ such that every point $x \in M$ is within a distance at most K from a geodesic ray starting at o . A Riemannian manifold M of dimension n has *bounded local geometry* provided about each $x \in M$, there is a geodesic ball $B(x, r)$ (with r independent of x) and a diffeomorphism $F : B(x, r) \rightarrow \mathbb{R}^n$ with

$$\frac{1}{c} \cdot d(y, z) \leq \|F(y) - F(z)\| \leq c \cdot d(y, z)$$

for all $y, z \in B(x, r)$, where c is independent of x . It is worth mentioning that M has bounded local geometry if M has Ricci curvature and injectivity radius bounded from below. Following the terminology of A. Ancona, M is called *coercive* if it has bounded local geometry and if the bottom $\lambda_1(M)$ of the spectrum is positive. We say that a complete, simply connected Riemannian manifold M satisfies condition (\clubsuit) if in addition M is coercive, roughly starlike and Gromov hyperbolic.

We first prove a local Fatou Theorem for Riemannian manifolds satisfying conditions (\clubsuit). Let U be an open set in M and let us denote by ∂M the geometric boundary of M . We say that a point $\theta \in \partial M$ is *tangential for U* if for all $c > 0$, the set $\Gamma_c^\theta \setminus U$ is bounded.

Theorem 1.1. *Let M be a manifold satisfying conditions (\clubsuit) and let U be an open subset of M . If u is a non-negative harmonic function on U , then for μ -almost all θ that is tangential for U , the function u converges non-tangentially at θ .*

The measure μ on ∂M is the *harmonic measure*. The proof follows the approach by F. Mouton [Mou07]. In Mouton's proof, geometry comes in at some key points by use of comparison Theorems in pinched negative curvature. As a Corollary of Theorem 1.1, we deduce that a harmonic function converges non-tangentially at almost all points where it is non-tangentially bounded from below (or above).

Corollary 1.2. *Let M be a manifold satisfying conditions (\clubsuit) and let u be a harmonic function on M . Then, for μ -almost all $\theta \in \partial M$, the following properties are equivalent:*

- (1) *The function u converges non-tangentially at θ .*
- (2) *The function u is non-tangentially bounded from below at θ .*
- 90 (3) *There exists $c > 0$ such that u is bounded from below on Γ_c^θ .*

Then we focus on the density of energy. In [Bro88], J. Brossard proved that for a harmonic function u on the Euclidean half-space, at almost all points of the boundary, u converges non-tangentially if and only if the supremum over $r \in \mathbb{R}$ of the density of area integral on the level set $\{u = r\}$ is finite. In [Mou07], F. Mouton
 95 proved that for a harmonic function u on a Riemannian manifold of pinched negative curvature, u converges non-tangentially at almost all points of the boundary where the density of energy on the level set $\{u = 0\}$ is finite, providing a partial geometric analogue of Brossard's Theorem. We focus here on Gromov hyperbolic manifolds and prove an analogue for the density of energy of Brossard's result. It generalizes
 100 and strengthens Mouton's Theorem.

Theorem 1.3. *Let M be a manifold satisfying conditions (\clubsuit) , $c > 0$, and let u be a harmonic function on M . Then, for μ -almost all $\theta \in \partial M$, the following properties are equivalent:*

- (1) *The function u converges non-tangentially at θ .*
- 105 (2) $\sup_{r \in \mathbb{R}} D_c^r(\theta) < +\infty$.
- (3) $D_c^0(\theta) < +\infty$.

As a Corollary, we prove the criterion of the finiteness of the non-tangential energy.

Corollary 1.4. *Let M be a manifold satisfying conditions (\clubsuit) and let u be a harmonic function on M . Then, for μ -almost all $\theta \in \partial M$, the following properties are equivalent:*

- (1) *The function u converges non-tangentially at θ .*
- (2) *The function u has finite non-tangential energy at θ .*

Corollaries 1.2 and 1.4 together yield in particular the Calderón-Stein Theorem
 115 on Gromov hyperbolic manifolds.

This paper is organized as follows. In section 2, we present the geometric framework of the results, recalling briefly some properties of Gromov hyperbolic metric spaces, the definition of a roughly starlike manifold and of a coercive manifold. In section 3 we discuss some basics related to Brownian motions needed later on, in
 120 particular the martingale property and the Doob's h-process method, allowing to condition Brownian motion to exit the manifold at a fixed point of the boundary. In section 4, we recall the different Harnack inequalities needed later on. Section 5 is devoted to the proofs of several lemmas ensuing Harnack inequalities and crucial in the proofs of the main results. Finally, in section 6 we prove Theorem 1.1 and
 125 Corollary 1.2 and in section 7, we prove Theorem 1.3 and Corollary 1.4.

2. PRELIMINARIES

From now on, M denotes a complete simply connected Riemannian manifold of dimension $n \geq 2$ and d denotes the usual Riemannian distance on M . Let Δ denote the Laplace-Beltrami operator on M , by G the associated Green function.
 130 A function $u : M \rightarrow \mathbb{R}$ is called *harmonic* if $\Delta u = 0$. The Green function G is

finite outside the diagonal, positive, symmetric and for every $y \in M$, the function $x \mapsto G(x, y)$ is harmonic on $M \setminus \{y\}$. We will make additional geometric assumptions on M , which will be described in the following paragraphs.

2.1. **Gromov hyperbolic spaces.** Gromov hyperbolic spaces have been introduced by M. Gromov in the 80's (see for instance [Gro81, Gro87]). These spaces are naturally equipped with a geometric boundary. There exists a wide literature on Gromov hyperbolic spaces, see [GDLH90, BH99] for nice introductions. We introduce here only the properties of these spaces which will be used in the following.

Let (X, d) denote a metric space. The *Gromov product* of two points $x, y \in X$ with respect to a basepoint $o \in X$ is defined by

$$(x, y)_o := \frac{1}{2} [d(o, x) + d(o, y) - d(x, y)].$$

Notice that $0 \leq (x, y)_o \leq \min\{d(o, x), d(o, y)\}$ and that if $o' \in X$ is another basepoint, then for every $x, y \in X$,

$$|(x, y)_o - (x, y)_{o'}| \leq d(o, o').$$

Definition 2.1. A metric space (X, d) is called *Gromov hyperbolic* if there exists $\delta \geq 0$ such that for every $x, y, z \in X$ and every basepoint $o \in X$,

$$(2.1) \quad (x, z)_o \geq \min\{(x, y)_o, (y, z)_o\} - \delta.$$

For a real $\delta \geq 0$, we say that (X, d) is δ -*hyperbolic* if inequality (2.1) holds for all $x, y, z, o \in X$.

Remark 2.2. From now on, when considering a Gromov hyperbolic metric space, we will always assume, without loss of generality, that inequality (2.1) holds with δ an integer greater than or equal to 3.

The definition of Gromov hyperbolicity makes sense in every metric space. When the metric space is Gromov hyperbolic and geodesic, the Gromov product $(x, y)_o$ may be seen as a rough measure of the distance between o and any geodesic segment between x and y . More precisely, if γ is a geodesic segment between x and y , we have

$$d(o, \gamma) - 2\delta \leq (x, y)_o \leq d(o, \gamma).$$

We now describe the geometric boundary of a Gromov hyperbolic space. Let (X, d) be a δ -hyperbolic metric space and fix a basepoint $o \in X$. A sequence $(x_i)_i$ in X *converges at infinity* if

$$\lim_{i, j \rightarrow +\infty} (x_i, x_j)_o = +\infty.$$

This condition is independent of the choice of the basepoint. Two sequences $(x_i)_i$ and $(y_j)_j$ converging at infinity are called *equivalent* if $\lim_{i \rightarrow +\infty} (x_i, y_i)_o = +\infty$. This defines an equivalence relation on sequences converging at infinity. The *geometric boundary* ∂X is the set of equivalence classes of sequences converging at infinity. In order to fix an appropriate topology on $\bar{X} := X \cup \partial X$, we extend the Gromov product to the boundary. Let us say that for a point $x \in X$, a sequence $(x_i) \in X^{\mathbb{N}}$ is in the class of x if $x_i \rightarrow x$. We then define

$$(x, y)_o := \sup \liminf_{i, j \rightarrow \infty} (x_i, y_j)_o,$$

where the supremum is taken over all sequences (x_i) in the class of $x \in \overline{X}$ and (y_j) in the class of $y \in \overline{X}$. The inequality

$$(2.2) \quad (x, z)_o \geq \min\{(x, y)_o, (y, z)_o\} - 2\delta$$

holds for every $x, y, z \in \overline{X}$. If in addition (X, d) is geodesic, then for every $x \in X$, $\xi \in \partial X$ and every geodesic ray γ from o to ξ , we have

$$(2.3) \quad d(x, \gamma) - 2\delta \leq (o, \xi)_x \leq d(x, \gamma) + 2\delta.$$

150 For a real $r \geq 0$ and a point $\xi \in \partial X$, denote $V_r(\xi) := \{y \in \overline{X} \mid (\xi, y)_o \geq r\}$. We then equip \overline{X} with the unique topology containing open sets of X and admitting the sets $V_r(\xi)$ with $r \in \mathbb{Q}_+$ as a neighborhood base at any $\xi \in \partial X$. This provides a compactification \overline{X} of X .

2.2. **Roughly starlike manifolds.** We will assume the manifold M to be roughly
155 starlike. From now on, fix a basepoint $o \in M$.

Definition 2.3. A complete Riemannian manifold M is called *roughly starlike with respect to the basepoint $o \in M$* if there exists $K \geq 0$ such that for every point $x \in M$, there exists a geodesic ray γ starting at o and within a distance at most K from x .

We will abbreviate to roughly starlike if there is no risk of ambiguity. Let us
160 notice that if M is δ -hyperbolic and K -roughly starlike with respect to o , then M is K' -roughly starlike with respect to o' , with $K' = K'(d(o, o'), \delta, K)$. The "roughly starlike" assumption has previously been used by A. Ancona [Anc88] and by M. Bonk, J. Heinonen and P. Koskela [BHK01].

165 Recall that a complete manifold M is said to have a *quasi-pole* in a compact set $\Omega \subset M$ if there exists $C > 0$ such that each point of M lies in a C -neighborhood of some geodesic ray emanating from Ω . If M is roughly starlike with respect to o , then M has a quasi-pole at o .

2.3. **Coercive manifolds.** As explained in the introduction, a manifold M of dimension n has *bounded local geometry* provided about each $x \in M$, there is a geodesic ball $B(x, r)$ (with r independent of x) and a diffeomorphism $F : B(x, r) \rightarrow \mathbb{R}^n$ with

$$\frac{1}{c}d(y, z) \leq \|F(y) - F(z)\| \leq cd(y, z)$$

for all $y, z \in B(x, r)$, where c is independent of x .

The manifold M is *coercive* if it has bounded local geometry and if the bottom $\lambda_1(M)$ of the spectrum is positive. Recall that the bottom of the spectrum of the Laplacian Δ is defined by

$$\lambda_1(M) := \inf_{\phi} \frac{\int_M \|\nabla \phi\|^2}{\int_M \phi^2},$$

170 where ϕ ranges over all smooth functions with compact support on M . Notice that for a manifold of bounded local geometry, $\lambda_1(M) > 0$ if and only if its Cheeger constant is positive (see [Bus82]).

175 It is worth mentioning that in general, Gromov hyperbolicity does not imply positivity of the bottom of the spectrum. In [Cao00], J. Cao gave conditions for a Gromov hyperbolic, roughly starlike manifold with bounded local geometry to have positive bottom of the spectrum.

2.4. **Comments.** Recall that we say that a complete, simply connected Riemannian manifold M satisfies condition (\clubsuit) if in addition M is coercive, roughly starlike and Gromov hyperbolic. On one hand, when the manifold M is Gromov hyperbolic, we can consider its geometric boundary as defined above. On the other
 180 hand, we can also consider its Martin boundary, which is natural when dealing with non-negative harmonic functions. In [Anc90], A. Ancona proved that for a manifold satisfying conditions (\clubsuit) (and even without roughly starlike assumption), the geometric compactification and the Martin compactification are homeomorphic.

The following Proposition is a consequence of conditions (\clubsuit) . It yields a uniformity in the behaviour of the Green function G and will be useful in the following.
 185

Proposition 2.4 ([Anc90], page 92). *If M satisfies conditions (\clubsuit) , there exist two positive constants $C_1 = C_1(M)$ and $c_1 = c_1(M)$ such that for every $x, y \in M$ with $d(x, y) \geq 1$, we have*

$$G(x, y) \leq C_1 \exp(-c_1 d(x, y)).$$

3. BROWNIAN MOTION AND CONDITIONING

Following the philosophy of J. Brossard [Bro88], our methods use Brownian motion and the connection between harmonic functions and Brownian motion given by the martingale property. We describe here the "Brownian material" needed in
 190 the proofs.

3.1. **Brownian motion.** The *Brownian motion* (X_t) on M is defined as the diffusion process associated with the Laplace-Beltrami operator Δ . If M satisfies condition (\clubsuit) , Brownian motion is defined for every $t \in \mathbb{R}_+$ ([Anc90], page 60). Choosing $\Omega := \mathcal{C}(\mathbb{R}_+, M)$ as the probability space, for every $t \in \mathbb{R}_+$, X_t is a
 195 random variable on Ω , with values in M , and for every $\omega \in \Omega$, $t \mapsto X_t(\omega)$ is a continuous function, that is a path in M . If we consider Brownian motion starting at a fixed point $x \in M$, we obtain a probability \mathbb{P}_x on Ω .

An important property of the Martin boundary (which in our case coincides with the geometric boundary, see section 2.4) is that for \mathbb{P}_x -almost every trajectory $\omega \in \Omega$, there exists a boundary point $\theta \in \partial M$ such that $\lim_{t \rightarrow +\infty} X_t(\omega) = \theta$. Let us denote by $X_\infty(\omega)$ the ∂M -valued random variable such that Brownian motion converges \mathbb{P}_x -almost surely to X_∞ for all $x \in M$. The *harmonic measure at x* , denoted by μ_x , is the distribution of X_∞ when Brownian motion starts at x . All the measures $\mu_x, x \in M$ on ∂M are equivalent. This gives rise to a notion of μ -negligibility. Defining the *Poisson kernel* $K(x, \theta)$ as limit of the Green kernels $\lim_{y \rightarrow \theta} \frac{G(x, y)}{G(o, y)}$, the Radon-Nikodym derivative of harmonic measure is given by

$$K(x, \theta) = (d\mu_x / d\mu_o)(\theta).$$

The *martingale property* (see [Dur84]) is a crucial tool in our methods: for a function f of class C^2 ,

$$f(X_t) + \frac{1}{2} \int_0^t \Delta f(X_s) ds$$

is a local martingale with respect to probabilities $(\mathbb{P}_x)_x$. Hence if u is harmonic, $(u(X_t))$ is a local martingale.

200 **3.2. Conditioning.** As claimed above, Brownian motion converges almost surely to a boundary point. *Doob's h-process method* [Doo57] allows to condition Brownian motion to "exit" the manifold at a fixed point $\theta \in \partial M$. For every $x \in M$, we obtain a new probability \mathbb{P}_x^θ on Ω , whose support is contained in the set of trajectories starting at x and converging to θ (see [Bro78, Mou94]). This probability satisfies a
 205 *strong Markov property* and an *asymptotic zero-one law*. For all $N \in \mathbb{N}$, denote by τ_N the exit time of the ball $B(o, N)$ and by \mathcal{F}_{τ_N} the associated σ -algebra. Let \mathcal{F}_∞ be the σ -algebra generated by \mathcal{F}_{τ_N} , $N \in \mathbb{N}$. We can reconstruct the probability \mathbb{P}_x with the conditioned probabilities: for a \mathcal{F}_∞ -measurable random variable F ,

$$(3.1) \quad \mathbb{E}_x[F] = \int_{\partial M} \mathbb{E}_x^\theta[F] d\mu_x(\theta).$$

3.3. Stochastic convergence. The behaviour of a harmonic function along trajectories of Brownian motion is easily studied by means of martingale theorems. For a function f on M , let us define the following event:

$$\mathcal{L}_f^{**} := \{\omega \in \Omega \mid \lim_{t \rightarrow \infty} f(X_t(\omega)) \text{ exists and is finite}\};$$

The asymptotic zero-one law implies that the quantity $\mathbb{P}_x^\theta(\mathcal{L}_f^{**})$ does not depend
 210 on x and has value 0 or 1. In the second case, we say that f *converges stochastically at θ* . In the same way we say that f is *stochastically bounded at θ* if \mathbb{P}_o^θ -a.s., $f(X_t)$ is bounded, and that f is of *finite stochastic energy at θ* if \mathbb{P}_o^θ -a.s., $\int_0^{+\infty} |\nabla f(X_t(\omega))|^2 dt < +\infty$. By the martingale property and martingale theorems, F. Mouton [Mou95] proved that for a harmonic function, the set of points
 215 $\theta \in \partial M$ where there is respectively stochastic convergence, stochastic boundedness and finiteness of stochastic energy, are μ -almost equivalent, that is they differ by a set of μ -measure zero.

When the harmonic function u is bounded, non-tangential and stochastic convergences at μ -almost all points of the boundary are automatic ([Anc90]):

Lemma 3.1. *A bounded harmonic function u on M converges non-tangentially and stochastically at μ -almost all points $\theta \in \partial M$ and the unique function $f \in L^\infty(\partial M, \mu)$ such that*

$$u(x) = \int_{\partial M} f(\theta) d\mu_x(\theta) = \mathbb{E}_x[f(X_\infty)]$$


220 *is μ -a.e. the non-tangential and stochastic limit of u .*

4. HARNACK INEQUALITIES

We will use comparison theorems between non-negative harmonic functions several times. The first one is the usual *Harnack inequality on balls*, sometimes called uniform Harnack (see [Anc90] page 21 and [CY75]):

Theorem 4.1 (Harnack on balls). *Let $r > 0$ and $R > 0$ such that $r < R$. There exists a constant $C > 0$ such that for all points $x \in M$ and all non-negative harmonic functions u on $B(x, R)$, we have*

$$\sup_{y \in B(x, r)} u(y) \leq C \cdot \inf_{y \in B(x, r)} u(y).$$


 FIGURE 4.1. Sets U_i^γ and points a_i^γ

225 The *Harnack principle at infinity* is a key principle of potential theory in hyperbolic geometry. It was established by A. Ancona ([Anc87]) in a very general framework using the concept of ϕ -chains. It can be stated simply in the Gromov hyperbolic framework:

Theorem 4.2 (Submultiplicativity of the Green function). *There exists a constant $C > 0$ such that for all pairs of points $(x, z) \in M^2$ and all points $y \in M$ on a geodesic segment between x and z with $\min\{d(x, y), d(y, z)\} \geq 1$, the Green function G satisfies*

$$C^{-1} \cdot G(x, y)G(y, z) \leq G(x, z) \leq C \cdot G(x, y)G(y, z).$$

We will also need another formulation of this principle. Let γ be a geodesic ray starting at $z \in M$ and denote

$$a_i^\gamma := \gamma(4i\delta), \quad i \in \mathbb{N} \setminus \{0\}$$

$$\text{and } U_i^\gamma := \{x \in M \mid (x, a_i^\gamma)_z > 4i\delta - 2\delta\}.$$

Let us point out that for all i , $a_i^\gamma \in U_i^\gamma \setminus U_{i+1}^\gamma$ (see figure 4.1). Let us also notice
230 that the decreasing sequence of sets (U_i^γ) and the sequence of points (a_i^γ) provide a ϕ -chain in the sense of A. Ancona ([Anc90], page 93). Then, the Harnack principle at infinity can be stated as follow:

Theorem 4.3 ([Anc88], page 12). *There exists a constant $C > 0$ such that for all $\theta \in \partial M$ and for all geodesic rays γ from o to θ , the following properties are*
235 *satisfied:*

- (1) *If u and v are two non-negative harmonic functions on U_i^γ , v does not vanish and u "vanishes" at $\overline{U_{i+1}^\gamma} \cap \partial M$, then*

$$\forall x \in U_{i+1}^\gamma, \quad \frac{u(x)}{v(x)} \leq C \frac{u(a_{i+1}^\gamma)}{v(a_{i+1}^\gamma)}.$$

- (2) *If u and v are two non-negative harmonic functions on $M \setminus U_{i+1}^\gamma$, v does not vanish and u "vanishes" at $\partial M \setminus \overline{U_{i+1}^\gamma}$, then*

$$\forall x \in U_i^\gamma, \quad \frac{u(x)}{v(x)} \leq C \frac{u(a_i^\gamma)}{v(a_i^\gamma)}.$$




FIGURE 5.1. Geometric Lemma 5.1

5. NON-TANGENTIAL BEHAVIOUR OF BROWNIAN MOTION

In this section, we gather several lemmas ensuing Harnack inequalities. They provide key ingredients in the proofs of the main results of the paper.

5.1. A geometric Lemma. The following geometric lemma is one of the main tools in the ensuing proofs and in particular in the proof of Theorem 1.1. In [Mou94], it is achieved by use of comparison theorems in pinched negative curvature. For a borelian set $E \subset \partial M$ and a real $c > 0$, denote

$$\Gamma_c(E) := \bigcup_{\theta \in E} \Gamma_c^\theta.$$

Lemma 5.1. *There exist $\eta > 0$ and $c_0 > 0$ such that for all borelian sets $E \subset \partial M$ and all $c > c_0$, one has*

$$\forall x \notin \Gamma_c(E), \mu_x(E) \leq 1 - \eta.$$

Figure 5.1 illustrates Lemma 5.1. The proof follows [Pet12]. We decompose it in two technical lemmas.

For a point $x \in M$, a point $\theta \in \partial M$ and a real $\alpha > 0$, denote

$$A_{x,\alpha}^\theta = \{\xi \in \partial M \mid (\xi, \theta)_x \geq \alpha\}.$$

Lemma 5.2. *There exist two constants $C_1 > 0$ and $d_1 > 0$ depending only on α and δ such that for all $\xi \in \partial M \setminus A_{x,\alpha}^\theta$ and all points y on a geodesic ray from x to θ with $d(x, y) \geq d_1$,*

$$\frac{d\mu_y}{d\mu_x}(\xi) \leq C_1 \cdot G(y, x).$$

Proof. Let $\xi \in \partial M \setminus A_{x,\alpha}^\theta$. Denote by γ a geodesic ray from x to ξ . Choose i such that $d(x, a_i^\gamma) - 3\delta = 4i\delta - 3\delta > \alpha + 4\delta$. By the hyperbolicity inequality (2.2),

$$\alpha > (\xi, \theta)_x \geq \min\{(\xi, y)_x, (y, \theta)_x\} - 2\delta$$

and if y lies on a geodesic ray from x to θ , there exists d_2 depending only on α such that $d(x, y) \geq d_2$ implies $(y, \theta)_x > \alpha + 2\delta$. Thus for such a point y , $(\xi, y)_x \leq \alpha + 2\delta$. Using once again the hyperbolicity inequality,

$$(5.1) \quad \alpha + 2\delta \geq (\xi, y)_x \geq \min\{(\xi, a_i^\gamma)_x, (a_i^\gamma, y)_x\} - 2\delta.$$




FIGURE 5.2. Proof of Lemma 5.2

We have $\xi \in \overline{U_i^\gamma}$, thus $(\xi, a_i^\gamma)_x \geq d(x, a_i^\gamma) - 3\delta > \alpha + 4\delta$. Combining with inequality
 245 (5.1), we obtain $(a_i^\gamma, y)_x \leq \alpha + 4\delta$ and thus $y \notin \overline{U_i^\gamma}$.

Let z be a point of γ in $\overline{U_{i+1}^\gamma}$. It is an exercise using hyperbolicity (see [Anc90] page 85 for details) to verify that the distance between a_i^γ and a geodesic segment between y and z is at most 50δ (see figure 5.2). Since i is fixed, the distance between x and a geodesic segment between y and z is bounded from above by a constant depending only on δ . Thus, the submultiplicativity of the Green function on geodesic segments (Theorem 4.2) associated with the Harnack inequality on balls (Theorem 4.1) give a constant C_1 depending only on δ such that $G(y, z) \leq C_1 \cdot G(y, x)G(x, z)$. Since we have

$$\frac{d\mu_y}{d\mu_x}(\xi) = \lim_{z \rightarrow \xi} \frac{G(y, z)}{G(x, z)},$$

letting $z \rightarrow \xi$, $z \in \overline{U_{i+1}^\gamma}$, we obtain

$$\frac{d\mu_y}{d\mu_x}(\xi) \leq C_1 \cdot G(x, y).$$

□

Lemma 5.3. *Given $\alpha > 0$, there exists a constant $\eta > 0$ such that for all points $x \in M$ and all $\theta \in \partial M$,*

$$\mu_x(A_{x,\alpha}^\theta) \geq \eta.$$

Proof. Fix $\alpha > 0$. We first prove that there exists $d = d(\alpha) > 0$ such that for all $x \in M$, all $\theta \in \partial M$ and all points y on a geodesic ray from x to θ with $d(x, y) \geq d$, we have

$$\mu_y(A_{x,\alpha}^\theta) > \frac{1}{2}.$$

Note that we have

$$(5.2) \quad \mu_y(\partial M \setminus A_{x,\alpha}^\theta) = \int_{\partial M \setminus A_{x,\alpha}^\theta} \frac{d\mu_y}{d\mu_x}(\xi) d\mu_x(\xi).$$

We deduce, from Lemma 5.2 and formula (5.2), that for all points y on a geodesic ray from x to θ with $d(x, y) \geq d_1$, $\mu_y(\partial M \setminus A_{x,\alpha}^\theta) \leq C_1 \cdot G(x, y)$. Since the Green




FIGURE 5.3. Proof of Lemma 5.1

function G has a uniform exponential decay at infinity (Proposition 2.4), there exists a d depending only on α and δ such that for all point y on a geodesic ray from x to θ , with $d(x, y) \geq d$,

$$\mu_y(A_{x,\alpha}^\theta) > \frac{1}{2}.$$

We conclude by Harnack inequality on balls (Theorem 4.1): if $x, y \in M$ with $d(x, y) = d$, we have

$$\mu_x(A_{x,\alpha}^\theta) \geq C(\alpha, \delta) \cdot \mu_y(A_{x,\alpha}^\theta) \geq \eta > 0$$

and the lemma is proved. \square

We can now prove Lemma 5.1.

Proof of Lemma 5.1. Fix $c_0 := K + 6\delta$, where K denotes the constant coming from the roughly starlike assumption on M . Let $c \geq c_0$, E be a borelian set in ∂M , and $x \notin \Gamma_c(E)$. Choose a geodesic ray $\bar{\gamma}$ with origin o such that $d(x, \bar{\gamma}) \leq K$, and denote by $\theta \in \partial M$ the endpoint of $\bar{\gamma}$. Since $x \notin \Gamma_c(E)$, $\theta \notin E$ (see figure 5.3). We prove that there exists a constant $\alpha > 0$ depending only on δ and K such that $A_{x,\alpha}^\theta \subset \partial M \setminus E$. We want to bound the quantity $(\xi, \theta)_x$ uniformly from above for $\xi \in E$. Fix $\xi \in E$. Inequality (2.2) gives

$$\min\{(\xi, \theta)_x, (\xi, o)_x\} \leq (\theta, o)_x + 2\delta.$$

On one hand, since by inequality (2.3), $(\theta, o)_x \leq d(x, \bar{\gamma}) + 2\delta \leq K + 2\delta$, we have $\min\{(\theta, \theta)_x, (\xi, o)_x\} \leq K + 4\delta$. On the other hand, denoting by γ a geodesic ray from o to ξ , we have $(\xi, o)_x \geq d(x, \gamma) - 2\delta \geq c - 2\delta \geq K + 4\delta$. We thus deduce that $(\xi, \theta)_x \leq K + 4\delta$. Since this holds for all $\xi \in E$, we obtain

$$A_{x,K+5\delta}^\theta \cap E = \emptyset.$$

By Lemma 5.3, there exists an $\eta > 0$ depending only on δ and K such that

$$\mu_x(E) \leq 1 - \eta,$$

250 which concludes the proof of Lemma 5.1. \square

For a borelian set $E \subset \partial M$, denote by $v_E(x) := \mu_x(E) = \mathbb{P}_x(X_\infty \in E)$. Let U be an open set in M . Recall that a point $\theta \in \partial M$ is called *tangential for U* if for all $c > 0$, the set $\Gamma_c^\theta \setminus U$ is bounded. The following corollary of Lemma 5.1 asserts that for almost every point θ that is tangential for an open set U in M , Brownian motion "ends its life \mathbb{P}_o^θ -almost surely" in U . 255

Corollary 5.4. *Let U be an open set in M . Then for μ -almost all θ that are tangential for U , \mathbb{P}_o^θ -almost surely, $X_t \in U$ for t large enough.*

Proof. Let $c > c_0$, where c_0 is the constant given in Lemma 5.1. Denote by T the set tangential points for U and, for $N \in \mathbb{N}$, let

$$T_N := \{\theta \in \partial M \mid \Gamma_c^\theta \setminus U \subset B(o, N)\}.$$

By countable union, it is sufficient to prove, for each $N \in \mathbb{N}$, that for μ -almost all $\theta \in T_N$, $X_t \in U$ for t large enough. Fix $N \in \mathbb{N}$. On one hand, since v_{T_N} is a bounded harmonic function, Lemma 3.1 asserts that for μ -almost all $\theta \in T_N$, \mathbb{P}_o^θ -almost surely,

$$\lim_{t \rightarrow \infty} v_{T_N}(X_t) = \mathbf{1}_{T_N}(\theta).$$

On the other hand, by Lemma 5.1,

$$\forall x \notin \Gamma_c(T_N), v_{T_N}(x) \leq 1 - \eta.$$

Thus, for μ -almost all $\theta \in T_N$, \mathbb{P}_o^θ -almost surely, $X_t \in \Gamma_c(T_N)$ for t large enough. Notice that for such a point $\theta \in T_N$ Brownian motion leaves \mathbb{P}_o^θ -almost surely the ball $B(o, N)$ and that $\Gamma_c(T_N) \setminus B(o, N) \subset U$ by definition of T_N . This proves the corollary. 260 □

Corollary 5.5. *Let $c > c_0$ and E be a Borelian subset of ∂M . Every $\theta \in \partial M$ such that v_E converges non-tangentially to 1 at θ is tangential for $\Gamma_c(E)$. In particular, μ -almost all $\theta \in E$ is tangential for $\Gamma_c(E)$.* 265

Proof. Let $\theta \in \partial M$ be such that v_E converges non-tangentially to 1 at θ and let Γ_e^θ be a non-tangential tube with vertex θ . Assume that $\Gamma_e^\theta \setminus \Gamma_c(E)$ is not bounded. Then there exists a sequence $(x_k)_k$ of points in $\Gamma_e^\theta \setminus \Gamma_c(E)$ such that $d(o, x_k) > k$. We thus have $v_E(x_k) \rightarrow 1$. However, by Lemma 5.1, $v_E(x_k) \leq 1 - \eta$, which gives a contradiction and proves the main statement of the corollary. In addition, by Lemma 3.1, v_E converges non-tangentially to 1 at μ -almost all $\theta \in E$ and μ -almost every $\theta \in E$ is tangential for $\Gamma_c(E)$. 270 □

5.2. Behaviour of Green functions. The next Lemma yields an estimate for the increasing rate of the minimal harmonic function $K(\cdot, \theta)$ along non-tangential tubes with vertex θ . The proof is a straightforward application of Theorem 4.2 and Theorem 4.1 (see [Anc90] page 99). 275

Lemma 5.6. *For all $c > 0$, there exist $0 < C < 1$ and $R > 0$ such that for all $\theta \in \partial M$, all $x \in \Gamma_c^\theta$, and all $y \in \Gamma_c^\theta \setminus B(o, R)$,*

$$C \leq G(o, x)K(x, \theta) \text{ and } G(o, y)K(y, \theta) \leq C^{-1}.$$

For an open set $U \subset M$, denote by G_U the Green function of U . The next lemma allows to compare G and G_U for a class of subsets $U \subset M$. This will be useful in the proof of Theorem 1.3.

Lemma 5.7. Fix large $c > e > 0$ and $\theta \in \partial M$. Let U be an open subset of M containing Γ_e^θ and denote by τ the exit time of U . Then we have

$$\lim_{x \rightarrow \theta, x \in \Gamma_e^\theta} \frac{G_U(o, x)}{G(o, x)} = \mathbb{P}_o^\theta(\tau = +\infty).$$

280 *Proof.* Since $G(\cdot, x)$ vanishes at infinity,

$$\begin{aligned} G_U(o, x) &= G(o, x) - \mathbb{E}_o[G(X_\tau, x)] \\ &= G(o, x) \left(1 - \mathbb{E}_o \left[\frac{G(X_\tau, x)}{G(o, x)} \cdot \mathbf{1}_{\tau < +\infty} \right] \right). \end{aligned}$$

Recall that for $\tau < +\infty$, $\lim_{x \rightarrow \theta} \frac{G(X_\tau, x)}{G(o, x)} = K(X_\tau, \theta)$. Hence, provided changing the order of limit and expectation is justified, we have

$$\lim_{x \in \Gamma_e^\theta, x \rightarrow \theta} \mathbb{E}_o \left[\frac{G(X_\tau, x)}{G(o, x)} \cdot \mathbf{1}_{\tau < +\infty} \right] = \mathbb{E}_o [K(X_\tau, \theta) \cdot \mathbf{1}_{\tau < +\infty}] = \mathbb{P}_o^\theta(\tau < +\infty)$$

and the lemma follows. It remains to justify changing the order of limit and expectation, which will be achieved by proving the following property:

There exists a constant $C > 0$ such that

$$(5.3) \quad \forall x \in \Gamma_e^\theta \setminus B(o, c), \forall z \notin \Gamma_c^\theta, \frac{G(z, x)}{G(o, x)} \leq C \cdot K(z, \theta).$$

To prove (5.3), we will apply Theorem 4.3 several times with $u = G(\cdot, y)$ for
285 a point $y \in M$ and $v = K(\cdot, \theta)$. The function $G(\cdot, y)$ is positive harmonic on $M \setminus \{y\}$, vanishes at infinity, and the function $K(\cdot, \theta)$ is positive harmonic. The assumptions of Theorem 4.3 will thus always be satisfied. In the rest of the proof, the constants depend only on the Gromov hyperbolicity constant δ , on the roughly starlike constant K and on c and e .

290 Let $z \notin \Gamma_c^\theta$, $x \in \Gamma_e^\theta \setminus B(o, c)$ and let γ be a geodesic ray starting at o and converging to θ such that $d(x, \gamma) < e$. First, remark that by Theorem 4.1, we can assume $x \in \gamma \setminus B(o, c)$. Indeed, if $x' \in \gamma \setminus B(o, c)$ is such that $d(x, \gamma) = d(x, x')$, then $\frac{G(z, x)}{G(o, x)} \leq C_0 \cdot \frac{G(z, x')}{G(o, x')}$.

Denote, for $i \in \mathbb{N}^*$, $a_i := a_i^\gamma = \gamma(4i\delta)$ and $U_i := U_i^\gamma = \{y \in M \mid (y, a_i)_o >$
295 $d(o, a_i) - 2\delta\}$. We split the proof in different cases:

Case 1: $z \notin U_3$.

By Theorem 4.3, there exists $C_1 > 0$ such that

$$\frac{G(z, x)}{K(z, \theta)} \leq C_1 \cdot \frac{G(a_2, x)}{K(a_2, \theta)}.$$

By definition of a_2 , $d(o, a_2) = 8\delta$ and using once again Theorem 4.1, there exists $C_2 > 0$ such that

$$\frac{G(z, x)}{K(z, \theta)} \leq C_2 \cdot \frac{G(o, x)}{K(o, \theta)} = C_2 \cdot G(o, x),$$

which gives (5.3) in case 1.

Case 2: $z \in U_3$.

By definition of U_3 , $d(o, z) > d(a_3, z) + 8\delta$. Denote by o' a point in γ such that $d(z, o') = \min_{z' \in \gamma} d(z, z')$. Since $d(a_3, z) \geq d(z, o')$, we have $d(o, o') \geq 8\delta$. Denote by γ' a geodesic ray starting at o' and within a distance at most K from z (recall




FIGURE 5.4. Proof of Lemma 5.7

that M is K -roughly starlike). If c is large enough (depending on δ and K), it is an easy exercise to prove that $z \in U_3^{\gamma'}$. We can thus apply Theorem 4.3 to have

$$\frac{G(z, x)}{K(z, \theta)} \leq C_3 \cdot \frac{G(\gamma'(8\delta), x)}{K(\gamma'(8\delta), \theta)}.$$

Hence it is sufficient to prove (5.3) for a point z within distance at most 8δ from γ . Let z be such a point and denote again by $o' \in \gamma$ a point so that $d(z, o') = \min_{z' \in \gamma} d(z, z')$. There are two cases, illustrated by Figure 5.4.

Case 2(a): $d(x, o') > 16\delta$.

In that case, Theorem 4.1 yields that it is sufficient to prove

$$(5.4) \quad \frac{G(o', x)}{G(o, x)} \leq C_4 \cdot K(o', \theta).$$

Recall that the three points o , o' and x lie on the geodesic ray γ .

– If o' is between o and x , we apply Theorem 4.3 with base point x and with $a_i, i = 1, 2$ the points of γ such that $d(o, x) = d(o, a_i) + 4i\delta$ and get

$$\frac{G(o', x)}{K(o', \theta)} \leq C_5 \cdot \frac{G(a_1, x)}{K(a_1, \theta)}.$$

Using Theorem 4.1, we obtain (5.4).

– If x is between o and o' , we apply Theorem 4.2 (recall that $K(\cdot, \theta) = \lim_{y \rightarrow \theta} \frac{G(\cdot, y)}{G(o, y)}$) and obtain

$$\frac{K(x, \theta)}{K(o', \theta)} \leq C_6 \cdot G(x, o').$$

Since $G(o, x)K(x, \theta) \geq C$ (Lemma 5.6) and since $G(x, o') \leq C_7$ (Proposition 2.4), we obtain (5.4).

Case 2(b): $d(x, o') \leq 16\delta$.

Since $8\delta \leq d(x, z) \leq 24\delta$, $K(x, \theta) \leq C_8 \cdot K(z, \theta)$ and $G(x, z) \leq C_9$. Combining these two inequalities with $G(o, x)K(x, \theta) \geq C$, we get property (5.3) in case 2(b). Changing the order of limit and expectation is justified and the proof is complete. \square




FIGURE 5.5. Brownian motion and non-tangential balls

5.3. **Brownian motion and non-tangential sets.** Harnack principles allow to
 315 prove the following lemma (see Figure 5.5 for an illustration), which helps build
 connections between stochastic properties and non-tangential properties. A. An-
 cona stated it in a potential theory terminology ([Anc90], Lemma 6.4) and used it
 to prove a Fatou's Theorem.

Lemma 5.8. Consider a sequence of balls of fixed positive radius whose centers
 320 converge non-tangentially to a point $\theta \in \partial M$, that is, converge to θ staying in a
 non-tangential cone Γ_c^θ for some $c > 0$. Then Brownian motion meets \mathbb{P}_o^θ -almost
 surely infinitely many of these balls.

We end this section by proving the following lemma:

Lemma 5.9. Let U be a connected open subset of M such that $o \in U$, and let τ
 325 denote the exit time of U . For every $\theta \in \partial M$ such that \mathbb{P}_o^θ -almost surely, $X_t \in U$
 for t large enough, we have $\mathbb{P}_o^\theta(\tau = +\infty) > 0$.

Remark 5.10. By Corollary 5.4, the conclusion holds in particular at μ -almost every point θ tangential for U .

Proof of Lemma 5.9. Let $\theta \in \partial M$ be such that \mathbb{P}_o^θ -almost surely, $X_t \in U$ for t large enough. Denote by h the non-negative harmonic function on U defined by $h(x) := K(x, \theta) \mathbb{P}_x^\theta(\tau = +\infty)$. By the maximum principle, h is either positive, or identically zero. We have

$$1 = \lim_{N \rightarrow \infty} \mathbb{P}_o^\theta(\forall t \geq \tau_N, X_t \in U),$$

where τ_N denotes the exit time of $B(o, N)$. Let N be large enough so that $\mathbb{P}_o^\theta(\forall t \geq \tau_N, X_t \in U) > 0$. By the strong Markov property,

$$\mathbb{P}_o^\theta(\forall t \geq \tau_N, X_t \in U) = \mathbb{E}_o^\theta[\mathbb{P}_o^\theta(\forall t \geq \tau_N, X_t \in U)|\tau_N] = \mathbb{E}_o^\theta[\varphi(X_{\tau_N})],$$

330 where $\varphi(x) := \mathbb{P}_x^\theta(\tau = +\infty)$ if $x \in U$ and $\varphi(x) := 0$ otherwise. The function φ , and therefore h , is not identically zero. The function h is thus positive and $\mathbb{P}_o^\theta(\tau = +\infty) > 0$, which proves the Lemma. \square

6. LOCAL FATOU THEOREM

The aim of this section is to prove Theorem 1.1. The proof is similar to the proof of Theorem 2 in [Mou07], and based upon the use of Lemma 5.9, which is achieved using Lemma 5.1. Although the main difference with [Mou07] lies in Lemma 5.1, we give here a detailed proof.

Proof of Theorem 1.1. We can assume, without loss of generality, that U is connected (since U is open, it has a countable number of connected components) and that $o \in U$. Denote again by τ the exit time of U . Let u be a non-negative harmonic function on U . The martingale property asserts that $(u(X_{t \wedge \tau}))$ is a non-negative local martingale and therefore converges \mathbb{P}_o -almost surely. By formula (3.1), for μ -almost all $\theta \in \partial M$, $(u(X_{t \wedge \tau}))$ converges \mathbb{P}_o^θ -almost surely.

By Lemma 5.9, for μ -almost all θ that is tangential for U , we have

$$\mathbb{P}_o^\theta(\tau = +\infty \text{ and } (u(X_t)) \text{ converges}) > 0.$$

Let θ be such a point. Denoting by $\tilde{u}(x) = u(x)$ for $x \in U$ and $\tilde{u}(x) = 0$ otherwise, the asymptotic zero-one law asserts that \tilde{u} converges stochastically at θ . Denote by ℓ the stochastic limit of \tilde{u} at θ and assume that \tilde{u} (and therefore u) does not converge non-tangentially to ℓ at θ . We will obtain a contradiction with Lemma 5.8. These step is standard (see for instance [BD63] page 403 and [Anc90] page 100). There exist $c > 0$, $\varepsilon > 0$ and a sequence $(y_k)_k$ of points in $\Gamma_c^\theta \setminus B(o, R)$ converging to θ such that for every k , $|u(y_k) - \ell| > 2\varepsilon$, where $R > 0$ is such that $\Gamma_{c+1}^\theta \setminus B(o, R) \subset U$. By Harnack inequalities, we have, even replacing 2ε by ε , the same inequality on $B(y_k, \lambda)$ for a $0 < \lambda < 1$ independent of k . By Lemma 5.8, Brownian motion meets \mathbb{P}_o^θ -almost surely infinitely many of the balls $B(y_k, \lambda)$. Let ω be a generic trajectory such that $(X_t(\omega))_t$ meets infinitely many of these balls, $\tau(\omega) = +\infty$ and $\lim_{t \rightarrow +\infty} u(X_t(\omega)) = \ell$. There exists t_0 such that for all $t \geq t_0$, $|u(X_t(\omega)) - \ell| \leq \varepsilon$. By compactness, $(X_t(\omega))_{t \geq t_0}$ meets at least one of the balls $B(y_k, \lambda)$, that is there exists $t_1 \geq t_0$ such that $X_{t_1}(\omega) \in B(y_k, \lambda)$ for some k . Then

$$0 < \varepsilon < |u(X_{t_1}(\omega)) - \ell| \leq \varepsilon,$$

which yields a contradiction. The theorem is proved. \square

We end this section by proving Corollary 1.2.

Proof of Corollary 1.2. Let u be a harmonic function on M . We have to prove that u converges non-tangentially at μ -almost all points $\theta \in \partial M$ where it is non-tangentially bounded from below. Fix $c > c_0$ (where c_0 comes from Lemma 5.1) and for $m \in \mathbb{N}$, let

$$A_c^m := \{\theta \in \partial M \mid \forall x \in \Gamma_c^\theta, u(x) \geq -m\}.$$

It is sufficient to prove that for every $m \in \mathbb{N}$, u converges non-tangentially at μ -almost all $\theta \in A_c^m$. Let $m \in \mathbb{N}$ and $U := \Gamma_c(A_c^m)$. The function $u + m$ is non-negative harmonic on U . By Theorem 1.1, it converges non-tangentially at μ -almost all points θ tangential for U and so the same holds for the function u . By corollary 5.5, μ -almost all $\theta \in A_c^m$ is tangential for U and the proof is complete. \square

350

7. DENSITY OF ENERGY

In this section, we prove Theorem 1.3 and Corollary 1.4. Let us define, for u harmonic on M , $\theta \in \partial M$ and $c > 0$ the *density of energy*

$$D_c^r(\theta) := -\frac{1}{2} \int_{\Gamma_c^\theta} \Delta|u - r|(dx).$$

We refer to [Bro88, Mou07] for introductions to the density of area integral and to the density of energy, respectively. Notice that by Sard's Theorem, for almost all $r \in \mathbb{R}$, $D_c^r(\theta) = \int_{\Gamma_c^\theta} |\nabla u(x)| \sigma_r(dx)$, where σ_r is the hypersurface measure on $\{u = r\}$. In addition, by the coarea formula, the *non-tangential energy* equals

$$J_c^\theta := \int_{\Gamma_c^\theta} |\nabla u|^2 d\nu_M = \int_{r \in \mathbb{R}} D_c^r(\theta) dr.$$

Proof of Theorem 1.3. In order to prove Theorem 1.3, we have to prove that for all $c > 0$:

Step 1: u converges non-tangentially at μ -almost all $\theta \in \partial M$ where $D_c^0(\theta) < +\infty$.

355 **Step 2:** $\sup_{r \in \mathbb{R}} D_c^r(\theta) < +\infty$ for μ -almost all $\theta \in \partial M$ where u converges non-tangentially;

Step 1: the proof goes as in the main Theorem of [Mou07], proved in the framework of manifold of pinched negative curvature. Thus, we give only the main ideas of the proof. The proof is based upon Theorem 1.1, Lemma 5.1 and Lemma 360 5.6.

For $m \in \mathbb{N}$, denote

$$\mathcal{D}_c^m := \{\theta \in \partial M \mid D_c^0(\theta) \leq m\}$$

and $\Gamma := \Gamma_c(\mathcal{D}_c^m)$. It is sufficient to prove that for all $m \in \mathbb{N}$, u converges non-tangentially at μ -a.e. $\theta \in \mathcal{D}_c^m$. Fix $m \in \mathbb{N}$ and recall that $v_{\mathcal{D}_c^m}(x) = \mathbb{P}(X_\infty \in \mathcal{D}_c^m)$. First we prove, using Lemmas 5.1 and 5.6 that there exists $\alpha \in (0, 1)$ such that $\{v_{\mathcal{D}_c^m} \geq \alpha\} \subset \Gamma$ and

$$I := - \int_{\{v_{\mathcal{D}_c^m} \geq \alpha\}} G(o, x) \Delta|u|(dx) < +\infty.$$

Then we prove that for an increasing sequence of compact regular domains V_n such that $\bigcup_n V_n = \{v_{\mathcal{D}_c^m} \geq \alpha\}$,

$$\sup_n \mathbb{E}_o[|u(X_{\tau_n})|] \leq |u(o)| + I,$$

where τ_n is the exit time of V_n . This allows us to decompose u as the difference of two non-negative harmonic functions on $\{v_{\mathcal{D}_c^m} \geq \alpha\}$ (see [Bro88]). Applying Theorem 1.1 to both functions, we get that u converges non-tangentially at μ -almost all tangential θ for $\{v_{\mathcal{D}_c^m} \geq \alpha\}$. By Lemma 3.1, $v_{\mathcal{D}_c^m}$ converges non-tangentially to 1 at μ -almost all $\theta \in \mathcal{D}_c^m$. Such a θ is thus tangential for $\{v_{\mathcal{D}_c^m} \geq \alpha\}$. Hence 365 u converges non-tangentially at μ -almost all $\theta \in \mathcal{D}_c^m$ and the proof of Step 1 is complete.

Step 2: For $m \in \mathbb{N}$ and $c > 0$, denote

$$\mathcal{N}_c^m := \{\theta \in \partial M \mid \sup_{x \in \Gamma_c^\theta} |u(x)| \leq m\}.$$

It is sufficient to show that for all $m \in \mathbb{N}$ and all $c > e > 0$, $\sup_{r \in \mathbb{R}} D_e^r(\theta) < +\infty$ for μ -a.e. $\theta \in \mathcal{N}_c^m$. Fix $c > e > 0$ and $m \in \mathbb{N}$. Let $\Gamma := \Gamma_c(\mathcal{N}_c^m)$ and let τ be the exit time of Γ . Let Γ_n be an increasing sequence of bounded domains such that $\bigcup_n \Gamma_n = \Gamma$ and let τ_n be the exit time of Γ_n . The local martingale $(u(X_{t \wedge \tau}))$ is bounded by m and thus by Barlow-Yor inequalities ([BY81]), $\mathbb{E}_o[\sup_{r \in \mathbb{R}} L_\tau^r] < +\infty$, where L_t^r denotes the local time in r of the local martingale $(u(X_t))$. Formula (3.1) gives that for μ -almost every $\theta \in \partial M$, $\mathbb{E}_o^\theta[\sup_{r \in \mathbb{R}} L_\tau^r] < +\infty$ and in particular, $\sup_n \mathbb{E}_o^\theta[\sup_{r \in \mathbb{R}} L_{\tau_n}^r] < +\infty$.

We now use the following Lemma, whose proof works exactly as Proposition 2 in [Bro88].

Lemma 7.1. *Let u be a harmonic function on M , $r \in \mathbb{R}$, and let L_t^r denote the local time in r of the local martingale $(u(X_t))$. Let also U be a bounded domain in M and τ be the exit time of U . We have*

$$\mathbb{E}_o^\theta[L_\tau^r] = - \int_U G_U(o, x) K(x, \theta) \Delta |u - r|(dx).$$

By Lemma 7.1, for μ -almost every $\theta \in \partial M$,

$$\sup_n \sup_{r \in \mathbb{R}} - \int_{\Gamma_n} G_{\Gamma_n}(o, x) K(x, \theta) \Delta |u - r|(dx) < +\infty.$$

Since for every $n \in \mathbb{N}$, $G_{\Gamma_n}(o, x) \mathbf{1}_{\Gamma_n}(x) \leq G_{\Gamma_{n+1}}(o, x) \mathbf{1}_{\Gamma_{n+1}}(x)$, by the monotone convergence theorem, we have for μ -almost all $\theta \in \partial M$,

$$(7.1) \quad \sup_{r \in \mathbb{R}} - \int_{\Gamma} G_{\Gamma}(o, x) K(x, \theta) \Delta |u - r|(dx) < +\infty.$$

On the other hand, by Lemma 5.9, for μ -almost every $\theta \in \mathcal{N}_c^m$, $\mathbb{P}_o^\theta(\tau = +\infty) > 0$. Hence by Lemmas 5.6 and 5.7, for μ -almost every $\theta \in \mathcal{N}_c^\theta$, there exist $R > 0$ and $C > 0$ such that

$$(7.2) \quad \forall x \in \Gamma_e^\theta \setminus B(o, R), G_{\Gamma}(o, x) K(x, \theta) \geq C.$$

Combining with (7.1), we obtain that for μ -almost every $\theta \in \mathcal{N}_c^m$,

$$2 \sup_{r \in \mathbb{R}} D_e^r(\theta) = \sup_{r \in \mathbb{R}} - \int_{\Gamma_e^\theta} \Delta |u - r|(dx) < +\infty$$

and Theorem 1.3 is proved. □

Proof of Corollary 1.4. Recall that the non-tangential energy at $\theta \in \partial M$ is

$$J_c^\theta = \int_{r \in \mathbb{R}} D_c^r(\theta) dr.$$

Note that if u converges non-tangentially at $\theta \in \partial M$, $D_c^r(\theta) = 0$ for $|r|$ large enough and therefore Theorem 1.3 implies that u has finite non-tangential energy at μ -almost all $\theta \in \partial M$ where u converges non-tangentially.

For $m \in \mathbb{N}$ and $c > 0$, denote

$$\mathcal{J}_c^m := \left\{ \theta \in \partial M \mid \int_{\Gamma_c^\theta} |\nabla u|^2 d\nu_M \leq m \right\}.$$

It is sufficient to prove that for all $m \in \mathbb{N}$, u converges non-tangentially at μ -almost all $\theta \in \mathcal{J}_c^m$. We have

$$\int_{\mathbb{R}} \int_{\mathcal{J}_c^m} D_c^r(\theta) d\mu_o(\theta) dr = \int_{\mathcal{J}_c^m} J_c^\theta d\mu_o(\theta) \leq m.$$

Then for almost every $r \in \mathbb{R}$, we have $\int_{\mathcal{J}_c^m} D_c^r(\theta) d\mu_o(\theta) < +\infty$. For such a real $r \in \mathbb{R}$, we have thus that for μ -almost all $\theta \in \mathcal{J}_c^m$, $D_c^r(\theta) < +\infty$ and by Theorem 1.3, for μ -almost all $\theta \in \mathcal{J}_c^m$, u converges non-tangentially at θ . □

390

REFERENCES

- [Anc87] A. Ancona. Negatively curved manifolds, elliptic operators, and the Martin boundary. *Ann. of Math. (2)*, 125(3):495–536, 1987.
- [Anc88] A. Ancona. Positive harmonic functions and hyperbolicity. In *Potential theory—surveys and problems (Prague, 1987)*, volume 1344 of *Lecture Notes in Math.*, pages 1–23. Springer, Berlin, 1988.
- [Anc90] A. Ancona. Théorie du potentiel sur les graphes et les variétés. In *École d’été de Probabilités de Saint-Flour XVIII—1988*, volume 1427 of *Lecture Notes in Math.*, pages 1–112. Springer, Berlin, 1990.
- [AP08] L. Atanasi and M.A. Picardello. The Lusin area function and local admissible convergence of harmonic functions on homogeneous trees. *Trans. Amer. Math. Soc.*, 360(6):3327–3343, 2008.
- [AS85] M.T. Anderson and R. Schoen. Positive harmonic functions on complete manifolds of negative curvature. *Ann. of Math. (2)*, 121(3):429–461, 1985.
- [BD63] M. Brelot and J. L. Doob. Limites angulaires et limites fines. *Ann. Inst. Fourier (Grenoble)*, 13(fasc. 2):395–415, 1963.
- [BH99] M.R. Bridson and A. Haefliger. *Metric spaces of non-positive curvature*, volume 319 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1999.
- [BHK01] M. Bonk, J. Heinonen, and P. Koskela. Uniformizing Gromov hyperbolic spaces. *Astérisque*, (270):viii+99, 2001.
- [Bro78] J. Brossard. Comportement “non-tangentiel” et comportement “brownien” des fonctions harmoniques dans un demi-espace. Démonstration probabiliste d’un théorème de Calderon et Stein. In *Séminaire de Probabilités, XII (Univ. Strasbourg, Strasbourg, 1976/1977)*, volume 649 of *Lecture Notes in Math.*, pages 378–397. Springer, Berlin, 1978.
- [Bro88] J. Brossard. Densité de l’intégrale d’aire dans \mathbf{R}_+^{n+1} et limites non tangentielles. *Invent. Math.*, 93(2):297–308, 1988.
- [Bus82] P. Buser. A note on the isoperimetric constant. *Ann. Sci. École Norm. Sup. (4)*, 15(2):213–230, 1982.
- [BY81] M. T. Barlow and M. Yor. (Semi-) martingale inequalities and local times. *Z. Wahrsch. Verw. Gebiete*, 55(3):237–254, 1981.
- [Cal50a] A.P. Calderón. On a theorem of Marcinkiewicz and Zygmund. *Trans. Amer. Math. Soc.*, 68:55–61, 1950.
- [Cal50b] A.P. Calderón. On the behaviour of harmonic functions at the boundary. *Trans. Amer. Math. Soc.*, 68:47–54, 1950.
- [Cao00] J. Cao. Cheeger isoperimetric constants of Gromov-hyperbolic spaces with quasi-poles. *Commun. Contemp. Math.*, 2(4):511–533, 2000.
- [CY75] S.Y. Cheng and S.T. Yau. Differential equations on Riemannian manifolds and their geometric applications. *Comm. Pure Appl. Math.*, 28(3):333–354, 1975.
- [Doo57] J.L. Doob. Conditional Brownian motion and the boundary limits of harmonic functions. *Bull. Soc. Math. France*, 85:431–458, 1957.

- [Dur84] R. Durrett. *Brownian motion and martingales in analysis*. Wadsworth Mathematics Series. Wadsworth International Group, Belmont, CA, 1984.
- 435 [Fat06] P. Fatou. Séries trigonométriques et séries de Taylor. *Acta Math.*, 30(1):335–400, 1906.
- [GDLH90] É. Ghys and P. De La Harpe. Espaces métriques hyperboliques. In *Sur les groupes hyperboliques d'après Mikhael Gromov (Bern, 1988)*, volume 83 of *Progr. Math.*, pages 27–45. Birkhäuser Boston, Boston, MA, 1990.
- [Gro81] M. Gromov. Hyperbolic manifolds, groups and actions. In *Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, N.Y., 1978)*, volume 97 of *Ann. of Math. Stud.*, pages 183–213. Princeton Univ. Press, Princeton, N.J., 1981.
- 440 [Gro87] M. Gromov. Hyperbolic groups. In *Essays in group theory*, volume 8 of *Math. Sci. Res. Inst. Publ.*, pages 75–263. Springer, New York, 1987.
- [Gun83] R.F. Gundy. The density of the area integral. In *Conference on harmonic analysis in honor of Antoni Zygmund, Vol. I, II (Chicago, Ill., 1981)*, Wadsworth Math. Ser., pages 138–149. Wadsworth, Belmont, CA, 1983.
- [Mou94] F. Mouton. *Convergence Non-Tangentielle des Fonctions Harmoniques en Courbure Négatives*. PhD thesis, Université Joseph Fourier, 1994.
- 450 [Mou95] F. Mouton. Comportement asymptotique des fonctions harmoniques en courbure négative. *Comment. Math. Helv.*, 70(3):475–505, 1995.
- [Mou00] F. Mouton. Comportement asymptotique des fonctions harmoniques sur les arbres. In *Séminaire de Probabilités, XXXIV*, volume 1729 of *Lecture Notes in Math.*, pages 353–373. Springer, Berlin, 2000.
- 455 [Mou07] F. Mouton. Local Fatou theorem and the density of energy on manifolds of negative curvature. *Rev. Mat. Iberoam.*, 23(1):1–16, 2007.
- [Mou10] F. Mouton. Non-tangential, radial and stochastic asymptotic properties of harmonic functions on trees. *arXiv:1004.4416v1[math.MG]*, 2010.
- [MZ38] J. Marcinkiewicz and A. Zygmund. A theorem of Lusin. Part I. *Duke Math. J.*, 4(3):473–485, 1938.
- 460 [Pet12] C. Petit. Harmonic functions on hyperbolic graphs. *Proc. Amer. Math. Soc.*, 140(1):235–248, 2012.
- [Pic10] M.A. Picardello. Local admissible convergence of harmonic functions on non-homogeneous trees. *Colloq. Math.*, 118(2):419–444, 2010.
- 465 [Pri16] J. Priwaloff. Sur les fonctions conjuguées. *Bull. Soc. Math. France*, 44:100–103, 1916.
- [Spe43] D.C. Spencer. A function-theoretic identity. *Amer. J. Math.*, 65:147–160, 1943.
- [Ste61] E.M. Stein. On the theory of harmonic functions of several variables. II. Behavior near the boundary. *Acta Math.*, 106:137–174, 1961.
- [Yau75] S.T. Yau. Harmonic functions on complete Riemannian manifolds. *Comm. Pure Appl. Math.*, 28:201–228, 1975.
- 470