



**HAL**  
open science

## **In vitro biosynthesis of a universal t6A tRNA modification in Archaea and Eukarya.**

Ludovic Perrochia, Estelle Crozat, Arnaud Hecker, Wenhua Zhang, Joseph Bareille, Bruno Collinet, Herman van Tilbeurgh, Patrick Forterre, Tamara Basta

► **To cite this version:**

Ludovic Perrochia, Estelle Crozat, Arnaud Hecker, Wenhua Zhang, Joseph Bareille, et al.. In vitro biosynthesis of a universal t6A tRNA modification in Archaea and Eukarya.. *Nucleic Acids Research*, 2013, 41 (3), pp.1953-64. 10.1093/nar/gks1287 . hal-00822760

**HAL Id: hal-00822760**

**<https://hal.science/hal-00822760>**

Submitted on 23 May 2013

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# *In vitro* biosynthesis of a universal t<sup>6</sup>A tRNA modification in Archaea and Eukarya

Ludovic Perrochia<sup>1</sup>, Estelle Crozat<sup>1</sup>, Arnaud Hecker<sup>1</sup>, Wenhua Zhang<sup>2</sup>, Joseph Bareille<sup>2</sup>, Bruno Collinet<sup>2,3</sup>, Herman van Tilbeurgh<sup>2</sup>, Patrick Forterre<sup>1,\*</sup> and Tamara Basta<sup>1,\*</sup>

<sup>1</sup>Institut de Génétique et Microbiologie, Université Paris-Sud, IFR115, UMR8621-CNRS, 91405 Orsay, France, <sup>2</sup>Institut de Biochimie et de Biophysique Moléculaire et Cellulaire, Université Paris-Sud, IFR115, UMR8619-CNRS, 91405 Orsay, France and <sup>3</sup>UFR Sciences de la Vie, Université Pierre et Marie Curie UPMC, Sorbonne Universités, 75005 Paris, France

Received September 19, 2012; Revised November 9, 2012; Accepted November 11, 2012

## ABSTRACT

*N*<sup>6</sup>-threonylcarbamoyladenine (t<sup>6</sup>A) is a modified nucleotide found in all transfer RNAs (tRNAs) decoding codons starting with adenosine. Its role is to facilitate codon–anticodon pairing and to prevent frameshifting during protein synthesis. Genetic studies demonstrated that two universal proteins, Kae1/YgjD and Sua5/YrdC, are necessary for t<sup>6</sup>A synthesis in *Saccharomyces cerevisiae* and *Escherichia coli*. In Archaea and Eukarya, Kae1 is part of a conserved protein complex named kinase, endopeptidase and other proteins of small size (KEOPS), together with three proteins that have no bacterial homologues. Here, we reconstituted for the first time an *in vitro* system for t<sup>6</sup>A modification in Archaea and Eukarya, using purified KEOPS and Sua5. We demonstrated binding of tRNAs to archaeal KEOPS and detected two distinct adenosine triphosphate (ATP)-dependent steps occurring in the course of the synthesis. Our data, together with recent reconstitution of an *in vitro* bacterial system, indicated that t<sup>6</sup>A cannot be catalysed by Sua5/YrdC and Kae1/YgjD alone but requires accessory proteins that are not universal. Remarkably, we observed interdomain complementation when bacterial, archaeal and eukaryotic proteins were combined *in vitro*, suggesting a conserved catalytic mechanism for the biosynthesis of t<sup>6</sup>A in nature. These findings shed light on the reaction mechanism of t<sup>6</sup>A synthesis and evolution of molecular

systems that promote translation fidelity in present-day cells.

## INTRODUCTION

All naturally occurring cellular RNA species carry numerous modified nucleosides. Of 100 different modified nucleosides reported to date, >80 are present in mature transfer RNAs (tRNAs). Each tRNA molecule has some subset of these modifications, altering nucleoside identity at an average of 11 positions. Modified nucleosides favour correct folding and stability of tRNA, and some of them are important for the efficiency and fidelity of translation (1–5). Notably, some tRNA positions are almost always modified. This is the case for the nucleotide at the position 37, which is modified in >70% of all tRNAs (6). In a vast majority of cases, the modifications introduced at position 37 are the universal *N*<sup>1</sup>-methyl-guanosine (m<sup>1</sup>G) and *N*<sup>6</sup>-threonylcarbamoyladenine (t<sup>6</sup>A), or their derivatives. t<sup>6</sup>A is found exclusively at the position 37, 3' adjacent to the third base of the anticodon, in all tRNAs reading ANN codons (where N corresponds to four canonical nucleotides A, C, G and T), tRNA<sup>Ile</sup>, tRNA<sup>Thr</sup>, tRNA<sup>Asn</sup>, tRNA<sup>Lys</sup>, tRNA<sup>Ser</sup> and tRNA<sup>Arg</sup> (Figure 1). The t<sup>6</sup>A modification facilitates correct anticodon–codon pairing by enhanced base stacking and by preventing intraloop base pairing of U<sub>33</sub> • A<sub>37</sub>, resulting in enhanced translation fidelity (7,8).


t<sup>6</sup>A modification has been discovered, and its chemical structure was elucidated >40 years ago (9). Initial studies of t<sup>6</sup>A biosynthesis performed *in vivo* or with *Escherichia coli* cellular extracts demonstrated that the reaction required adenosine triphosphate (ATP), carbonate,

\*To whom correspondence should be addressed. Tel: +33 1 69 15 74 89; Fax: +33 1 69 15 66 78; Email: patrick.forterre@igmors.u-psud.fr  
Correspondence may also be addressed to Tamara Basta. Tel: +33 1 69 15 46 28; Fax: +33 1 69 15 78 08; Email: tamara.basta@igmors.u-psud.fr  
Present address:  
Arnaud Hecker, Université de Lorraine, UMR1136 INRA Université de Lorraine 'Interactions Arbres/Micro-organismes', IFR 110 EFABA, Faculté des Sciences BP 239, 54506 Vandoeuvre Cedex, France.

The authors wish it to be known that, in their opinion, the first two authors should be regarded as joint First Authors

© The Author(s) 2012. Published by Oxford University Press.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits non-commercial reuse, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact [journals.permissions@oup.com](mailto:journals.permissions@oup.com).



**Figure 1.**  $t^6A$  modification. On the left is a schematic representation of an anticodon loop of tRNA carrying NNU anticodon and an mRNA molecule carrying ANN codon. Adenosine at position 37 is indicated in white circle. On the right is the chemical structure of  $t^6A$ -modified nucleoside. Chemical groups originating from threonine or carbonate are indicated in grey and bold, respectively.

magnesium and threonine (10,11). However, the proteins involved in the catalysis of this multistep reaction were not identified until recently. The breakthrough came when de Crecy-Lagard *et al.* made the link between the universality of  $t^6A$  and a previous comparative genomic analysis that identified YrdC and YgjD and their archaeal and eukaryotic homologues (Sua5 and Kae1, respectively) as universal proteins of unknown function (12). In particular, they noticed that proteins HypF and NodU/CmcH, which catalyse chemical reactions similar to the ones expected in  $t^6A$  biosynthesis, are formed by the fusion of domains homologous to YrdC and YgjD. Using genetic experiments, they demonstrated that YrdC/Sua5 and YgjD/Kae1 are responsible for the formation of  $t^6A$  *in vivo* in *E. coli* and *Saccharomyces cerevisiae* (13,14).

Earlier this year, Deutsch *et al.* (15) successfully reconstituted the  $t^6A$  synthesis reaction *in vitro* by combining YrdC, YgjD and two additional *E. coli* proteins, YeaZ and YjeE, in the presence of ATP,  $Mg^{2+}$ , threonine and carbonate. Subsequently, the reconstitution of a *Bacillus subtilis* system using purified YwlC (YrdC), YdiB (YjeE), YdiC (YeaZ) and YdiE (YgjD) was reported (16). Notably, YeaZ and YjeE have no detectable orthologues in Archaea and Eukarya. YeaZ is a paralogue of YgjD (thus homologous but distantly related to Kae1), whereas YjeE is specific to Bacteria. This suggested either that Sua5 and Kae1 are sufficient to perform the reaction in these two domains of life or that the YeaZ and YjeE functions are performed by functional analogues evolutionarily unrelated to their bacterial counterparts. Obvious candidate proteins to fulfil this role are the partner proteins of Kae1 (the YgjD homologue) in the KEOPS (kinase, endopeptidase, and other proteins of small size) or EKC (endopeptidase-like and kinase associated with transcribed chromatin) complex. In addition to Kae1, the KEOPS/EKC complex (thereafter called KEOPS) contains three proteins conserved between

Archaea and Eukarya, the protein kinase Bud32, Cgi121 and Pcc1 (17,18). Genetic and transcriptomic studies in *S. cerevisiae* demonstrated that, in addition to Kae1, Bud32 and Pcc1 are required for the formation of  $t^6A$  modification (19,20). Recent genetic investigation of the KEOPS complex in halophilic archaeon *Haloferax volcanii* showed that Kae1, Bud32 and Cgi121 are essential, whereas Pcc1 knockout strain presented a decreased growth rate and reduced levels of  $t^6A$  modification (21). Furthermore, using a whole cell yeast extract for *in vitro*  $t^6A$  synthesis, Srinivasan *et al.* (20) showed that, as in the case of the bacterial system, the reaction requires Sua5, Kae1, threonine, bicarbonate and ATP.

We have previously characterized the archaeal homologues of Kae1 and Bud32, with the aim to get more insight into functions that were probably already present in the last universal common ancestor (LUCA; Kae1/YgjD) or in the last common ancestor of Archaea and Eukarya (Bud32). We have shown that Kae1 is not an *O*-sialoglycoprotein endopeptidase, as initially reported (22), but rather a metalloprotein that binds ATP and DNA *in vitro*, suggesting a role in some unknown universal function related to nucleic acid. In the course of our work, we have purified the archaeal KEOPS complex from the hyperthermophilic archaeon *Pyrococcus abyssi* (PaKEOPS, this work) and the *S. cerevisiae* KEOPS complex (ScKEOPS) (23) for structural and functional studies. Here, we report for the first time the *in vitro* formation of  $t^6A_{37}$ -modified tRNA catalysed by the archaeal and yeast complexes. We present in more detail the enzymatic properties of the more active PaKEOPS complex. The efficiency of the reconstituted *in vitro* system indicates that KEOPS and Sua5 are both necessary and sufficient for the reaction. Notably, PaKEOPS, but not Sua5, strongly binds tRNA *in vitro*, suggesting a role for KEOPS as a tRNA-binding platform in the  $t^6A$  modification reaction. Finally, we identified two distinct ATP-dependent catalytic steps associated with Sua5 and KEOPS. Our data, together with results obtained with the bacterial system (15), show that the universal proteins Sua5/YrdC and Kae1/YgjD cannot synthesize  $t^6A$  by themselves but require additional non-universal accessory proteins. Using interdomain complementation experiments, we demonstrated that the KEOPS complex and bacterial proteins YeaZ, YgjD and YjeE are functional analogues, suggesting mechanistic similarity of  $t^6A$  synthesis in the three domains of life. We discuss evolutionary implications raised by these observations and alternative models they suggested on the nature of LUCA.

## MATERIALS AND METHODS

### Cloning procedures

A polycistronic sequence containing the genes encoding the PaKEOPS subunits PAB1159 (Kae1), PAB1047 (Bud32), PAB1522 (Cgi121) and PAB3073 (Pcc1) was synthesized by GenScript and cloned into pET28(a) plasmid (Novagen). PAB3073 (Pcc1) was fused to a hexa-histidine tag at its 3'-end. The gene PAB1159

(Kae1) was cloned into the pET28(a) as previously described (24).

The gene PAB1302 encoding PaSua5 was amplified by polymerase chain reaction (PCR) using total genomic DNA from the *P. abyssi* GE5 strain as template and cloned into pET26b (Novagen). The ScSua5 gene was amplified by PCR using genomic DNA from *S. cerevisiae* S288C strain (ATCC). A hexa-histidine tag was added at the 3'-end of the gene, which was subsequently cloned into pET21d vector (Novagen).

*E. coli* genes *ygjD* and *yeaZ* were cloned in pET9a with a histidine tag at their 3'-end; *yrdC* was cloned in pET26b with a histidine tag at its 3'-end. *yjeE* gene was synthesized (GenScript) and cloned in pET28a with a Strep-tag II on its 3'-end.

Media were supplemented with 50 µg/ml kanamycin, 34 µg/ml chloramphenicol or 100 µg/ml ampicillin when needed.

### Recombinant protein expression and purification

Recombinant proteins were expressed in Rosetta2 (DE3) pLysS *E. coli* strains (Novagen). Protein overexpression was induced in Overnight Express Instant TB medium (Novagen) supplemented with 10% glycerol.

Cells were collected by centrifugation, resuspended in lysis buffer (Lbf) and sonicated on ice. After centrifugation and in the case of archaeal protein purification, the supernatant was heated at 65°C for 15 min to precipitate bacterial proteins. His-tagged proteins from the soluble fraction were purified by gravity-flow chromatography on an Ni-NTA resin column (Qiagen) according to the manufacturer's recommendations. Proteins carrying a strep-tag II were purified on a Strep-tactin Sepharose resin (IBA) following manufacturer advice. Fractions of interest were pooled, concentrated and injected on a Superdex™200 column (GE Healthcare) for KEOPS or Superdex™75 column for Sua5 and bacterial proteins. Fractions containing pure proteins were concentrated and stored at -20°C in Lbf without β-mercaptoethanol and 10% glycerol.

### tRNA expression plasmid construction

*P. abyssi* (Pa) and *E. coli* (Ec) tRNA gene sequences PaVal (GAC), PaLys (UUU), PaIle (GAU), PaAsn (GUU), PaMet (CAU), EcVal (GAC), EcLys (UUU), EcIle (GAU) and EcAsn (GUU) were retrieved from the Genomic tRNA Database (<http://gtrnadb.ucsc.edu/>) (25). Two complementary oligonucleotides corresponding to the forward and reverse sequence of tRNA genes were synthesized (IDT), annealed and cloned in a pBlue-Script vector yielding pB-tRNA, to allow their expression under the control of T5 promoter/*lac* operator sequence. All sequences are available on request.

### tRNA production and purification

XL1 Blue *E. coli* cells (Agilent Technologies) were transformed with pB-tRNA, and tRNAs were overexpressed in Overnight Express Instant TB medium (Novagen). After centrifugation, total RNAs were extracted with Trizol (Sigma) using the manufacturer's procedure.

Run-off transcription of tRNA genes was performed according to Mougin *et al.* (26): Pa\_tRNA<sup>Lys</sup> and Ec\_tRNA<sup>Asn</sup> genes were cloned in pB-tRNA and amplified by PCR using primers containing the sequence of the T7 promoter upstream of the gene to be transcribed. The transcription reaction was carried out in transcription buffer supplemented with 200 nmol of each rNTP, 40 units of RNase (Thermo Scientific), 300 ng of PCR product and 0.2 µM of his-T7 RNA polymerase expressed and purified according to Arnaud *et al.* (27). Reaction occurred at 37°C for 4h. Template DNA was then digested with DNase I (Thermo Scientific) for 30 min at 37°C.

For purification, RNAs were loaded on an 8 M urea, 12% polyacrylamide gel, and the dominant band corresponding to overexpressed tRNA was cut out. tRNA was eluted from gel in elution buffer, precipitated and resuspended in water. To favour correct folding of tRNAs, the solution was heated at 70°C, cooled slowly to room temperature and stored at -80°C.

### In vitro assay for the synthesis of t<sup>6</sup>A-modified tRNA

The reaction was performed in a final volume of 50 µl containing archaeal (1 µM or 5 µM for complementation experiments), yeast (5 µM) or bacterial (5 µM) proteins, tRNA (2 µM) and 18.2 µM <sup>14</sup>C-L-threonine (0.05 µCi, 55 Ci/mol, Hartmann Analytic) in reaction buffer. After 30-min incubation at 50°C (or other incubation times when indicated), macromolecules were precipitated by the addition of 1 ml of trichloroacetic acid 15% (TCA), and incubated at 4°C for 1h. Precipitated material was applied on pre-wet glass microfibre filters GF/F (Whatmann) using vacuum apparatus (Millipore). Filters were washed with 2 ml of 5% TCA and 3 ml of 95% EtOH and dried. Radioactivity was recorded as average counts per minute (CPM) for 2 min, with a Packard liquid scintillation analyser.

The amount of incorporated threonine was calculated from a standard curve obtained with different concentrations of <sup>14</sup>C-L-threonine spotted on filters (1 pmol = 102.8 CPM).

Yield was calculated as [CPM (assay) - CPM (background)]/102.8/(pmol tRNA) × 100.

### 2D thin layer chromatography for detection of t<sup>6</sup>A-modified nucleosides

tRNAs obtained after *in vitro* t<sup>6</sup>A reaction were extracted with phenol/chloroform and ethanol precipitated. The tRNA pellet was dissolved in 10 µl of 100 mM ammonium acetate supplemented with 1 µg of P<sub>1</sub> nuclease from *Penicillium citrinum* (Sigma) and incubated overnight at 37°C. Digested tRNAs (2 µl) were mixed with 20 µg of cold 5'P-monomonucleosides pA, pU, pG and pC and spotted on a CEL-300 cellulose plate (Merck). 5'P-monomonucleosides were separated using chromatographic solvents A, B or C as previously described (28). First dimension chromatography was performed in solvent A, and the second in solvent B or C. The positions of the four major mononucleosides, pA, pG, pU and pC, were revealed by UV shadowing. The position of t<sup>6</sup>A nucleoside

carrying radiolabelled threonine was revealed by phosphorimaging and compared with that of reference maps obtained under identical experimental conditions.

#### ATP hydrolysis assay

ATP hydrolysis reactions were carried out in a final volume of 25  $\mu$ l containing archaeal or yeast proteins (2  $\mu$ M, if not otherwise specified) in reaction buffer, supplemented with 100  $\mu$ M of cold ATP and 1  $\mu$ Ci of  $\alpha$ -<sup>32</sup>P-ATP (PerkinElmer). When indicated, tRNA, single- or double-strand random sequence RNA (1  $\mu$ M) and other amino acids (50  $\mu$ M) were added to the reaction.

Reaction was performed at 50°C during 30 min, stopped on ice, and 1  $\mu$ l of the mix was spotted on a PEI cellulose plate (Merck), pre-run in distilled water. Radioactive nucleotides were separated by thin layer chromatography (TLC) using 0.5 M KH<sub>2</sub>PO<sub>3</sub><sup>-</sup>, pH 3.5, as the solvent. Plates were dried and revealed by phosphorimaging.

#### Electrophoretic mobility shift assay

The binding reactions were carried out in a final volume of 20  $\mu$ l containing 0.05 pmol (10 nM) of RNA probes and purified PaKEOPS or PaSua5 (0.5 nM to 50  $\mu$ M) in binding buffer. Incubations were performed for 15 min at 4°C, the products were loaded onto a 10% non-denaturing acrylamide gel and electrophoresis was performed in 1  $\times$  TGE buffer at 4°C for 4 h at 7.5 V/cm. The RNA–protein complexes were visualized by autoradiography and phosphorimaging.

## RESULTS

### *P. abyssi* KEOPS binds tightly to tRNAs

The whole PaKEOPS complex (Kae1, Bud32, Cgi121 and Pcc1) was purified from an *E. coli* overproducing strain by exploiting the polyhistidine tag of the Pcc1 subunit. The complex eluted as a single peak with an apparent molecular mass of ~200 kDa after gel filtration (Supplementary Figure S1A). The theoretical mass of PaKEOPS was 90.5 kDa; therefore, the apparent molecular mass of this peak could correspond to a dimer, in agreement with the 3D structural model proposed previously (29). Interestingly, PaKEOPS co-purified with nucleic acids that were resistant to DNaseI digestion but sensitive to RNaseA (not shown) and displayed a similar electrophoretic migration pattern as tRNAs isolated from *E. coli* expression cells (Supplementary Figure S1B). These observations suggested that purified recombinant PaKEOPS was bound to *E. coli* tRNAs. In contrast, recombinant PaSua5 did not co-purify with nucleic acids.

Electrophoretic mobility shift assays of PaSua5 and PaKEOPS in the presence of substrate, non-substrate tRNA and random sequence 65mer RNA showed binding of all RNAs only by PaKEOPS, whereas PaSua5 did not bind tRNA even at a high protein to tRNA molar ratio (Supplementary Figure S2). This suggested that the discrimination between substrate and non-substrate tRNAs occurs after the binding of tRNA to the complex and that specific tRNA-like structural and

sequence features are not required for the binding of RNA by KEOPS. Taken together, these observations suggested that PaKEOPS might be mainly responsible for the binding of tRNA substrates during t<sup>6</sup>A synthesis.


### PaSua5 and PaKEOPS together catalyse the biosynthesis of t<sup>6</sup>A

To determine whether PaSua5 and PaKEOPS were able to catalyse the biosynthesis of t<sup>6</sup>A, we needed an appropriate tRNA substrate lacking this modification. We initially anticipated that additional modifications might also be important for the formation of t<sup>6</sup>A because a previous *in vivo* study using *Xenopus* oocytes showed that t<sup>6</sup>A and several nucleotide modifications (D47, m5C48, m1A58) were formed almost simultaneously (30). To obtain tRNA substrates that most closely satisfied these criteria, we designed a recombinant strategy for tRNA production *in vivo* using *E. coli* cells that overexpressed various *P. abyssi* (Pa\_tRNA) or *E. coli* (Ec\_tRNA) tRNA genes under the control of an inducible promoter. Under these conditions, it is known that the resulting overexpressed tRNA is often hypomodified, thus generating a collection of so-called modivariants (RNA molecules that differ only by their modified nucleoside content) (31,32). Using this system, we typically obtained a major band on denaturing acrylamide gel corresponding to overexpressed, possibly undermodified tRNAs, species (Supplementary Figure S3A).

We incubated PaSua5 and PaKEOPS in presence of ATP, Pa\_tRNA<sup>Lys</sup>, carbonate and <sup>14</sup>C-threonine, and measured the incorporation of <sup>14</sup>C-threonine into TCA-precipitated material. We observed a significant increase of radioactivity when both PaSua5 and PaKEOPS were present in the reaction mixture (Figure 2A). In contrast, no incorporation was observed with PaSua5 or PaKEOPS alone, indicating that both are necessary for the reaction to take place. Notably, PaSua5 and PaKae1 together do not catalyse the incorporation of threonine, indicating that PaKae1 cannot substitute for PaKEOPS. Threonine incorporation also required the addition of ATP and tRNA but, surprisingly, not carbonate. We hypothesized that carbonate was still present in the reaction mixtures despite boiling and extensive degassing of reaction solutions. For comparison, we performed the complete t<sup>6</sup>A assay using *E. coli* proteins (as explained later). As shown in Supplementary Figure S7C, added carbonate had a stimulating effect on the *E. coli* system, indicating that at least some of atmospheric carbonate was removed from reaction solutions. Thus, it appears that archaeal and also eukaryal systems (as explained later) function at lower carbonate concentrations as compared with bacterial systems.

The optimal temperature for threonine incorporation was 50°C with 65% of activity still retained at 73°C (Supplementary Figure S3C). Because the optimal growth temperature of *P. abyssi* GE5 is 96°C, it is likely that the reaction is limited by thermal denaturation of tRNA in non-physiological *in vitro* conditions.

Importantly, the threonine incorporation test was negative for non-ANN decoding tRNA<sup>Val</sup> from *E. coli*



**Figure 2.** *In vitro* reconstitution of t<sup>6</sup>A-modified tRNAs. (A) Incorporation of radiolabelled threonine in tRNAs: reaction mixtures composed of the reactants indicated under the graph were precipitated with TCA, and the amount of incorporated <sup>14</sup>C-threonine was measured. The tRNA species from *P. abyssi* (Pa) and *E. coli* (Ec) (purified from *in vivo* overexpression) are indicated. In one case, Kae1 was used instead of the whole KEOPS complex. HCO<sub>3</sub><sup>-</sup> was added (+) or not (-) to the reaction. Error bars indicate standard deviation from the mean obtained for three independent experiments. (B) Detection of t<sup>6</sup>A-modified nucleotide using 2D TLC. Upper line shows chromatographic mobility in solvent system A + B, lower line in solvent system A + C. Left panels: abundant, non-modified nucleotides were detected by ultraviolet (UV) shadowing. Centre panel: <sup>14</sup>C-threonine incorporated in modified nucleotides was detected by phosphorimaging, and the positions of unmodified nucleotides were superimposed by aligning with the UV image (white circles). Right panel: reference maps [from (28)] showing the theoretical position of t<sup>6</sup>A nucleoside in both solvent systems. (C) Incorporation of radiolabelled threonine using different tRNA species as substrates. Legend is the same as in (A); tRNAs were overexpressed *in vivo*, or produced *in vitro* when stated. fMet, bacterial initiator *N*-formyl-methionine; AsnC37 is Asn tRNA where A37 was mutated into a cytosine; Sc tRNAs are tRNAs purified from a  $\Delta$ sua5 *S. cerevisiae* strain. The percentage indicated on top of each bar is the yield of t<sup>6</sup>A-modified tRNA obtained for each reaction.


and *P. abyssi*, thus confirming that only ANN-decoding tRNA molecules are valid substrates. Moreover, Ec\_tRNA<sup>Asn</sup> failed to act as a substrate when adenosine at position 37 was replaced by cytosine, showing that t<sup>6</sup>A is specifically added on the adenosine 37.

The incorporation of threonine into tRNAs was confirmed by purification of tRNAs after t<sup>6</sup>A reaction. As expected, only tRNAs from the reaction containing PaSua5 and PaKEOPS were radioactive (Supplementary Figure S3B).

Finally, we performed 2D TLC to demonstrate that the incorporation of threonine into tRNA corresponded to t<sup>6</sup>A formation. For this, tRNAs were isolated from the reaction mixture, hydrolysed to individual nucleotides and separated by 2D TLC. The position of the radioactive

spot on the plate was reproducible and consistent with the position of the t<sup>6</sup>A-modified nucleotide according to previously published reference maps (33) (Figure 2B).

Next, it was important to verify that the *in vitro* assay we established can generate a significant amount of t<sup>6</sup>A-modified tRNA. Standard assay conditions yielded 25–32% of modified Pa\_tRNA<sup>Lys</sup> (i.e. mole t<sup>6</sup>A<sub>37</sub> per mole of tRNA present in the incubation mixture). This result indicated that the *in vitro* reaction could still be optimized or that the enzymatic machinery may work better *in vivo* under the physiological conditions of the cell. To optimize the reaction, we added manganese divalent ions, which were reported to be required for optimal activity of Bud32 (34). Indeed, manganese had a stimulating effect because we could increase the yield of



**Figure 3.** ATPase activity of Sua5 and KEOPS. (A) *P. abyssi* proteins and (B) *S. cerevisiae* proteins. Produced radiolabelled ( $\alpha$ - $^{32}$ P) ADP and AMP were separated by TLC and visualized by phosphorimaging. Presence (+) or absence (-) of different reactants in reaction mixtures is indicated. Pa\_tRNA<sup>Lys</sup> produced in overexpressing *E. coli* strain was used as a substrate.

the reaction to 50% for archaeal system and to 38% for the *S. cerevisiae* system (as explained later) (Supplementary Figure S5).

However, the yield reported is the lowest estimate for our assay conditions because, as noted previously, it is possible that a subpopulation of the *in vivo* produced Pa\_tRNAs<sup>Lys</sup> already carried the t<sup>6</sup>A modification. The yield could not be improved by adding a fresh aliquot of PaSua5 and PaKEOPS or ATP into the reaction mixture (not shown), which indicated that the amount of substrate tRNA was the limiting factor.

To further characterize the range of substrate tRNAs used by PaKEOPS and PaSua5, we also tested overexpressed *E. coli* tRNAs, total tRNAs isolated from a *S. cerevisiae* Sua5 deletion mutant (19) and *in vitro* transcribed tRNAs. All of these acted as equivalent substrates (Figure 2C), demonstrating that (i) universally conserved structural and/or sequence features of tRNAs decoding ANN codons are recognized by PaKEOPS and PaSua5; and (ii) presence of modified nucleotides on ANN-decoding tRNA is not required for the t<sup>6</sup>A<sub>37</sub> modification to take place.

### Two distinct ATP-dependent reactions occur in the course of t<sup>6</sup>A synthesis by PaSua5 and PaKEOPS

To investigate the ATP-dependent activation steps undertaken by PaKEOPS and PaSua5, we assayed the ATPase activity using  $\alpha$ - $^{32}$ P-ATP and TLC to visualize the resulting products (Figure 3A).

When all the proteins and substrates were put together in a reaction that leads to t<sup>6</sup>A synthesis and under saturating concentration of ATP, we observed that both AMP and ADP were produced. The production of AMP occurred when PaSua5 was present in the reaction mixture alone. This activity was significantly increased when threonine and carbonate were added but did not require tRNA (Figure 3A and Supplementary Figure S7A and S7B). In contrast, PaSua5 activity was not stimulated by glycine, phenylalanine and homoserine, an isomer of threonine (Supplementary Figure S4B).

In contrast to PaSua5, incubation of PaKEOPS alone with ATP generated ADP (Figure 3A). This basal activity is detected in the absence of tRNA; however, the addition of substrate tRNA or non-substrate Ec\_tRNA<sup>Asn</sup> (C37) and Pa\_tRNA<sup>Val</sup> significantly increased this activity

(Figure 3A and Supplementary Figure S4A). Notably, such activation was not observed when tRNA was replaced by random sequence 65mer DNA or RNA, although this RNA was bound by KEOPS with an affinity similar to that observed for tRNA (see Supplementary Figures S2C and S4A).

The KEOPS-dependent ATP hydrolysis into ADP seems to be required for t<sup>6</sup>A synthesis because the t<sup>6</sup>A reaction was completely inhibited when ATP was replaced by AMP-PNP, an ATP analogue that can be hydrolysed into AMP (data not shown).

Taken together, our data indicate that two distinct ATP-dependent reactions, one generating AMP and another generating ADP, are catalysed by PaSua5 and PaKEOPS, respectively, in the course of t<sup>6</sup>A synthesis.

#### *In vitro* synthesis of t<sup>6</sup>A by Sua5 and KEOPS from *S. cerevisiae*


The *S. cerevisiae* KEOPS (ScKEOPS) complex was heterologously expressed and purified as previously described (23). In contrast to PaKEOPS, ScKEOPS was not associated with tRNAs when purified from *E. coli* cells (not shown). Consistent with observations made for PaSua5, the purified recombinant yeast homologue ScSua5 did not co-purify with tRNA either.

To test whether ScSua5 and ScKEOPS could also catalyse the biosynthesis of the t<sup>6</sup>A modification, we used the same experimental approach as for *P. abyssi* proteins except that the assay was performed at 30°C. As for *P. abyssi* proteins, we observed the incorporation of <sup>14</sup>C-threonine into TCA-precipitated material, and 2D TLC analysis confirmed that t<sup>6</sup>A was formed (Figure 4B). This reaction was ATP and tRNA dependent and required the presence of ScSua5 and ScKEOPS. Similar as for the *P. abyssi* system, the omission of carbonate did not have an effect on the amount of incorporated threonine (Supplementary Figure S7C). The amount of threonine incorporation was similar for endogenous tRNAs and *E. coli* tRNAs, indicating that, like *P. abyssi* proteins, the *S. cerevisiae* proteins recognize features common to all ANN-decoding tRNA molecules.

We next examined the ATPase activity of ScSua5 and ScKEOPS using TLC (Figure 3B). We observed weak ATPase activity of ScSua5 of which the product was AMP. This activity was threonine dependent and carbonate dependent (Supplementary Figure S7A and S7B) but did not require tRNA, and addition of tRNA did not have a stimulating effect. Thus, the ScSua5 behaved in the same manner as the PaSua5. In contrast, the behaviour of ScKEOPS was different from that of PaKEOPS because we observed the production of ADP and AMP in a tRNA-independent manner. Note, however, that we cannot exclude a contaminating ATPase activity from the *E. coli* expression strain.


#### Interdomain exchange of t<sup>6</sup>A machinery components

The similar behaviour of archaeal, yeast and bacterial t<sup>6</sup>A machineries suggested that the function of KEOPS/YgjD-YjeE-YeaZ (hereafter called DEZ) and Sua5/YrdC has been conserved throughout evolution. To test this



**Figure 4.** *In vitro* synthesis of t<sup>6</sup>A modification using *S. cerevisiae* KEOPS and Sua5 proteins. (A) Incorporation of radiolabelled threonine in tRNAs: the amount of incorporated <sup>14</sup>C-threonine was measured as in Figure 2A. Total tRNAs were purified from a  $\Delta$ Sua5 strain; for comparison, *in vitro* transcribed *E. coli* tRNA<sup>Lys</sup> was used when indicated. Error bars indicate standard deviation from the mean obtained from three independent experiments. (B) Detection of t<sup>6</sup>A-modified nucleotide using 2D TLC. The chromatography was done as described in Figure 2B.

hypothesis, we purified bacterial proteins and set up a functional *in vitro* system to perform complementation assays. We combined Sua5/YrdC proteins and KEOPS/DEZ complexes. TLC analysis confirmed that the mixed systems produce t<sup>6</sup>A-modified tRNA (Supplementary Figure S6). Remarkably, despite the non-optimal temperatures used in the assays, the amount of t<sup>6</sup>A produced by heterologous combinations was comparable with the amount produced by the homologous systems at the respective reaction temperatures (43°C for Pa and Ec proteins, and 35°C for Sc proteins) (Figure 5). Lower values were obtained in complementation with PaKEOPS, owing to suboptimal temperatures because at 43°C and 35°C, PaKEOPS retains 40% and 25% of its activity (Supplementary Figure S3C). Nevertheless, those values were reproducible and are still fairly above the background signal. These results indicate that Sua5/YrdC



**Figure 5.** Functional complementations assays between bacterial, archaeal and eukaryotic proteins. Complementation assays were tested using PaKEOPS, ScKEOPS or EcDEZ combined with PaSua5, ScSua5 or EcYgjD. Incorporation of radiolabelled threonine was measured as in Figure 2A. Activities of archaeal (A), bacterial (B) and eukaryotic (E) complexes were measured at 43°C (A, B and A/B) and 35°C (E, B/E and A/E). The dotted line indicates the background level, obtained without ATP.

and KEOPS/DEZ components can be exchanged, suggesting that  $t^6A$ -synthesizing machineries from the three domains of life share a common catalytic mechanism.

## DISCUSSION

### Archaeal and eukaryal Sua5 and KEOPS together catalyse the synthesis of $t^6A$ -modified tRNA

In the present study, we established an *in vitro* system for enzymatic synthesis of a universal and essential  $t^6A$  tRNA modification by archaeal and eukaryotic KEOPS and Sua5 proteins. This work thus provides for the first time direct evidence for the biological function of KEOPS and Sua5 in archaeal and eukaryotic cells and confirms the results of previous genetic and transcriptome studies in *S. cerevisiae* and *H. volcanii* (13,14,19,20,21). In the light of these findings, it is possible that the pleiotropic phenotypes described for yeast KEOPS and Sua5 deletion mutants (17,18,35) are indirect effects owing to the translational defects occurring in cells lacking  $t^6A$  modification. This is well illustrated by a recent study that showed that the diverse cellular dysfunctions linked to the development of type 2 diabetes in mice can be attributed to abnormal protein synthesis caused by the lack of 2-methylthio- $N^6$ -threonylcarbamoyladenine ( $ms^2t^6A$ )-modified tRNA<sup>Lys</sup> (36).


Purified proteins and a reconstituted *in vitro* system allowed us to get the first insights onto the structural and mechanistic features of the  $t^6A$  modification process. We made two important observations: firstly, PaKEOPS, but not PaSua5, binds tRNA tightly; and secondly, the reaction requires the hydrolysis of ATP into AMP by Sua5 and the hydrolysis of ATP into ADP by KEOPS.

### The KEOPS complex is a tRNA binding platform

PaKEOPS binds tRNA of *E. coli* *in vivo* and *in vitro*, and this binding is strong because it is preserved throughout the purification process. It therefore seems plausible, suggesting that the KEOPS complex is responsible for tRNA binding in the course of the  $t^6A$  synthesis. Interestingly, PaKEOPS binds tRNA in a positive cooperative manner (Supplementary Figure S2). It is unlikely that PaKEOPS binds more than one molecule of tRNA simultaneously because of the large size of tRNA. An alternative explanation is that one tRNA molecule is bound by each subunit of a PaKEOPS homodimer. We have previously shown that KaeI from *P. abyssi* binds DNA *in vitro* (24). This suggested that KaeI could have a general affinity for nucleic acids. Indeed, we could show using electrophoretic mobility shift assays that purified *P. abyssi* KaeI can bind ssRNA and tRNA, albeit with lower affinity as compared with KEOPS (unpublished data). This suggests that tRNA interacts with KaeI in the KEOPS/tRNA complex. However, because purified KaeI is not bound to *E. coli* tRNA, it is likely that other proteins of the complex are also involved in tRNA binding. Interestingly, the archaeal/eukaryal KEOPS and bacterial  $t^6A$  machinery can use as substrate tRNA from the three domains of life. This again suggests that KaeI is directly involved in tRNA binding because it is the only KEOPS protein with a bacterial homologue (YgjD). This also indicates that universal structural and/or sequence features of tRNA are recognized by all systems, suggesting a similar mode of interaction between tRNA and the proteins.

### The catalytic mechanism for $t^6A$ synthesis seems to be conserved in the three domains of life

It was suggested that  $t^6A$  synthesis, which is a complex modification, required at least two ATP-dependent activation steps (37). In agreement with this proposal, Deutsch *et al.* reported that both AMP and ADP were produced when all four proteins of the bacterial system were mixed (15). They observed that YrdC alone hydrolysed ATP into AMP in a threonine-dependent reaction, and that YgjD, YjeE and YeaZ (DEZ) generated ADP. These results were confirmed by a recent study that reconstituted *in vitro* the  $t^6A$  biosynthesis by *B. subtilis* proteins YwIC (YrdC), YdiB (YjeE), YdiC (YeaZ) and YdiE (YgjD) (16). One common point between the bacterial system and our archaeal/eukaryal systems is the production of both AMP and ADP in the reaction that generates  $t^6A$ -modified tRNA. The conservation of this feature across the three domains of life corroborates the hypothesis of at least two distinct ATP-dependent steps in the course of  $t^6A$  synthesis, suggesting a conserved catalytic mechanism for the biosynthesis of  $t^6A$  in nature, despite the fact that some of the proteins are unrelated. The second common point of all three systems is the carbonate and threonine dependence of AMP generation by YrdC/Sua5. This observation is in line with the universal character of these proteins and indicates that they perform the same catalytic role in the three domains of life.



**Figure 6.** Proposed reaction scheme for  $t^6A$  synthesis. Different catalytic steps are catalysed by proteins indicated above arrows. Activation of bicarbonate to carboxy-AMP [1]; condensation of carboxy-AMP with threonine to form carbamoylthreonine [2]; activation of carbamoylthreonine to threonylcarbamoyladenylate [3]; transfer of threonylcarbamoyladenylate to Kae1/YgjD active site where second condensation reaction occurs to form  $t^6A$ -modified tRNA [4]. The participation of Sua5 in the last catalytic step is currently unclear. Further details are in the text.

The three systems diverge significantly in terms of complexes involved in the ADP-generating reaction step. In Bacteria, this reaction is catalysed by DEZ, whereas it is catalysed by KEOPS in Archaea and Eukarya. Kae1/YgjD is the only common protein between these different systems, suggesting that it might be responsible for this activity.

Finally, using *in vitro* complementation experiments, we have demonstrated that Sua5 can replace YrdC and *vice versa* and that KEOPS is interchangeable with DEZ. These data corroborate the notion of a common catalytic mechanism for  $t^6A$  synthesis catalysed by universal proteins Sua5/YrdC and Kae1/YgjD.

#### A putative mechanism for $t^6A$ synthesis

The threonine-dependent hydrolysis of ATP into AMP by Sua5 and YrdC suggests the formation of an adenylate reaction intermediate, such as threonylcarbamoyladenylate. The fact that we do not directly observe threonylcarbamoyladenylate is probably because of its unstable nature under acidic conditions used for TLC analysis. This intermediate was isolated using HPLC under neutral pH conditions with *B. subtilis* Yw1C (YrdC), which catalysed the conversion of L-threonine, bicarbonate/ $CO_2$  and ATP to give L-threonylcarbamoyl-AMP intermediate and pyrophosphate as products (16). Moreover, L-threonylcarbamoyl-AMP was found in the 3D crystal structure of Sua5 from *Sulfolobus tokodaii*, and a recent study of TobZ protein also suggested a putative model for  $t^6A$  synthesis involving a threonylcarbamoyladenylate intermediate (38,39). This protein, which synthesizes the antibiotic nebramycin, consists of two domains homologous to YrdC and Kae1 fused into a single polypeptide. In the proposed model for the reaction catalysed by TobZ, the YrdC-like domain catalyses the ATP-dependent formation of the carbamoyladenylate intermediate, followed by the transfer of the carbamoyl moiety to tobramycin to form nebramycin with

concomitant release of AMP. This final step is performed by the Kae1-like domain of TobZ, which positions the adenylate intermediate and the sugar substrate in the correct orientation. By analogy, Parthier *et al.* (39) suggested that Sua5 catalyses the formation of threonylcarbamoyladenylate, which is then accommodated in the active site of Kae1, together with the anticodon loop of tRNA, to form  $t^6A$ -modified tRNA. Our results are compatible with this model. However, Parthier *et al.* further suggested that N-carbamoylthreonine, the precursor for the synthesis of the threonylcarbamoyladenylate intermediate, is generated from bicarbonate and threonine by an unknown enzymatic activity. Our data prove that it is not the case because Sua5 and KEOPS are sufficient to synthesize  $t^6A$ . Moreover, the observation that ATPase activity of PaSua5 [and of YrdC and Yw1C in bacterial systems (15,16)] is carbonate and threonine dependent strongly suggests the involvement of Sua5/YrdC in the early steps of the catalysis, leading to the formation of threonylcarbamoyladenylate. In addition, our data imply that KEOPS is involved in the last step of the catalysis, which requires the presence of tRNA. This is supported by the finding that the reaction of tRNA with L-threonylcarbamoyl-AMP can be catalysed by the *B. subtilis* YdiB (YjeE), YdiC (YeaZ) and YdiE (YgjD) complex in the absence of Yw1C (YrdC) (16). Based on these data and previous studies, we propose a novel mechanism for the synthesis of  $t^6A$  (Figure 6). In the first step, bicarbonate is activated by Sua5/YrdC to carboxy-AMP [1], which can then condense with the threonine to form carbamoylthreonine [2]. Sua5/YrdC uses a second ATP to activate this product and generates threonylcarbamoyladenylate [3]. Finally, this intermediate is transferred to Kae1/YgjD active site where second condensation reaction occurs to form  $t^6A$ -modified tRNA. This proposal is compatible with the data reported by Lauhon (16), but it is in disagreement with the work of El Yacoubi *et al.* (14), who proposed the activation of

bicarbonate to carboxyphosphate catalysed by Kae1. We do observe generation of ADP by KEOPS, and this activity is mandatory. However, KEOPS activity is sensible to neither carbonate nor threonine, suggesting strongly that the complex, and thus Kae1, is not involved in early steps of catalysis. In favour of our proposal, YrdC-homologous domain of TobZ protein was shown to catalyse formation of carbamoyl-AMP, a compound that is structurally reminiscent of carboxy-AMP. Our model is compatible with the data obtained for the bacterial t<sup>6</sup>A systems, and carboxy-AMP was included in the proposed mechanism scheme as an alternative to the formation of carboxyphosphate (15). The present data do not allow assigning convincingly the ADP-generating ATPase activity of KEOPS/DEZ to a particular process. However, in archaeal system, we observed that the active site responsible for ATP hydrolysis is not in a catalytically competent conformation until tRNA is bound, indicating that a set of mutually induced conformational changes occur between PaKEOPS and tRNA.

Therefore, we propose that KEOPS and DEZ must undergo conformational changes to attain a catalytically relevant conformation and that ATP hydrolysis is needed to compensate for the free energy cost of rearrangements. Further experimental work is needed to verify this hypothesis.

#### **Possible role of archaeal Sua5 and KEOPS in the synthesis of N<sup>6</sup>-hydroxynorvalylcarbamoyladenine (hn<sup>6</sup>A)**

Two other modified adenosines occur at position 37 and contain carbamoyl and amino acid moieties: N<sup>6</sup>-glycylcarbamoyladenine (g<sup>6</sup>A) and N<sup>6</sup>-hydroxynorvalylcarbamoyladenine (hn<sup>6</sup>A) (3,40), but the enzymes catalysing the synthesis of these modified nucleosides have not been identified yet. g<sup>6</sup>A is present in certain cytoplasmic tRNAs of eukaryotes, whereas hn<sup>6</sup>A is found in tRNAs of Bacteria and Archaea (41). In agreement with this phylogenetic distribution, we found that PaSua5 hydrolyses ATP to AMP in presence of hydroxynorvaline, but not in presence of glycine (Supplementary Figure S4B). This suggested that Sua5 proteins could be responsible for the synthesis of the hydroxynorvalylcarbamoyladenylyl and that, together with KEOPS, they may be responsible for the formation of hn<sup>6</sup>A in Archaea. Further experimental work is needed to confirm this hypothesis, which if true, would extend the role of archaeal KEOPS and Sua5 proteins to the synthesis of another tRNA modification.

#### **Evolutionary implications**

The present work and the previous work on bacterial system (15) revealed that the universal proteins Sua5/YrdC and Kae1/YgjD alone are not sufficient to perform t<sup>6</sup>A modification but require two sets of non-universal proteins, one specific to Bacteria (YeaZ, YjeE) and the other common to Archaea and Eukarya (Bud32, Cgi121 and Pcc1). A similar situation is found in many other fundamental molecular systems, e.g. translation initiation and transcription, in which a core of universally

conserved proteins is supplemented by factors specific for Bacteria on one side and for Archaea/Eukarya on the other. It is often argued that such a situation suggests a simple LUCA [a progenote *sensu* (42) and subsequent independent refinement of molecular processes (i.e. translation)] in Bacteria and in Archaea/Eukarya. In case of t<sup>6</sup>A, it seems unlikely that LUCA lacked this modification. Indeed, several studies identified t<sup>6</sup>A as a part of a minimal tRNA modifications set required for life (43–45). Moreover, the genes responsible for the synthesis of t<sup>6</sup>A are resistant to loss in insect symbionts and mollicutes that have reduced genomes, suggesting that these genes emerged early during cellular evolution, and once acquired became essential for the cell (45,46). Finally, it has been argued convincingly that t<sup>6</sup>A modification is necessary for the accurate reading of the three letters genetic code (7,8,19). Because this code is universal, it was likely present in LUCA, and its accuracy was probably already optimal (47).

If t<sup>6</sup>A was already present in LUCA, one should imagine evolutionary scenarios to explain how an ancestral tRNA modification mechanism evolved into two distinct modern versions. The idea that these two versions evolved from a common mechanism is supported by the existence of two universal proteins in both versions, by the universality of the two distinct ATP-dependent steps and by the interdomain functional complementations. One possible scenario is that the core proteins (Sua5/YrdC and Kae1/YgjD) are the remnants of an ancient synthetase operating in LUCA, which could have performed the complete reaction without accessory proteins. After the separation of the three domains, additional proteins were added independently in Bacteria and in ancestors of Archaea and Eukarya to improve speed and accuracy of protein translation in increasingly complex cells. In this scenario, we propose that these proteins were selected to provide additional surfaces for tRNA binding. This suggestion fits well with the observation that Kae1 alone binds tRNA less tightly as compared with KEOPS (as explained previously). Alternatively, LUCA might have possessed either DEZ or KEOPS to work with the ancestor of Sua5/YrdC; later on, YeaZ and YjeE were replaced by Bud32, Pcc1 and Cgi121 in Archaea and Eukarya or *vice versa*, and KEOPS was replaced by DEZ in Bacteria. Our complementation experiments render this hypothesis plausible because we show that both KEOPS and DEZ can complement *in vitro* the Sua5/YrdC protein of the three different domains, suggesting that such situation could have happened *in vivo* in ancestors of modern organisms.

The study of t<sup>6</sup>A synthesis in mitochondria could contribute to the comprehension of t<sup>6</sup>A machinery evolution. Eukaryotic genomes encode mitochondrial homologue of Kae1 called Qri7 in *S. cerevisiae*. Genetic data from two independent studies indicated that *S. cerevisiae* Qri7 functions in mitochondrial t<sup>6</sup>A synthesis. Interestingly, Sua5 is required for both cytoplasmic and mitochondrial t<sup>6</sup>A synthesis, indicating that it is imported into mitochondria despite the lack of a recognizable import signal (14,20). It remains to be investigated if Qri7 and Sua5 alone could catalyse the formation of t<sup>6</sup>A-modified tRNA or if other

proteins are required. In the case of the second hypothesis, the proteins of KEOPS are the obvious candidates. The first hypothesis, if true, would substantially argue in favour of the scenario in which the two universal proteins formed the ancient t<sup>6</sup>A synthetic machinery.

In conclusion, this study established an *in vitro* system for the synthesis of essential t<sup>6</sup>A tRNA modification by eukaryotic and archaeal t<sup>6</sup>A synthetases. Characterization of these systems allowed us to propose a novel reaction scheme for the synthesis of t<sup>6</sup>A. Our data suggest that the catalytic mechanism for t<sup>6</sup>A synthesis is conserved in nature. Our work paves the way for further mechanistic and structural studies of these enzymatic complexes. The full comprehension of how this ancient modification is synthesized should result in better understanding of evolutionary mechanisms that contributed to increased translation efficiency and fidelity in present day cells.

## SUPPLEMENTARY DATA

Supplementary Data are available at NAR Online: Supplementary Figures 1–7, Supplementary Methods.

## ACKNOWLEDGEMENTS

The authors are indebted to Henri Grosjean for his encouragement, stimulating discussions and critical reading of the manuscript. The authors thank Domenico Libri and Marie-Claire Daugeron for helpful discussions and for providing the Sua5 yeast deletion strain; Dorian Guetta, Philippe Bouloc, Chantal Bohn and Emmanuelle Schmitt for sharing protocols and for helpful suggestions. B.C. wishes to warmly thank Magali Blaud for her helpful advices and protocols and François Mallet for his generous gift of pMRT-7 wt.

## FUNDING

Agence Nationale de Recherche [ANR-09-BLAN-0349 to P.F. and H.vT.]; Marie Curie postdoctoral fellowship (to E.C.); ENS Lyon (PhD fellowship to L.P.). Funding for open access charge: Agence Nationale de Recherche [ANR-09-BLAN-0349].

*Conflict of interest statement.* None declared.

## REFERENCES

- Grosjean, H., Sprinzl, M. and Steinberg, S. (1995) Posttranscriptionally modified nucleosides in transfer RNA: their locations and frequencies. *Biochimie*, **77**, 139–141.
- Gustilo, E.M., Vendeix, F.A. and Agris, P.F. (2008) tRNA's modifications bring order to gene expression. *Curr. Opin. Microbiol.*, **11**, 134–140.
- Grosjean, H. Nucleic acids are not boring long polymers of only four types of nucleotides. *DNA and RNA Modification Enzymes: Structure, Mechanism, Function and Evolution*. Landes Biosciences, Austin, Texas, USA, pp. 1–18.
- Juhling, F., Morl, M., Hartmann, R.K., Sprinzl, M., Stadler, P.F. and Putz, J. (2009) tRNAdb 2009: compilation of tRNA sequences and tRNA genes. *Nucleic Acids Res.*, **37**, D159–D162.
- Giegé, R., Juhling, F., Pütz, J., Stadler, P., Sauter, C. and Florentz, C. (2012) Structure of transfer RNAs: similarity and variability. *Wiley Interdiscipl. Rev.RNA*, **3**, 37–61.
- Auffinger, P. and Westhof, E. Location and distribution of modified nucleotides in tRNA. In: Grosjean, H. and Benne, R. (eds), *Modification and editing of RNA*. ASM press, Washington DC, USA, pp. 103–112.
- Stuart, J.W., Gdaniec, Z., Guenther, R., Marszałek, M., Sochacka, E., Malkiewicz, A. and Agris, P.F. (2000) Functional anticodon architecture of human tRNA<sup>Lys3</sup> includes disruption of intraloop hydrogen bonding by the naturally occurring amino acid modification, t<sup>6</sup>A. *Biochemistry*, **39**, 13396–13404.
- Murphy, F.V., Ramakrishnan, V., Malkiewicz, A. and Agris, P.F. (2004) The role of modifications in codon discrimination by tRNA(Lys)UUU. *Nat. StructMol. Biol.*, **11**, 1186–1191.
- Chheda, G.B., Hall, R.H., Mozejko, J., Magrath, D.I., Schweizer, M.P., Stasiuk, L. and Taylor, P.R. (1969) Aminoacyl nucleosides. VI. Isolation and preliminary characterization of threonyladenine derivatives from transfer ribonucleic acid. *Biochemistry*, **8**, 3278–3282.
- Chheda, G., Hong, C. and Piskorz, C. (1972) Biosynthesis of *N*-(purin-6-ylcarbamoyl)-L-threonine riboside. Incorporation of L-threonine in vivo into modified nucleoside of transfer ribonucleic acid. *Biochem. J.*, **127**, 515–519.
- Körner, A. and Söll, D. (1974) *N*-(purin-6-ylcarbamoyl)threonine: biosynthesis in vitro in transfer RNA by an enzyme purified from *Escherichia coli*. *FEBS Lett.*, **39**, 301–306.
- Galperin, M.Y. (2004) Conserved hypothetical proteins: prioritization of targets for experimental study. *Nucleic Acids Res.*, **32**, 5452–5463.
- El Yacoubi, B., Lyons, B., Cruz, Y., Reddy, R., Nordin, B., Agnelli, F., Williamson, J.R., Schimmel, P., Swairjo, M.A. and de Crecy-Lagard, V. (2009) The universal YrdC/Sua5 family is required for the formation of threonylcarbamoyladenine in tRNA. *Nucleic Acids Res.*, **37**, 2894–2909.
- El Yacoubi, B., Hatin, I., Deutsch, C., Kahveci, T., Rousset, J., Iwata-Reuyl, D., Murzin, A.G. and de Crecy-Lagard, V. (2011) A role for the universal Kae1/Qri7/YgjD (COG0533) family in tRNA modification. *EMBO J.*, **30**, 882–893.
- Deutsch, C., El Yacoubi, B., de Crecy-Lagard, V. and Iwata-Reuyl, D. (2012) Biosynthesis of Threonylcarbamoyl Adenosine (t<sup>6</sup>A), a Universal tRNA Nucleoside. *J Biol. Chem.*, **287**, 13666–13673.
- Lauhon, C. (2012) Mechanism of N<sup>6</sup>-threonylcarbamoyladenine (t<sup>6</sup>a) biosynthesis: isolation and characterization of the intermediate threonylcarbamoyl-AMP. *Biochemistry*, **51**, 8950–8963.
- Downey, M., Houlsworth, R., Maringe, L., Rollie, A., Brehme, M., Galicia, S., Guillard, S., Partington, M., Zubko, M.K., Krogan, N.J. *et al.* (2006) A Genome-wide screen identifies the evolutionarily conserved KEOPS complex as a telomere regulator. *Cell*, **124**, 1155–1168.
- Kisseleva-Romanova, E., Lopreiato, R., Baudin-Baillieu, A., Rousselle, J., Ilan, L., Hofmann, K., Namane, A., Mann, C. and Libri, D. (2006) Yeast homolog of a cancer-testis antigen defines a new transcription complex. *EMBO J.*, **25**, 3576–3585.
- Daugeron, M., Lenstra, T.L., Frizzarin, M., El Yacoubi, B., Liu, X., Baudin-Baillieu, A., Lijnzaad, P., Decourty, L., Saveanu, C., Jacquier, A. *et al.* (2011) Gcn4 misregulation reveals a direct role for the evolutionary conserved EKC/KEOPS in the t(6)A modification of tRNAs. *Nucleic Acids Res.*, **39**, 1–13.
- Srinivasan, M., Mehta, P., Yu, Y., Prugar, E., Koonin, V.E., Karzai, A.W. and Sternlanz, R. (2011) The highly conserved KEOPS/EKC complex is essential for a universal tRNA modification, t<sup>6</sup>A. *EMBO J.*, **30**, 873–881.
- Naor, A., Thiaville, P.C., Altman-Price, N., Cohen-Or, I., Allers, T., de Crécy-Lagard, V. and Gophna, U. (2012) A genetic investigation of the KEOPS complex in halophilic Archaea. *PLoS One*, **7**, e43013.
- Abdullah, K.M., Lo, R.Y. and Mellors, A. (1991) Cloning, nucleotide sequence, and expression of the Pasteurella haemolytica A1 glycoprotease gene. *J. Bacteriol.*, **173**, 5597–5603.
- Collinet, B., Friberg, A., Brooks, M.A., Elzen, T., Henriot, V., Dziembowski, A., Graille, M., Dur, D., Leulliot, N., Saint

- André, C.S. *et al.* (2011) Strategies for the structural analysis of multi-protein complexes: lessons from the 3D-Repertoire project. *J. Struct. Biol.*, **175**, 147–158.
24. Hecker, A., Leulliot, N., Gabelle, D., Graille, M., Justome, A., Dorlet, P., Brochier, C., Quevillon-Cheruel, S., Le Cam, E., Le Cam, E. *et al.* (2007) An archaeal orthologue of the universal protein Kae1 is an iron metalloprotein which exhibits atypical DNA-binding properties and apurinic-endonuclease activity in vitro. *Nucleic Acids Res.*, **35**, 6042–6051.
25. Lowe, T.M. and Eddy, S.R. (1997) tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.*, **25**, 955–964.
26. Mougin, A., Grégoire, A., Banroques, J., Ségault, V., Fournier, R., Brulé, F., Chevrier-Miller, M. and Branlant, C. (1996) Secondary structure of the yeast *Saccharomyces cerevisiae* pre-U3A snoRNA and its implication for splicing efficiency. *RNA*, **2**, 1079–1093.
27. Arnaud, N., Cheynet, V., Oriol, G., Mandrand, B. and Mallet, F. (1997) Construction and expression of a modular gene encoding bacteriophage T7 RNA polymerase. *Gene*, **199**, 149–156.
28. Grosjean, H., Droogmans, L., Roovers, M. and Keith, G. (2007) Detection of enzymatic activity of transfer RNA modification enzymes using radiolabeled tRNA substrates. *Methods Enzymol.*, **425**, 55–101.
29. Mao, D.Y.L., Neculai, D., Downey, M., Orlicky, S., Haffani, Y.Z., Ceccarelli, D.F., Ho, J.S.L., Szilard, R.K., Zhang, W., Ho, C.S. *et al.* (2008) Atomic structure of the KEOPS complex: an ancient protein kinase-containing molecular machine. *Mol. Cell*, **32**, 259–275.
30. Morin, A., Auxilien, S., Senger, B., Tewari, R. and Grosjean, H. (1998) Structural requirements for enzymatic formation of threonylcarbamoyladenine (t6A) in tRNA: an in vivo study with *Xenopus laevis* oocytes. *RNA*, **4**, 24–37.
31. Madore, E., Florentz, C., Giegé, R., Sekine, S., Yokoyama, S. and Lapointe, J. (1999) Effect of modified nucleotides on *Escherichia coli* tRNA<sup>Glu</sup> structure and on its aminoacylation by glutamyl-tRNA synthetase. Predominant and distinct roles of the mnm5 and s2 modifications of U34. *Eur. J. Biochem.*, **266**, 1128–1135.
32. Tisné, C., Rigourd, M., Marquet, R. and Ehresmann, C. (2000) NMR and biochemical characterization of recombinant human tRNA (Lys) 3 expressed in *Escherichia coli*: identification of posttranscriptional nucleotide modifications. *RNA*, **6**, 1403–1412.
33. Grosjean, H., Keith, G. and Droogmans, L. (2004) Detection and quantification of modified nucleotides in RNA using thin-layer chromatography. *Methods Mol. Biol.*, **265**, 357–391.
34. Stocchetto, S., Marin, O., Carignani, G. and Pinna, L.A. (1997) Biochemical evidence that *Saccharomyces cerevisiae* YGR262c gene, required for normal growth, encodes a novel Ser/Thr-specific protein kinase. *FEBS Lett.*, **414**, 171–175.
35. Meng, F., Chen, X., Hu, Y., Tang, H., Dang, W. and Zhou, J. (2010) Sua5p is required for telomere recombination in *Saccharomyces cerevisiae*. *Cell Res.*, **20**, 495–498.
36. Wei, F., Suzuki, T., Watanabe, S. and Kimura, S. (2011) Deficit of tRNA<sup>Lys</sup> modification by Cdk11 causes the development of type 2 diabetes in mice. *J. Clin. Invest.*, **121**, 3598–3608.
37. Garcia, G.A. and Goodenough-Lashua, D.M. Mechanisms of RNA modifying and editing enzymes. In: Dans Grosjean, H. and Benne, R. (eds), *Modification and Editing of RNA*. ASM Press, Washington DC, pp. 135–168.
38. Kuratani, M., Kasai, T., Akasaka, R., Higashijima, K., Terada, T., Kigawa, T., Shinkai, A., Bessho, Y. and Yokoyama, S. (2011) Crystal structure of *Sulfolobus tokodaii* sua5 complexed with L-threonine and AMPNP. *Proteins: Structure, Function, and Bioinformatics*, **79**, 2065–2075.
39. Parthier, C., Görlich, S., Jaenecke, F., Breithaupt, C., Bräuer, U., Fandrich, U., Clausnitzer, D., Wehmeier, U.F., Böttcher, C., Scheel, D. *et al.* (2012) The O-Carbamoyltransferase TobZ catalyzes an ancient enzymatic reaction. *Angew. Chem. Int. Ed. Engl.*, **51**, 4046–4052.
40. Cantara, W.A., Crain, P.F., Rozenski, J., McCloskey, J.A., Harris, K.A., Zhang, X., Vendeix, F.A., Fabris, D. and Agris, P.F. (2011) The RNA Modification Database, RNAMDB: 2011 update. *Nucleic Acids Res.*, **39**, D195–D201.
41. Czerwoniec, A., Dunin-Horkawicz, S., Purta, E., Kaminska, K.H., Kasprzak, J.M., Bujnicki, J.M., Grosjean, H. and Rother, K. (2009) MODOMICS: a database of RNA modification pathways. 2008 update. *Nucleic Acids Res.*, **37**, D118–D121.
42. Woese, C.R. and Fox, G.E. (1977) Phylogenetic structure of the prokaryotic domain: the primary kingdoms. *Proc. Natl Acad. Sci. USA*, **74**, 5088–5090.
43. Björk, G.R., Ericson, J.U., Gustafsson, C.E., Hagervall, T.G., Jönsson, Y.H. and Wikström, P.M. (1987) Transfer RNA modification. *Annu. Rev. Biochem.*, **56**, 263–287.
44. Forster, A.C. and Church, G.M. (2006) Towards synthesis of a minimal cell. *Mol. Syst. Biol.*, **2**, doi: 10.1038/msb4100090.
45. de Crecy-Lagarde, V., Marck, C. and Grosjean, H. (2012) Decoding in *Candidatus Riesia pediculicola*, close to minimal tRNA modification set? *Trends Cell Mol. Biol.*, **7**, 11–34.
46. de Crecy-Lagarde, V., Marck, C., Brochier-Armanet, C. and Grosjean, H. (2007) Comparative RNomics and modomics in mollicutes: prediction of gene function and evolutionary implications. *IUBMB Life*, **59**, 634–658.
47. Vetsigian, K., Woese, C. and Goldenfeld, N. (2006) Collective evolution and the genetic code. *Proc. Natl Acad. Sci. USA*, **103**, 10696–10701.