



HAL
open science

SIMULTANEOUS RESOLUTION OF SINGULARITIES IN THE NASH CATEGORY: FINITENESS AND EFFECTIVENESS

Kartoue Mady Demdah

► **To cite this version:**

Kartoue Mady Demdah. SIMULTANEOUS RESOLUTION OF SINGULARITIES IN THE NASH CATEGORY: FINITENESS AND EFFECTIVENESS. *Revista Matematica Complutense*, 2015, 29 (1), pp.207-224. 10.1007/s13163-015-0176-8. hal-00962155v1

HAL Id: hal-00962155

<https://hal.science/hal-00962155v1>

Submitted on 20 Mar 2014 (v1), last revised 11 May 2016 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SIMULTANEOUS RESOLUTION OF SINGULARITIES IN THE NASH CATEGORY: FINITENESS AND EFFECTIVENESS

DEMDAH KARTOUE MADY

ABSTRACT. In this paper we present new proofs using real spectra of the finiteness theorem on Nash trivial simultaneous resolution and the finiteness theorem on Blow-Nash triviality for isolated real algebraic singularities. That is, we prove that a family of Nash sets in a Nash manifold indexed by a semialgebraic set always admits a Nash trivial simultaneous resolution after a partition of the parameter space into finitely many semialgebraic pieces and in the case of isolated singularities it admits a finite Blow-Nash trivialization. We also complement the finiteness results with recursive bounds.

INTRODUCTION

In [CS], M.Coste and M. Shiota proved Nash triviality in family, which can be seen as the Nash version of the Ehresmann's Lemma which gives a trivialization of proper submersion. The additional difficulty in obtaining Nash triviality is that integration of vector fields cannot be used. Several authors (Fukui, Koike, Shiota...) have obtained similar results which are useful for classification of singularities, for instance in the study of Blow-analytic equivalence. In this paper we give new proofs of such results using real spectra.

Let N be an *affine* Nash manifold (i.e. Nash submanifold of affine space \mathbb{R}^n) and J be a semialgebraic subset of \mathbb{R}^s . A mapping $F : N \times J \rightarrow \mathbb{R}^k$ is said to be semialgebraic family of Nash maps if F is semialgebraic and for any $t \in J$ the map $f_t : N \rightarrow \mathbb{R}^k$ defined by $f_t(x) = F(x, t)$ is Nash. For $Q \subset J$, we denote $F|_{N \times Q}$ by F_Q . We say that the family of zero-sets $F_Q^{-1}(0)$ admits a *Nash trivial simultaneous resolution* along Q if there exists a desingularization of the *Nash set* defined by $F_Q = 0$ which is Nash trivial. We say also that the family of zero-sets $F_Q^{-1}(0)$ admits a *Blow-Nash trivialization* along Q if there exist a desingularization of the *Nash set* defined by $F_Q = 0$ and a Nash triviality upstairs which induces a semialgebraic trivialization of the zero-set $F_Q^{-1}(0)$ (for precise definitions, see Sect.1). Here are the results we prove in this paper.

Theorem (I). *Let $F : N \times J \rightarrow \mathbb{R}^k$ be semialgebraic family of Nash maps from N to \mathbb{R}^k . Then there exists a finite partition $J = J_1 \cup \dots \cup J_p$ which satisfies the following conditions:*

- (1) *Each J_i is Nash manifold which is Nash diffeomorphic to an open simplex in some Euclidean space.*
- (2) *For each i , $F_{J_i}^{-1}(0)$ admits a Nash trivial simultaneous resolution along J_i .*

2010 *Mathematics Subject Classification.* 14P10 - 14P20.

Key words and phrases. Nash manifolds, Resolution, Singularities, Real spectrum, Blow-Nash. The author is partially supported by CONFOFOR/TCHAD fund.

When for all $t \in J$, $f_t^{-1}(0)$ has only isolated singularities, the following result holds true.

Theorem (II). *Let $F : N \times J \rightarrow \mathbb{R}^k$ be semialgebraic family of Nash maps from N to \mathbb{R}^k such that for all $t \in J$, $f_t^{-1}(0)$ has only isolated singularities. Then there exists a finite partition $J = J_1 \cup \dots \cup J_p$ which satisfies the following conditions:*

- (1) *Each J_i is Nash manifold which is Nash diffeomorphic to an open simplex in some Euclidean space.*
- (2) *For each i , $F_{J_i}^{-1}(0)$ admits a Blow-Nash trivialization along J_i .*

In the setting of Nash germs, these theorems were first proved by Fukui, Koike and Shiota (cf. [FKS], [K1]). In this paper, we obtain the same results, but in the global case with N an affine Nash manifold not necessary compact using the techniques of real spectra. First, we verify that the desingularization theorem holds true for the category of Nash sets over an arbitrary real closed field, equipped with a sheaf of Nash functions. The precise definition of this category needs the use of the real spectrum and of finite sheaves of ideals of Nash functions over this real spectrum (see Sect.2). That is, we check that the above category verifies the list of the criteria given by Bierston and Milman (cf. [BM1] p.234) to apply the desingularization Theorem. We apply the desingularization Theorem for the fiber of the family at a point of the real spectrum of the parameter space. This fiber is defined over a real closed field which is not \mathbb{R} a priori, which explains the need to have a desingularisation valid over any real closed field. The finiteness follows from the compactness property of the real spectrum of the parameter space.

The paper is organized as follows. In the first section we recall useful results and give the definition of Blow-Nash triviality. In the second section, we recall the notion of real spectrum and some related results which will be used. In section three, we check that the desingularization theorem holds true over an arbitrary real closed field. In particular, we prove that glueing local Blowing-up of an affine Nash manifold gives an affine Nash manifold (we need always to work with affine Nash manifolds). As consequence of the section three, we prove in section four the finiteness of Nash trivial simultaneous resolution (**Theorem (I)**). Finally, in the last section we prove the **Theorem (II)**.

Acknowledgment - I would like to thank Michel Coste for his advice during the preparation of this paper.

1. USEFUL RESULTS

In this paper, a *Nash manifold* is a semialgebraic subset of \mathbb{R}^n (or of R^n , where R is a real closed field) which is also a C^∞ submanifold. For more details see [S] or [BCR] for the use of an arbitrary real closed field. We shall work in this paper only with *affine* Nash manifolds to avoid some pathological situation. We define a *Nash subset* (or a *Nash variety*) of a Nash manifold M as a subset of M which is the zero set of a Nash function over M .

Let us now introduce the notion of normal crossing that we will use in this paper. Let M be a manifold of dimension m . Let N_1, \dots, N_r be Nash proper

submanifolds in M . Let $P \in \bigcap_{i=1}^r N_i$. The submanifolds N_1, \dots, N_r are said to have *simultaneous normal crossings at a point P* when there exist (x_1, \dots, x_m) a local coordinates system for M at P , J_1, \dots, J_r disjoint subsets of $\{1, \dots, m\}$ such that $N_i = \{x_j = 0 \text{ for } j \in J_i\}$ for $i = 1, \dots, r$.

Now consider N_1, \dots, N_u a finite family of Nash proper submanifolds of M . The manifolds N_1, \dots, N_u are said to have *simultaneous only normal crossings* if for all $P \in M$, the submanifolds N_i which contain P have simultaneous normal crossings at P .

Let M and M' be Nash manifolds, and let \mathcal{V} be a Nash subset of M . A map $\Pi : M' \rightarrow M$ is said to be a *Nash resolution* of \mathcal{V} in M if there exists a finite sequence of blowing-ups $\sigma_{j+1} : M_{j+1} \rightarrow M_j$ with smooth centers C_j (where σ_j, M_j and C_j are of Nash class) such that:

- Π is the composite of σ_j 's;
- the critical set of Π is a union of smooth Nash divisors D_1, \dots, D_r ;
- the strict transform of \mathcal{V} in M by Π (denoted by \mathcal{V}') is a Nash submanifold of M ;
- $\mathcal{V}', D_1, \dots, D_r$ simultaneously have only normal crossings.

It is folklore that the desingularization theorem (cf.[H1], [H2], [BM1] and [BM2]) apply in the Nash category to give:

Theorem 1.1. (*Desingularization*)

For a Nash subset V of a Nash manifold M , there exists a Nash resolution $\Pi : M' \rightarrow M$ of V in M .

We shall make precise the Nash category in the following section, working over an arbitrary real closed field.

Now let M, J be Nash manifolds and \mathcal{V} a Nash subset of M . Let $\Pi : \mathcal{M} \rightarrow M$ be a Nash resolution of \mathcal{V} and $q : M \rightarrow J$ be an onto Nash submersion. For any $t \in J$, let $\mathcal{M}_t = (q \circ \Pi)^{-1}(t)$, $M_t = q^{-1}(t)$ and $V_t = \mathcal{V} \cap M_t$. The map Π is said to be a *Nash simultaneous resolution of \mathcal{V} in M over J* if

- $q \circ \Pi|_{\mathcal{V}} : \mathcal{V}' \rightarrow J$,
- $q \circ \Pi|_{D_{j_1} \cap \dots \cap D_{j_s}} : D_{j_1} \cap \dots \cap D_{j_s} \rightarrow J$,
- $q \circ \Pi|_{\mathcal{V}' \cap D_{j_1} \cap \dots \cap D_{j_s}} : \mathcal{V}' \cap D_{j_1} \cap \dots \cap D_{j_s} \rightarrow J$ ($1 \leq j_1 < \dots < j_s \leq r$)

are onto submersions;

Now let $\Pi : \mathcal{M} \rightarrow M$ be a Nash simultaneous resolution of \mathcal{V} over J , and let $t_0 \in J$. We say that Π gives a *Nash trivial simultaneous resolution of \mathcal{V} in M over J* if there is a Nash diffeomorphism $\Phi : \mathcal{M} \rightarrow \mathcal{M}_{t_0} \times J$ such that

- (1) $(q \circ \Pi) \circ \Phi^{-1} : \mathcal{M}_{t_0} \times J \rightarrow J$ is the natural projection,
- (2) $\Phi(\mathcal{V}') = \mathcal{V}'_{t_0} \times J$, $\Phi(D_i) = D_{i,t_0} \times J$,
- (3) $\Phi^{-1}(x, t_0) = x$, for every $x \in \mathcal{M}_{t_0}$.

When the Nash trivialization Φ induces the semialgebraic trivialization of \mathcal{V} in M over J , we say that (M, \mathcal{V}) admits a *Π -Blow-Nash trivialization along J* . This is equivalent to saying that there is a semialgebraic homeomorphism ϕ such that the

following diagram commutes:

$$\begin{array}{ccccc}
\mathcal{M} & \xrightarrow{\Pi_i} & M & \xrightarrow{q} & J \\
\Phi \downarrow & & \phi \downarrow & & \parallel \\
\mathcal{M}_{t_0} \times J & \xrightarrow{\Pi_{|\mathcal{M}_{t_0}} \times id_J} & M_{t_0} \times J & \xrightarrow{p} & J
\end{array}$$

where $p(x, t) = t$ for $(x, t) \in M_{t_0} \times J$.

2. REAL SPECTRUM, FAMILY OF NASH FUNCTIONS

In order to work over any real closed field, real spectrum is necessary. So, in this section we recall certain facts about Nash sheaves and real spectrum.

Let M be a Nash manifold defined over a real closed field R . In general, one cannot define the sheaf of Nash functions over M by the order topology, because this topology can be totally disconnected. Consider for example the locally Nash function f over \mathbb{R}_{alg} defined by $f = 0$ over $\mathbb{R}_{alg} \cap (-\infty, \pi)$ and $f = 1$ over $\mathbb{R}_{alg} \cap (\pi, +\infty)$: it is not a Nash function over \mathbb{R}_{alg} since it is not semialgebraic over \mathbb{R}_{alg} . For $R = \mathbb{R}$ the sheaf \mathcal{N} of rings of Nash functions germs on M is well defined, but in the non-compact case another problem may appear. The sheaf \mathcal{N} is coherent, but in [LT], the authors observe that there is a coherent sheaf of ideals of germs of Nash functions which is not generated by its global sections. For instance, the sheaf $\mathcal{I}_{\mathbb{Z}}$ of germs of Nash functions vanishing on $\mathbb{Z} \subset \mathbb{R}$ is not generated by global Nash functions.

Let us recall now the notion of real spectrum which will be used in this paper. Let us consider $M \subset R^v$ a semialgebraic subset. Define \widetilde{M} to be the set of ultrafilters of the Boolean algebra of semialgebraic subsets of M . To any $x \in M$, we associate α_x the principal ultrafilter of semialgebraic subsets of M containing x . If S is a semialgebraic subset of M , \widetilde{S} may be identified with the subset of \widetilde{M} consisting of those ultrafilters $\alpha \in \widetilde{M}$ such that $S \in \alpha$. The real spectrum has the following compactness property: for any covering $\widetilde{M} = \bigcup_{i \in I} \widetilde{S}_i$ where S_i are semialgebraic

subset of M , there exists a finite subset J of I such that $\widetilde{M} = \bigcup_{i \in J} \widetilde{S}_i$. We define a

topology on \widetilde{M} by taking as basis of open subsets all \widetilde{U} where U is a semialgebraic open subset of M . Denote it \mathcal{T}^{sa} .

Now, let M be a Nash submanifold of R^n . There exists a unique sheaf of rings \mathcal{N} on \widetilde{M} such that for every semialgebraic open subset U of M , $\mathcal{N}(\widetilde{U}) = \mathcal{N}(U)$ the ring of Nash functions on U . Let be $\alpha \in \widetilde{M}$, the stalk of \mathcal{N} at α is defined by $\mathcal{N}_\alpha = \varinjlim_{\alpha \in \widetilde{U}} \mathcal{N}(U)$.

The proof of the existence and the uniqueness of the sheaf \mathcal{N} can be found in [[BCR], Proposition 8.8.2] and the fact that, for any $\alpha \in \widetilde{M}$, the stalk \mathcal{N}_α is a henselian local ring. Its residue field is denoted by $k(\alpha)$ which is a real closed

extension of R . If $\alpha = x \in M$, of course $\mathcal{N}_\alpha = \mathcal{N}_x$ the ring of germs of Nash functions at x and $k(x) = R$ (cf. [BCR], Proposition 8.8.3). Recall that the dimension of α is the smallest dimension of U for $U \in \alpha$. Consider the completion $\widehat{\mathcal{N}}_\alpha$ of the ring $\mathcal{N}_{\widetilde{M},\alpha}$, then $\widehat{\mathcal{N}}_{\widetilde{M},\alpha} \cong k(\alpha)[[X_1, \dots, X_r]]$ with $r = \dim(M) - \dim(\alpha)$.

Let us denote by \mathbf{Nash}_R the Nash category over a real closed field R . On the Nash manifold \widetilde{M} (a smooth space in the sense that for all $\alpha \in \widetilde{M}$, \mathcal{N}_α is a regular local ring as consequence of above paragraph) let us consider a locally finite sheaf of ideals $\mathcal{I} \subset \mathcal{N}$, that is, a sheaf of ideals locally (for the topology \mathcal{T}^{sa}) generated by finitely many Nash sections. Then there exists an ideal I of the ring of Nash function $\mathcal{N}(M)$ such that $\mathcal{I} = I \cdot \mathcal{N}_{\widetilde{M}}$ and $\mathcal{N}/\mathcal{I}(M) = \mathcal{N}(M)/I$ (see [CRS]). Thus one can note that the problems indicated in the beginning of this section are settled with the use of the real spectrum. So, we have that $Z(\mathcal{I}) = \widetilde{Z(I)}$ is a Nash subset of \widetilde{M} . Then in \mathbf{Nash}_R , $(\widetilde{Z(I)}, \mathcal{N}/\mathcal{I})$ is a closed subspace of \widetilde{M} . The family of Nash sets on \widetilde{M} defines a Zariski topology which is noetherian.

Now, let us fix some notations about semialgebraic family which will be used. Let S be a semialgebraic subset of R^s and $U \subset R^n \times S$ be a semialgebraic set. Then one can write

$$U = \{(x, t) \in R^n \times S : \Phi(x, t)\}$$

with $\Phi(x, t)$ a first order formula in theory of real closed field with parameter in R . Denote $t(\alpha) = (t_1(\alpha), \dots, t_s(\alpha)) \in k(\alpha)^s$ where $t_i(\alpha)$, with $i = 1, \dots, s$, are images of coordinate functions in $k(\alpha)$. Then for $\alpha \in \widetilde{S}$ set

$$U_\alpha = \{x \in k(\alpha)^n : \Phi(x, t(\alpha))\}$$

which is a semialgebraic subset of $k(\alpha)^n$ (the fiber of the semialgebraic family U at α).

Reciprocally, if $\Omega \subset k(\alpha)^n$ is a semialgebraic set, there is a semialgebraic set S with $\alpha \in \widetilde{S}$ and a semialgebraic set $U \subset R^n \times S$ such that $U_\alpha = \Omega$. We have also that $U_\alpha = V_\alpha$ if and only if there exists a semialgebraic set S with $\alpha \in \widetilde{S}$ such that $U_S = V_S$.

Hence, any semialgebraic subset of $k(\alpha)^n$ can be represented as a fiber at α of a semialgebraic family of semialgebraic subsets of R^n . Two families of semialgebraic sets have the same fiber in α if and only if they coincide over a semialgebraic set S with $\alpha \in \widetilde{S}$ (idem for the semialgebraic functions). After that, we make an abuse of notation writing U_α the semialgebraic subsets defined over $k(\alpha)$ (in $k(\alpha)^n$).

The philosophy behind the use of fibers at points of the real spectrum is that any property which can be written in the first order language of real closed fields is true at α if and only if it is true at all $t \in S$ for some semialgebraic set such that $\alpha \in \widetilde{S}$. This applies for instance for the property of being the graph of a semialgebraic function, and so one can define the fiber f_α for a semialgebraic family of semialgebraic functions $(f_t)_{t \in S}$, for any $\alpha \in \widetilde{S}$.

The following theorems will be used to prove our main results.

Theorem 2.1. ([BCR], Proposition 8.10.1)

Let $S \subset R^s$ be a semialgebraic set and $\alpha \in \tilde{S}$. Let U be an open semialgebraic subset of $R^m \times S$, and $F : U \rightarrow R$, a semialgebraic family of functions parametrized by S . If, for every $t \in S$ the fiber $f_t : U_t \rightarrow R$ is a Nash function, then for any $\alpha \in \tilde{S}$, the fiber $f_\alpha : U_\alpha \rightarrow k(\alpha)$ is also a Nash function.

By [CS], the ultrafilter $\alpha \in \tilde{R}^s$ are generated by Nash submanifolds of R^s (not closed), Nash diffeomorphic to open simplices. Then we have the following theorems.

Theorem 2.2. ([BCR], Proposition 8.10.3)

Let $\alpha \in \tilde{R}^s$, and let U_α be an open semialgebraic subset of $k(\alpha)^m$, and $f_\alpha : U_\alpha \rightarrow k(\alpha)$, a Nash function. Then there exist a Nash submanifold $Q \subset R^s$, with $\alpha \in \tilde{Q}$, an open semialgebraic subset U_Q of $R^m \times Q$ and a semialgebraic family of functions $f_Q : U_Q \rightarrow R$ parametrized by Q , such that f_Q is a Nash mapping.

Theorem 2.3. ([CS], Proposition 2.2)

Let $B \subset R^s$ be a semialgebraic set, and let X be a semialgebraic subset of $R^n \times B$. Let $\alpha \in \tilde{B}$ be such that X_α is a Nash submanifold of $k(\alpha)^n$. Then there is a Nash submanifold Q of R^s , $Q \subset B$, such that $\alpha \in \tilde{Q}$, X_Q is a Nash submanifold of $R^n \times Q$, and the projection $p : X_Q \rightarrow Q$ is a submersion.

We prove the following technical lemma.

Lemma 2.4. Let $\alpha \in \tilde{R}^s$, and let $N_{1\alpha}$ and $N_{2\alpha}$ be closed Nash submanifolds of an affine Nash manifold M_α . Then

- (i) If there is $S \subset R^s$ semialgebraic subset with $\alpha \in \tilde{S}$ such that for all $t \in S$, N_{1t} and N_{2t} are closed Nash submanifolds of M_t with only normal crossings, then $N_{1\alpha}$ and $N_{2\alpha}$ have only normal crossings.
- (ii) If $N_{1\alpha}$ and $N_{2\alpha}$ have only normal crossings, then there exists a Nash submanifold $Q \subset R^s$, with $\alpha \in \tilde{Q}$ such that N_{1Q} and N_{2Q} have only normal crossings and the projections $N_{1Q} \rightarrow Q$, $N_{2Q} \rightarrow Q$ and $N_{1Q} \cap N_{2Q} \rightarrow Q$ are Nash submersions.

Proof. (i) Assume that there is a semialgebraic subset $S \subset R^s$ with $\alpha \in \tilde{S}$ such that for all $t \in S$, N_{1t} and N_{2t} are closed Nash submanifolds of M_t with only normal crossings. For $t \in S$, note that N_{1t} and N_{2t} have only normal crossings is equivalent to saying that for any $x \in N_{1t} \cap N_{2t}$, the codimension of $T_x N_{1t} \cap T_x N_{2t}$ is equal to the sum of the codimensions of $T_x N_{1t}$ and $T_x N_{2t}$ which can be expressed by a first order formula of the theory of real closed fields. In fact, being a tangent vector at x and the linear independence of vectors can be expressed into a first order formula of the theory of real closed fields. It follows that $N_{1\alpha}$ and $N_{2\alpha}$ have only normal crossings.

- (ii) Using the first part of the proof and Theorem 2.3, we deduce that there exists a Nash submanifold $Q \subset R^s$, with $\alpha \in \tilde{Q}$ such that N_{1Q} and N_{2Q} have only

normal crossings and the projections $N_{1Q} \rightarrow Q$, $N_{2Q} \rightarrow Q$ and $N_{1Q} \cap N_{2Q} \rightarrow Q$ are Nash submersions. This completes the proof. \square

Remark 2.5. One can easily verify the Lemma 2.4 in general case with an arbitrary finite number of Nash submanifolds.

Now, let us recall the theorem of Nash Isotopy. Let M be a Nash manifold and $\varpi : M \rightarrow \mathbb{R}^s$ be a Nash submersion. For $Z \subset M$, $Q \subset \mathbb{R}^s$ and $t \in \mathbb{R}^s$ let us denote by $Z_Q = Z \cap \varpi^{-1}(Q)$ and $Z_t = Z \cap \varpi^{-1}(t)$. The following result is proved in [K2]:

Theorem 2.6. ([K2], Theorem I)

Let N_1, \dots, N_r be Nash submanifolds of M which have only normal crossings. Let $\varpi : M \rightarrow \mathbb{R}^s$, $s > 0$, be Nash submersion such that for every $0 \leq i_1 < \dots < i_q \leq r$, $\varpi|_{N_{i_1} \cap \dots \cap N_{i_q}} : N_{i_1} \cap \dots \cap N_{i_q} \rightarrow \mathbb{R}^s$ is submersion (unless $N_{i_1} \cap \dots \cap N_{i_q} = \emptyset$). Then there exists a finite partition of \mathbb{R}^s into Nash simplices Q_j and $t_j \in Q_j$ for $j = 1, \dots, u$ and for any j there exists a Nash diffeomorphism

$$\varphi_j = (\varphi'_j, \varpi_j) : M_{Q_j} \rightarrow M_{t_j} \times Q_j$$

such that $\varphi|_{M_{t_j}} = id_{M_{t_j}}$ and $\varphi_j(N_{i_{Q_j}}) = N_{i_{t_j}} \times Q_j$, for $i = 1, \dots, r$.

Let us recall that a semialgebraic subset of \mathbb{R}^n is said of *complexity* at most (p, q) if it admits a description as follows

$$\bigcup_{i=1}^s \bigcap_{j=1}^{k_i} \{x \in \mathbb{R}^n | f_{ij}(x) *_{ij} 0\},$$

where $f_{ij} \in R[X_1, \dots, X_n]$, and $*_{ij} \in \{<, >, =\}$, $\sum_{i=1}^s k_i \leq p$, $\deg(f_{ij}) \leq q$ for $i = 1, \dots, s$ and $j = 1, \dots, r_i$.

We call *Nash NC-system* any family $(M; N_1, \dots, N_r)$ such that M is an affine Nash manifold and N_i for $i = 1, \dots, r$ are closed Nash submanifolds of M having simultaneously only normal crossings. The *complexity* of a Nash NC-system $(M; N_1, \dots, N_r)$ is bounded by (p, q) if M and N_1, \dots, N_r have complexities bounded by (p, q) .

As consequence of Nash isotopy theorem, we prove that there is finitely and effectively many models of Nash NC-systems with bounded complexity.

Theorem 2.7. *Given the integers n, p, q and r , there exist a triplet of integers (u, v, w) such that for each $i = 1, \dots, u$ there is a Nash NC-system $(M_i, N_{i_1}, \dots, N_{i_r})$ of complexity bounded by (p, q) and for any Nash NC-system (M, N_1, \dots, N_r) of complexity bounded by (p, q) there is $i \in \{1, \dots, u\}$ and a Nash diffeomorphism Φ between M and M_i whose graph has complexity bounded by (v, w) and such that $\Phi(N_j) = N_{i_j}$ for $j = 1, \dots, r$.*

Moreover, u, v, w are bounded by recursive functions of n, p, q and r .

Proof. Using the technique in [D] we can parametrize the Nash NC-systems of complexity bounded by (p, q) by a semialgebraic subset defined by a first order formula of the theory of real closed fields with coefficients in \mathbb{Z} . Indeed we know

that the fact to be a Nash manifold of complexty bounded by (p, q) can be translated into a first order formula of the theory of real closed fields (see [R], Proposition 3.5 and [D], Proposition 5.7 (i)) and by the proof of Lemme 2.4 the normal crossing property also can be translated into a first order formula of the theory of real closed fields.

Applying [CS] Corollary 2.3 and the Theorem 2.6 to the projection of the family of Nash NC-systems of complexity bounded by (p, q) on its parameter space, there is a triplet of integers (u, v, w) such that for each $i = 1, \dots, u$ there is a Nash NC-system $(M_i, N_{i_1}, \dots, N_{i_r})$ of complexity bounded by (p, q) and for any Nash NC-system (M, N_1, \dots, N_r) of complexity bounded by (p, q) there is $i \in \{1, \dots, u\}$ and a Nash diffeomorphism Φ between M and M_i whose graph have the complexity bounded by (v, w) and such that $\Phi(N_j) = N_{i_j}$ for $j = 1, \dots, r$.

Now, fixing n, p, q and r , by the first part of the proof, there exists $(u, v, w) \in \mathbb{N}^3$ such that the following formula holds:

$$\Psi(n, p, q, r, u, v, w) :=$$

“There are u Nash NC-systems $(M_i, N_{i_1}, \dots, N_{i_r})$, $i = 1, \dots, u$, with complexity bounded by (p, q) such that for every Nash NC-system (M, M, M_1, \dots, M_r) in \mathbb{R}^n of complexity at most (p, q) there are $i \in \{1, \dots, u\}$ and a Nash diffeomorphism Φ between M and M_i whose graph have the complexity bounded by (v, w) and such that $\Phi(N_j) = N_{i_j}$ for $j = 1, \dots, r$.”

There is an algorithm with input n, p, q, r, u, v, w and output the formula $\Psi(n, p, q, r, u, v, w)$. So the decidability of the first order theory of real-closed fields implies that the function

$$(n, p, q, r) \mapsto (u(n, p, q, r), v(n, p, q, r), w(n, p, q, r)),$$

is bounded by a recursive function. This achieves the proof. \square

3. DESINGULARIZATION ON THE NASH CATEGORY OVER AN ARBITRARY REAL CLOSED FIELD

In [BM1] and [BM2], the authors prove that the desingularization theorem holds true for the objects X belonging to the category \mathcal{A} of locally ringed spaces over a field k of characteristic zero which satisfies the four properties that we shall directly verify for the category \mathbf{Nash}_R which interest us in this paper.

- (i) Every locally ringed space $(\tilde{X}, \mathcal{N}_{\tilde{X}})$ in the category \mathbf{Nash}_R is a closed subspace $(Z(\mathcal{I}), \mathcal{N}_{\tilde{M}}/\mathcal{I})$ of a Nash manifold $(\tilde{M}, \mathcal{N}_{\tilde{M}})$ which is a smooth space (all its local rings are regular local rings).
- (ii) For $(\tilde{M}, \mathcal{N}_{\tilde{M}})$ a smooth space in the category \mathbf{Nash}_R , there is a neighbourhood basis given by (the support of) Nash coordinate charts \tilde{U} . That is, there is coordinates system (x_1, \dots, x_m) on \tilde{U} consisting of Nash functions on \tilde{U} . Indeed each $x_i \in \mathcal{N}_{\tilde{M}}(\tilde{U})$ and the partial derivatives $\frac{\partial^{|\beta|}}{\partial x^\beta} = \frac{\partial^{\beta_1 + \dots + \beta_m}}{\partial x_1^{\beta_1} \dots \partial x_m^{\beta_m}}$ make sense as transformations $\mathcal{N}_{\tilde{M}}(\tilde{U}) \rightarrow \mathcal{N}_{\tilde{M}}(\tilde{U})$.

Furthermore, for each $\alpha \in \tilde{U}$, the residue fields $k(\alpha)$ of \tilde{M} at α , by [BCR] p.162, there is an injective mapping $T_\alpha : \mathcal{N}_{\tilde{M}, \alpha} \rightarrow k(\alpha)[[X_1, \dots, X_r]]$ which maps each element of $\mathcal{N}_{\tilde{M}, \alpha}$ to its Taylor expansion, such that T_α induces an isomorphism from the completion $\hat{\mathcal{N}}_{\tilde{M}, \alpha}$ to $k(\alpha)[[X_1, \dots, X_r]]$ where $r = m - \dim(\alpha)$. One can verify that T_α commutes with differentiation: $T_\alpha \circ (\frac{\partial^{|\beta|}}{\partial x^\beta}) = \frac{\partial^{|\beta|}}{\partial x^\beta} \circ T_\alpha$, for all $\beta \in \mathbb{N}^r$.

- (iii) For each $\tilde{X} \in \mathbf{Nash}_R$, $\mathcal{N}_{\tilde{X}}$ is a coherent sheaf of ideals of Nash functions and \tilde{X} , in this category, as shown in the section 2, is noetherian. Every decreasing sequence of closed subspaces of \tilde{X} stabilizes since $\mathcal{N}_{\tilde{X}}(\tilde{X})$ is a noetherian ring.

It remains to show that \mathbf{Nash}_R satisfies the last condition which required it to be closed under blowing-ups. It is clear that the blowing-up of a Nash manifold gives a Nash manifold. But in our case we consider only the *affine* Nash manifolds, and it is not clear that the Nash manifold obtained by blowing-up (which is locally defined) is affine. For instance consider \mathbb{R}/\mathbb{Z} and the Nash open sets $(0, 1)$ and $(\frac{1}{2}, \frac{3}{2})$. We define a Nash structure on \mathbb{R}/\mathbb{Z} by glueing the charts $(0, 1)$ and $(\frac{1}{2}, \frac{3}{2})$ in the following way:

$$(0, 1) \supset (\frac{1}{2}, 1) \xrightarrow{Id} (\frac{1}{2}, 1) \subset (\frac{1}{2}, \frac{3}{2}) \text{ and}$$

$$(0, 1) \supset (0, \frac{1}{2}) \xrightarrow{\sim} (1, \frac{3}{2}) \subset (\frac{1}{2}, \frac{3}{2})$$

$$t \mapsto t + 1,$$

Consider $\varphi \in \mathcal{N}_{\mathbb{R}/\mathbb{Z}}$. The canonical mapping $f : \mathbb{R} \rightarrow \mathbb{R}/\mathbb{Z}$ is a Nash map. Then we have $\varphi \circ f$ is constant because it is periodic and a periodic Nash map is constant. That implies that φ is constant. Then \mathbb{R}/\mathbb{Z} is non-affine.

In the following subsection we prove that the blowing-up of an affine Nash manifold gives an affine Nash manifold.

Global definition of Blowing-up in Nash category

For the sake of simplicity, although we work on the category \mathbf{Nash}_R , we shall write here the objects without the tilde symbol \sim . So, let $C \subset M$ be a closed Nash submanifold of an affine Nash manifold M of codimension c . Let \mathcal{I}_C be the sheaf of ideals of germs of Nash functions vanishing on C . It is known that the sheaf \mathcal{I}_C is a finite sheaf (cf.[S]). It follows from [CRS] that there are finitely many $f_1, \dots, f_r \in \mathcal{N}_M(M)$ such that $\mathcal{I}_C = (f_1, \dots, f_r)\mathcal{N}_M$. Consider the semialgebraic subset $M' \subset M \times \mathbb{P}^{r-1}$ defined as follows:

$$M' = \text{clos}(\{(x, [\xi_1, \dots, \xi_r]) \in (M \setminus C) \times \mathbb{P}^{r-1} : f_i(x)\xi_j = f_j(x)\xi_i; i, j = 1, \dots, r\}),$$

where clos defines the closure respect to the usual topology. Let us show that M' is a blowing-up of M with center C . Consider the projection $\Pi : M' \rightarrow M$. It follows from the construction of M' that $\Pi|_{\Pi^{-1}(M \setminus C)} : \Pi^{-1}(M \setminus C) \rightarrow M \setminus C$ is a Nash diffeomorphism.

Now for any $\alpha_0 \in C$, there is a semialgebraic open neighbourhood U of α_0 in M such that c of the f_i belong to a coordinates system over U , say $f_1, \dots, f_c, x_{c+1}, \dots, x_m$ for simplicity, such that for $c < j \leq r$ there are $h_{j,k} \in \mathcal{N}_M(U)$ with $f_j = h_{j,1}f_1 + \dots + h_{j,c}f_c$ and $k = 1, \dots, c$. By the usual blowing-up of U with center $U \cap C = \{f_1 = \dots = f_c = 0\}$, we get:

$$U' = \{(x, [\xi_1, \dots, \xi_c]) \in U \times \mathbb{P}^{c-1} : f_i(x)\xi_j = f_j(x)\xi_i, 1 \leq i, j \leq c\}.$$

It is clear that U' is a Nash submanifold of $U \times \mathbb{P}^{c-1}$. Let us show that U' is Nash diffeomorphic to $M' \cap \Pi^{-1}(U)$. Let us consider the embedding $\Phi : U \times \mathbb{P}^{c-1} \rightarrow U \times \mathbb{P}^{r-1}$ defined by:

$$(x, [\xi_1, \dots, \xi_c]) \mapsto (x, [\xi_1, \dots, \xi_c, \sum_{i=1}^c h_{c+1_i}(x)\xi_i, \dots, \sum_{i=1}^c h_{r_i}(x)\xi_i]).$$

We *claim* that: $\Phi(U') = M' \cap \Pi^{-1}(U)$. It follows that M' is the blowing-up of M with center C and M' is an affine Nash manifold.

Proof of claim: It is clear that $\Phi(U' \setminus (C \times \mathbb{P}^{c-1})) = \Pi^{-1}(U \setminus C)$.

The set $U' \setminus (C \times \mathbb{P}^{c-1})$ is open and dense in U' , and $\Phi(U')$ is a closed submanifold of $U \times \mathbb{P}^{r-1}$. It follows that $\Pi^{-1}(U)$, which is the closure of $\Pi^{-1}(U \setminus C)$ in $C \times \mathbb{P}^{r-1}$, is equal to $\Phi(U')$. This completes the proof.

Which verifies that **Nash_R** is closed under Blowing-ups. We conclude that the desingularization theorem is valid in the category **Nash_R**.

4. NASH TRIVIAL SIMULTANEOUS RESOLUTION

In this section, we prove the global case of the result of Koike ([K], Theorem I, p.317). In other words, we prove the global case of the finiteness theorem of Nash trivial simultaneous resolution.

Let us first prove the following technical lemma:

Lemma 4.1. *Let $\alpha \in \widetilde{R}^s$, and let N_α be a Nash submanifold of $k(\alpha)^m$. Let $\sigma_\alpha : N_{1\alpha} \rightarrow N_\alpha$ be the Blowing-up of N_α with center C_α a Nash submanifold. Then there exist a Nash submanifold $Q \subset R^s$, with $\alpha \in \widetilde{Q}$, Nash manifolds N_{1Q} , N_Q and $C_Q \subset N_Q$, and a Nash map $\sigma_Q : N_{1Q} \rightarrow N_Q$ the Blowing-up of N_Q with center C_Q such that $N_Q \subset R^m \times Q$ and the projections $N_Q \rightarrow Q$ and $C_Q \rightarrow Q$ are Nash submersions.*

Proof. Since σ_α is a Nash map, its graph Γ_{σ_α} is a semialgebraic set. Then we may write it as follows

$$\Gamma_{\sigma_\alpha} = \{(x, y) \in N_{1\alpha} \times N_\alpha : \Phi(x, y)\}$$

where $\Phi(x, y)$ is a first order formula in the theory of real closed fields with coefficients $a = (a_1, \dots, a_n) \in k(\alpha)^n$ for some $n \in \mathbb{N}$. We may also write $C_\alpha = \{x \in N_\alpha : \Psi(x)\}$ where $\Psi(x)$ is a first order formula in the theory of real closed fields with coefficients $b = (b_1, \dots, b_c) \in k(\alpha)^c$ for some $c \in \mathbb{N}$.

The fact that the Nash map $\sigma_\alpha : N_{1\alpha} \rightarrow N_\alpha$ is the blowing-up of N_α with center C_α can be expressed into a first order formula of theory of real closed fields (see

the construction of Global definition of Blowing-up in Nash category in Sect.2). It follows that there exist a Nash submanifold $Q \subset \mathbb{R}^s$ with $\alpha \in \tilde{Q}$, a Nash submanifold $C_Q \subset N_Q$ and $\sigma_Q : N_{1Q} \rightarrow N_Q$ is the blowing-up of N_Q with smooth center C_Q . Applying also Theorem 2.3, shrinking Q if necessary, the projection $N_{1Q} \rightarrow Q$, $N_Q \rightarrow Q$, and $C_Q \rightarrow Q$ are Nash submersions. This ends the proof. \square

Now we will prove the **Theorem (I)**. Let N be an affine Nash manifold and J a semialgebraic subset of \mathbb{R}^s . Let $F : N \times J \rightarrow \mathbb{R}^k$ be semialgebraic family of Nash maps $f_t : N \rightarrow \mathbb{R}^k$ ($t \in J$).

Proof. of Theorem (I)

As seen in the third section (Sect.3), the Theorem 1.1 is valid over any real closed fields. Let $\alpha \in \tilde{J}$, by the Theorem 2.1, f_α is a Nash map. It follows that the fiber $f_\alpha^{-1}(0) \subset N_{k(\alpha)}$ is a Nash set. By the Nash desingularization Theorem, there exists a Nash resolution of $f_\alpha^{-1}(0)$ in $N_{k(\alpha)}$, $\Pi_\alpha : \mathcal{M}_\alpha \rightarrow N_{k(\alpha)}$. The properties that Π_α must satisfy to be Nash resolution can be translated in a first order formula of the theory of real closed fields (That is a consequence of the construction of Global definition of Blowing-up in Nash category in Sect.3). Consider $D_{1\alpha}, \dots, D_{k\alpha}$ the Nash divisors and V'_α the strict transform of $V_\alpha = f_\alpha^{-1}(0)$ respected to Π_α . By the definition of Π_α , we have that, $D_{1\alpha}, \dots, D_{k\alpha}$ and V'_α have only normal crossings.

So by the Theorem 2.2, Theorem 2.3, Lemma 4.1 and Lemma 2.4 there exists a Nash submanifold $Q \subset J$, which can be supposed Nash diffeomorphic to an open simplex in an euclidean space, such that $\alpha \in \tilde{Q}$ and the following diagram commutes:

$$\begin{array}{ccc} \mathcal{M}_Q & \longrightarrow & Q \\ \Pi \downarrow & & \parallel \\ N \times Q & \xrightarrow{p} & Q \end{array}$$

where $p \circ \Pi$ is a Nash submersion and the map Π is a Nash resolution of $V = F_Q^{-1}(0)$ in $N \times Q$ and V' (the strict transform of V), D_1, \dots, D_k have only normal crossings and

- $p \circ \Pi|_V : V' \rightarrow Q$,
- $p \circ \Pi|_{D_{j_1} \cap \dots \cap D_{j_s}} : D_{j_1} \cap \dots \cap D_{j_s} \rightarrow Q$,
- $p \circ \Pi|_{V' \cap D_{j_1} \cap \dots \cap D_{j_s}} : V' \cap D_{j_1} \cap \dots \cap D_{j_s} \rightarrow Q$ ($1 \leq j_1 < \dots < j_s \leq k$)

are Nash submersions.

Applying Theorem 2.6 to $p \circ \Pi$, there is a finite partition of Q into Nash open simplices Q_j , $j = 1, \dots, b$, and for any j , there exists a Nash diffeomorphism $\Phi_j : M_{Q_j} \rightarrow M_{t_j} \times Q_j$ such that $\Phi(V'_j) = V'_{t_j} \times Q_j$, $\Phi(D_i) = D_{it_j} \times Q_j$, $i = 1, \dots, r$ and

the following diagram is commutative:

$$\begin{array}{ccc} \mathcal{M}_{Q_i} & \xrightarrow{\Phi_j} & \mathcal{M}_{t_j} \times Q_j \\ \Pi_j \downarrow & & \downarrow p \\ N \times Q_j & \xrightarrow{p} & Q_i \end{array} ,$$

where $\mathcal{M}_{t_j} = (p \circ \Pi)^{-1}(t_j)$, $V'_{t_j} = V'_j \cap (p \circ \Pi)^{-1}(t_j)$ and $D_{it_j} = D_i \cap (p \circ \Pi)^{-1}(t_j)$ with $t_j \in Q_j$. It follows that Π_j gives a Nash trivial simultaneous resolution of V_j in $N \times Q_j$ over Q_j . There is $j_0 \in \{1, \dots, b\}$ such that $\alpha \in \widetilde{Q}_{j_0}$. Let us write it \widetilde{Q}^α .

Since $\widetilde{J} = \bigcup_{\alpha \in \widetilde{J}} \widetilde{Q}^\alpha$ with $Q^\alpha \subset J$ a Nash submanifold diffeomorphic to an open simplex in an Euclidean space over which there is a Nash trivial simultaneous resolution of $F_{Q^\alpha}^{-1}(0)$ in $N \times Q^\alpha$, by compactness property of \widetilde{J} , we get $\widetilde{J} = \bigcup_{i=1}^p \widetilde{Q}^{\alpha_i}$. Set $J_i = Q^{\alpha_i}$, and this ends the proof. \square

Using the same technique for the proof of Theorem 2.7 and **Theorem (I)**, we get the following effective result.

Theorem 4.2. *Given the integers n, p and q , there exist a triplet of integers (u, v, w) such that for each $i = 1, \dots, u$ there is a Nash subset V_i of \mathbb{R}^n of complexity bounded by (p, q) with a Nash resolution $\Pi_i : M_i \rightarrow \mathbb{R}^n$ and for any Nash subset V of \mathbb{R}^n of complexity bounded by (p, q) with a Nash resolution $\Pi : M \rightarrow \mathbb{R}^n$, there is $i \in \{1, \dots, u\}$ and a Nash diffeomorphism Φ between M and M_i whose graph have the complexity bounded by (v, w) which sends strict transform of V to the strict transform of V_i and exceptional divisors to exceptional divisors.*

Moreover, u, v and w are recursive functions of n, p and q .

5. FINITENESS THEOREM ON BLOW-NASH TRIVIALITY

Now, we can prove our second result. Let N be an affine Nash manifold and J a semialgebraic subset of \mathbb{R}^s . Let $F : N \times J \rightarrow \mathbb{R}^k$ be semialgebraic family of Nash maps $f_t : N \rightarrow \mathbb{R}^k$ such that $f_t^{-1}(0)$ for all $t \in J$ has only isolated singularities.

Proof. of **Theorem (II)**

First of all, let us note that the number of singular points of $f_t^{-1}(0)$ for $t \in J$ is uniformly bounded. This implies that, given $\alpha \in \widetilde{J}$, the Nash set $f_\alpha^{-1}(0)$ has finitely many isolated singular points, say $s_{1,\alpha}, \dots, s_{\nu,\alpha}$. By the Nash desingularization Theorem, there exists a Nash resolution of $f_\alpha^{-1}(0)$ in $N_{k(\alpha)}$, $\Pi_\alpha : \mathcal{M}_\alpha \rightarrow N_{k(\alpha)}$. We index the Nash exceptional divisors as $D_{i,j\alpha}$ for $i = 1, \dots, \nu$ and $j = 1, \dots, \mu(i)$ with the property $\Pi_\alpha(D_{i,j\alpha}) = \{s_{i,\alpha}\}$ and V'_α the strict transform of $V_\alpha = f_\alpha^{-1}(0)$ with respect to Π_α . By the definition of Π_α , we have that, $D_{i,j\alpha}$ for $i = 1, \dots, \nu$, $j = 1, \dots, \mu(i)$ and V'_α have only normal crossings. So by the Theorem 2.2, Theorem 2.3, Lemma 4.1 and Lemma 2.4 there exists a Nash submanifold $Q \subset J$, which can be supposed Nash diffeomorphic to an open simplex in an euclidean space, such that $\alpha \in \widetilde{Q}$ and the following diagram commutes:

$$\begin{array}{ccc} \mathcal{M}_Q & \longrightarrow & Q \\ \Pi \downarrow & & \parallel \\ N \times Q & \xrightarrow{p} & Q \end{array}$$

where $p \circ \Pi$ is a Nash submersion and the map Π is a Nash resolution of $V = F_Q^{-1}(0)$ in $N \times Q$ such that V' (the strict transform of V) and $D_{i,j}$ the Nash divisors for $i = 1, \dots, \nu$, $j = 1, \dots, \mu(i)$ simultaneously have only normal crossings and

- $p \circ \Pi|_{V'} : V' \rightarrow Q$,
- $p \circ \Pi|_{D_{i,j_1} \cap \dots \cap D_{i,j_s}} : D_{i,j_1} \cap \dots \cap D_{i,j_s} \rightarrow Q$,
- $p \circ \Pi|_{V' \cap D_{i,j_1} \cap \dots \cap D_{i,j_s}} : V' \cap D_{i,j_1} \cap \dots \cap D_{i,j_s} \rightarrow Q$ ($i = 1, \dots, \nu$ and $1 \leq j_1 < \dots < j_s \leq \mu(i)$)

are Nash submersions. There are Nash maps $s_i : Q \rightarrow N$ for $i = 1, \dots, \nu$ such that $\Pi(D_{i,j}) \subset \Gamma_{s_i}$ for $j = 1, \dots, \mu(i)$ where Γ_{s_i} is the graph of the map s_i .

Now, applying Theorem 2.6 to $p \circ \Pi$, we may shrink Q so that $\alpha \in \tilde{Q}$ and there exist $t_0 \in Q$ and a Nash diffeomorphism $\Phi : \mathcal{M}_Q \rightarrow \mathcal{M}_{t_0} \times Q$ such that $\Phi(V') = V'_{t_0} \times Q$, $\Phi(D_{i,j}) = D_{i,j,t_0} \times Q$, $i = 1, \dots, \nu$, $j = 1, \dots, \mu(i)$ and the following diagram is commutative:

$$\begin{array}{ccc} \mathcal{M}_Q & \xrightarrow{\Phi} & \mathcal{M}_{t_0} \times Q \\ \Pi \downarrow & & \downarrow p \\ N \times Q & \xrightarrow{p} & Q \end{array}$$

where $\mathcal{M}_{t_0} = (p \circ \Pi)^{-1}(t_0)$, $V'_{t_0} = V' \cap (p \circ \Pi)^{-1}(t_0)$ and $D_{i,j,t_0} = D_{i,j} \cap (p \circ \Pi)^{-1}(t_0)$. It follows that Π gives a Nash trivial simultaneous resolution of V in $N \times Q$ over Q .

Let us construct a semialgebraic homeomorphism $\phi : N \times Q \rightarrow N \times Q$ such that the following diagram commutes:

$$\begin{array}{ccccc} \mathcal{M}_Q & \xrightarrow{\Pi} & N \times Q & \xrightarrow{p} & Q \\ \Phi \downarrow & & \phi \downarrow & & \parallel \\ \mathcal{M}_{t_0} \times Q & \xrightarrow{\Pi|_{\mathcal{M}_{t_0}} \times id_Q} & N \times Q & \xrightarrow{p} & Q \end{array}$$

So, let us set the map ϕ as follows: $\phi(x, t) = (y, t)$ such that

- $\phi(x, t) = (\Pi|_{\mathcal{M}_{t_0}} \times id_Q) \circ \Phi \circ \Pi^{-1}(x, t)$ for $x \notin \{s_1(t), \dots, s_\nu(t)\}$
- $\phi(s_i(t), t) = (s_i(t_k), t_0)$ for $i = 1, \dots, \nu$.

Let us show that ϕ as constructed is a semialgebraic homeomorphism. Set $S = \{(s_i(t), t) : i = 1, \dots, \nu \text{ and } t \in Q\}$ the singular set of $V = F_Q^{-1}(0)$. Of course, $\phi|_{(N \times Q) \setminus S}$ is a Nash diffeomorphism as composition of Nash diffeomorphisms and $\phi|_S$ is a semialgebraic homeomorphism. So, ϕ is a semialgebraic map. It remains to show that ϕ is continuous. Indeed, let be $(s_i(t), t) \in S$. Consider a semialgebraic path $\gamma : [0, 1] \rightarrow N \times Q$ with $\gamma([0, 1]) \subset (N \times Q) \setminus S$ and $\gamma(1) = (s_i(t), t)$. Consider

a semialgebraic lifting $\gamma' : [0, 1] \rightarrow \mathcal{M}_Q$ of γ in \mathcal{M}_Q by Π (γ' exists because Π is proper). It follows that $\gamma'(1) \in \bigcup_j^{\mu(i)} D_{i,j_t}$ where $D_{i,j_t} = D_{i,j} \cap (p \circ \Pi)^{-1}(t)$. For $s \in [0, 1)$, by definition of ϕ we get

$$\phi(\gamma(s)) = (\Pi|_{\mathcal{M}_t} \times id_Q) \circ \Phi \circ \Pi|_{(N \times Q) \setminus S}^{-1}(\gamma(s)).$$

It is also clear that $\phi(\gamma(1)) = (\Pi|_{\mathcal{M}_{t_0}} \times id_Q) \circ \Phi(\gamma'(1)) = (s_i(t), t_0)$ since $\gamma'(1) \in \bigcup_j^{\mu(i)} D_{i,j_t}$. It follows that ϕ is a semialgebraic homeomorphism. Hence $(N \times Q, V)$ admits a Π -Blow-Nash trivialization over Q . Let us write this Q by Q^α . Since $\tilde{J} = \bigcup_{\alpha \in \tilde{J}} \tilde{Q}^\alpha$ with $Q^\alpha \subset J$ a Nash submanifold diffeomorphic to an open simplex in an Euclidean space over which $(N \times Q^\alpha, V)$ admits a Π -Blow-Nash triviality, by compactness property of \tilde{J} , we get $\tilde{J} = \bigcup_{i=1}^p \tilde{Q}^{\alpha_i}$. Set $J_i = Q^{\alpha_i}$, and this completes the proof. \square

To end this paper, using the same technique of the proof of Theorem 2.7 and **Theorem (II)**, we get that there are finitely many Blow-Nash equivalence classes of Nash sets with only isolated singularities and bounded complexity. We get also that the result is effective.

Theorem 5.1. *Given the integers n, p and q , there exist a triplet of integers (u, v, w) such that for each $i = 1, \dots, u$ there is a Nash subset V_i of \mathbb{R}^n of complexity bounded by (p, q) having only isolated singularities such that for any Nash subset V of \mathbb{R}^n of complexity bounded by (p, q) having only isolated singularities there are $i \in \{1, \dots, u\}$ and a Nash diffeomorphism Φ between M and M_i and a semialgebraic homeomorphism ϕ between \mathbb{R}^n whose graphs have the complexities*

$$M \xrightarrow{\Phi} M_i$$

bounded by (v, w) making the following diagram commutative

$$\begin{array}{ccc} M & & M_i \\ \Pi \downarrow & & \downarrow \Pi_i \\ \mathbb{R}^n & \xrightarrow{\phi} & \mathbb{R}^n \end{array}$$

where $\Pi : M \rightarrow \mathbb{R}^n$ and $\Pi_i : M_i \rightarrow \mathbb{R}^n$ are respectively Nash resolutions of singularity of V and V_i .

Moreover, u, v and w are bounded by recursive functions of n, p and q .

REFERENCES

- [BCR] J. Bochnak, M. Coste, and M-F. Roy. *Real algebraic geometry*, volume 36 of *Ergebnisse der Mathematik und ihrer Grenzgebiete(3)* [Results in Mathematics and related Area(3)]. Springer-Verlag, Berlin, 1998. Translated from the 1987 French original, Revised by the authors.
- [BM1] E. Bierstone and P.D. Milman *Uniformization of analytic spaces*, Jour.Amer. Math. Soc. 2 (1989), 801-836.
- [BM2] E. Bierstone and P.D. Milman *Canonical desingularization in characteristic zero by blowing-up the maximum strata of a local invariant*, Invent. math. 128 (1997), 207-302.
- [CRS] M. Coste, J.M Ruiz and M. Shiota *Global Problem on Nash Functions* Rev. Mat. Complut. 2004,17;Nùm 1, 83-115.

- [CS] M. Coste and M. Shiota *Nash triviality in family of Nash manifolds* Invent. Math. 108, 349-368 (1992).
- [CS1] M. Coste and M. Shiota *Thom's first isotopy lemma: a semialgebraic version, with uniform bound*, Real Analytic and Algebraic Geometry (Ed. F. Broglia, M. Galbiati, A. Tognoli), Walter de Gruyter, Berlin 83-101, 1995.
- [D] Demdah K.Mady *h-cobordism and s-cobordism theorems: Transfer over Semialgebraic and Nash categories, Uniform bound and Effectiveness* Annales de l'institut Fourier, t. 61, n.5(2011), p. 1573-1597.
- [FKS] T. Fukui, S. Koike and M. Shiota *Modified Nash triviality of family of zero-sets of real polynomial mappings* Annales de l'institut Fourier, t. 48, n.5, p. 1395-1440, 1998.
- [H1] H. Hironaka *Resolution of singularities of an algebraic variety over a field of characteristic zero*, I, II, Ann. of math. 79 (1964) 109-302.
- [H2] H. Hironaka *Idealistic exponents of singularity*, J.J. Sylvester Sympos., Johns Hopkins Univ., Baltimore (1976) 52-125.
- [K1] S. Koiké *Nash Trivial simultaneous resolution for a family of zero-sets of Nash mappings*, Mathematisch Zeitschrift.237,313-336 (2000).
- [K2] S. Koiké *Finiteness Theorem for Blow-semialgebraic Triviality of a family of three-dimensional Algebraic sets*, Proc. London. Math. Soc. 105 (3),506-540 (2012).
- [LT] F. Lazerri, A. Tognoli *Alcune proprietà degli spazi algebrici*, Ann. Scuola Norm.Sup. Pisa 24, 597-632 (1970).
- [P] A. Prestel *Model theory for the real algebraic geometer*, Dottorato di Ricerca in Matematica, Dip. Mat. Univ. Pisa. istituti Editoriali Poligrafici Internazionali 1998.
- [R] R. Ramanakorasina *Complexité des fonctions de Nash*, Commun. Algebra 17 (n.6),1395-1406, 1989.
- [S] M. Shiota *Nash Manifolds*. Lecture Notes in Mathematics 1269 . Springer-Verlag, Berlin , 1980.

DEMDAH KARTOUE MADY
 Département de Mathématiques
 Faculté des Sciences Exactes et Appliquées
 Université de Ndjamena
 BP:1027, 1 Route de Farcha
 email:kartoue@hotmail.com
 kartoue.demdah@univ-rennes1.fr