



**HAL**  
open science

# The Gauß-Bonnet operator of an infinite graph

Colette Anné, Nabila Torki-Hamza

► **To cite this version:**

Colette Anné, Nabila Torki-Hamza. The Gauß-Bonnet operator of an infinite graph. 2014. hal-00768827v3

**HAL Id: hal-00768827**

**<https://hal.science/hal-00768827v3>**

Preprint submitted on 28 Apr 2014 (v3), last revised 11 Sep 2014 (v4)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# THE GAUSS-BONNET OPERATOR OF AN INFINITE GRAPH

COLETTE ANNÉ AND NABILA TORKI-HAMZA

ABSTRACT. — We propose a general condition, to ensure essential self-adjointness for the Gauß-Bonnet operator  $D = d + \delta$ , based on a notion of completeness as Chernoff. This gives essential self-adjointness of the Laplace operator both for functions or 1-forms on infinite graphs. This is used to extend Flanders result concerning solutions of Kirchhoff's laws.

RÉSUMÉ. Nous proposons une condition générale qui assure le caractère essentiellement auto-adjoint de l'opérateur de Gauss-Bonnet  $D = d + \delta$ , basée sur une notion de complétude comme Chernoff. Comme conséquence, l'opérateur de Laplace agissant sur les fonctions ou les 1-formes de graphes infinis est essentiellement auto-adjoint. Nous utilisons ce cadre pour étendre le résultat de Flanders à propos des solutions des lois de Kirchhoff.

## 1. INTRODUCTION

Operators on infinite graphs are of large interest and a lot of recent works deals with this subject. One approach can be to study how technics of spectral geometry can be extended on graphs regarded as one-dimensional simplicial complexes. We refer to Dodziuk [D84, DK87] for general presentation of this approach and to [CdV98, CTT11] for the geometric point of view, and also [CdV91] for the relation between Kirchhoff's laws and Hodge theory.

We consider here only connected locally finite infinite graphs and we study Kirchhoff's laws. Flanders has first studied this question on infinite graphs seen as infinite electric networks, see [F71]. Several authors have clarified and extended Flanders work on electric networks, see for instance Thomassen [T90], Soardi [S94], Doyle & Snell [DS99], Zemanian [Z08], Georgakopoulos [G10], Carmesin [Cm12] and also the book of Jorgensen & Pearse [JP14] for a general approach.

Flanders main result is that there exists a unique current flow in an infinite network with a finite number of sources which is the limit of finite flows.

In our paper, this question is approached by the study of a Dirac type operator: *the Gauß-Bonnet operator*  $D = d + \delta$ , introduced on an infinite graph considered as a one-dimensional simplicial complex. Indeed, this operator is a generalisation of the Dirac operator studied on  $\mathbb{Z}$  by Golenia & Haugomat in [GH12]. We give a general condition on the graph by defining the notion of  $\chi$ -completeness, see Section 3.2. One of the main results is to prove essential self-adjointness of the Gauß-Bonnet operator, when the graph is  $\chi$ -complete (or *complete homogeneous*). This condition

---

*Date:* April 28, 2014 *File:* graf-10-2-2014.tex

2010 *Mathematics Subject Classification.* 39A12, 05C63, 47B25, 05C12, 05C50.

*Key Words and Phrases.* infinite graph,  $\chi$ -completeness, difference operator, coboundary operator, Dirac type operator, Gauß-Bonnet operator, essential self-adjointness.

covers the situations of [HKMW13], [M09], and [T10] (or [T12]), and it is a discrete version of a result of Chernoff, see [Ch73], in the case of manifolds. One of the applications in his paper concludes that, on a complete manifold, every power of the Dirac operator  $d + \delta$  is essentially self-adjoint. In particular, for every power of the Laplace-Beltrami operator, essential self-adjointness is true.

In Section 4.3, we define the property of *positivity at infinity* for Dirac type operators. And by adding this assumption on our Gauß-Bonnet operator, we prove that its range is closed and consequently the Hodge property holds, in a similar result as Anghel's for compact Riemannian manifold, see [A93]. This situation permits us to enlarge the conditions on the current source and the voltage source in the Flanders problem. In Section 5, we give new examples of infinite graphs where it applies.

## 2. PRELIMINARIES

**2.1. Definitions on Graphs.** ( cf. [LP14]) A graph  $K$  is a simplicial complex of dimension one. We denote by  $\mathcal{V}$  the set of vertices and  $\mathcal{E}$  the set of *oriented edges*, considered as a subset of  $\mathcal{V} \times \mathcal{V}$ . We assume that  $\mathcal{E}$  is symmetric without loops :

$$v \in \mathcal{V} \Rightarrow (v, v) \notin \mathcal{E}, \quad (v_1, v_2) \in \mathcal{E} \Rightarrow (v_2, v_1) \in \mathcal{E}.$$

Choosing an orientation of the graph consists of defining a partition of  $\mathcal{E}$  :

$$\begin{aligned} \mathcal{E}^+ \sqcup \mathcal{E}^- &= \mathcal{E} \\ (v_1, v_2) \in \mathcal{E}^+ &\iff (v_2, v_1) \in \mathcal{E}^-. \end{aligned}$$

For  $e = (v_1, v_2) \in \mathcal{E}$ , let's set

$$e^+ = v_2, \quad e^- = v_1, \quad -e = (v_2, v_1).$$

$e^+$  and  $e^-$  are called boudary points of the edge  $e$ .

2.1.1. A *path* between two vertices  $x, y$  in  $\mathcal{V}$  is a finite set of edges  $e_1, \dots, e_n, n \geq 1$  such that

$$e_1^- = x, \quad e_n^+ = y \quad \text{and, if } n \geq 2, \quad \forall j, \quad 1 \leq j \leq (n-1) \Rightarrow e_j^+ = e_{j+1}^-.$$

Notice that each path has a beginning and an end, and that an edge is a path.

Let us denote  $\Gamma_{xy}$  the set of the paths from the vertex  $x$  to the vertex  $y$ .

2.1.2. The graph is *connected* if two vertices are always related by a path, *ie.* if  $\Gamma_{xy}$  is non empty for all  $x, y$  in  $\mathcal{V}$ .

2.1.3. The graph is *locally finite* if each vertex belongs to a finite number of edges. The *degree* or *valence* of a vertex  $x \in \mathcal{V}$  is the cardinal of the set  $\{e \in \mathcal{E}; e^+ = x\}$ .

2.1.4. A *subgraph* of a graph  $K$  is a graph  $K_0 = (\mathcal{V}_0, \mathcal{E}_0)$  such that  $\mathcal{V}_0 \subset \mathcal{V}$  and  $\mathcal{E}_0 \subset \mathcal{E}$ .

**Remark 1.** *All the graphs we shall consider on the sequel will be connected, locally finite, so with countably many vertices.*

**2.2. Functions and forms.** The 0-cochains are just scalar functions on  $\mathcal{V}$ , we denote their set  $C^0(K)$ .

The 1-cochains or forms are odd scalar functions on  $\mathcal{E}$  we denote their set  $C^1(K)$ . Thus we have

$$\begin{aligned} C^0(K) &= \mathbb{C}^{\mathcal{V}}, \\ C^1(K) &= \{\varphi : \mathcal{E} \rightarrow \mathbb{C}, \varphi(-e) = -\varphi(e)\}. \end{aligned}$$

The sets of cochains with finite support are denoted by  $C_0^0(K)$ ,  $C_0^1(K)$ . To obtain Hilbert spaces we need weights, let's give

$$c : \mathcal{V} \rightarrow \mathbb{R}_+^*,$$

and

$$r : \mathcal{E} \rightarrow \mathbb{R}_+^* \text{ even}$$

so  $r(-e) = r(e)$ .

They define scalar products :

$$\begin{aligned} \forall f, g \in C_0^0(K); \quad \langle f, g \rangle &= \sum_{v \in \mathcal{V}} c(v) f(v) \bar{g}(v) \\ \forall \varphi, \psi \in C_0^1(K); \quad \langle \varphi, \psi \rangle &= \frac{1}{2} \sum_{e \in \mathcal{E}} r(e) \varphi(e) \bar{\psi}(e) \end{aligned} \quad (1)$$

**Remark 2.** As the products  $r(e)\varphi(e)\bar{\psi}(e)$ ,  $e \in \mathcal{E}$  in (1) are even, the term  $\frac{1}{2}$  allows to recover the usual definition.

**Remark 3.** In the context of electric networks, our weight on edges would play the role of the conductance, the intensity would be on  $e \in \mathcal{E}$  :  $I(e) = r(e)\varphi(e)$  and the energy  $\|\varphi\|^2 = \frac{1}{2} \sum_{e \in \mathcal{E}} \frac{1}{r(e)} I(e)^2$ . So, indeed,  $\frac{1}{r(e)}$  is the resistance of the edge  $e$ !

Let us finally define the Hilbert spaces

$$\begin{aligned} L_2(\mathcal{V}) &= \overline{C_0^0(K)}, \\ L_2(\mathcal{E}) &= \overline{C_0^1(K)}. \end{aligned}$$

### 2.3. Operators.

2.3.1. *The difference operator.* It is the operator

$$d : C_0^0(K) \rightarrow C_0^1(K),$$

given by

$$d(f)(e) = f(e^+) - f(e^-), \quad (2)$$

for  $f \in C_0^0(K)$ ,  $e \in \mathcal{E}$ .

2.3.2. *The coboundary operator.* It is  $\delta$  the formal adjoint of  $d$ . Thus it satisfies

$$\langle df, \varphi \rangle = \langle f, \delta\varphi \rangle \quad (3)$$

for all  $f \in C_0^0(K)$  and  $\varphi \in C_0^1(K)$ .

**Lemma 4.** *The coboundary operator  $\delta : C_0^1(K) \rightarrow C_0^0(K)$ , is defined by the formula*

$$\delta(\varphi)(x) = \frac{1}{c(x)} \sum_{e, e^+=x} r(e)\varphi(e). \quad (4)$$

*Proof.* — Using the equation (3), we have

$$\frac{1}{2} \sum_{e \in \mathcal{E}} r(e) (f(e^+) - f(e^-)) \bar{\varphi}(e) = \frac{1}{2} \sum_{x \in \mathcal{V}} f(x) \left( \sum_{e^+=x} r(e)\varphi(e) - \sum_{e^-=x} r(e)\varphi(e) \right)$$

But  $r\varphi$  is odd and  $\mathcal{E}$  symmetric, so

$$\sum_{e^-=x} r(e)\varphi(e) = - \sum_{e^+=x} r(e)\varphi(e).$$

We remark that the sum entering in the formula (4) of  $\delta$  is finite due to the hypothesis that the graph is locally finite.  $\square$

**Remark 5.** *The operator  $d$  is defined by (2) in all  $C^0(K)$ , but to define  $\delta$  in all  $C^1(K)$ , we need an hypothesis on  $K$  : we suppose that the graph is locally finite. This hypothesis could be weakened by assuming that the edge weights  $r(e)$ ,  $e \in \mathcal{E}$  are summable around each vertex as considered in [KL12].*

With these two operators we can form the following two operators.

2.3.3. *The Gauß-Bonnet operator.* It is the endomorphism

$$D = d + \delta : C_0^0(K) \oplus C_0^1(K) \curvearrowright$$

given by

$$D(f, \varphi) = \delta\varphi + df$$

for all  $f \in C_0^0(K)$  and  $\varphi \in C_0^1(K)$ .

This operator is symmetric and of Dirac type.

2.3.4. *Laplacian.* By definition, it is

$$\Delta = D^2 : C_0^0(K) \oplus C_0^1(K) \curvearrowright.$$

This operator preserves the direct sum  $C_0^0(K) \oplus C_0^1(K)$ , so we can write

$$\Delta = \Delta_0 \oplus \Delta_1.$$

2.4. **Metrics.** A *metric* is an even function

$$a : \mathcal{E} \rightarrow \mathbb{R}_+^*,$$

it defines a distance on the graph  $K$  in the following way.

One first defines the *length of a path* : for  $\gamma = (e_1, \dots, e_n)$

$$l_a(\gamma) = \sum_{j=1}^n \sqrt{a(e_j)}.$$

Then the *metric distance* between two vertices  $x, y$  is given by

$$d_a(x, y) = \inf_{\gamma \in \Gamma_{xy}} l_a(\gamma).$$

### 3. CLOSABILITY AND SELF-ADJOINTNESS

#### 3.1. Closability.

**Lemma 6.** *If the graph  $K$  is connected and locally finite the operators  $d$  and  $\delta$  are closable.*

*Proof.* — Let us suppose that there exists a sequence  $(f_n)_{n \in \mathbb{N}}$  in  $C_0^0(K)$  such that  $\|f_n\| \rightarrow 0$  and  $(d(f_n))_n$  converges. Let us denote by  $\varphi$  this limit.

We have to show that  $\varphi = 0$ . If

$$\|f_n\| + \|d(f_n) - \varphi\| \rightarrow 0,$$

then for each vertex  $v$ ,  $f_n(v)$  converges to 0 and for each edge  $e$ ,  $d(f_n)(e)$  converges to  $\varphi(e)$ . But by the first statement and the expression of  $d$ , for each edge  $e$ ,  $d(f_n)(e)$  converges to 0.

The same can be done for  $\delta$  : convergence in norm to 0 of a sequence  $(\varphi_n)_n$  implies pointwise convergence to 0 which implies pointwise convergence of  $\delta(\varphi_n)$  to 0, because of local finiteness of the graph ; if  $\delta(\varphi_n)$  converges in norm, it must be to 0.  $\square$

Thus, we can consider different extensions of these operators in the framework of Hilbert spaces (see [RS80]).

The smallest extension is the closure, denoted  $\bar{d} = d_{min}$  (resp.  $\bar{\delta} = \delta_{min}$  and  $\bar{D} = D_{min}$ ) has the domain

$$\text{Dom}(d_{min}) = \left\{ f \in L_2(\mathcal{V}); \exists (f_n)_{n \in \mathbb{N}}, f_n \in C_0^0(K), L_2\text{-}\lim_{n \rightarrow \infty} f_n = f, \right. \\ \left. L_2\text{-}\lim_{n \rightarrow \infty} d(f_n) \text{ exists} \right\} \quad (5)$$

for such an  $f$ , one puts

$$d_{min}(f) = \lim_{n \rightarrow \infty} d(f_n).$$

The largest is  $d_{max} = \delta^*$ , the adjoint operator of  $\delta_{min}$ , (resp.  $\delta_{max} = d^*$ , the adjoint operator of  $d_{min}$ .)

#### 3.2. A sufficient condition for self-adjointness of $D$ .

### 3.2.1. Geometric hypothesis for the graph $K$ .

**Definition 7.** The graph  $K$  is  $\chi$ -complete if there exists a increasing sequence of finite sets  $(B_n)_{n \in \mathbb{N}}$  such that  $\mathcal{V} = \dot{\cup} B_n$  and there exist related functions  $\chi_n$  satisfying the following three conditions :

- (i)  $\chi_n \in C_0^0(K)$ ,  $0 \leq \chi_n \leq 1$
- (ii)  $v \in B_n \Rightarrow \chi_n(v) = 1$
- (iii)  $\exists C > 0, \forall n \in \mathbb{N}, x \in \mathcal{V}, \frac{1}{c(x)} \sum_{e, e^\pm = x} r(e) d\chi_n(e)^2 \leq C$ .

For this type of graphs one has

$$\forall p \in \mathbb{N}, \exists n_p, n \geq n_p \Rightarrow \forall e \in \mathcal{E}, \text{ such that } e^+ \text{ or } e^- \in B_p, d\chi_n(e) = 0 \quad (6)$$


$$\mathcal{E} = \dot{\cup} \mathcal{E}_n \text{ if } \mathcal{E}_n = \{e \in \mathcal{E}, e^+ \in B_n \text{ or } e^- \in B_n\} \quad (7)$$

$$\forall f \in L_2(\mathcal{V}), \lim_{n \rightarrow \infty} \langle \chi_n f, f \rangle = \|f\|^2 \quad (8)$$

$$\forall \varphi \in L_2(\mathcal{E}), \|\varphi\|^2 = \lim_{n \rightarrow \infty} \frac{1}{2} \sum_{e \in \mathcal{E}} r(e) \chi_n(e^+) |\varphi(e)|^2 \quad (9)$$

$$\text{and } \lim_{n \rightarrow \infty} \sum_{e \in \text{supp}(d\chi_n)} r(e) |\varphi(e)|^2 = 0. \quad (10)$$

**Example 8.** Let's consider an infinite tree with increasing valence:



Taking constant weights on vertices and edges, this graph is  $\chi$ -complete. Indeed, one can define *generations* of vertices on such a graph : the considered origine vertex  $c$  is of generation 0 and valence 2, it is related to two vertices which are of generation 1 and valence 3, and more generally there are  $2n!$  vertices of generation  $n$  and valence  $(n+2)$ .

One defines  $B_n, n \in \mathbb{N}$ , as the set of vertices of generation less than  $n^2$  and  $\chi_n$  constant on each generation of vertices:

$$x \text{ of generation } p \Rightarrow \chi_n(x) = \left( \frac{(n+1)^2 - p}{2n+1} \wedge 1 \right) \vee 0.$$

So,  $p \leq n^2 \Rightarrow \chi_n(x) = 1$  and  $p \geq (n+1)^2 \Rightarrow \chi_n(x) = 0$  while  $|d\chi_n(e)| \leq 1/(2n+1)$  is in fact supported on edges between generations larger than  $n^2$  and less than

$(n + 1)^2$ . To verify the condition (iii), one has to calculate for these generations,  $(p + 2)/(2n + 1)^2 \leq ((n + 1)^2 + 2)/(2n + 1)^2$  which is bounded independantly on  $n$ .

**Remark 9.** *The condition of  $\chi$ -completeness covers many situations that have been already studied. Particularly it covers the situation studied in [HKMW13]. In this work, the authors define a notion of intrinsic pseudo metric.*

**Lemma 9. 1.** *If the graph admits an intrinsic path metric  $d$  such that  $(\mathcal{V}, d)$  is complete, then the graph is  $\chi$ -complete.*

*Proof.* The hypothesis means that our infinite, connected, locally finite, weighted graph admits a metric  $a$  as defined in section 2.4 such that

$$\forall x \in \mathcal{V}, \frac{1}{c(x)} \sum_{e \in \mathcal{E}, e^+ = x} r(e)a(e) \leq 1$$

(the relation between our notations and those of [HKMW13] is :  $\sigma^2 = a$ ). We suppose also that the metric distance  $d_a$  defines  $(\mathcal{V}, d_a)$  as a complete metric space.

We then define the functions  $\chi_n$  as follows. Fix  $O$  a vertex in  $\mathcal{V}$  and put

$$\forall n \in \mathbb{N}, B_n = \{x \in \mathcal{V}; d_a(O, x) \leq n\}, \chi_n(x) = \sup\{(1 - d_a(x, B_n)), 0\} \quad (11)$$

As pointed in [HKMW13] completeness of  $(\mathcal{V}, d_a)$  gives that the  $B_n$  are finite. We verify that

- (i) The support of  $\chi_n$  is finite : it is included in  $\{x; d_a(x, B_n) \leq 1\} \subset B_{n+1}$ .
- (ii)  $x \in B_n \Rightarrow d_a(x, B_n) = 0 \Rightarrow \chi_n(x) = 1$
- (iii) finally, by the triangle inequality,  $d\chi_n(e)^2 \leq a(e)$ ; then the condition of *intrinsic metric* gives :

$$\forall x \in \mathcal{V}, \frac{1}{c(x)} \sum_{e \in \mathcal{E}, e^+ = x} r(e)d\chi_n(e)^2 \leq 1.$$

□

**Remark 10.** *If we consider the metric already introduced in [CTT11] (but to study non complete situation)*

$$a(e) = \frac{\min(c(e^+), c(e^-))}{r(e)} \quad (12)$$

and with bounded valence :

$$\exists A > 0, \forall v \in \mathcal{V}, \#\{e \in \mathcal{E}, e^+ = v\} \leq A.$$

then, if the graph is complete for this metric,  $\chi$ -completeness is also satisfied. Indeed,  $\frac{a}{A}$  is an intrinsic metric, because :

$$\forall x \in \mathcal{V}, \frac{1}{c(x)} \sum_{e \in \mathcal{E}, e^+ = x} r(e)a(e) \leq A.$$

It is also the case in the situation of [M09] where the hypothesis taken give that

$$\sup_{B_n} \frac{1}{c(x)} \sum_{e, e^\pm = x} r(e)d\chi_n(e)^2 = o(1)$$

for some  $\chi_n$  satisfying  $d\chi_n(e)^2 = O(n^{-2})$ .

**Theorem 1.** *Let  $K$  be a connected, locally finite graph. If  $K$  is  $\chi$ -complete, then the operator  $D$  is essentially self-adjoint.*

*Proof.* — Remark first that if  $d_{min} = d_{max}$  and  $\delta_{min} = \delta_{max}$  then  $D$  is essentially self-adjoint.

Indeed,  $D$  is a direct sum and if  $F = (f, \varphi) \in \text{Dom}(D^*)$  then  $\varphi \in \text{Dom}(d^*)$  and  $f \in \text{Dom}(\delta^*)$  and then, by hypothesis,  $\varphi \in \text{Dom}(\delta_{min})$  and  $f \in \text{Dom}(d_{min})$ , thus  $F \in \text{Dom}(\bar{D})$ .

For the following we need some formulas taken in [M09]. First we set, for each  $f \in C^0(K)$

$$\tilde{f}(e) = \frac{1}{2}(f(e^+) + f(e^-)). \quad (13)$$

The function  $\tilde{f}$  is even on the edges. We have

$$\begin{aligned} \forall f, g \in C^0(K), \forall e \in \mathcal{E}, \quad d(fg)(e) &= f(e^+)dg(e) + df(e)g(e^-) \\ &= \tilde{f}(e)dg(e) + \tilde{g}(e)df(e) \end{aligned} \quad (14)$$

$$\forall f \in C^0(K) \varphi \in C^1(K), \forall v \in \mathcal{V}, \quad \delta(\tilde{f}\varphi)(v) = f(v)\delta\varphi(v) - \frac{1}{c(v)} \sum_{e^+=v} r(e)df(e)\varphi(e). \quad (15)$$

3.2.2. *If  $f \in \text{Dom}(d_{max})$  then  $\|(f - \chi_n f)\| + \|d(f - \chi_n f)\| \rightarrow 0$  when  $n \rightarrow \infty$ . This will show that  $d_{min} = d_{max}$ .*

Let  $f \in \text{Dom}(d_{max})$ , we can then calculate

$$\|(f - \chi_n f)\|^2 \leq \sum_{v \notin B_n} c(v)|f(v)|^2 \xrightarrow{n \rightarrow \infty} 0$$

because  $f \in L_2(\mathcal{V})$ . For the second term, the relation (14) gives

$$d(f - \chi_n f)(e) = (1 - \chi_n)(e^+)d(f)(e) - f(e^+)d(\chi_n)(e).$$

Because of (9) (and with an abuse of notation),

$$\lim_{n \rightarrow \infty} \|(1 - \chi_n)(e^+)d(f)(e)\| = 0$$

On the other hand,

$$\begin{aligned} \|f(e^+)d(\chi_n)(e)\|^2 &= \sum_{e \in \mathcal{E}} r(e)|f(e^+)|^2 |d(\chi_n)(e)|^2 \\ &= \sum_{x \in \mathcal{V}} |f(x)|^2 \sum_{e^+=x} r(e) |d(\chi_n)(e)|^2 \\ &\leq \sum_{x \in \mathcal{V}, \exists e \in \text{supp}(d\chi_n), e^+=x} Cc(x)|f(x)|^2 \end{aligned}$$

by the hypothesis (iii). The property (7) permits to conclude that this term tends to 0 as  $n \rightarrow \infty$ .

3.2.3. If  $\varphi \in \text{Dom}(\delta_{max})$  then  $\|(\varphi - \tilde{\chi}_n\varphi)\| + \|\delta(\varphi - \tilde{\chi}_n\varphi)\| \rightarrow 0$  when  $n \rightarrow \infty$ . This will show that  $\delta_{min} = \delta_{max}$ .

Let  $\varphi \in \text{Dom}(\delta_{max})$ , by the properties (6) and (7) we know that

$$\forall p \in \mathbb{N}, \forall n \geq n_p, \quad \|\varphi - \tilde{\chi}_n\varphi\|^2 \leq \sum_{e \in \mathcal{E}_p^c} r(e)|\varphi(e)|^2$$

so  $\lim_{n \rightarrow \infty} \|\varphi - \tilde{\chi}_n\varphi\| = 0$ .

On the other hand, by (15)

$$\begin{aligned} \delta(\varphi - \tilde{\chi}_n\varphi)(v) &= \delta\left(\widetilde{(1 - \chi_n)\varphi}\right)(v) \\ &= (1 - \chi_n)(v)\delta\varphi(v) + \frac{1}{c(v)} \sum_{e^+=v} r(e)d\chi_n(e)\varphi(e) \end{aligned}$$

Clearly

$$\lim_{n \rightarrow \infty} \|(1 - \chi_n)\delta\varphi\| = 0$$

because  $\delta\varphi \in L_2(\mathcal{V})$ . For the second term, we use (iii) and the Cauchy-Schwarz inequality :

$$\begin{aligned} \forall v \in \mathcal{V}, \left| \sum_{e^+=v} r(e)d\chi_n(e)\varphi(e) \right|^2 &\leq \sum_{e^+=v} r(e)|d\chi_n(e)|^2 \sum_{e \in \text{supp}(d\chi_n), e^+=v} r(e)|\varphi(e)|^2 \\ &\leq Cc(v) \sum_{e \in \text{supp}(d\chi_n), e^+=v} r(e)|\varphi(e)|^2 \end{aligned}$$

$$\begin{aligned} \text{so, } \sum_{v \in \mathcal{V}} c(v) \left| \frac{1}{c(v)} \sum_{e^+=v} r(e)d\chi_n(e)\varphi(e) \right|^2 &\leq C \sum_{v \in \mathcal{V}} \sum_{e \in \text{supp}(d\chi_n), e^+=v} r(e)|\varphi(e)|^2 \\ &\leq C \sum_{e \in \text{supp}(d\chi_n)} r(e)|\varphi(e)|^2. \end{aligned}$$

This term tends to 0 by properties (6) and (7).

**Corollary 11.** *Let  $K$  be a connected, locally finite graph. If  $K$  is  $\chi$ -complete, then the operator  $\Delta$  is essentially self-adjoint.*

*Proof.* — If  $D$  is essentially self-adjoint, then  $\text{Im}(D \pm i)$  is dense and  $(\bar{D} \pm i)$  are invertible. This is a result for essentially self-adjoint operators (Corollary of Theorem VIII.3 in [RS80]). By the second property we know that

$$\exists C_2 > 0, \forall F \in \text{Dom}(\bar{D}), \|F\|_{L_2} \leq C_2 \|(D \pm i)(F)\|_{L_2}. \quad (16)$$

Remark also that

$$D(C_0^0(K) \oplus C_0^1(K)) \subset C_0^0(K) \oplus C_0^1(K).$$

Now, by the theorem of von Neumann,  $(\bar{D})^2 = D^*\bar{D}$  is self-adjoint when  $D^* = \bar{D}$  and it is an extension of  $\Delta$ ; then its domain contains the domain of  $\bar{\Delta}$ , the closure of  $\Delta$ . But

$$\text{Dom}(\bar{\Delta}) \subset \text{Dom}((\bar{D})^2) \Rightarrow \text{Dom}((\bar{D})^2) \subset \text{Dom}(\Delta^*).$$

In fact, we have also  $\text{Dom}(\Delta^*) \subset \text{Dom}((\bar{D})^2)$ : let  $\Psi \in \text{Dom}(\Delta^*)$ , then

$$\exists C_1 > 0, \forall F \in C_0^0(K) \oplus C_0^1(K), \quad | \langle (\Delta + 1)(F), \Psi \rangle | \leq C_1 \|F\|_{L_2}.$$

We now consider the linear form defined on  $C_0^0(K) \oplus C_0^1(K)$ , by

$$G \longmapsto \langle (D - i)G, \Psi \rangle$$

For all  $G \in \text{Im}(D + i)$ ,  $\exists F \in C_0^0(K) \oplus C_0^1(K)$ , such that  $G = (D + i)(F)$  so  $G \in C_0^0(K) \oplus C_0^1(K)$  and, using (16)

$$| \langle (D - i)G, \Psi \rangle | = | \langle (\Delta + 1)F, \Psi \rangle | \leq C_1 \|F\|_{L_2} \leq C_1 C_2 \|G\|_{L_2}$$

Hence

$$\exists C > 0, \forall G \in \text{Im}(D + i), \quad | \langle (D - i)G, \Psi \rangle | \leq C \|G\|_{L_2}. \quad (17)$$

But  $\text{Im}(D + i)$  is dense, it means that the considered linear form extends continuously on  $L_2$  or that  $(D + i)\Psi \in L_2$ . Thus  $\Psi \in \text{Dom}(\bar{D})$  because  $\bar{D}$  is self-adjoint. It is then clear that  $D(\Psi) \in \text{Dom}(\bar{D})$  :

$$\forall F \in C_0^0(K) \oplus C_0^1(K), \quad | \langle D(F), D(\Psi) \rangle | = | \langle \Delta(F), \Psi \rangle | \leq (C_1 + \|\Psi\|_{L_2}) \|F\|_{L_2}.$$

So, we have proved

$$\text{Dom}(\Delta^*) \subset \text{Dom}((\bar{D})^2) \Rightarrow \text{Dom}((\bar{D})^2) \subset \text{Dom}(\bar{\Delta})$$

because  $\Delta^{**} = \bar{\Delta}$ , and finally

$$\text{Dom}(\bar{\Delta}) = \text{Dom}((\bar{D})^2)$$

and then  $\bar{\Delta} = (\bar{D})^2$  is self-adjoint, see also [Ch73].  $\square$

**3.2.4. The converse is also true :** if  $\Delta$  is essentially self-adjoint then  $D$  is also essentially self-adjoint. Indeed, this is an easy consequence of the Corollary of Theorem VIII.3 in [RS80] : if  $\Delta$  is essentially self-adjoint then  $\text{Im}(\Delta + 1)$  is dense but

$$\Delta + 1 = (D + i)(D - i) = (D - i)(D + i) \Rightarrow \text{Im}(\Delta + 1) \subset \text{Im}(D \pm i).$$

Thus  $\text{Im}(D \pm i)$  are both dense and  $D$  is essentially self-adjoint.

**Remark 12.** — *The case studied in [T10] namely a complete graph for the metric*

$$a(e) = \frac{\sqrt{c(e^+)c(e^-)}}{r(e)}$$

*and with a valence bounded by  $A$  can be proved directly with the same kind of calculus. Indeed the condition satisfied now is*

$$\exists C > 0, \forall e \in \mathcal{E}, n \in \mathbb{N}, \quad r(e)d\chi_n(e)^2 \leq C\sqrt{c(e^+)c(e^-)}$$

*We write*

$$\begin{aligned} \sum_{e \in \mathcal{E}} r(e)f(e^-)d\chi_n(e)\bar{\varphi}(e) &= \frac{1}{2} \sum_{e \in \mathcal{E}} r(e)(f(e^+) + f(e^-))d\chi_n(e)\bar{\varphi}(e) \\ &\leq \frac{1}{2} \sqrt{\sum_{e \in \text{supp}(d\chi_n)} r(e)|\varphi(e)|^2} \sqrt{\sum_{e \in \text{supp}(d\chi_n)} r(e)|f(e^+) + f(e^-)|^2 d\chi_n(e)^2} \end{aligned}$$

$$\begin{aligned}
& \text{and } \sum_{e \in \text{supp}(d\chi_n)} r(e) |f(e^+) + f(e^-)|^2 d\chi_n(e)^2 \\
&= \sum_{e \in \text{supp}(d\chi_n)} r(e) \left[ |(f(e^+) - f(e^-))|^2 + 4 \operatorname{Re} \left( f(e^+) \bar{f}(e^-) \right) \right] d\chi_n(e)^2 \\
&= \sum_{e \in \text{supp}(d\chi_n)} r(e) |d(f)(e)|^2 + 4 \operatorname{Re} \left( \sum_{x \in \mathcal{V}} f(x) \sum_{e^+=x} r(e) \bar{f}(e^-) d\chi_n(e)^2 \right)
\end{aligned}$$

the first term tends to 0 by completeness and the second is bounded as follows

$$\begin{aligned}
& \operatorname{Re} \left( \sum_{x \in \mathcal{V}} |f(x)| \sum_{e^+=x} r(e) |f(e^-)| d\chi_n(e)^2 \right) \\
& \leq C \sum_{x \in \mathcal{V}} |f(x)| \sum_{e \in \text{supp } d\chi_n, e^+=x} |f(e^-)| \sqrt{c(e^+)c(e^-)} \\
& \leq AC \sum_{x \in \mathcal{V}, \exists e \in \text{supp } d\chi_n, e^+=x} c(x) |f(x)|^2
\end{aligned}$$

because, as  $\mathcal{E}$  is symmetric, one has

$$\sum_{x \in \mathcal{V}, \exists e \in \text{supp } d\chi_n, e^+=x} c(x) |f(x)|^2 = \sum_{x \in \mathcal{V}, \exists e \in \text{supp } d\chi_n, e^-=x} c(x) |f(x)|^2$$

So the second term also tends to 0, because of completeness and bounded valence.

#### 4. FLANDERS THEOREM

**4.1. Flanders problem.** In 1971, Flanders published a very nice result [F71] concerning resistive networks. The problem is the following : Let  $i$  be a finite current source, *i.e.* an element of  $C_0^0(K)$ , and  $E'$  a finite voltage source, *i.e.* an element of  $C_0^1(K)$ ,

is there a resulting current flow, and is it unique?

*i.e.* find solutions  $I$  of the problem (Kirchhoff's laws):

$$\begin{cases} \text{(Kirchhoff's current law)} & \delta(I) + i = 0, \\ \text{(Kirchhoff's voltage law)} & \forall Z, \partial Z = 0, \quad \int_Z E' = \int_Z I, \end{cases} \quad (18)$$

Here  $Z$  is a cycle, *i.e.* a 1-chain (a formal finite sum of oriented edges) with no boundary.

Formally we write

$$Z = \sum_{e \in \mathcal{E}^+} z_e e, \quad z_e \in \mathbb{Z}$$

or

$$Z = \frac{1}{2} \sum_{e \in \mathcal{E}} z_e e, \quad z_e \in \mathbb{Z}, \quad \text{with } z_e = -z_{-e} \quad (19)$$

and the boundary  $\partial$  of a 1-chain is an operator defined on the edges by

$$\partial(e) = e^+ - e^-,$$

so

$$\partial Z = \sum_{x \in \mathcal{V}} \left( \sum_{e^+ = x} z_e \right) x. \quad (20)$$

The integral in (18) has to be understood in the simplicial framework :

$$\int_Z I = \frac{1}{2} \sum_{e \in \mathcal{E}} z_e I(e) \quad (21)$$

Flanders studies this problem for an infinite graph with weight  $c = 1$  on vertices (Remark that our weight on edges is in fact the inverse of the resistances  $r$  introduced by Flanders, so our unknown  $I$  corresponds to  $r.I$  in the notations of Flanders). He shows that this problem has a unique  $L_2$ -solution which is the limit of finite flows (*ie.* solutions on an increasing sequence of finite subgraphs) if  $i$  has zero mean value  $\sum_{v \in \mathcal{V}} i(v) = 0$ .

**4.2. Flanders type Theorem.** In the framework we have introduced in Section 2, this question is related to the question of the Hodge decomposition. Indeed, the second condition tells us that the period of  $I$  are given by those  $E'$ , this determine the harmonic component of  $I$ , *ie.* the orthogonal projection of  $I$  on  $\text{Ker}(\delta)$ , while the complementary must be sent by  $\delta$  on  $-i$ . So we have to look for  $I = E_0 + I_0$  such that  $E_0$  is the harmonic component of  $E'$  and  $I_0$  satisfies  $-i = \delta(I_0)$  and  $\int_Z I_0 = 0$  on cycles.

**Remark 13.** *With this choice of  $E_0$ , we will find the solution with minimal energy. Indeed, the problem of uniqueness has been studied very carefully. It appears that, at least with finite source current and no voltage current, the two general solutions are the free current which is the solution of Flanders, and the wired current which is our solution. In this context, the uniqueness problem is to find conditions where these solutions coincide. See [LP14] for a precise presentation.*

**Lemma 14.** *Any cycle  $Z$  defines a 1-cochain  $E_Z$  in  $\text{Ker } \delta$  by the formula*

$$Z = \sum_{e \in \mathcal{E}^+} z_e e, z_e \in \mathbb{Z} \quad \Rightarrow \quad E_Z = \sum_{e \in \mathcal{E}^+} \frac{z_e}{r_e} e^*.$$

where the cochain  $e^*$  is defined by  $e^*(e) = 1$  and  $e^*(e') = 0$  if  $e' \neq \pm e$ .

The fact that  $E_Z \in \text{Ker } \delta$  is a simple consequence of (20).

**Definition 15.** *An  $L_2$ -cycle  $Z$  is an (infinite) cycle such that  $E_Z \in L_2(\mathcal{E})$ .*

**Lemma 16.** *For any  $E \in L_2(\mathcal{E})$  orthogonal to  $\text{Ker } \delta$  and any  $L_2$ -cycle  $Z$*

$$\int_Z E = 0.$$

Indeed, for any  $L_2$ -cycle,  $E_Z \in L_2(\mathcal{E})$  and

$$\int_Z E = \langle E, E_Z \rangle .$$

**Theorem 2.** *Let  $K$  be a connected, locally finite graph. We suppose that it is  $\chi$ -complete such that the operator  $D$  defined on  $C_0^0(K) \oplus C_0^1(K)$  is essentially self-adjoint. Then for any  $i \in C_0^0(K)$  satisfying  $\sum_{v \in \mathcal{V}} c(v)i(v) = 0$  and for any  $E' \in L_2(\mathcal{E})$  there exists a unique solution of minimal energy  $I \in L_2(\mathcal{E})$  of the problem :*

$$\delta(I) + i = 0, \text{ and } \forall Z, L_2\text{-cycle} \quad \int_Z E' = \int_Z I \quad (22)$$

*Proof.* — The space  $\text{Ker } \delta$  is closed in  $L_2(\mathcal{E})$ , so any element  $I \in L_2(\mathcal{E})$  can be written  $I = E_0 + I_0$  with  $E_0 \in \text{Ker } \delta$  and  $I_0$  in its orthogonal. By Lemma 16,  $E_0$  must be the orthogonal projection of  $E'$  on  $\text{Ker } \delta$  : if  $Z$  is a cycle, then  $E_Z \in \text{Ker } \delta$  and, as a consequence of Lemma 16,

$$\forall Z L_2\text{-cycle}, \quad \int_Z E' = \int_Z E_0.$$

Now, the existence of  $I_0$  is related to the property of  $-i$  to be in the range of  $\Delta$ . In the case where  $i$  has finite support, we can do as follows : let  $K_0$  be a finite connected subgraph of  $K$  (see 2.1.4, vertex of  $K_0$  are vertex of  $K$  and edges of  $K_0$  are edges of  $K$ ). We suppose that the support of  $i$  is included in  $K_0$ . Denote by  $d_0$  the difference operator of  $K_0$ . The Laplacian  $\Delta_0$  of  $K_0$  is self-adjoint and  $\text{Im } \Delta_0 = \text{Ker } \Delta_0^\perp$ . Thus, as  $\text{Ker } \Delta_0 = \mathbb{R}$  consists of constant functions

$$\langle i, 1 \rangle = 0 \Rightarrow \exists f \in C^0(K_0), \quad -i = \Delta_0(f).$$

Let  $\varphi \in C_0^1(K)$  be the extension of  $d_0 f$  by 0 on the edges which don't belong to  $K_0$ . This form is certainly different from  $df$  but  $\delta\varphi = -i$ .

We define now  $I_0$  as the orthogonal projection of  $\varphi$  on the orthogonal of  $\text{Ker } \delta$ , it means that  $I_0$  differs from  $\varphi$  by an element of  $\text{Ker } \delta$  and that  $I_0 \in \text{Ker } \delta^\perp$ . Using Lemma 16, we conclude that :

$$\delta I_0 = -i \text{ and } \forall Z L_2\text{-cycle}, \quad \int_Z I_0 = 0.$$

Thus  $I_0 + E_0$  is a solution of the problem, it is clearly the unique one.  $\square$

**Remark 17.** — *In the case of Flanders,  $E'$  has finite support, and we only take care of finite cycles, but the proof extends easily to  $E' \in L_2(\mathcal{E})$  if we consider only  $L_2$ -cycles. The question is how extend on more general  $i$ . It is related to the property of closeness of  $\text{Im}(\Delta)$ , what we explore below.*

**4.3. Anghel's hypothesis.** In [A93], N. Anghel shows that a Dirac type operator  $D$  defined on a complete manifold is Fredholm if and only if  $D^2$  is *positive at infinity*.

Let us define the complementary of a subgraph of a graph.

**Definition 18.** *For a subgraph  $K_0$  of a graph  $K$ , we define the complementary graph  $K_0^c = (\mathcal{V}^c, \mathcal{E}^c)$  as follows*

$$\mathcal{V}^c = \mathcal{V} \setminus \mathcal{V}_0, \quad \mathcal{E}^c = \{e \in \mathcal{E} \setminus \mathcal{E}_0, \partial(e) \subset \mathcal{V}^c\}.$$

**Remark 19.** (1) *In particular boundary points of edges in  $\mathcal{E}_0$  belong to  $\mathcal{V}_0$ .*

- (2) As a consequence of the definition,  $\mathcal{E}^c$  avoids the edges with one end in  $\mathcal{V}^c$  and one in  $\mathcal{V}_0$ .

Following [KL10], we define the *boundary* of a subgraph  $K_0$  to be its edge boundary :

$$\partial(K_0) = \mathcal{E} \setminus (\mathcal{E}_0 \cup \mathcal{E}^c).$$

**Definition 20.** We say that a Dirac type operator is positive at infinity if there exists a finite connected subgraph  $K_0 = (\mathcal{V}_0, \mathcal{E}_0)$  of  $K$  such that

$$\exists C > 0, \quad \forall (f, \varphi) \in L_2(\mathcal{V}^c) \times L_2(\mathcal{E}^c) \cap \text{Dom}(D), \quad \|(f, \varphi)\| \leq C \|D(f, \varphi)\|. \quad (23)$$

(Remark that this definition gives rather positivity of  $\Delta$ .)

**Theorem 3.** If the graph (connected and locally finite) is  $\chi$ -complete and if its Gauß-Bonnet operator

$$D = d + \delta$$

(which is essentially self-adjoint) is positive at infinity, then  $\text{Im}(D)$  is closed and, as a consequence, the Hodge property holds :

$$L_2(\mathcal{E}) = \text{Ker } \delta \oplus \text{Im}(d), \quad L_2(\mathcal{V}) = \text{Ker } d \oplus \text{Im}(\delta). \quad (24)$$

*Proof.* — The condition (23) implies that the closed restriction operator  $D^c$  of  $D$  on  $K_0^c$  :

$$D^c : \text{Dom}(D^c) \subset L_2(\mathcal{V}^c) \times L_2(\mathcal{E}^c) \rightarrow L_2(\mathcal{V}^c) \times L_2(\mathcal{E}^c)$$

is continuous (for the operator norm), injective and with closed image. By the inversion theorem, there exists

$$P : L_2(\mathcal{V}^c) \times L_2(\mathcal{E}^c) \rightarrow \text{Dom}(D^c)$$

such that  $P \circ D = \mathbb{I}$ , and  $\mathbb{I} - D \circ P$  is the orthogonal projector on the subspace  $\text{Im}(D^c)^\perp$ .

Let now  $\psi \in \overline{\text{Im}(D)}$ . It means :

$$\exists \text{ a sequence } (\sigma_n)_{n \in \mathbb{N}} \text{ in } \text{Dom}(D), \quad \sigma_n \in \text{Ker}(D)^\perp, \text{ and } \lim_{n \rightarrow \infty} D(\sigma_n) = \psi.$$

The sequence  $(\sigma_n)$  is bounded. If not, by extraction we can construct

$$\varphi_n = \frac{\sigma_n}{\|\sigma_n\|}$$

a subsequence tending to  $+\infty$  which satisfies

$$\|\varphi_n\| = 1, \quad \lim_{n \rightarrow \infty} D(\varphi_n) = 0.$$

Then the restriction of  $D(\varphi_n)$  to  $K_0^c$  also converge to 0 in  $L_2(\mathcal{V}^c) \times L_2(\mathcal{E}^c)$ .

But the set of vertices not in  $\mathcal{V}^c$  and the set of edges not in  $\mathcal{E}^c$  are finite. As  $\varphi_n$  is bounded, we can by extraction suppose that all their values in these finite sets converge, and by the same argument we can suppose that the value of  $\varphi_n$  on the vertices which are boundary points of edges in  $\partial(K_0)$  converge. By local finiteness we conclude that  $D(\varphi_n|_{K_0^c})$  converges.

By (23), then also  $\varphi_n|_{K_0^c}$  converges, thus finally  $\varphi_n$  converges, let  $\varphi$  be the limit, it satisfies

$$\|\varphi\| = 1, \quad \varphi \in \text{Ker}(D)^\perp, \quad D(\varphi) = 0 \quad \text{absurd.}$$

So we can suppose that  $(\sigma_n)$  is bounded, then by the same kind of reasoning, we show that  $(\sigma_n)_n$  admits a subsequence which converges, let  $\sigma$  be this limit. As  $D$  is closed and  $D(\sigma_n)$  converges, then  $\sigma \in \text{Dom}(D)$  and  $D(\sigma) = \psi$ .  $\square$

We see that the reasoning is separated for 0-forms and 1-forms. This gives :

**Corollary 21.** *Let  $K$  be a graph (connected and locally finite)  $\chi$ -complete so its Gauß-Bonnet operator  $D = d + \delta$  is essentially self-adjoint. If  $d$  satisfies the condition*

$$\exists C > 0, \quad \forall f \in L_2(\mathcal{V}^c) \cap \text{Dom}(d), \quad \|f\| \leq C\|df\| \tag{25}$$


for the complementary of some finite graph, then  $\text{Im } d$  is closed and

$$L_2(\mathcal{E}) = \text{Ker } \delta \oplus \text{Im}(d).$$

And there exists a similar statement for  $\delta$ .

### 5. EXAMPLES

It is clear that if  $K$  possesses infinitely many cycles (as infinite ladders, or infinite grids) it does not work. A family of examples could be a graph with finite geometry: there exists a finite subgraph  $K_0$  such that  $K_0^c$  is a disconnected (finite) union of branches.



**Proposition 22.** *If the connected graph  $K$  admits a finite subgraph such that its complementary is a finite union of trees with constant valence larger than 3, then, considered with the weights constant equal to 1 on vertex and edges, it is  $\chi$ -complete and  $\text{Im } d$  is closed.*

*Proof.* — We will prove that  $d$  is positive at infinity, *ie.* on each tree. Let  $U$  be a tree with a base point and valence  $p + 1$ ,  $p \geq 2$ . We apply Corollary 17 of [KL10], taking the notations of this paper (in particular  $\sharp$  denotes the cardinality): in our case  $D_U = p + 1$  is finite, so it suffices to show that the isoperimetric constant  $\alpha_U$  is

positive.

Recall that

$$\alpha_U = \inf_{W \subset U, \text{finite}} \frac{\sharp(\partial W)}{\sharp W}. \quad (26)$$

For a tree, one has a notion of *height*: the base point is of height 0, and for another point its height is the necessary number of edges to join it to the base point.

Let  $W$  be a finite set of vertices of  $U$ , we shall show by recurrence on  $\sharp W$  that

$$\sharp(\partial W) \geq \sharp W.$$

If  $\sharp W = 1$ , then  $\sharp(\partial W) = p+1$ . If  $\sharp W = n \geq 1$ , let  $x \in W$  be a point of highest height in  $W$  and  $y$  is the point just below. Then define  $W' = W - \{x\}$  so  $\sharp W' = \sharp W - 1$  and

$$\begin{aligned} y \in W &\Rightarrow \sharp(\partial W) = p - 1 + \sharp(\partial W') \\ y \notin W &\Rightarrow \sharp(\partial W) = p + 1 + \sharp(\partial W') \end{aligned}$$

In all cases, applying the recurrence hypothesis, we get:

$$\sharp(\partial W) \geq p - 1 + \sharp(\partial W') \geq p - 1 + \sharp W - 1 \geq \sharp W.$$

□

**Corollary 23.** *Such a graph (as in the proposition 22) satisfies also that  $\text{Im } \delta$  is closed and  $\text{Ker } d = \{0\}$  (because constants are not in  $L_2$ ), so  $\delta$  is surjective. As a consequence, for such a graph Flanders problem (18) has always a unique solution.*

*Proof.* — Indeed, if (25) is satisfied, then

$$\forall f \in \text{Dom}(\Delta^c) \subset L_2(\mathcal{V}^c), \quad \|f\| \leq C^2 \|\Delta(f)\|. \quad (27)$$

Thus, by the same reasoning as before the range of  $\Delta$  acting on functions is closed. Now if  $(\varphi_n)_n$  is a sequence of 1-forms such that  $\delta(\varphi_n)$  converges, we can apply the Hodge decomposition (24) at  $\varphi_n$ , because of the Proposition 22:

$$\exists f_n \in \text{Dom}(d) \text{ such that } \delta \circ d(f_n) \in L_2(\mathcal{V}) \text{ and converges.}$$

But we can extract a subsequence of  $(f_n)_n$  which converges, because of (27). □

**Proposition 24.** *If the connected graph  $K$  admits a finite subgraph such that its complementary is a finite union of trees with valence larger than 3, then, considered with the weights equal to the valence on vertices and constant equal to 1 on edges, it is  $\chi$ -complete and  $\text{Im } d$  is closed.*

*Proof.* — It is clear that such a graph satisfies the condition of  $\chi$ -completeness. The fact that  $d$  is positive at infinity is again a consequence of the results of [KL10]. Indeed, by hypothesis we have  $\forall v \in \mathcal{V}, m(v) = \sharp\{e \in \mathcal{E}, e^+ = v\}$  at least on the "tree-part", thus is it equal to the function  $n$  introduced in [KL10] and their  $d$  is constant equal to 1. By their Proposition 15, the quadratic form on a part  $U$  is

bounded from below by  $1 - \sqrt{1 - \alpha_U^2}$  if  $\alpha_U$  is the isoperimetric constant introduced in (26) but now with the volumes  $|\cdot|$  defined by the weights :

$$\alpha_U = \inf_{W \subset U, \text{finite}} \frac{|\partial W|}{|W|}.$$

Let  $W$  be a finite part of a tree. Its number of (oriented) edges is  $\sum_{v \in W} m(v) = |W|$ . But, because it is in a tree the number of interior edges is at most  $2 \cdot \sharp(W)$ . Thus


$$\frac{|\partial W|}{|W|} \geq \frac{\sum_{v \in W} (m(v) - 2)}{\sum_{v \in W} m(v)} \geq \frac{1}{3}$$

because  $m(v) \geq 3$ . □

The same Corollary as before holds, for the same reasons.

**Corollary 25.** *Such a graph (as in the Proposition 24) satisfies also that  $\text{Im } \delta$  is closed and  $\text{Ker } d = \{0\}$  (because constants are not in  $L_2$ ), so  $\delta$  is surjective. As a consequence, for such a graph, Flanders problem (18) has always a unique solution with minimal energy.*

**Remark 26.** *Take care to the fact that in these situations  $\text{Ker } \delta$  can be non trivial : on a tree of valence 3, with all the weights equal to 1, fix a point  $O$ , it has at least two edges which go to infinity :  $(x, O)$  and  $(y, O)$ .*



Let  $\varphi$  be the form such that

$$\varphi(x, O) = 1, \quad \varphi(y, O) = -1$$

at the  $n$ -level on the branch emanating from  $x$  we put the value of  $\varphi$  to be  $\frac{1}{2^n}$ , and

at the  $n$ -level on the branch emanating from  $y$  we put the value of  $\varphi$  to be  $\frac{-1}{2^n}$ .

Elsewhere, we put  $\varphi(e) = 0$ .

It is easy to verify that such a  $\varphi$  is in  $L_2$  and satisfies  $\delta(\varphi) = 0$ , see also [Ay13].

**Remark 27.** *In these two last cases the Laplacian is bounded, and the non zero spectrum is bounded from below because the isoperimetric constant  $\alpha_U$  admits a bound independant on  $U$ .*

**Acknowledgements** Part of this work was done while the author N.T-H was visiting the University of Nantes. She would like to thank the Laboratoire de Mathématiques Jean Leray (LMJL) for its hospitality. She is greatly indebted to the research unity (UR / 13 Z S 47) for its continuous support.

This work was supported by Grants through both Géanpyl project (FR 2962 du CNRS Mathématiques des Pays de Loire) and PHC-Utique (13 G 15-01) "Graphes, géométrie et théorie spectrale".

The authors thank Sylvain Golenia, Matthias Keller and Ognjen Milatovic for their reading with great interest and for their remarks.

#### REFERENCES

- [A93] N. Anghel, *An abstract index theorem on noncompact Riemannian manifolds*, Houston J. Math. **19** no. 2 (1993), 223–237.
- [Ay13] H. Ayadi, *Semi-Fredholmness of the discrete Gauß-Bonnet operator*, preprint, (2013).
- [Cm12] J. Carmesin, *A characterization of the locally finite networks admitting non-constant harmonic functions*, Potential Anal., **37**, (2012), 229–245.
- [Ch73] Paul R. Chernoff, *Essential self-adjointness of powers of generators of hyperbolic equations.*, J. Funct. Anal. **12** (1973), 401–414.
- [CdV91] Y. Colin de Verdière, *Théorème de Kirchhoff et théorie de Hodge*, Séminaire de théorie spectrale et géométrie, Chambéry-Grenoble (1990–1991), 89–94.
- [CdV98] Y. Colin de Verdière, *Spectres de graphes*, Cours Spécialisés [Specialized Courses], **4**, Société Mathématique de France, Paris, (1998).
- [CTT11] Y. Colin de Verdière, N. Torki-Hamza, F. Truc, *Essential Self-adjointness for combinatorial Schrödinger Operators II- Metrically non complete graphs*, Math Phys Anal Geom **14** (2011), 21–38.
- [D84] J. Dodziuk, *Difference equations, isoperimetric inequality and transience of certain random walks*, Trans. Amer. Math. Soc. **284** no. 2 (1984), 787–794.
- [DK87] J. Dodziuk and L. Karp, *Spectral and function theory for combinatorial Laplacians*, Geometry of random motion (Ithaca, N.Y., 1987) 25–40, Contemp. Math., 73, Amer. Math. Soc., Providence, RI, (1988).
- [DS99] P.G. Doyle and J.L. Snell, *Random walks and electric networks*, the Carus Mathematical Monographs, **22**, (1999).
- [F71] H. Flanders, *Uniqueness of electrical currents in a network of finite total resistance*, J. London Math. Soc. **82** no. 2 (2010), 256–272.
- [G10] A. Georgakopoulos, *Infinite networks : I- Resistive Networks*, IEEE Trans. Circuit Theory **CT-18** no. 3 (1971), 326–331
- [GH12] S. Golénia and T. Haugomat, *On the A.C. spectrum of 1D discrete Dirac operator* ArXiv 1207.3516, (2012) to appear in Meth. Funct. An. Top.
- [HKMW13] X. Huang, M. Keller, J. Masamune, R.K. Wojciechowski, *A note on self-adjoint extensions of the Laplacian on weighted graphs*, J. Funct. Anal. **265** no. 8 (2013), 1556–1578.
- [JP14] P.E.T. Jorgensen and E.P.J. Pearse, *Operator theory of electrical resistance networks*, 380 pages, to appear in Springer’s Universitext series, ArXiv 0806.3881.

- [KL10] M. Keller, D. Lenz, *Unbounded Laplacians on graphs: basic spectral properties and the heat equation*, Math. Model. Nat. Phenom. **5** no. 4 (2010), 198–224.
- [KL12] M. Keller, D. Lenz, *Dirichlet forms and stochastic completeness of graphs and subgraphs*, J. reine angew. Math., **666**, (2012), 189–223. arXiv:0904.2985 [math.FA]
- [LP14] R. Lyons, Y. Peres, *Probability on Trees and Networks*, Cambridge University Press (2014), In preparation. Current version available at <http://mypage.iu.edu/~rdlyons/>.
- [M09] J. Masamune, *A Liouville property and its application to the Laplacian of an infinite graph*, Contemporary Mathematics **484** (2009), 103–115.
- [RS80] M. Reed, B. Simon, *Methods of Modern Mathematical Physics I*, Academic Press (1980).
- [S94] P.M. Soardi, *Potential theory on infinite networks*, Lecture Notes in Mathematics **1590**, Springer (1994).
- [T90] C. Thomassen, *Resistances and currents in infinite networks*, J. Comb. Th. **B 49**, (1990), 87–102.
- [T10] N. Toriki-Hamza, *Laplaciens de graphes infinis I - Graphes métriquement complets*, Confluentes Mathematici, **2** no. 3 (2010), 333–350.
- [T12] N. Toriki-Hamza, *Essential Self-adjointness for combinatorial Schrödinger Operators I- Metrically complete graphs*, ArXiv:1201.4644 , 1–22, Translation of [T10] with some add, correction and update.
- [Z08] A. H. Zemanian, *Infinite electrical networks*, Cambridge Tracts in Mathematics ( **101**), ( 2008), 324 p.

LABORATOIRE DE MATHÉMATIQUES JEAN LERAY, UNIVERSITÉ DE NANTES, CNRS, FACULTÉ DES SCIENCES, BP 92208, 44322 NANTES, FRANCE

*E-mail address:* `colette.anne@univ-nantes.fr`

UNITÉ DE RECHERCHES MATHÉMATIQUES ET APPLICATIONS, UR / 13 Z S 47, FACULTÉ DES SCIENCES DE BIZERTE DE L'UNIVERSITÉ DE CARTHAGE, 7021-BIZERTE; ISIG-K, UNIVERSITÉ DE KAIROUAN, 3100-KAIROUAN; TUNISIE

*E-mail address:* `nabila.torki-hamza@fsb.rnu.tn; natorki@gmail.com`