



HAL
open science

On full groups of non ergodic probability measure preserving equivalence relations

François Le Maître

► **To cite this version:**

François Le Maître. On full groups of non ergodic probability measure preserving equivalence relations. 2014. hal-00992639v1

HAL Id: hal-00992639

<https://hal.science/hal-00992639v1>

Preprint submitted on 19 May 2014 (v1), last revised 15 Feb 2024 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On full groups of non ergodic probability measure preserving equivalence relations

François Le Maître

Abstract

This article generalizes the results in [LM14] by giving a formula relating the topological rank of the full group of an aperiodic pmp equivalence relation to the cost of its ergodic components. Furthermore, we obtain examples of full groups having a dense free subgroup whose rank is equal to the topological rank of the full group, using a Baire category argument. We then study the automatic continuity property for full groups of aperiodic equivalence relations, and find a connected metric for which they have the automatic continuity property. This allows us to give an algebraic characterization of aperiodicity for pmp equivalence relations, namely the non-existence of homomorphisms from their full groups into totally disconnected separable groups. A simple proof of the extreme amenability of full groups of hyperfinite pmp equivalence relations is also given, generalizing to the non ergodic case a result of Giordano and Pestov [GP07, thm. 5.7].

Contents

1	Introduction	2
1.1	Topological rank for full groups	4
1.2	(Dense) free groups in full groups	5
1.3	Automatic continuity for a finer metric	6
1.4	Extreme amenability	8
1.5	Notations and preliminaries	8
2	Conditional expectation and non ergodic equivalence relations	9
2.1	Conditional expectation and conditional measure	9
2.2	A non ergodic analogue of Zimmer's lemma	10
2.3	Conditional cost	11
3	The hyperfinite ergodic equivalence relation \mathcal{R}_0	12
3.1	Dyadic permutations	12
3.2	The odometer T_0	13
4	Two topological generators for the full group of $\mathcal{R}_0^Y = \Delta_Y \times \mathcal{R}_0$	14
4.1	Some subgroups which topologically generate $[\mathcal{R}_0^Y]$	14
4.2	Two generators for many finite subgroups	15
4.3	Two topological generators for $[\mathcal{R}_0^Y]$	16

5	Consequences	20
5.1	Computation of the topological rank for full groups	20
5.2	A dense G_δ set of topological generators	21
6	Free subgroups in full groups	22
7	A finer metric on full groups	25
7.1	Automatic continuity for $([\mathcal{R}], d_C)$	25
7.2	Connectedness	27
8	Extreme amenability	28

1 Introduction

When considering dynamical systems given by a measure preserving bijection on a standard probability space, one has at hand various invariants which sometimes allow to classify them up to conjugacy. For instance, the entropy is a complete invariant for Bernoulli shifts, meaning that two Bernoulli shifts are conjugate iff they share the same entropy. An easier to define invariant is the partition of the probability space induced by the orbits of the dynamical systems. This partition turns out to be as far as possible from a complete invariant: a result of Dye asserts that up to a measure preserving bijection, all ergodic dynamical systems induce the same partition of the space almost everywhere [Dye59]. In other words, they are all **orbit equivalent**.

Orbit equivalence is thus trivial in the realm of classical ergodic theory, that is, ergodic theory of probability measure preserving \mathbb{Z} -action. We then move to the setting of probability measure preserving (pmp) actions of arbitrary countable groups Γ , where the picture gets much richer. Indeed, whenever Γ is a non amenable group, Epstein has shown that it admits a continuum of non orbit equivalent pmp ergodic free actions [Eps08]. On the other hand, generalizing the aforementioned result of Dye, Ornstein and Weiss have shown that any two pmp ergodic actions of any two amenable groups are orbit equivalent [OW80].

Given these results, it is desirable to have a more precise understanding of the situation for pmp ergodic free actions of non amenable groups. But given a non-amenable group Γ , there are currently no invariants which yield an infinite family of non orbit equivalent pmp free ergodic Γ -actions. However, if one wants to distinguish the partitions induced by pmp free ergodic actions of two different non-amenable groups, there is an invariant which might do the trick, namely the cost.

Introduced by Levitt in 1995 [Lev95], the cost of a pmp action is roughly the minimal number of elements needed to generate the partition of the space into orbits induced by this action. In 2000, Gaboriau showed that all the pmp free actions of the free group on n generators have cost n [Gab00], so that free groups of different ranks can never have orbit equivalent pmp free actions. As we will see later, the cost is deeply linked to another natural invariant of pmp actions: their full group, which we now define.

Let us fix a standard probability space (X, μ) and denote by $\text{Aut}(X, \mu)$ the group of its measure preserving bijections, identified up to null sets. Given a non trivial pmp action of a countable group Γ on (X, μ) , one associates to it a **pmp equivalence relation** \mathcal{R}_Γ defined by $x\mathcal{R}_\Gamma y$ iff $x \in \Gamma y$. Because the equivalence classes of \mathcal{R}_Γ are precisely the Γ -orbits, orbit equivalence can be rephrased as the study of such pmp equivalence relations up to isomorphism. Now if \mathcal{R} is a pmp equivalence relation, its **full group**, denoted by $[\mathcal{R}]$ is defined by

$$[\mathcal{R}] = \{T \in \text{Aut}(X, \mu) : \forall x \in X, T(x)\mathcal{R}x\}.$$

The full group of a pmp equivalence relation is a Polish group when equipped with the uniform metric $d_u(T, T') = \mu(\{x \in X : T(x) \neq T'(x)\})$, homeomorphic as a topological space to an infinite dimensional Hilbert space [KT10, thm. 1.5].

Suppose that $T : X \rightarrow X$ is an isomorphism between two pmp equivalence relations \mathcal{R} and \mathcal{R}' , i.e. a class measure preserving bijection of (X, μ) such that for almost every $x \in X$, $T([x]_{\mathcal{R}}) = [T(x)]_{\mathcal{R}'}$. Then T conjugates the full group of \mathcal{R} to the full group of \mathcal{R}' , which are consequently isomorphic as topological groups. In other words, the full group is an invariant of orbit equivalence.

We thus leave the realm of orbit equivalence and are led to study these full groups as topological groups in their own right, which was the main motivation for the already mentioned article of Kittrell and Tsankov [KT10]. Furthermore, Dye's reconstruction theorem implies that full groups are a complete invariant of orbit equivalence: any abstract group isomorphism between full groups must descend from a class measure preserving orbit equivalence between the corresponding equivalence relations, as long as one of the two pmp equivalence relations has no finite orbits.

Let us now give examples where one can see concretely how the full group "remembers" the pmp equivalence relation.

Theorem 1.1 (Eigen [Eig81], Dye [Dye63, prop. 5.1]). *Let \mathcal{R} be a pmp equivalence relation. Then \mathcal{R} is ergodic iff $[\mathcal{R}]$ is simple. Furthermore, when \mathcal{R} is not ergodic, the closed normal subgroups of its full group are of the form*

$$[\mathcal{R}_A] = \{T \in [\mathcal{R}] : \text{supp } T \subseteq A\},$$

where A is a \mathcal{R} -invariant subset of X .

Theorem 1.2 (Giordano-Pestov [GP07, thm. 5.7]). *Let \mathcal{R} be an ergodic equivalence relation. Then \mathcal{R} is amenable iff $[\mathcal{R}]$ is extremely amenable.*

Theorem 1.3 (Le Maître [LM14, thm. 1]). *Let \mathcal{R} be a pmp ergodic equivalence relation. Then topological rank¹ $t([\mathcal{R}])$ of the full group of \mathcal{R} is related to the integer part of the cost of \mathcal{R} by the formula*

$$t([\mathcal{R}]) = \lfloor \text{Cost}(\mathcal{R}) \rfloor + 1.$$

This article is an attempt to understand further algebraic and topological properties of full groups, especially in the non ergodic case. The results are extracted from the author's thesis.

¹By definition, the topological rank $t(G)$ of a topological group G is the minimal number of elements needed to generate a dense subgroup of G .

1.1 Topological rank for full groups

Our main result is a direct continuation of [LM14], where the question of the topological rank of the full group of a pmp ergodic equivalence relation \mathcal{R} was addressed, using previous works of Kittrell, Tsankov [KT10] and Matui [Mat13]. We may first note that when a pmp equivalence relation \mathcal{R} has only finite orbits, its full group cannot be topologically generated by a finite number of elements, for it is a locally finite group (i.e. all finitely generated subgroups are finite). Moreover, if there is a positive \mathcal{R} -invariant set A onto which \mathcal{R} has only finite equivalence classes, the full group of \mathcal{R} factors continuously onto the full group of its restriction to A , so that the full group of \mathcal{R} is not topologically finitely generated. So we can restrict ourselves to **aperiodic** pmp equivalence relations, that is, having only infinite classes.

In this article, we compute the topological rank of full groups of pmp aperiodic equivalence relations, using the same line of ideas as in [LM14], but with some modifications which we now briefly explain. First, in [LM14] we used a result of Matui [Mat13, thm. 3.2] which gives two topological generators for the full group of the *ergodic* hyperfinite equivalence relation², and so we had to generalize Matui's theorem to non-ergodic hyperfinite equivalence relations (theorem 4.1).

Second, instead of working with the ergodic decomposition, we chose a more “functional” approach which was already present in Dye's founding paper [Dye59]. If \mathcal{R} is a pmp equivalence relation, we have a conditional expectation which turns every measurable map into an $M_{\mathcal{R}}$ -measurable map, where $M_{\mathcal{R}}$ is the σ -algebra of all \mathcal{R} -invariant sets. This yields what we call the **\mathcal{R} -conditional measure** of a set A , defined to be the conditional expectation of its characteristic function χ_A . Such a conditional measure can of course be understood as $x \mapsto \mu_x(A)$, where $(\mu_x)_{x \in X}$ is the ergodic decomposition of \mathcal{R} . Also note that when \mathcal{R} is ergodic, the \mathcal{R} -conditional measure equals μ . Every element of $[\mathcal{R}]$ preserves the \mathcal{R} -conditional measure and conversely, if $A, B \subseteq X$ have the same \mathcal{R} -conditional measure, then there is $T \in [\mathcal{R}]$ such that $T(A) = B$ (proposition 2.3). More details can be found in subsection 2.1.

In the same spirit, we introduce a notion of cost for non-ergodic equivalence relations which can be understood as a function mapping an ergodic component to its cost, which we refer to as conditional cost. Again rather than using the ergodic decomposition, we see the conditional cost of a pmp equivalence relation \mathcal{R} as an $M_{\mathcal{R}}$ -measurable function from X to $\mathbb{R}^+ \cup \{+\infty\}$. Also note that when \mathcal{R} is ergodic, the conditional cost is the usual cost. For more details, cf. subsection 2.3.

We can now state our main theorem:

Theorem 1.4. *Let \mathcal{R} be an aperiodic pmp equivalence relation. Then the topological rank $t([\mathcal{R}])$ of the full group of \mathcal{R} is related to the conditional cost of \mathcal{R} by the relation*

$$t([\mathcal{R}]) = \operatorname{ess\,sup}_{x \in X} [\text{C-Cost}(\mathcal{R})(x)] + 1.$$

Because every countable product of full groups can itself be seen as a full group, we have the following corollary.

²Note that Andrew Marks has now a much shorter proof of Matui's result, available as a note on his website.

Corollary 1.5. *Let $(\mathcal{R}_i)_{i \in I}$ be a countable family of pmp equivalence relations. Then*

$$t \left(\prod_{i \in I} [\mathcal{R}_i] \right) = \sup_{i \in I} t([\mathcal{R}_i]).$$

Remark. Theorem 1.4 also allows us to remove the ergodicity assumption in corollaries 2 and 3 of [LM14].

1.2 (Dense) free groups in full groups

Recall that $T \in \text{Aut}(X, \mu)$ is called **aperiodic** if all its orbits are infinite, and that the set of such elements, denoted by APER , is closed for the uniform metric. Our next result goes into the direction of a better understanding of free groups in full groups, and generalizes [Kec10, thm. 3.7] (cf. theorem 6.2).

Theorem 1.6. *Let \mathcal{R} be a pmp equivalence relation, and let $T \in [\mathcal{R}]$ have infinite order. Then for all $n \in \mathbb{N}$ we have the following:*

(1) *The set of n -uples (U_1, \dots, U_n) of elements of $[\mathcal{R}]$ such that*

$$\langle T, U_1, \dots, U_n \rangle \simeq \mathbb{F}_{n+1}$$

is a dense G_δ in $[\mathcal{R}]^n$.

(2) *If furthermore \mathcal{R} is aperiodic, then the set of n -uples of aperiodic elements (U_1, \dots, U_n) of $[\mathcal{R}]$ such that*

$$\langle T, U_1, \dots, U_n \rangle \simeq \mathbb{F}_{n+1}$$

is a dense G_δ in $([\mathcal{R}] \cap \text{APER})^n$.

Combining this theorem with a density result for topological generators in aperiodic pmp equivalence relations of cost one (proposition 5.1), we get a good understanding of topological generators for full groups of aperiodic pmp equivalence relations of cost one (theorem 6.3).

Theorem 1.7. *Let \mathcal{R} be any pmp equivalence relation, and consider the set*

$$G = \{(T, U) \in (\text{APER} \cap [\mathcal{R}]) \times [\mathcal{R}] : \overline{\langle T, U \rangle} = [\mathcal{R}] \text{ and } \langle T, U \rangle \simeq \mathbb{F}_2\}.$$

Then the following assertions are equivalent.

(1) *\mathcal{R} is aperiodic of cost one.*

(2) *G is a dense G_δ in $(\text{APER} \cap [\mathcal{R}]) \times [\mathcal{R}]$.*

Note that the aperiodicity restriction on the first coordinate is necessary. Indeed, topological generators of $[\mathcal{R}]$ have to generate the equivalence relation \mathcal{R} , and if D is any dense subset of $[\mathcal{R}] \times [\mathcal{R}]$, it contains elements of arbitrarily small support. Then, by definition of the cost some elements of D do not generate the equivalence relation \mathcal{R} , hence cannot be topological generators of $[\mathcal{R}]$.

When \mathcal{R} is generated by a single pmp automorphism³, the set $\text{GEN}(\mathcal{R})$ of generators of \mathcal{R} is a dense G_δ in $\text{APER} \cap [\mathcal{R}]$, so that in this case we have the following reformulation of theorem 1.7.

Corollary 1.8. *Let \mathcal{R} be any aperiodic pmp singly-generated equivalence relation. Then the set*

$$\{(T, U) \in \text{GEN}(\mathcal{R}) \times [\mathcal{R}] : \overline{\langle T, U \rangle} = [\mathcal{R}] \text{ and } \langle T, U \rangle = \mathbb{F}_2\}$$

is a dense G_δ in $\text{GEN}(\mathcal{R}) \times [\mathcal{R}]$.

Using the Kuratowski-Ulam theorem, we see that for \mathcal{R} singly-generated aperiodic, the set of $T \in \text{GEN}(\mathcal{R})$ such that

$$\{U \in [\mathcal{R}] : \overline{\langle T, U \rangle} = [\mathcal{R}] \text{ and } \langle T, U \rangle = \mathbb{F}_2\}$$

is itself a dense G_δ in $\text{GEN}(\mathcal{R})$. But it might actually be equal to $\text{GEN}(\mathcal{R})$, which yields the following general question.

Question 1.9. Let T be an aperiodic element of $\text{Aut}(X, \mu)$, let \mathcal{R}_T be the pmp equivalence relation it generates. Does there exist $U \in [\mathcal{R}_T]$ such that $\langle T, U \rangle = [\mathcal{R}_T]$? If yes, is the set of such U 's a dense G_δ ?

Remark. In the author's thesis, it is shown that the answer to both these questions is yes for every $T \in \text{Aut}(X, \mu)$ of rank one, constructing a $U \in [\mathcal{R}_T]$ very similar to the one we have here for the odometer (theorem 4.1). But this idea seems to fail for infinite rank transformations like Bernoulli shifts.

When \mathcal{R} is aperiodic of cost one, theorem 1.7 also gives a positive answer to the following question, which was suggested to the author by Gelfander.

Question 1.10. Let \mathcal{R} be an aperiodic pmp equivalence relation. Does it have a dense free subgroup of rank $t([\mathcal{R}])$?

We do not even have an *explicit* dense free subgroup of rank 2 in the full group of the hyperfinite ergodic equivalence relation. Let us end this section by mentioning a very interesting conjecture of Thom about free groups in full groups.

Conjecture (Thom). There are no discrete free subgroups of rank 2 in the full group of the hyperfinite ergodic equivalence relation.

1.3 Automatic continuity for a finer metric

Dye's reconstruction theorem [Dye63, thm. 2] states that whenever a pmp equivalence relation \mathcal{R} is aperiodic, any abstract group isomorphism ψ between the full groups of \mathcal{R} and another pmp equivalence relation \mathcal{R}' is implemented by a class measure preserving orbit equivalence $T \in \text{Aut}^*(X, \mu)$ between \mathcal{R} and \mathcal{R}' , i.e. we have $\psi(g) = TgT^{-1}$ for all $g \in [\mathcal{R}]$ and $T \times T(\mathcal{R}) = \mathcal{R}'$. Moreover, if \mathcal{R} is ergodic, T

³By a theorem of Dye [Dye59, thm. 1], a pmp equivalence relation \mathcal{R} is singly-generated iff it is hyperfinite (cf. definition 1.16), which in turn is equivalent to the amenability of \mathcal{R} [CFW81].

is automatically measure preserving. So in the ergodic case Dye's theorem implies that any abstract group isomorphism between full groups is an isometry, while in the non ergodic case it states that it is a homeomorphism. This was the motivation for Kittrell and Tsankov to study the automatic continuity property for full groups.

Definition 1.11. Let G be a topological group. It satisfies the **automatic continuity property** if every homomorphism from G to a separable topological group is continuous.

Theorem 1.12 ([KT10, thm. 1.2]). *Let \mathcal{R} be a pmp ergodic equivalence relation. Then the full group of \mathcal{R} endowed with the uniform metric satisfies the automatic continuity property.*

It is a classical fact that whenever G is a Polish group having the automatic continuity property, every isomorphism between G and a Polish group H is a homeomorphism (cf. for instance [Ros09]). The preceding theorem can thus be seen as a partial generalization of Dye's reconstruction theorem, and also has the following important consequence.

Corollary 1.13 ([KT10, thm. 1.2]). *Every full groups of a pmp ergodic equivalence relations carries a unique Polish group topology, namely the uniform topology.*

Note that whenever \mathcal{R} is not aperiodic and non trivial, its full group does not have the automatic continuity property, for it is connected but admits a non trivial morphism into $\mathbb{Z}/2\mathbb{Z}$ (cf. corollary 7.6 and the paragraph following it). So a natural question is: what happens in the aperiodic case⁴?

We only provide a very partial answer to this question (theorem 7.3), namely we define a finer connected metric d_C on the full group of \mathcal{R} such that $([\mathcal{R}], d_C)$ has the automatic continuity property whenever $[\mathcal{R}]$ is aperiodic. This metric can be thought of as uniform convergence along ergodic components, so that in the ergodic case it is the uniform metric, but in general it is not separable. Our result is thus a generalization of theorem 1.12, but the proof is very similar, and we think the metric d_C is actually the right setting for Kittrell and Tsankov's result.

We use this result to give a characterization of aperiodic equivalence relations in terms of their full groups (corollary 7.6).

Theorem 1.14. *Let \mathcal{R} be a pmp equivalence relation. Then the following assertions are equivalent.*

- (1) \mathcal{R} is aperiodic.
- (2) $[\mathcal{R}]$ has no nontrivial morphisms into $\mathbb{Z}/2\mathbb{Z}$.
- (3) $[\mathcal{R}]$ has no nontrivial morphisms into totally disconnected separable groups.

⁴When \mathcal{R} has finitely many ergodic components, its full group is the finite product of the full groups of its restrictions to these components. But by theorem 1.12, each of these has the automatic continuity property, hence the full group of \mathcal{R} itself has the automatic continuity. So we may restrict ourselves to aperiodic pmp equivalence relations having either a continuous or a countably infinite ergodic decomposition.

1.4 Extreme amenability

Our last result (theorem 8.4) generalizes a result of T. Giordano and V. Pestov on the relationship between the amenability of a pmp equivalence relation and the extreme amenability of its full group.

Definition 1.15. A topological group G is **extremely amenable** if every continuous G -action on a compact space K admits a fixed point.

Extreme amenability is a very strong property, specific to “infinite dimensional groups”, for no non-trivial locally compact group can be extremely amenable [Vee77]. The terminology is justified by the following definition of amenability for topological groups: a topological group G is **amenable** if every continuous G -action *by affine isometries* on a compact *convex subset of a locally convex topological vector space* admits a fixed point.

In the discrete case, amenability admits a lot of different reformulations, some of which yield different characterizations of amenability for equivalence relations, and we refer the reader to [Kai97] for these. For our purposes, the following definition is the most relevant, and turns out to be equivalent to amenability by a celebrated result of Connes, Feldman and Weiss [CFW81]. Recall that a pmp equivalence relation is **finite** if all its classes are finite.

Definition 1.16. A pmp equivalence relation \mathcal{R} is **hyperfinite** if it can be written as a countable increasing union of finite pmp equivalence relations.

Giordano and Pestov showed in [GP07, prop. 5.3] that an ergodic pmp equivalence relation is hyperfinite iff its full group is extremely amenable. We generalize this to the non ergodic setting and give an easier proof (cf. theorem 8.4). One should note however that our proof does not yield that the full group of an amenable equivalence relation is a Levy group, while their does. Also, we still rely on the concentration of measure phenomenon via a theorem of Glasner [Gla98, theorem 1.3], and it would be nice to have a proof which does not.

1.5 Notations and preliminaries

Everything will take place “modulo sets of measure zero”. A **cycle** is a measure-preserving automorphism of (X, μ) who has only finite orbits. A cycle is **odd** when all its orbits have an odd cardinality. We keep the same notations and definitions as in [LM14, sec. 1 and 2], which we now briefly recall.

If (X, μ) is a standard probability space, and A, B are Borel subsets of X , a **partial isomorphism** of (X, μ) of **domain** A and **range** B is a Borel bijection $f : A \rightarrow B$ which is measure-preserving for the measures induced by μ on A and B respectively. We denote by $\text{dom } f = A$ its domain, and by $\text{rng } f = B$ its range. Note that in particular, $\mu(\text{dom } f) = \mu(\text{rng } f)$. A **graphing** is a countable set of partial isomorphisms of (X, μ) , denoted by $\Phi = \{\varphi_1, \dots, \varphi_k, \dots\}$ where the φ_k 's are partial isomorphisms. It **generates** a pmp equivalence relation \mathcal{R}_Φ , defined to be the smallest equivalence relation containing $(x, \varphi(x))$ for every $\varphi \in \Phi$ and $x \in \text{dom } \varphi$.

The **pseudo full group** of \mathcal{R} , denoted by $[[\mathcal{R}]]$, consists of all partial isomorphisms φ such that $\varphi(x) \mathcal{R} x$ for all $x \in \text{dom } \varphi$.

Let $p \in \mathbb{N}$. A **pre- p -cycle** is a graphing $\Phi = \{\varphi_1, \dots, \varphi_{p-1}\}$ such that the following two conditions are satisfied:

- (i) $\forall i \in \{1, \dots, p-1\}, \text{rng } \varphi_i = \text{dom } \varphi_{i+1}$.
- (ii) The following sets are all disjoint:

$$\text{dom } \varphi_1, \text{dom } \varphi_2, \dots, \text{dom } \varphi_{p-1}, \text{rng } \varphi_{p-1}.$$

A **p -cycle** is an element $C \in \text{Aut}(X, \mu)$ whose orbits have cardinality 1 or p . So according to the terminology introduced at the beginning of this section, a p -cycle is a cycle which is odd iff p is odd.

Given a pre- p -cycle $\Phi = \{\varphi_1, \dots, \varphi_{p-1}\}$, we can extend it to a p -cycle $C_\Phi \in \text{Aut}(X, \mu)$ as follows:

$$C_\Phi(x) = \begin{cases} \varphi_i(x) & \text{if } x \in \text{dom } \varphi_i \text{ for some } i < p, \\ \varphi_1^{-1} \varphi_2^{-1} \cdots \varphi_{p-1}^{-1}(x) & \text{if } x \in \text{rng } \varphi_{p-1}, \\ x & \text{otherwise.} \end{cases}$$

The following theorem will again prove very useful.

Theorem 1.17 ([KT10], theorem 4.7). *Let $\mathcal{R}_1, \mathcal{R}_2, \dots$ be measure-preserving equivalence relations on (X, μ) , and let \mathcal{R} be their join (i.e. the smallest equivalence relation containing all of them). Then $\langle \bigcup_{n \in \mathbb{N}} [\mathcal{R}_n] \rangle$ is dense in $[\mathcal{R}]$.*

2 Conditional expectation and non ergodic equivalence relations

2.1 Conditional expectation and conditional measure

Given a pmp equivalence relation \mathcal{R} , one may look at the measure subalgebra of the \mathcal{R} -invariant sets, i. e. the set of all $A \in \text{MAlg}(X, \mu)$ such that for any $\varphi \in [\mathcal{R}]$, $\varphi^{-1}(A) = A$. We denote this closed subalgebra by $M_{\mathcal{R}}$, and by $L^2_{\mathcal{R}}(X)$ the set of square integrable $M_{\mathcal{R}}$ -measurable real functions.

The orthogonal projection $\mathbb{E}_{\mathcal{R}}$ from the Hilbert space $L^2(X)$ onto the closed subspace $L^2_{\mathcal{R}}(X)$ satisfies the following equality, which defines it uniquely: for any $f \in L^2(X)$ and $g \in L^2_{\mathcal{R}}(X)$,

$$\int_X fg = \int_X \mathbb{E}_{\mathcal{R}}(f)g.$$

$\mathbb{E}_{\mathcal{R}}$ is called a **conditional expectation**. When A is a subset of X , its characteristic function is an element of $L^2(X)$, and we call $\mathbb{E}_{\mathcal{R}}(\chi_A)$ the **\mathcal{R} -conditional measure** of A , denoted by $\mu_{\mathcal{R}}(A)$. Because $\mathbb{E}_{\mathcal{R}}$ is a contraction for the L^∞ norm, $\mu_{\mathcal{R}}(A)$ is an $M_{\mathcal{R}}$ -measurable function taking values in $[0, 1]$. Proposition 2.2 roughly states that any $M_{\mathcal{R}}$ -measurable function $f : (X, \mu) \rightarrow [0, 1]$ is equal to $\mu_{\mathcal{R}}(A)$ for

some $A \subseteq X$.

The following proposition is due to Dye [Dye59]; we give a simple proof, based on the marker lemma which we now recall.

Lemma 2.1 ([KM04], lemma 6.7). *Let \mathcal{R} be an aperiodic equivalence relation, then there exists a decreasing sequence of $A_n \subseteq X$ which intersect every \mathcal{R} class and such that $\bigcap_n A_n = \emptyset$.*

We call such A_n 's a sequence of **markers**.

Proposition 2.2 ([Dye59], Maharam's lemma). *\mathcal{R} is aperiodic iff for any $A \subseteq X$, and for any $M_{\mathcal{R}}$ -measurable function f such that $0 \leq f \leq \mu_{\mathcal{R}}(A)$, there exists $B \subseteq A$ such that the \mathcal{R} -conditional measure of B equals f .*

Proof. First note that we can restrict ourselves to the case $0 < f \leq \mu_{\mathcal{R}}(A)$. By a maximality argument, it suffices to show that we can find $B \subseteq A$ such that $0 < \mu_{\mathcal{R}}(B) \leq f$. Let A_n be a sequence of markers for $\mathcal{R}_{\upharpoonright A}$, note that $\mu(A_n) = \int \mu_{\mathcal{R}}(A_n) \rightarrow 0$, so up to taking a subsequence we can assume $\mu_{\mathcal{R}}(A_n)$ converges pointwise to 0. Now define B to be the set of $x \in X$ such that there exists an integer n for which $x \in A_n$ and $\mu_{\mathcal{R}}(A_n)(x) < f(x)$.

Conversely, if $\mathcal{R}_{\upharpoonright A}$ is finite, one can further restrict A and suppose all its classes are of size n . Now the function $\frac{1}{2n}$ cannot arise as the conditional measure of $B \subseteq A$. \square

Remark. The above proposition has the following nice consequence: given $A \subseteq X$, we can split it "equally among ergodic components": one puts $f = \mu_{\mathcal{R}}(A)/2$ and gets $B \subseteq A$ such that $\mu_{\mathcal{R}}(B) = \mu_{\mathcal{R}}(A \setminus B) = \mu_{\mathcal{R}}(A)/2$ (in particular, $\mu(B) = \frac{\mu(A)}{2}$). And of course one can also split A in any finite number of pieces, all which have the same conditional measure.

Proposition 2.3. *Let \mathcal{R} be a pmp equivalence relation. Then if two sets A and B have the same \mathcal{R} -conditional measure, there exists $\varphi \in [[\mathcal{R}]]$ whose domain is A and whose range is B .*

Proof. This is a standard maximality argument. One can for instance use the construction in [KM04, lemma 7.10], and check that if $\mu(A \setminus A') > 0$, then the \mathcal{R} -conditional measures of $A \setminus A'$ and $B \setminus B'$ are the same and positive. But then the \mathcal{R} -saturation of $A \setminus A'$ contains $B \setminus B'$, so that there exists a minimal n such that $\gamma_n^{-1}(B \setminus B') \cap (A \setminus A')$ is non null, a contradiction. \square

2.2 A non ergodic analogue of Zimmer's lemma

Recall that every closed subalgebra of $\text{MAlg}(X, \mu)$ arises as the inverse image of $\text{MAlg}(Y, \nu)$ by some measure preserving $\pi : (X, \mu) \rightarrow (Y, \nu)$, where (Y, ν) is a standard probability space possibly with atoms. So if \mathcal{R} is a measure preserving equivalence relation, one can find a standard probability space $(Y_{\mathcal{R}}, \nu_{\mathcal{R}})$, possibly with atoms, and a measure preserving map $\pi_{\mathcal{R}} : (X, \mu) \rightarrow (Y_{\mathcal{R}}, \nu_{\mathcal{R}})$ such

that $\pi_{\mathcal{R}}^{-1}(\text{MAlg}(Y_{\mathcal{R}}, \nu_{\mathcal{R}})) = M_{\mathcal{R}}$. This yields an identification between $L^2_{\mathcal{R}}(X)$ and $L^2(Y_{\mathcal{R}}, \nu_{\mathcal{R}})$.

The following proposition is essentially due to Dye [Dye59, thm. 4 and 5], and generalizes the well-known fact often attributed to Zimmer that any ergodic equivalence relation contains an ergodic subequivalence relation generated by a single automorphism.

Proposition 2.4 (Dye). *Let \mathcal{R} be a pmp aperiodic equivalence relation on (X, μ) , let $\pi_{\mathcal{R}} : (X, \mu) \rightarrow (Y_{\mathcal{R}}, \nu_{\mathcal{R}})$ be the measure preserving map associated with the algebra of \mathcal{R} -invariant subsets of X . Then there exists an isomorphism $\varphi : (X, \mu) \rightarrow (Y_{\mathcal{R}}, \nu_{\mathcal{R}}) \otimes (2^{\mathbb{N}}, \lambda)$ such that $\text{id}_{Y_{\mathcal{R}}} \times \mathcal{R}_0$ is mapped into \mathcal{R} by φ^{-1} , and the following diagram commutes:*

$$\begin{array}{ccc} (X, \mu) & \xrightarrow{\varphi} & (Y_{\mathcal{R}} \times 2^{\mathbb{N}}, \nu_{\mathcal{R}} \otimes \lambda) \\ & \searrow \pi_{\mathcal{R}} & \swarrow \pi \\ & (Y_{\mathcal{R}}, \nu_{\mathcal{R}}) & \end{array}$$

2.3 Conditional cost

Let \mathcal{R} be an aperiodic equivalence relation. As orbit equivalence remembers the ergodic decomposition of \mathcal{R} , the following notion makes sense as an invariant for non ergodic equivalence relations.

Definition 2.5. The **C-cost** (conditionnal cost) of a pmp equivalence relation \mathcal{R} is the infimum over all graphings $\Phi = \{\varphi_i\}_{i \in \mathbb{N}}$ which generate \mathcal{R} of the functions

$$\text{C-Cost}(\Phi) = \sum_{i \in \mathbb{N}} \mu_{\mathcal{R}}(\varphi_i).$$

Let us check that this definition makes sense. The functions $\mu_{\mathcal{R}}(\Phi)$ are positive elements of the von Neumann algebra $L^{\infty}_{\mathcal{R}}(X, \mu) = L^{\infty}(Y_{\mathcal{R}}, \nu_{\mathcal{R}})$, to show that they have an infimum we must check that they constitute a directed set. But if Φ and Φ' are two graphings which generate \mathcal{R} , the set $A = \{x \in X : \text{C-Cost}(\Phi)(x) < \mu_{\mathcal{R}}(x)\}$ is \mathcal{R} -invariant, and one can define a new graphing Ψ made of the restrictions of the elements of Φ to A and the restrictions of the elements of Φ' to $X \setminus A$. Such a Ψ still generates \mathcal{R} , and we have

$$\text{C-Cost}(\Psi) = \inf(\text{C-Cost}(\Phi), \text{C-Cost}(\Phi')).$$

Using a maximality argument, one can then see that for every $M_{\mathcal{R}}$ -measurable everywhere positive function f , there exists a graphing Φ which generates \mathcal{R} such that its C-cost is less than $f + \text{C-Cost } \mathcal{R}$.

We now give a non ergodic version of lemma III.5 in [Gab00].

Lemma 2.6. *Let \mathcal{R} be an aperiodic pmp equivalence relation, let $\mathcal{R}_0 \subseteq \mathcal{R}$ be hyperfinite with the same algebra of invariant sets as \mathcal{R} (cf. proposition 2.4), and fix a graphing Ψ_0 of C-cost 1 which generates \mathcal{R}_0 . Let f be an $M_{\mathcal{R}}$ -measurable function, everywhere positive. Then there exists a graphing Φ whose C-cost is everywhere less than $\text{C-Cost}(\mathcal{R}) - 1 + f$, such that $\Phi_0 \cup \Phi$ generates \mathcal{R} .*

Proof. Begin with a graphing Φ which generates \mathcal{R} , and whose C-Cost is less than $f/2 + \text{C-Cost}(\mathcal{R})$. By proposition 2.2, we can find $A \subseteq X$ whose conditional measure is everywhere less than $f/2$. Now the induction procedure as described in [Gab00, lemme II.8] yields a treeing Φ_0 of \mathcal{R}_0 whose C-cost is $(1 - \mu_{\mathcal{R}}(A))$ and a graphing $\tilde{\Phi}$ of \mathcal{R} whose C-cost is $(\text{C-Cost}(\Phi) - (1 - \mu_{\mathcal{R}}(A)))$ such that Φ_0 and $\tilde{\Phi}$ generate \mathcal{R} . Then in particular $\Psi_0 \cup \tilde{\Phi}$ generates \mathcal{R} . \square

The relation between the C-cost and the cost of the ergodic components is very simple, as shown by the following proposition.

Proposition 2.7. *Let \mathcal{R} be a pmp equivalence relation. Then the conditionnal cost of \mathcal{R} is the function which associates to $x \in X$ the cost of \mathcal{R} for the \mathcal{R} -ergodic measure whose support contains x .*

Proof. This is a simple reformulation of the proof of proposition 18.4. in [KM04]. \square

3 The hyperfinite ergodic equivalence relation \mathcal{R}_0

Let $2^{\mathbb{N}} = \{0, 1\}^{\mathbb{N}}$, and for $n \in \mathbb{N}$, $2^n = \{0, 1\}^n$. Given $s \in 2^n$, we define the basic clopen set

$$N_s = \{x \in 2^{\mathbb{N}} : x_i = s_i \text{ for } 1 \leq i \leq n\}.$$

We can see elements $a \in 2^n$ and $b \in 2^{\mathbb{N}} \cup \bigcup_{n \in \mathbb{N}} 2^n$ as words in $\{0, 1\}$, and denote their concatenation by $a \frown b$. For $\epsilon \in \{0, 1\}$ and $n \in \mathbb{N}$, ϵ^n is the word $(x_i)_{i=1}^n \in 2^n$ defined by $x_i = \epsilon$.

3.1 Dyadic permutations

Let $n \in \mathbb{N}$. We view \mathfrak{S}_{2^n} as the group of permutations of the set 2^n . This defines a natural inclusion $\alpha_n : \mathfrak{S}_{2^n} \hookrightarrow \mathfrak{S}_{2^{n+1}}$ given by

$$\alpha_n(\sigma)(x_1, \dots, x_{n+1}) = (\sigma(x_1, \dots, x_n), x_{n+1})$$

for $\sigma \in \mathfrak{S}_{2^n}$ and $(x_1, \dots, x_{n+1}) \in 2^{n+1}$. Let \mathfrak{S}_{2^∞} be the inductive limit of these groups, called the group of **dyadic permutations**.

Let us now define a function $\sqrt{\cdot} : \mathfrak{S}_{2^p} \rightarrow \mathfrak{S}_{2^{p+1}}$ such that for any $\sigma \in \mathfrak{S}_{2^p}$, $\sigma = (\sqrt{\sigma})^2$ and $\sqrt{\sigma}$ has same support as σ . The idea is just to make $\sqrt{\sigma}$ “twice as slow”, as the following picture shows:

To be more precise, if $A \subseteq 2^p$ is the support of σ , define $\sqrt{\sigma}$ by

$$\sqrt{\sigma}(s \frown \epsilon) = \begin{cases} s \frown 1 & \text{if } s \in A \text{ and } \epsilon = 0 \\ \sigma(s) \frown 0 & \text{if } s \in A \text{ and } \epsilon = 1 \\ s \frown \epsilon & \text{if } s \notin A \end{cases} .$$

It is easily checked that $\sqrt{\sigma^2} = \sigma$, and that $\sqrt{\sigma}$ has the same support as σ .

This allows us to define functions $\sqrt[q]{\cdot} : \mathfrak{S}_{2^p} \rightarrow \mathfrak{S}_{2^{p+q}}$ inductively on $q \in \mathbb{N}$ by $\sqrt[q+1]{T} = \sqrt{\sqrt[q]{T}}$.




Figure 1: A transposition $\sigma \in \mathfrak{S}_2$ and its square root $\sqrt{\sigma} \in \mathfrak{S}_4$.

The key feature of \mathfrak{S}_{2^∞} is that it acts in a measure preserving way on $(2^\mathbb{N}, \lambda)$ by, for $\sigma \in \mathfrak{S}_{2^n}$, $s \in 2^n$ and $x \in 2^\mathbb{N}$,

$$\sigma(s \frown x) = \sigma(s) \frown x.$$

Let \mathcal{R}_0 be the measure preserving equivalence relation generated by this countable group. It is immediate that

$$(x_i)_{i \in \mathbb{N}} \mathcal{R}_0 (y_i)_{i \in \mathbb{N}} \iff \exists n \in \mathbb{N} \text{ such that } \forall i \geq n, x_i = y_i.$$

To avoid confusion, when we see \mathfrak{S}_{2^n} as a subgroup of $[\mathcal{R}_0]$, we will denote it by $\tilde{\mathfrak{S}}_{2^n}$. We still have square root functions $\sqrt[\cdot]{\cdot} : \tilde{\mathfrak{S}}_{2^p} \rightarrow \tilde{\mathfrak{S}}_{2^{p+q}}$.

The following proposition belongs to the folklore, for a proof see e.g. [Kec10, prop. 3.8].

Proposition 3.1. *The group of dyadic permutations is dense in the full group of \mathcal{R}_0 .*

3.2 The odometer T_0

The **odometer** is the map $T_0 \in \text{Aut}(2^\mathbb{N}, \lambda)$ defined by

$$(x_i)_{i \in \mathbb{N}} \in 2^\mathbb{N} \mapsto 1^n \frown (x_i)_{i > n},$$

where n is the first integer such that $x_n = 0$ (note that this is well defined on a set of full measure). This can be understood as adding $(1, 0, 0, \dots)$ to $(x_i)_{i \in \mathbb{N}}$ with right carry.

The map T_0 belongs to $\text{Aut}(2^\mathbb{N}, \lambda)$, an explicit inverse being given by

$$T_0^{-1} : (x_i) \mapsto 1^{n-1}0 \frown (x_i)_{i > n}$$

where n is the first integer such that $x_n = 1$. This inverse can of course be understood as subtraction with right borrow. Moreover, one can check that T_0 generates \mathcal{R}_0 .

Let $n \in \mathbb{N}$, then we define a “finite odometer” $\sigma_n \in \mathfrak{S}_{2^n}$ by

$$\sigma_n((s_i)_{i=1}^n) = \begin{cases} 0^n & \text{if } (s_i) = 1^n \\ 1^k \frown (s_i)_{i>k} & \text{else, where } k \text{ is the first integer such that } s_k = 0. \end{cases}$$

We denote by T_n the corresponding element in $\tilde{\mathfrak{S}}_{2^n}$. Note that by definition, T_n and T_0 coincide on $2^{\mathbb{N}} \setminus N_{1^n}$.

4 Two topological generators for the full group of $\mathcal{R}_0^Y = \Delta_Y \times \mathcal{R}_0$

In this section, our aim is to prove the following theorem, which gives two topological generators for non necessarily ergodic aperiodic hyperfinite equivalence relations, the second topological generator having an arbitrarily small support.

Theorem 4.1. *Let (Y, ν) be a standard probability space, possibly with atoms, and let $(X, \mu) = (Y \times 2^{\mathbb{N}}, \nu \otimes \lambda)$ where λ is the standard Bernoulli measure.. Let $(Y_k)_{k \in \mathbb{N}}$ be a partition of Y . Finally, let $\mathcal{R}_0^Y = \Delta_Y \times \mathcal{R}_0$ be the pmp equivalence relation on $(X, \mu) = (Y \times 2^{\mathbb{N}}, \nu \otimes \lambda)$ defined by*

$$(x, x') \mathcal{R} (y, y') \iff x = y \text{ and } x' \mathcal{R}_0 y'.$$

Then there exists $U \in [\mathcal{R}_0^Y]$ whose support is contained in $Z = \bigsqcup_{k \in \mathbb{N}} Y_k \times (N_{0^{k+1}} \cup N_{1^{k+1}})$ such that the following properties are satisfied for all $m \geq 0$:

- (I) *The full group of \mathcal{R} is topologically generated by $T = \text{id}_Y \times T_0$ and U^{2^m} , where T_0 is the odometer.*
- (II) *For every odd cycle C with disjoint support from U^{2^m} , the closed group generated by $(CU)^{2^m}$ contains U^{2^m}*

The following two sections are devoted to basic facts about the full group of \mathcal{R} , which will be used in the proof of theorem 4.1 that we give afterwards.

4.1 Some subgroups which topologically generate $[\mathcal{R}_0^Y]$

Let \mathcal{R}_0^Y be as in the statement of the previous theorem. By definition, the full group of \mathcal{R}_0^Y is isomorphic to the group of measurable maps from (Y, ν) to $[\mathcal{R}_0]$, denoted by $L^0(Y, \nu, [\mathcal{R}_0])$. This identification is isometric when we put on $L^0(Y, \nu, [\mathcal{R}_0])$ the L^1 metric d_1 defined by

$$\forall f, g : Y \rightarrow [\mathcal{R}_0], \quad d_1(f, g) = \int_Y d_u(f(y), g(y)) d\nu(y).$$

Now given a non-null subset A of Y , we can define an injective group morphism $\iota_A : [\mathcal{R}_0] \rightarrow [\mathcal{R}_0^Y]$ by letting $\iota_A(V)$ be the element of $L^0(Y, \nu, [\mathcal{R}_0])$ which sends every $y \in A$ to V , and every $y \notin A$ to $\text{id}_{2^{\mathbb{N}}}$. This morphism is bi-Lipschitz, and if A, B are

disjoint subsets of Y and $V, W \in [\mathcal{R}_0]$, the (commuting) product $i_A(V)i_B(W)$ can be understood as the map

$$y \in Y \mapsto \begin{cases} V & \text{if } y \in A, \\ W & \text{if } y \in B, \\ \text{id}_{2^{\mathbb{N}}} & \text{otherwise.} \end{cases} .$$

Since $[\mathcal{R}_0]$ is separable, the measurable maps from (Y, ν) to $[\mathcal{R}_0]$ with countable range are dense in $L^0(Y, \nu, [\mathcal{R}_0])$, which in turn yields that the measurable maps with finite range are dense in $L^0(Y, \nu, [\mathcal{R}_0])$. But every such map f can be decomposed as a product

$$f = \iota_{A_1}(V_1) \cdots \iota_{A_n}(V_n)$$

where $V_i \in [\mathcal{R}_0]$ and the A_i 's partition Y . Suppose now that \mathcal{B} is a dense subalgebra of $\text{MAlg}(Y, \nu)$, we may then find \tilde{f} arbitrarily close to f , where

$$\tilde{f} = \iota_{B_1}(V_1) \cdots \iota_{B_n}(V_n)$$

and the B_i 's belong to \mathcal{B} . We have proved the following lemma:

Lemma 4.2. *The full group of \mathcal{R}_0^Y is topologically generated by the set*

$$\bigcup_{B \in \mathcal{B}} \iota_B([\mathcal{R}_0]),$$

where \mathcal{B} is any dense subalgebra of $\text{MAlg}(Y, \nu)$.

4.2 Two generators for many finite subgroups

Let $n \geq 2$, and define $\tau_n \in \mathfrak{S}_{2^n}$ to be the transposition which exchanges $0^{n-1}1$ and $1^{n-1}0$. Let U_n be the corresponding element of $\tilde{\mathfrak{S}}_{2^n}$, that is, the group element implementing the action of τ_n on $2^{\mathbb{N}}$. Note that the support of U_n is $N_{0^{n-1}1} \cup N_{1^{n-1}0}$, so that the supports of the U_n 's are all disjoint.

Lemma 4.3. *For all $A \subseteq Y$ non null, $\iota_A(\tilde{\mathfrak{S}}_{2^n})$ is generated by $\iota_A(U_n)$ and T . Moreover, there exists $\kappa(n) \in \mathbb{N}$, independent of A , such that every element of $\iota_A(\tilde{\mathfrak{S}}_{2^n})$ can be written as a word in T and $\iota_A(U_n)$ of size less than $\kappa(n)$.*

Proof. Recall that $T_n \in \tilde{\mathfrak{S}}_{2^n}$ is obtained from the ‘‘finite odometer’’ $\sigma_n \in \mathfrak{S}_{2^n}$, which acts transitively on 2^n . And U_n corresponds to a transposition τ_n which permutes $1^{n-1}0$ and $0^{n-1}1 = \sigma_n(1^{n-1}0)$. For $i \in \{1, \dots, 2^n\}$, let $a_i = \sigma_n^{i-1}(0^n)$, and note that $a_{2^n} = 1^n$. For $i \in \{1, \dots, 2^n - 1\}$, let ρ_i be the transposition exchanging a_i and a_{i+1} , and let V_i be the corresponding element of $\tilde{\mathfrak{S}}_{2^n}$. Note that by construction the ρ_i 's generate \mathfrak{S}_{2^n} , so that the V_i 's generate $\tilde{\mathfrak{S}}_{2^n}$. Also, since $a_{2^{n-1}} = 1^{n-1}0$, we have $\rho_{2^{n-1}} = \tau_n$.

Then observe that, for $j \in \{-2^{n-1} + 1, \dots, 2^{n-1} - 1\}$, we have the conjugation relation

$$\sigma_n^j \rho_{2^{n-1}} \sigma_n^{-j} = \rho_{2^{n-1}+j}.$$

Going back to $[\mathcal{R}_0]$, this yields

$$T_n^j U_n T_n^{-j} = V_{2^{n-1+j}}.$$

But because T and $\iota_A(T_n)$ coincide on $A \times (2^{\mathbb{N}} \setminus N_{1^n})$, and $\iota_A(U_n), \iota_A(V_{2^{n-1+j}})$ act trivially outside of $A \times 2^{\mathbb{N}}$, the previous formula yields

$$T^j \iota_A(U_n) T^{-j} = \iota_A(V_{2^{n-1+j}}).$$

So the group generated by T and $\iota_A(U_n)$ contains all the $\iota_A(V_i)$ for $1 \leq i \leq 2^n - 1$. Since the V_i 's generate $\tilde{\mathfrak{S}}_{2^n}$, the group generated by T and $\iota_A(U_n)$ contains $\iota_A(\tilde{\mathfrak{S}}_{2^n})$. And the second part of the lemma follows from the finiteness of $\tilde{\mathfrak{S}}_{2^n}$ and the fact that the construction is independent from the subset A of Y we chose. \square

We now fix once and for all a function $n \mapsto \kappa(n)$ given by the previous lemma.

4.3 Two topological generators for $[\mathcal{R}_0^Y]$

We now begin the proof of theorem 4.1. Let $(X, \mu) = (Y \times 2^{\mathbb{N}}, \nu \otimes \lambda)$ and $\mathcal{R}_0^Y = \Delta_Y \times \mathcal{R}_0$ be the equivalence relation $(x, y) \mathcal{R}_0^Y (x', y')$ iff $x = x'$ and $y \mathcal{R}_0 y'$. Note that by Dye's theorem, every aperiodic singly generated equivalence relation is orbit equivalent to such an \mathcal{R}_0^Y .

We also define $T = \text{id}_Y \times T_0 \in [\mathcal{R}_0^Y]$, where T_0 is the odometer.

We first focus on property (I) in the case $m = 0$, and we will see that the constructed U also satisfies properties (I) and (II) for all $m \in \mathbb{N}$. So we fix a partition $(Y_k)_{k \in \mathbb{N}}$ of Y , and we want to find U such that its support is a subset of

$$Z = \bigsqcup_{k \in \mathbb{N}} Y_k \times (N_{0^{k+1}} \cup N_{1^{k+1}}),$$

and the set $\{T, U\}$ topologically generates $[\mathcal{R}_0^Y]$. Note that the conditional measure of Z on $L^2(Y, \nu)$ is an arbitrary small positive function. To be more precise, if $f \in L^2(Y, \nu)$ is any function such that $1 \geq f(y) > 0$ for all $y \in Y$, define $Y_k = \{y \in Y : \frac{1}{2^k} \leq f(y) < \frac{1}{2^{k-1}}\}$, and note that the conditional measure of $Z = \bigsqcup_{k \in \mathbb{N}} Y_k \times (N_{0^{k+1}} \cup N_{1^{k+1}})$ is smaller than f .

Let \mathcal{A}_n be an increasing family of finite subalgebras of $\text{MAlg}(Y, \nu)$ with a dense reunion. Define $\mathcal{B}_n = \{A \cap (\bigcup_{k \leq n} Y_k) : A \in \mathcal{A}_n\}$ and let $\mathcal{B} = \bigcup_{n \in \mathbb{N}} \mathcal{B}_n$. Note that because $\nu(\bigcup_{k \leq n} Y_k) \rightarrow 1$ as $n \rightarrow +\infty$, the countable set \mathcal{B} is dense in $\text{MAlg}(Y, \nu)$, and we have the following additional property: for every $B \in \mathcal{B}$, there exists $K \in \mathbb{N}$ such that $B \times (N_{0^K} \cup N_{1^K}) \subseteq Z$.

First recall that for all $B \in \mathcal{B}$, there exists $K \in \mathbb{N}$ such that $B \times (N_{0^K} \cup N_{1^K}) \subseteq Z$. Since the support of $\iota_B(U_n)$ is $B \times N_{0^{n-1}} \cup N_{1^{n-1}}$, which is a subset of $B \times (N_{0^K} \cup N_{1^K})$ for all $n \geq K - 1$, we have the following lemma.

Lemma 4.4. *For every $B \in \mathcal{B}$ and n large enough, the support of $\iota_B(U_n)$ is a subset of Z .*

We now build an element U of the full group of \mathcal{R} such that the support of U is a subset of Z and $[\mathcal{R}_0^Y]$ is topologically generated by T and U .

Sketch of the construction. Let us briefly explain the idea behind the construction of U . Observe that for all $B, B' \in \mathcal{B}$ and $n \neq m \in \mathbb{N}$, $\iota_B(U_n)$ and $\iota_{B'}(U_m)$ have disjoint support (hence they commute). We fix an enumeration $(B_k)_{k \in \mathbb{N}}$ of \mathcal{B} such that each element of \mathcal{B} appears infinitely many times. Now start with $U = \iota_{B_1}(U_{n_1})$, where n_1 is such that the support of $\iota_{B_1}(U_{n_1})$ is a subset of Z . Then by lemma 4.3, the group generated by T and U contains $\iota_{B_1}(\tilde{\mathfrak{S}}_{2^{n_1}})$.

We now also want the group generated by T and U to contain $\iota_{B_2}(\tilde{\mathfrak{S}}_{2^{n_2}})$ for some n_2 such that the support of $\iota_{B_1}(U_{n_2})$ is a subset of Z . If we put $U = \iota_{B_1}(U_{n_1})\iota_{B_2}(\sqrt{U_{n_2}})$, observe that since U_{n_1} is an involution, $U^2 = \iota_{B_2}(U_{n_2})$ so that the group generated by T and U contains $\iota_{B_2}(\tilde{\mathfrak{S}}_{2^{n_2}})$. The problem is that now, the group generated by T and U does not contain $\iota_{B_1}(\tilde{\mathfrak{S}}_{2^{n_1}})$ anymore. But if we choose n_2 big enough, the support of $\iota_{B_2}(\sqrt{U_{n_2}})$ will be very small, so we will still have $\iota_{B_1}(\tilde{\mathfrak{S}}_{2^{n_1}})$ in the group generated by T and U , up to a small error. If we keep doing that, the group generated by T and U will contain up to an error converging to 0 every $\iota_{B_k}(\tilde{\mathfrak{S}}_{2^{n_k}})$, so by proposition 3.1 it will actually contain every $\iota_B([\mathcal{R}_0])$. Then lemma 4.2 yields that the group topologically generated by T and U contains the full group of \mathcal{R}_0^Y . \square

We now begin the actual construction, reverse-engineering the sketch above. Thanks to lemma 4.2, it is enough to find $U \in [\mathcal{R}_0^Y]$ such that the support of U is a subset of Z , and for every $B \in \mathcal{B}$, $\iota_B([\mathcal{R}_0])$ is contained in the closed group generated by T and U .

Fix a sequence of positive reals $(\epsilon_k)_{k \in \mathbb{N}}$ decreasing to zero. Since ι_B is bi-Lipschitz, by proposition 4.2 it suffices to find $U \in [\mathcal{R}_0]$ whose support is a subset of Z and an increasing sequence of integers $(n_k)_{k \in \mathbb{N}}$ such that

- (a) For all $B \in \mathcal{B}$ and infinitely many $k \in \mathbb{N}$, every element of $\iota_B(\tilde{\mathfrak{S}}_{2^{n_k}})$ is in the ϵ_k -neighborhood of $\langle T, U \rangle$.

We now fix an enumeration $(B_k)_{k \in \mathbb{N}}$ of \mathcal{B} such that each element of \mathcal{B} appears infinitely many times. This allows us to replace condition (a) by condition

- (a') For all $k \in \mathbb{N}$, every element of $\iota_{B_k}(\tilde{\mathfrak{S}}_{2^{n_k}})$ is in the ϵ_k -neighborhood of $\langle T, U \rangle$.

Finally, the definition of $\kappa(n)$ (cf. the end of section 4.2), the fact that d_u is biinvariant and lemma 4.3 yield that we can replace (a') by the weaker condition

- (a'') For all $k \in \mathbb{N}$, the element $\iota_{B_k}(U_{n_k})$ is in the $\frac{\epsilon_k}{\kappa(n_k)}$ -neighborhood of the group generated by U .

First choose $n_1 \in \mathbb{N}$ such that the support of $\iota_{B_1}(U_{n_1})$ is a subset of Z (such an n_1 exists by lemma 4.4). Then, given n_k , find $n_{k+1} > n_k$ such that

- (i) the support of $\iota_{B_{k+1}}(U_{n_{k+1}})$ is a subset of Z and

(ii)
$$\frac{1}{2^{n_{k+1}-2}} < \frac{\epsilon_k}{\kappa(p_k)}.$$

We are ready to define U as an infinite product of commuting elements:

$$U = \prod_{l \in \mathbb{N}^*} \iota_{B_l} ({}^{2^{l-1}}\sqrt{U_{n_l}}).$$

Let us check that condition (a'') is satisfied. If we fix $k \in \mathbb{N}^*$, we have

$$\begin{aligned} U^{2^k} &= \prod_{l \in \mathbb{N}^*} \iota_{B_l} ({}^{2^{l-1}}\sqrt{U_{n_l}}^{2^{k-1}}) \\ &= \prod_{l=1}^{k-1} \iota_{B_l} (U_{n_l}^{2^{k-l}}) \cdot U_{n_k} \cdot \prod_{l=k+1}^{+\infty} \iota_{B_l} ({}^{2^{l-k}}\sqrt{U_{n_l}}) \end{aligned}$$

Because the U_{n_l} 's are involution, the first product is equal to the identity, so that

$$U^{2^{k-1}} = \iota_{B_k} (U_{n_k}) \cdot \prod_{l=k+1}^{+\infty} \iota_{B_l} ({}^{2^{l-k}}\sqrt{U_{n_l}}). \quad (1)$$

We now check that the error term $W_k = \prod_{l=k+1}^{+\infty} \iota_{B_l} ({}^{2^{l-k}}\sqrt{U_{n_l}})$ is small. Because for every $l \in \mathbb{N}$, ${}^{2^{l-k}}\sqrt{U_{n_l}}$ has same support as U_{n_l} , the support of W_k has measure smaller than

$$\nu(B_l) \sum_{l=k+1}^{+\infty} \lambda(\text{supp } U_{n_l}) \leq \sum_{l=k+1}^{+\infty} \frac{1}{2^{n_l-1}} \quad (2)$$

Since $(n_l)_{l \in \mathbb{N}}$ is increasing, we have for all $l \geq k+1$,

$$\frac{1}{2^{n_l-1}} \leq \frac{1}{2^{n_{k+1}+(l-k-2)}}$$

We can now bound from above the sum in (2) and get the inequality

$$\begin{aligned} \mu(\text{supp } W_k) &\leq \frac{1}{2^{n_{k+1}}} \cdot \sum_{l=k+1}^{+\infty} \frac{1}{2^{l-k-2}} \\ &\leq \frac{4}{2^{n_{k+1}}} \\ &\leq \frac{\epsilon_k}{\kappa(n_k)} \end{aligned}$$

because condition (ii) states that $\frac{1}{2^{n_{k+1}-2}} < \frac{\epsilon_k}{\kappa(p_k)}$. In the end, because d_u is bi-invariant, formula (1) yields

$$\begin{aligned} d_u(U^{2^{k-1}}, \iota_{B_k}(U_{n_k})) &\leq \mu(\text{supp}(W_k)) \\ &\leq \frac{\epsilon_k}{\kappa(n_k)}. \end{aligned}$$

Condition (a'') is thus satisfied, so that T and U topologically generate the full group of \mathcal{R}_0^Y . In other words, property (I) holds for $m = 0$. Let now $m \geq 0$, and let

$U' = U^{2^m}$. By the definition of the square root function, and the fact that the U_n 's are involutions, we get

$$\begin{aligned} U' &= \prod_{l=m+1}^{+\infty} \iota_{B_l} ({}^{2^{l-m-1}}\sqrt{U_{n_l}}) \\ &= \prod_{l \in \mathbb{N}^*} \iota_{B_{m+l}} ({}^{2^{l-1}}\sqrt{U_{n_{l+m}}}). \end{aligned}$$

So if we put $B'_l = B_{m+l}$, the B'_l 's still enumerate \mathcal{B} in a way that every element of \mathcal{B} appears infinitely many times, and we can apply the previous proof to see that U' satisfies condition

(b'') For all $k \in \mathbb{N}^*$, the element $\iota_{B'_k}(U_{n_{k+m}})$ is in the $\frac{\epsilon_{k+m}}{\kappa(n_{k+m})}$ -neighborhood of the group generated by U' .

Using the same argument as for condition (a'') on U , one can see that this implies that T and U' topologically generate the full group of \mathcal{R}_0^Y . So for all $m \geq 0$, we have that T and U^{2^m} topologically generate the full group of \mathcal{R}_0^Y , in other words condition (I) is satisfied.

Let us now check that condition (II) is also satisfied for all $m \geq 0$, that is, let us show that if C is an odd cycle (an element whose orbits are all finite and have odd cardinality) with disjoint support from the support of U^{2^m} , then U^{2^m} is a cluster point of the sequence $((CU^{2^m})^k)_{k \in \mathbb{N}}$. For this, we just need the fact that $V = U^{2^m}$ only has orbits of cardinality 2^l for $l \in \mathbb{N}$.

Let the integers $(p_l)_{l \in \mathbb{N}}$ enumerate the cardinalities of the orbits of C , and let $P_k = \prod_{l \leq k} p_l$. We have $C^{P_k} \rightarrow \text{id}_X$ when $k \rightarrow +\infty$. For $k \in \mathbb{N}$, let X_k be the set of elements whose V -orbit has cardinality at most 2^k . Then, because all the orbits of V are finite, we have $\mu(X_k) \rightarrow 1$ as $k \rightarrow +\infty$. So, if we let

$$V_k(x) = \begin{cases} V(x) & \text{if } x \in X_k \\ x & \text{else} \end{cases},$$

we can write $V = V_k W_k$, where V_k and W_k have disjoint support and $W_k \rightarrow \text{id}_X$ as $k \rightarrow +\infty$.

Since P_k is odd we find $m_k \in \mathbb{N}$ such that $m_k P_k \equiv 1 \pmod{2^k}$. Observe that then, $m_k P_k \equiv 1 \pmod{2^l}$ for all $l \leq k$. This implies that $V_k^{m_k P_k} = V_k$, so that

$$\begin{aligned} d_u(V^{m_k P_k}, V_k) &= d_u(V_k^{m_k P_k} W_k^{m_k P_k}, V_k) \\ &= d_u(V_k W_k^{m_k P_k}, V_k) \\ &\leq \mu(\text{supp } W_k) \rightarrow 0 \quad [k \rightarrow +\infty]. \end{aligned}$$

In the end, because V and C have disjoint support,

$$(CV)^{m_k P_k} = (C^{P_k})^{m_k} V^{m_k P_k} \rightarrow V \quad [k \rightarrow +\infty],$$

so that condition (2) is satisfied and theorem 4.1 is proved.

5 Consequences

5.1 Computation of the topological rank for full groups

We now begin the proof of theorem 1.4. Let \mathcal{R} be a pmp aperiodic equivalence relation. We first prove the inequality

$$t([\mathcal{R}]) \geq \sup_{x \in X} \lfloor \text{C-Cost}(\mathcal{R})(x) \rfloor + 1.$$

Let A be a positive \mathcal{R} -invariant set and $n \in \mathbb{N}$ such that $\lfloor \text{C-Cost}(\mathcal{R})(x) \rfloor \geq n$ for all $x \in A$. Then the restriction to A yields a continuous morphism $[\mathcal{R}] \rightarrow [\mathcal{R}|_A]$. The normalised cost of $\mathcal{R}|_A$ is at least n , so that by an argument of Miller (cf. [LM14, proof of theorem 1]), $t([\mathcal{R}|_A]) \geq n + 1$. Because the restriction is a continuous morphism, we get $t([\mathcal{R}]) \geq n + 1$. By definition of the essential supremum, the first inequality is proved.

We now prove the reverse inequality. Let us first put ourselves in the situation given by proposition 2.4, so $X = Y_{\mathcal{R}} \times 2^{\mathbb{N}}$, $\mu = \nu_{\mathcal{R}} \otimes \lambda$, and \mathcal{R} contains the hyperfinite equivalence relation $\mathcal{R}_0 = \Delta_{Y_{\mathcal{R}}} \times \mathcal{R}_0$. The \mathcal{R} -conditional expectation is then the projection onto $L^2(Y_{\mathcal{R}}, \nu_{\mathcal{R}})$.

Let n be the essential supremum of the function $y \mapsto \lfloor \text{C-Cost}(\mathcal{R})(y) \rfloor$. We must find $n+1$ topological generators for $[\mathcal{R}]$. By definition, the C-cost of \mathcal{R} is everywhere less than $n + 1$.

Fix a graphing Φ_0 which generates \mathcal{R}_0 , and whose C-cost is 1. By lemma 2.6, we can find a graphing Φ of \mathcal{R} such that $\Phi_0 \cup \Phi$ generates \mathcal{R} and the C-Cost of Φ is everywhere lesser than n . Now define $f(y) = \frac{\text{C-Cost}(\Phi)(y)}{n} < 1$, and write $\Phi = \Phi_1 \cup \dots \cup \Phi_n$, where each Φ_i has C-cost f . We may find an odd valued function $q(y)$ such that for all $y \in Y$,

$$\frac{q(y) + 2}{q(y)} f(y) < 1.$$

Theorem 4.1 then provides elements T and U of the full group of \mathcal{R}_0 such that the support of U has \mathcal{R} conditional measure less than $1 - \frac{q(y)+2}{q(y)} f(y)$, and the following properties are satisfied:

- (1) The full group of \mathcal{R} is topologically generated by $T = \text{id}_Y \times T_0$ and U .
- (2) For every odd cycle C with disjoint support from U , the closed group generated by UC contains U .

We now let $Y_p = \{y \in Y : q(y) = p\}$ for $p \in \mathbb{N}$. Note that when p is even, Y_p is empty, and that the Y_p 's partition Y . Then by lemma 2.2, for every $p \in \mathbb{N}$ we may find disjoint subsets A_p^1, \dots, A_p^{p+2} of $(Y_p \times 2^{\mathbb{N}}) \setminus \text{supp } U$, each of the same \mathcal{R} conditional measure $\frac{f(y)}{p}$.

Fix $p \in \mathbb{N}$ as well as $i \in \{1, \dots, n\}$, and consider the restriction of Φ_i to $Y_p \times 2^{\mathbb{N}}$. Using proposition 2.3, we may assume, after cutting/gluing elements of Φ_i and pre/post-composing by partial isomorphisms of \mathcal{R}_0 , that the restriction of Φ_i to $Y_p \times 2^{\mathbb{N}}$ is a pre- $(p+1)$ -cycle.

Now for all $p \in \mathbb{N}$, fix $\psi_p \in [[\mathcal{R}_0]]$ whose domain is A_p^{p+1} and whose range is A_p^{p+2} . Put $\tilde{\Phi}_i = \Phi_i \cup \{\psi_p : p \in \mathbb{N}\}$.

Then, \mathcal{R} is still generated by $\Phi_0 \cup \tilde{\Phi}_1 \cup \dots \cup \tilde{\Phi}_n$, and every $\tilde{\Phi}_i$ is a pre- $(p+2)$ -cycle when restricted to $Y_p \times 2^{\mathbb{N}}$. Let C_i^p be the corresponding $(p+2)$ cycles, and let C_i be the odd cycle obtained by gluing together all the C_i^p . Finally, let $U_1 = UC_1$.

Claim. *The $(n+1)$ elements T, U_1, C_2, \dots, C_n topologically generate the full group of \mathcal{R} .*

Proof of the claim. Let G be the closed group generated by T, U_1, C_2, \dots, C_n . First, thanks to condition (2) and the fact that $U_1 = UC_1$ belongs to G , we have that U belongs to G . So by condition (1), G contains $[\mathcal{R}_0]$. It follows that G contains $U^{-1}U_1 = C_1$. Since every $\mathcal{R}_{\tilde{\Phi}_i}$ is generated by its restrictions to every $Y_p \times 2^{\mathbb{N}}$, which contain $\psi_p \in [[\mathcal{R}_0]]$, the arguments in [LM14, proposition 9] and the fact that \mathcal{R} is the join of $\mathcal{R}_0, \mathcal{R}_{\tilde{\Phi}_1}, \dots, \mathcal{R}_{\tilde{\Phi}_n}$ yield the conclusion. \square

5.2 A dense G_δ set of topological generators

Let $APER$ denote the set of elements of $\text{Aut}(X, \mu)$ whose orbits are all infinite. In this section we show the following proposition, which will be improved at the end of the next section (cf. theorem 6.3).

Proposition 5.1. *Let \mathcal{R} be any aperiodic pmp equivalence relation with $\text{Cost}(\mathcal{R}) = 1$. Then the set*

$$\{(T, U) \in (\text{APER} \cap [\mathcal{R}]) \times [\mathcal{R}] : \overline{\langle T, U \rangle} = [\mathcal{R}]\}$$

is a dense G_δ in $(\text{APER} \cap [\mathcal{R}]) \times [\mathcal{R}]$.

Proof. It is a standard fact that the set we are interested in is a G_δ , so we only have to show its density. Because of Rohlin's lemma, any aperiodic element of $[\mathcal{R}]$ has a dense conjugacy class in $[\mathcal{R}]$, so that we actually only have to find one aperiodic $T \in [\mathcal{R}]$ such that the set

$$\{U \in [\mathcal{R}] : \overline{\langle T, U \rangle} = [\mathcal{R}]\}$$

is dense in $[\mathcal{R}]$.

By proposition 2.4, we can suppose $X = Y \times 2^{\mathbb{N}}$ equipped with a product measure $\mu = \lambda \otimes \mathcal{B}(1/2)$, where (Y, λ) is a standard probability space (possibly with atoms) and $\mathcal{B}(1/2)$ is the standard Bernoulli measure. Moreover, we may assume that \mathcal{R} contains $\mathcal{R}_0 = \text{id}_Y \times \mathcal{R}_0$, which is generated by $T = \text{id}_Y \times T_0$. We will show that the set $\{U \in [\mathcal{R}] : \overline{\langle T, U \rangle} = [\mathcal{R}]\}$ is dense.

Recall that the set of cycles is dense in the full group of \mathcal{R} . One can then show that the set of cycles whose support does not intersect $Y \times (N_{0^{k+1}} \cup N_{1^{k+1}})$ for some k , and whose orbits have bounded cardinality is dense in $[\mathcal{R}]$.

Fix such a cycle C and $\epsilon > 0$, write the order of C as a product $2^K N$, where N is odd. Also fix $k \in \mathbb{N}$ such that $Y \times (N_{0^{k+1}} \cup N_{1^{k+1}})$ is disjoint from the support of C , and $\mu(Y \times (N_{0^{k+1}} \cup N_{1^{k+1}})) < \epsilon$, and find an odd number M coprime with N .

Theorem 4.1 provides us $U \in [\mathcal{R}]$ whose support is a subset of $Z = Y \times (N_{0^{k+2}} \cup N_{1^{k+2}})$, such that the full group of \mathcal{R} is topologically generated by T and U^{2^K} .

Now note that because \mathcal{R} is aperiodic, its conditional cost is greater or equal than one. Since \mathcal{R} has cost one, its conditional cost must then be constant equal to one. Using the same ideas as in the proof of theorem 1.4, we may find a cycle C' of order M supported on $Y \times (N_{0^{k+1}})$ (which is disjoint from the supports of U and C), such that each C' orbit contains two consecutive \mathcal{R}_0 related elements, and \mathcal{R} is generated by T_0 and C' . We define $U' = UCC'$, note that U' is ϵ -close to C so that the proof boils down to the following claim.

Claim. T and U' topologically generate $[\mathcal{R}]$.

Indeed, let G the the closed subgroup generated by T and U' . Since CC' and U have disjoint support and CC' is an odd cycle, condition (II) in theorem 4.1 ensures that U^{2^K} belongs to the group topologically generated by $U'^{2^K} = (CC'U)^{2^K}$, so G contains $[\mathcal{R}_0]$. Then G must contain $CC' = U'U^{-1}$, but by the chinese remainder theorem, because C and C' commute and their orders are coprime with each other, G contains C' . Now because each C' orbit contains two \mathcal{R}_0 related elements, the same proof as for theorem 4.1 yields that G contains $[\mathcal{R}_{C'}]$, hence G contains $[\mathcal{R}]$. \square

6 Free subgroups in full groups

This section is devoted to a generalization of the following theorem of Kechris, which we then use to find dense free subgroups in some full groups. Note that the set of tuples generating a free group is always a G_δ in a Polish group, so that in the next theorems, the key feature is the density of tuples generating a dense subgroup.

Theorem 6.1 ([Kec10, Theorem 3.9]). *Let \mathcal{R} be a pmp ergodic equivalence relation and $n \in \mathbb{N}$. Then the set of n -tuples of aperiodic elements of $[\mathcal{R}]$ generating a free subgroup is a dense G_δ in the set of n -tuples of aperiodic elements of $[\mathcal{R}]$.*

By the Kuratowski-Ulam theorem, for $n = 2$ the above statement is equivalent to having a dense G_δ of aperiodic elements T for which there is a dense G_δ of aperiodic elements U such that the subgroup generated by T and U is free. Our result yields a dense G_δ above *any* $T \in [\mathcal{R}]$ of infinite order.

Theorem 6.2. *Let \mathcal{R} be a pmp equivalence relation, and let $T \in [\mathcal{R}]$ have infinite order. Then for all $n \in \mathbb{N}$ we have the following:*

(1) *The set of n -uples (U_1, \dots, U_n) of elements of $[\mathcal{R}]$ such that*

$$\langle T, U_1, \dots, U_n \rangle \simeq \mathbb{F}_{n+1}$$

is a dense G_δ in $[\mathcal{R}]^n$.

(2) *If furthermore \mathcal{R} is aperiodic, then the set of n -uples of aperiodic elements (U_1, \dots, U_n) of $[\mathcal{R}]$ such that*

$$\langle T, U_1, \dots, U_n \rangle \simeq \mathbb{F}_{n+1}$$

is a dense G_δ in $([\mathcal{R}] \cap \text{APER})^n$.

Proof. For the sake of notational simplicity, we only give a detailed proof for $n = 1$. Whenever w is a reduced word in two letters t and u , we denote by $w(\varphi, \psi)$ the evaluation of this word on elements φ, ψ belonging to the pseudo-full group of \mathcal{R} . Note that $w(\varphi, \psi)$ may be nowhere defined, but that whenever φ and ψ belong to the full group of $[\mathcal{R}]$, $w(\varphi, \psi)$ also does.

Let $T \in \mathcal{R}$ have infinite order, which is equivalent to asking that T has unbounded orbits. We have to show that for every non empty reduced word w in t and u , the set of $U \in [\mathcal{R}]$ (respectively in $U \in \text{APER} \cap [\mathcal{R}]$) such that $w(T, U) \neq 1$ is dense in $[\mathcal{R}]$ (respectively in $\text{APER} \cap [\mathcal{R}]$). Indeed, every one of these sets is open, and their intersection is the set of U 's such that T and U generate a free group.

So we fix a reduced word w , $\delta > 0$ and $U \in [\mathcal{R}]$ (respectively $U \in \text{APER} \cap [\mathcal{R}]$). We want to find $U' \in [\mathcal{R}]$ (respectively $U' \in \text{APER} \cap [\mathcal{R}]$) such that $d_u(U, U') < \delta$ and $w(T, U') \neq \text{id}_X$. Because T has infinite order, we may restrict ourselves to the case when w contains at least one occurrence of u . Up to conjugating w , we may then suppose that w ends with the letter u or u^{-1} . Let n be the length of w .

Because T has unbounded orbits, there is $A \subseteq X$ such that $(T^j(A))_{j=0}^{2(n+1)^2}$ is a disjoint family of subsets of X . The map which associates to every $B \in \text{MAlg}(X, \mu)$ the first return map U_B induced by U on B is continuous. So, up to shrinking A , we may furthermore assume that

$$d_u(U, U_{X \setminus \bigsqcup_{j=0}^{2(n+1)^2} T^j(A)}) < \frac{\delta}{2},$$

and also that $\mu(\bigsqcup_{j=0}^{2(n+1)^2} T^j(A)) < \frac{\delta}{2}$.

For all $i \in \{0, \dots, n\}$ and $j \in \{-n, \dots, n\}$, let $A_{i,j} = T^{j+n+2(n+1)i}(A)$. We think of the $A_{i,j}$'s as squares on the plane whose center has coordinates (i, j) . In this picture, T mostly acts by vertical translation.

Write $w = w_n \cdots w_1$, with $w_k \in \{t, t^{-1}, u, u^{-1}\}$. Let us define by recurrence on $k \in \{0, \dots, n\}$ a sequence of pairs of integers $(i_k, j_k)_{k=0}^n$ by putting $(i_0, j_0) = (0, 0)$, and then

$$(i_{k+1}, j_{k+1}) = (i_k, j_k) + \begin{cases} (0, 1) & \text{if } w_k = t \\ (0, -1) & \text{if } w_k = t^{-1} \\ (1, 0) & \text{if } w_k = u^{\pm 1} \end{cases}.$$

Note that the sequence $(i_k, j_k)_{k=0}^n$ is injective because w is a reduced word. We now build up an element φ of the pseudo full group of \mathcal{R} such that $w(T, \varphi)$ "follows the same path" as the sequence (i_k, j_k) (cf. figure 2).

Let K^+ be the set of $k \in \{1, \dots, n\}$ such that $w_k = t$, and K^- the set of $k \in \{1, \dots, n\}$ such that $w_k = t^{-1}$. For all $i, j \in \{1, \dots, n\} \times \{-n, \dots, n\}$, we fix a partial isomorphism $\psi_{i,j} \in [[\mathcal{R}]]$ whose domain is $A_{i-1,j}$ and whose image is $A_{i,j}$ (for instance, one can choose $\psi_{i,j} = T_{|A_{i-1,j}}^{2n+1}$).

The partial isomorphism $\psi \in [[\mathcal{R}]]$ is defined by gluing together some of the $\psi_{i,j}$:

$$\psi = \bigsqcup_{k \in K^+} \psi_{i_k, j_k} \sqcup \bigsqcup_{k \in K^-} \psi_{i_k, j_k}^{-1}.$$

Note that ψ is well defined because an element of K^+ is never followed by an element of K^- and vice-versa, since w is a reduced word. By construction, for all




Figure 2: The construction of ψ for $w = t^2u^{-2}tut^{-2}u^{-1}$

$k \in \{1, \dots, n\}$, $w_k \cdots w_1(T, \varphi)$ has domain A_{i_0, j_0} and range A_{i_k, j_k} . In particular, the range of $w_n \cdots w_1(T, \varphi) = w(T, \psi)$ is disjoint from its domain.

We now seek $U' \in [\mathcal{R}]$ (respectively $U' \in \text{APER} \cap [\mathcal{R}]$) near to U and which extends φ . First recall that we have

$$d_u(U, U_{X \setminus \bigsqcup_{j=0}^{2(n+1)^2} T^j(A)}) < \frac{\delta}{2},$$

and that the domain as well as the range of ψ are subsets of $\bigsqcup_{j=0}^{2(n+1)^2} T^j(A)$.

We now consider the two cases of the theorem separately. In the first case, we extend ψ to an element \tilde{U} of the full group of the restriction of \mathcal{R} to $\bigsqcup_{j=0}^{2(n+1)^2} T^j(A)$. In the second case, because φ is obtained by gluing together pre- p -cycles with disjoint supports, the aperiodicity of \mathcal{R} enables us to extend it to an aperiodic element \tilde{U} of the full group of the restriction of \mathcal{R} to $\bigsqcup_{j=0}^{2(n+1)^2} T^j(A)$.

We may then define

$$U'(x) = \begin{cases} \tilde{U}(x) & \text{if } x \in \bigsqcup_{j=0}^{2(n+1)^2} T^j(A) \\ U_{X \setminus \bigsqcup_{j=-n}^n T^j(A)}(x) & \text{else.} \end{cases}$$

We clearly have $d_u(U, U') < \delta$, and since U' extends φ , we have $w(T, U') \neq \text{id}_X$. Finally, U' belongs to $[\mathcal{R}]$ (respectively to $\text{APER} \cap [\mathcal{R}]$). \square

We now apply this theorem to give a characterization of aperiodic pmp equivalence relations of cost one, using the results of the previous section.

Theorem 6.3. *Let \mathcal{R} be any pmp equivalence relation, and consider the set*

$$G = \{(T, U) \in (\text{APER} \cap [\mathcal{R}]) \times [\mathcal{R}] : \overline{\langle T, U \rangle} = [\mathcal{R}] \text{ and } \langle T, U \rangle \simeq \mathbb{F}_2\}.$$

Then the following assertions are equivalent.

- (1) \mathcal{R} is aperiodic of cost one.
- (2) G is a dense G_δ in $(\text{APER} \cap [\mathcal{R}]) \times [\mathcal{R}]$.

Proof. Suppose first that \mathcal{R} is a aperiodic. Then theorem 6.2 yields that the set

$$G_1 = \{(T, U) \in (\text{APER} \cap [\mathcal{R}]) \times [\mathcal{R}] : \langle T, U \rangle \simeq \mathbb{F}_2\}$$

is a dense G_δ , while proposition 5.1 asserts that the set

$$G_2 = \{(T, U) \in (\text{APER} \cap [\mathcal{R}]) \times [\mathcal{R}] : \overline{\langle T, U \rangle} = [\mathcal{R}]\}$$

is a dense G_δ . So $G = G_1 \cap G_2$ is a dense G_δ .

Conversely, if \mathcal{R} is not aperiodic, it is not topologically finitely generated, hence G_2 is empty, so in particular G is empty. And if \mathcal{R} is aperiodic but not of cost one, its cost has to be strictly greater than one, so suppose by contradiction that G_2 were dense in $(\text{APER} \cap [\mathcal{R}]) \times [\mathcal{R}]$. Its vertical section would contain U 's of arbitrarily small support, which along with any $T \in [\mathcal{R}]$ would then fail to generate the equivalence relation \mathcal{R} by the definition of the cost, so that they could not topologically generate the full group of \mathcal{R} , a contradiction. \square

7 A finer metric on full groups

The notion of \mathcal{R} -conditional measure induces a new metric d_C on the full group of \mathcal{R} , defined by

$$d_C(S, T) = \|\mu_{\mathcal{R}}(\text{supp } S^{-1}T)\|_\infty.$$

In this section, we study this bi-invariant complete metric, which is non separable as soon as \mathcal{R} has infinitely many ergodic components, and show that $([\mathcal{R}], d_C)$ satisfies the automatic continuity property as soon as \mathcal{R} is aperiodic.

First, remark that if \mathcal{R} has only finitely many ergodic components, this metric induces the uniform topology on $[\mathcal{R}]$, so that our automatic continuity result is a generalization of [KT10, theorem 3.1]. Note however that our proof follows the exact same outline as theirs.

7.1 Automatic continuity for $([\mathcal{R}], d_C)$

The following notion, introduced by Rosendal and Solecki [RS07], will be key in order to prove the automatic continuity property for full groups.

Definition 7.1. Let $n \in \mathbb{N}$. A topological group G is said to be n -**Steinhaus** if for every subset A of G containing the identity such that $A = A^{-1}$ and such that G is covered by countably many left translates of A , the set A^n of products of at most n elements of A contains a neighborhood of the identity.

For a survey on automatic continuity, including in particular a proof of the next proposition, see [Ros09].

Proposition 7.2 (Rosendal-Solecki). *Suppose a topological group G is n -Steinhaus for some fixed $n \in \mathbb{N}$. Then every homomorphism from G into a separable group is continuous.*

We may now state the main theorem of this section.

Theorem 7.3. *Let \mathcal{R} be a pmp aperiodic equivalence relation. Then $([\mathcal{R}], d_C)$ is 38-Steinhaus. In particular, every homomorphism from $([\mathcal{R}], d_C)$ into a separable group is continuous.*

Proof. Take $(B_n)_{n \in \mathbb{N}}$ which partition X , each of them having constant positive conditional measure. For any $B \subseteq X$, let $[\mathcal{R}]_B = \{T \in [\mathcal{R}] : \text{supp } T \subseteq B\}$. The proof of the first step goes verbatim as in [KT10], so we omit it.

Step 1. There exists $n \in \mathbb{N}$ such that

$$\forall T \in [\mathcal{R}]_{B_n} \exists S \in W^2, S_{\upharpoonright B_n} = T_{\upharpoonright B_n}$$

We fix n as in the previous lemma, and let $B = B_n$.

Step 2. W^2 contains an involution S whose support is contained in B , and has constant conditional expectation lesser than $\mu(B)/2$.

Fix disjoint $A_1, A_2 \subseteq B$ of constant conditional measure $\mu(B)/4$, and fix an involution T supported in $A_1 \cup A_2$, exchanging them. Now one can find an increasing family $(A_t)_{t \in (0, \mu(B)/4) \rightarrow}$ of subsets of A_1 where each A_t has constant conditional measure equal to t (for instance, lemma 2.2 provides such a family for dyadic t , and one can then use the completeness of the measure algebra to extend this uniquely for $t \in (0, \mu(B)/4)$).

Let $T_t = T_{\upharpoonright A_t \cup T(A_t)} \sqcup \text{id}_{X \setminus (A_t \cup T(A_t))}$, note that these involutions all belong to the commutative group $[\mathcal{R}_T]$, and as there are uncountably many such T_t 's, there is m such that $g_m W$ contains two distinct elements T_{t_1}, T_{t_2} with $t_1 < t_2$. Now $S = T_{t_1} T_{t_2} = T_{t_1}^{-1} T_{t_2}$ belongs to $W g_m^{-1} g_m W = W^2$ and has support $A_{t_2} \setminus A_{t_1} \sqcup T(A_{t_2} \setminus A_{t_1})$, which has constant conditional expectation lesser than $\mu(B)/2$.

We fix such an S , and a subset C of B such that $\text{supp } S \subseteq C$ and $\mu_R(C) = 2\mu_R(\text{supp } S)$.

Step 3. We have $[\mathcal{R}]_C \subseteq W^{36}$.

Indeed, let $T \in [\mathcal{R}]_C$ be an involution. We first construct an involution $U \in [\mathcal{R}]_C$ such that the support of the involution $S(USU)$ has conditional measure $f = \mu_R(\text{supp } T)$. By lemma 2.2, we may find $E \subseteq (C \setminus \text{supp } S)$ which has conditional measure $\frac{f}{2}$. Let F_1 be a subset of a fundamental domain of $S_{\upharpoonright \text{supp } S}$ having conditional measure $\frac{\mu(C) - f}{4}$. Put $F_2 = T(F_1)$, let $F = F_1 \sqcup F_2$ and let $\varphi \in [[\mathcal{R}]]$ be a partial

isomorphism of domain E and range F given by lemma 2.3. Define the involution $U \in [\mathcal{R}]_C$ by

$$U(x) = \begin{cases} \varphi(x) & \text{if } x \in E \\ \varphi^{-1}(x) & \text{if } x \in F \\ x & \text{else} \end{cases}$$

Then the support of $S(USU)$ is $E \cup F$, hence has conditional measure $f = \mu_{\mathcal{R}}(\text{supp } T)$. Thanks to step 1, we may find $\tilde{U} \in W^2$ which coincides with U on $\text{supp } S \subseteq B$. Then, $\tilde{U}S\tilde{U}^{-1} = USU$ so that $S(USU)$ belongs to W^8 . Because its support has the same conditional measure as the support of T , lemma 2.3 yields that $SUSU$ and T are conjugated by an involution V supported on C . Again, by step 1, we find \tilde{V} such that V and \tilde{V} coincide on C , and we deduce that $T = VSUSUV = \tilde{V}S\tilde{U}S\tilde{U}^{-1}\tilde{V}^{-1}$ belongs to W^{12} .

Now by a result of Ryzhikov [Ryz93] any element of $[\mathcal{R}]_C$ is the product of at most 3 involutions in $[\mathcal{R}]_C$, so $[\mathcal{R}]_C \subseteq W^{36}$.

We can now conclude the proof exactly as in [KT10]: let (T_n) be a sequence of elements of $[\mathcal{R}]$ such that $d_C(T_n, 1) \rightarrow 0$. Let $D_n = \text{supp } T_n$, and $D = \bigcup_m g_m D_m$. We only need to show that there exists m such that $T_m \in W^{98}$. Going to a subsequence if necessary, we may assume that $\sum \mu_{\mathcal{R}}(D_n) \leq \mu_{\mathcal{R}}(C)$, hence $\mu_{\mathcal{R}}(D) \leq \mu_{\mathcal{R}}(C)$. Now there is $A \subseteq C$ such that $\mu_{\mathcal{R}}(A) = \mu_{\mathcal{R}}(D)$. We may find $S \in [\mathcal{R}]$ such that $S(A) = D$; there is m such that $S \in g_m W$, and we put $T = g_m^{-1}S \in W$. We have $D_m \subseteq T(A)$, and so $T^{-1}T_m T \in [\mathcal{R}]_A \subseteq [\mathcal{R}]_C \subseteq W^{36}$. Thus T_m belongs to $WW^{36}W = W^{38}$, and we are done. \square

7.2 Connectedness

Whenever \mathcal{R} is of type I_n , the metric d_C induces the discrete topology. However, if \mathcal{R} is aperiodic, the induced topology is connected.

Lemma 7.4. *Let \mathcal{R} be an aperiodic pmp equivalence relation. Then $([\mathcal{R}], d_C)$ is connected.*

Proof. By a result of Ryzhikov [Ryz93], $[\mathcal{R}]$ is generated by involutions, so it suffices to connect involutions to the identity. But if T is an involution, let A be a fundamental domain of T . Since \mathcal{R} is aperiodic we may, as in step 2 of the proof of theorem 7.3, find an increasing family $(A_t)_{t \in [0,1]}$ of subsets of A such that $\mu_{\mathcal{R}}(A_t) = t\mu_{\mathcal{R}}(A)$. Then define the involution T_t by

$$T_t(x) = \begin{cases} T(x) & \text{if } x \in A_t \\ T^{-1}(x) & \text{if } x \in T(A_t) \\ x & \text{else} \end{cases}$$

Then $d_C(T_{t_1}, T_{t_2}) \leq |t_1 - t_2|$, $T_0 = \text{id}_X$ and $T_1 = T$, so that T is connected to the identity. \square

Putting together lemma 7.4 and theorem 7.3, we have the following consequence.

Corollary 7.5. *Let \mathcal{R} be a pmp aperiodic equivalence relation. Then every morphism from $[\mathcal{R}]$ to a zero dimensional separable group is trivial.*

Remark. This implies that the full group of an aperiodic pmp equivalence relation cannot act nontrivially on a Cantor set, in particular it can never be abstractly isomorphic to the full group of a Cantor dynamical system, answering a question suggested by Vincent Tassion (for a definition of full groups in the topological setting, cf. [GPS99]).

We can then characterize aperiodic pmp equivalence relations algebraically.

Corollary 7.6. *Let \mathcal{R} be a pmp equivalence relation. Then \mathcal{R} is aperiodic iff every morphism from its full group to a totally disconnected separable group is trivial. Moreover, when \mathcal{R} is periodic, it admits a non trivial morphism onto $\mathbb{Z}/2\mathbb{Z}$.*

Proof. By corollary 7.5, we only need to show that if \mathcal{R} is not aperiodic, then there exists a non trivial morphism $[\mathcal{R}] \rightarrow \mathbb{Z}/2\mathbb{Z}$. So let \mathcal{R} be a non aperiodic equivalence relation, then we find an \mathcal{R} -invariant positive set A and $n \in \mathbb{N}$ such that all the elements of A have \mathcal{R} -classes of cardinality n . Then we have a morphism $\pi_A : [\mathcal{R}] \rightarrow [\mathcal{R}]_A$ given by restriction. Let B be a fundamental domain of $\mathcal{R}|_A$, then we may identify the full group of $\mathcal{R}|_A$ with the group of measurable maps from B to the symmetric group on n elements. Using the signature and π_A , we get a non trivial morphism φ from $[\mathcal{R}]$ to the group $L^0(B, \mathbb{Z}/2\mathbb{Z})$ of measurable maps from B to $\mathbb{Z}/2\mathbb{Z}$. Such a group has only elements of order 2, so it is a $\mathbb{Z}/2\mathbb{Z}$ vector space. Fix a non trivial element y in the image of φ , then using the axiom of choice we find a projection $p : L^0(B, \mathfrak{S}_2) \rightarrow \mathbb{Z}/2\mathbb{Z}$ onto the vector space spanned by y . Then $p \circ \varphi$ is a non trivial morphism from $[\mathcal{R}]$ to $\mathbb{Z}/2\mathbb{Z}$. \square

Note that this theorem implies that if \mathcal{R} is not aperiodic, then its full group equipped with the *uniform* metric does not satisfy the automatic continuity property, as it is connected but admits a morphism onto $\mathbb{Z}/2\mathbb{Z}$. It would be interesting to characterize the full group of \mathcal{R} equipped with the uniform metric in terms of the automatic continuity property, namely to answer the following question.

Question 7.7. Let \mathcal{R} be a pmp aperiodic equivalence relation. Is every morphism from $([\mathcal{R}], d_u)$ to a separable group continuous?

8 Extreme amenability

This section is devoted to giving a simple proof of the following proposition, which was proved by Giordano and Pestov in the ergodic case [GP07, prop 5.3].

Proposition 8.1. *Let \mathcal{R} be a hyperfinite equivalence relation, then its full group is extremely amenable.*

The previous proposition is a straightforward application of the two following lemmas. The first one is an immediate consequence of theorem 1.17, but we give here a direct proof.

Lemma 8.2. *Let $\mathcal{R} = \bigcup_n \mathcal{R}_n$ be a pmp equivalence relation, where the \mathcal{R}_n 's are increasing finite equivalence relations. Then $\bigcup_n [\mathcal{R}_n]$ is dense in $[\mathcal{R}]$.*

Proof. Let $\varphi \in [\mathcal{R}]$ and $\epsilon > 0$, since the \mathcal{R}_n 's exhaust \mathcal{R} there exists $n \in \mathbb{N}$ such that $\mu(\{x \in X : \varphi(x) \mathcal{R}_n x\}) > 1 - \epsilon$. Now put $\psi(x) = \varphi(x)$ whenever $\varphi(x) \mathcal{R}_n x$. We can extend ψ to an element $\tilde{\psi}$ of the full group of \mathcal{R}_n . Indeed, ψ induces a partial bijection in every \mathcal{R}_n -class, and every such partial bijection can be extended in a Borel way to have full domain (using a Borel total order on X , one can for instance send the first element of each equivalence class not in the domain of ψ to the first element of the class not in the range of ψ , etc.). By definition, $d_u(\tilde{\psi}, \varphi) < \epsilon$. \square

Lemma 8.3. *Let \mathcal{R} be a finite equivalence relation. Then its full group is extremely amenable.*

Proof. For $n \in \mathbb{N}$, let X_n be the set of $x \in X$ whose \mathcal{R} -class has cardinality n . Then we have a topological group isomorphism between the full group of \mathcal{R} and the product $\prod_{n \in \mathbb{N}} [\mathcal{R}_{\upharpoonright X_n}]$. Let A_n be a transversal for $\mathcal{R}_{\upharpoonright X_n}$, then we can identify $[\mathcal{R}_{\upharpoonright X_n}]$ with the group $L^0(A_n, \mathfrak{S}_n)$ of measurable maps from A_n to \mathfrak{S}_n . This identification is an isometry if we put on $L^0(A_n, \mathfrak{S}_n)$ the L^1 metric d_1 defined by:

$$d_1(f, g) = \int_{A_n} d_{\text{Ham}}(f(x), g(x)) d\mu(x),$$

where d_{Ham} is the Hamming metric on \mathfrak{S}_n (i.e. $d_{\text{Ham}}(\sigma, \sigma') = \text{Card}(\{x \in \{1, \dots, n\} : \sigma(x) \neq \sigma'(x)\})$). Then, a theorem of Glasner ([Gla98, theorem 1.3], see also [GP07, theorem 2.20]) implies that $[\mathcal{R}_{\upharpoonright X_n}] = L^0(A_n, \mathfrak{S}_n)$ is extremely amenable since \mathfrak{S}_n is compact. We conclude that $[\mathcal{R}] = \prod_{n \in \mathbb{N}} [\mathcal{R}_{\upharpoonright X_n}]$ is also extremely amenable. \square

Remark. In order to prove proposition 8.1, one could also use Dye's theorem and identify the full group of the aperiodic part X_∞ of \mathcal{R} with $L^0(X_\infty, [\mathcal{R}_0])$, which is easily seen to be extremely amenable once we know that $[\mathcal{R}_0]$ is amenable. This also yields that the full group of any hyperfinite equivalence relation is actually a Levy group.

Using the theorem of Connes, Feldmann and Weiss, and the fact that the full group acts continuously by affine isometric transformations on the field of invariant means on \mathcal{R} , one can easily see that the converse of proposition 8.1 holds, and so we get the following corollary.

Theorem 8.4. *Let \mathcal{R} be a pmp equivalence relation. Then the following statements are equivalent.*

- (i) \mathcal{R} is hyperfinite.
- (ii) The full group of \mathcal{R} is amenable.
- (iii) The full group of \mathcal{R} is extremely amenable.

References

- [CFW81] A. Connes, J. Feldman, and B. Weiss. An amenable equivalence relation is generated by a single transformation. *Ergodic Theory Dynamical Systems*, 1(4):431–450 (1982), 1981.

- [Dye59] H. A. Dye. On groups of measure preserving transformation. I. *Amer. J. Math.*, 81:119–159, 1959.
- [Dye63] H. A. Dye. On groups of measure preserving transformations. II. *Amer. J. Math.*, 85:551–576, 1963.
- [Eig81] S. J. Eigen. On the simplicity of the full group of ergodic transformations. *Israel J. Math.*, 40(3-4):345–349 (1982), 1981.
- [Eps08] Inessa Epstein. *Some results on orbit inequivalent actions of non-amenable groups*. ProQuest LLC, Ann Arbor, MI, 2008. Thesis (Ph.D.)–University of California, Los Angeles.
- [Gab00] Damien Gaboriau. Coût des relations d’équivalence et des groupes. *Invent. Math.*, 139(1):41–98, 2000.
- [Gla98] Eli Glasner. On minimal actions of Polish groups. *Topology Appl.*, 85(1-3):119–125, 1998. 8th Prague Topological Symposium on General Topology and Its Relations to Modern Analysis and Algebra (1996).
- [GP07] Thierry Giordano and Vladimir Pestov. Some extremely amenable groups related to operator algebras and ergodic theory. *J. Inst. Math. Jussieu*, 6(2):279–315, 2007.
- [GPS99] Thierry Giordano, Ian F. Putnam, and Christian F. Skau. Full groups of Cantor minimal systems. *Israel J. Math.*, 111:285–320, 1999.
- [Kai97] Vadim A. Kaimanovich. Amenability, hyperfiniteness, and isoperimetric inequalities. *C. R. Acad. Sci. Paris Sér. I Math.*, 325(9):999–1004, 1997.
- [Kec10] Alexander S. Kechris. *Global aspects of ergodic group actions*, volume 160 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2010.
- [KM04] Alexander S. Kechris and Benjamin D. Miller. *Topics in orbit equivalence*, volume 1852 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2004.
- [KT10] John Kittrell and Todor Tsankov. Topological properties of full groups. *Ergodic Theory Dynam. Systems*, 30(2):525–545, 2010.
- [Lev95] Gilbert Levitt. On the cost of generating an equivalence relation. *Ergodic Theory Dynam. Systems*, 15(6):1173–1181, 1995.
- [LM14] François Le Maître. The number of topological generators for full groups of ergodic equivalence relations. *to appear in Inventiones Mathematicae*, 2014.
- [Mat13] Hiroki Matui. Some remarks on topological full groups of Cantor minimal systems II. *Ergodic Theory Dynam. Systems*, 33(5):1542–1549, 2013.

- [OW80] Donald S. Ornstein and Benjamin Weiss. Ergodic theory of amenable group actions. I. The Rohlin lemma. *Bull. Amer. Math. Soc. (N.S.)*, 2(1):161–164, 1980.
- [Ros09] Christian Rosendal. Automatic continuity of group homomorphisms. *Bull. Symbolic Logic*, 15(2):184–214, 2009.
- [RS07] Christian Rosendal and Sławomir Solecki. Automatic continuity of homomorphisms and fixed points on metric compacta. *Israel J. Math.*, 162:349–371, 2007.
- [Ryz93] V. V. Ryzhikov. Factorization of an automorphism of a full Boolean algebra into the product of three involutions. *Mat. Zametki*, 54(2):79–84, 159, 1993.
- [Vee77] William A. Veech. Topological dynamics. *Bull. Amer. Math. Soc.*, 83(5):775–830, 1977.