

Separating inequalities for nonnegative polynomials that are not sums of squares

Sadik Iliman^a, Timo de Wolff^b

^a*Goethe-Universität, FB 12 – Institut für Mathematik, Postfach 11 19 32, 60054 Frankfurt am Main, Germany*

^b*Universität des Saarlandes, Fachrichtung Mathematik, Postfach 15 11 50, 66041 Saarbrücken, Germany*

Abstract

Ternary sextics and quaternary quartics are the smallest cases where there exist nonnegative polynomials that are not sums of squares (SOS). A complete classification of the difference between these cones was given by G. Blekherman via analyzing the extreme rays of the corresponding dual cones. However, an exact computational approach in order to build separating extreme rays for nonnegative polynomials that are not sums of squares is a widely open problem. We provide a method substantially simplifying this computation for certain classes of polynomials on the boundary of the PSD cones. In particular, our method yields separating extreme rays for every nonnegative ternary sextic with at least seven zeros, which proves a slight variation of a conjecture by Blekherman for many instances. As an application, we compute rational certificates for some prominent polynomials.

Keywords: PSD, nonnegative polynomial, SOS, extreme ray, certificate, exact methods
2010 Mathematics Subject Classification. 11E20, 11E25, 14P99, 52A20

1. Introduction

We consider real polynomials in the vector space of all homogeneous polynomials in n variables of degree d , denoted by $H_{n,d}$. For every $p \in H_{n,d}$ we denote its real projective variety as $\mathcal{V}(p)$. Let $P_{n,d} \subset H_{n,d}$ be the cone of all nonnegative polynomials in n variables of degree d .

Inside $H_{n,2d}$, there are two full dimensional convex cones of special interest, the cone of nonnegative polynomials and the cone of sums of squares (for a general background about nonnegative polynomials and sums of squares see e.g. [11, 15, 17, 22]; for some metric and convexity properties of these cones see [2]).

$$\begin{aligned} P_{n,2d} &:= \{f \in H_{n,2d} : f(x) \geq 0, \text{ for all } x \in \mathbb{R}^n\} \\ \Sigma_{n,2d} &:= \left\{ f \in P_{n,2d} : f = \sum_i f_i^2 \text{ for some } f_i \in H_{n,d} \right\}. \end{aligned}$$

The investigation of the relationship between the cone of nonnegative polynomials and the cone of sums of squares began in the seminal work of Hilbert when he showed that the cone of nonnegative polynomials coincides with the cone of sums of squares exactly in the cases of bivariate forms ($n = 2$), quadratic forms ($2d = 2$) and ternary quartics ($n = 3, 2d = 4$) ([14]).

The Motzkin polynomial $m(x, y, z) = x^4y^2 + x^2y^4 - 3x^2y^2z^2 + z^6$ was the first explicitly known example for a nonnegative polynomial which is not a sum of squares. Most proofs for this fact are based on term by term inspections (see e.g. [18, 22]). In near past other proofs were found, e.g. using representation theory (see [9]).

In [3], Blekherman showed that for fixed dimension $2d \geq 4$ there are significantly more nonnegative polynomials than sums of squares as n tends to infinity. However, the question of precisely when nonnegative polynomials begin to significantly overtake sums of squares is much less understood. In the smallest cases where there exist nonnegative polynomials which are not sums of squares ($(n, 2d) = (3, 6), (4, 4)$) the general conjecture is that these two cones do not differ very much. This conjecture is supported by the following two facts: Firstly, the maximal dimensional difference between exposed faces of the cone of nonnegative polynomials and sums of squares is one (see [7]). Secondly, all extreme rays of the dual sums of squares cone $\Sigma_{3,6}^*$ have rank one or rank seven (see [4]).

Recently, in [6], it is shown that, except the discriminant, there is a unique component of the algebraic boundary of $\Sigma_{3,6}$ with degree 83200, which indicates the complicated structure of the SOS cone. But still, the geometry and the relationship between these two cones in the smallest cases are less understood.

In the smallest cases $(n, 2d) = (3, 6)$ and $(n, 2d) = (4, 4)$, Blekherman shows that it is precisely the Cayley-Bacharach relation that prevents sums of squares from filling out the cone of nonnegative polynomials. More precisely, in [4], it is shown that every separating extreme ray in the dual SOS cone for a given nonnegative polynomial that is not a sum of squares depends on an 9-point configuration for $(n, 2d) = (3, 6)$ resp. an 8-point configuration for $(n, 2d) = (4, 4)$ coming from the intersection of two cubic resp. three quadric polynomials. Furthermore, given an appropriate 9-point (resp. 8-point) configuration, one can write down an extreme ray of the dual SOS cone (see Theorems 2.6 and 2.7) corresponding to faces of maximal dimension of the SOS cone. In [5], Blekherman extends the investigation of the extreme rays of the dual sums of squares cones, especially for ternary forms.

A central problem in this area is how to determine the separating inequalities efficiently. This can always be done in a numerical way (see Section 2.2), but is widely open for exact methods currently. Of course, symbolic results are strongly preferred – not only since they provide algebraic certificates, which are exact. Furthermore, they can also be connected to the whole machinery of algebraic geometry and thus be used to tackle follow-up questions (like the semi-algebraic description of appropriate nine point configurations for the Motzkin polynomial, which we provide here; see e.g., Figure 5). Hence, finding constructive symbolic methods for computing these inequalities is one main research issue. Blekherman’s result does not provide an efficient symbolic way to obtain a proper 9-point

(resp. 8-point) configuration to solve this problem (see 2.2 for further details).

The key idea of this article is to construct a proper 9-point (resp. 8-point) configuration out of a given initial set of points. Specifically, we investigate nonnegative polynomials p which lie on the boundary of the cones $P_{3,6}$ and $P_{4,4}$ (which cover most of the explicitly known nonnegative polynomials that are not SOS, see [3]). Our main result, Theorem 3.1, provides a sufficient condition for using k zeros of p as a subset of a 9-point (resp. 8-point) configuration. The idea is to fill up the set of k zeros with $9-k$ (resp. $8-k$) points such that a genericity and a quadratic condition based on the Cayley-Bacharach relation hold (note that $k \leq 10$ for $p \in P_{3,6} \setminus \Sigma_{3,6}$ and $p \in P_{4,4} \setminus \Sigma_{4,4}$; see [6, 10]). Given these conditions, which are computationally easy to check, we can construct a separating extreme ray immediately. This method reduces the difficulty of constructing separating extreme rays via symbolic computation significantly. Furthermore, it yields rational certificates for rational point configurations and even for rational varieties $\mathcal{V}(p) \subset \mathbb{Q}^3$ resp. $\mathcal{V}(p) \subset \mathbb{Q}^4$.

We show that for $p \in P_{3,6} \setminus \Sigma_{3,6}$ and $k \geq 7$ (resp. $p \in P_{4,4} \setminus \Sigma_{4,4}$ and $k \geq 6$) almost every 9-point (resp. 8-point) configuration containing the seven (resp. six) zeros leads to a certificate for a nonnegative polynomial p to be not SOS. This proves a slightly modified version of Blekherman's Conjecture 3.6 for many instances.

In Section 2, we review some curve theoretical issues as, e.g., the Cayley-Bacharach relation and present Blekherman's results on the SOS cones $\Sigma_{3,6}$ and $\Sigma_{4,4}$. In Section 3, we state and prove our main Theorems 3.1 and 3.3 for ternary sextics and quaternary quartics and discuss exactness and rationality of our methods. Section 4 deals with the special case of polynomials with exactly seven zeros. We show that in this case our method generically yields a separating extreme ray (Theorem 4.1), which also proves the special instances in Blekherman's Conjecture 3.6. Finally, we discuss the difficulties of dropping zeros in our method in Section 5 by applying it to the Motzkin polynomial and constructing a rational extreme ray certificate for it. Furthermore, based on the Motzkin polynomial we investigate some geometric aspects of the set of appropriate point configurations for our method (see Figure 5).

2. Preliminaries

2.1. Curve theoretical background

We recall some classical results from algebraic geometry. From now on we consider every investigated polynomial to be homogeneous. We start with the Cayley-Bacharach relation. It exists in various formulations (see [13]); we use the ones given in [4].

Lemma 2.1. *Let $p_1, p_2 \in \mathbb{C}[x_1, x_2, x_3]$ be two ternary cubics intersecting transversely in nine projective points $\gamma_1, \dots, \gamma_9$. Let v_1, \dots, v_9 be affine representatives of γ_i . Then there is a unique linear relation on the values of any ternary cubic on v_i*

$$\sum_{j=1}^9 u_j f(v_j) = 0 \text{ for all ternary cubics } f \tag{2.1}$$

with nonzero $u_j \in \mathbb{C}$. Furthermore, if (2.1) is satisfied, then the following genericity condition holds

$$\text{no four of the } v_i \text{ lie on a line and no seven on a quadric.} \quad (2.2)$$

Analogously, we have for quaternary quadrics

Lemma 2.2. *Let $p_1, p_2, p_3 \in \mathbb{C}[x_1, \dots, x_4]$ be three quaternary quadrics intersecting transversely in eight projective points $\gamma_1, \dots, \gamma_8$. Let v_1, \dots, v_8 be affine representatives of γ_i . Then there is a unique linear relation on the values of any quaternary quadric on v_i*

$$\sum_{j=1}^8 u_j f(v_j) = 0 \text{ for all quaternary quadrics } f \quad (2.3)$$

with nonzero $u_j \in \mathbb{C}$. Furthermore, if (2.3) is satisfied, then the following genericity condition holds

$$\text{no five of the } v_i \text{ lie on a plane (i.e., a projective linear 2-space).} \quad (2.4)$$

For the genericity condition (2.4) see e.g. [14]. Note that, if all points v_j are real, then all Cayley-Bacharach coefficients u_j are real, too (see e.g. [4, Lemma 4.1]) and can be computed by solving a system of linear equations with the coefficients of forms in $H_{3,3}$ resp. $H_{4,2}$ as variables.

Each of the conditions (2.2) and (2.4) can be checked easily by investigating the minors of the matrix given by the vectors v_j .

For the description of separating extreme rays of $\Sigma_{3,6}^*$ yielded by Blekherman's Theorem 2.6 one needs to investigate 9-point configurations given as the intersection of two coprime ternary cubics. The following lemma shows that coprimality is the case generically (see [23]).

Lemma 2.3. *Suppose $A := \{v_1, \dots, v_8\}$ is a set of eight distinct points in \mathbb{R}^3 , no four on a line and no seven on a quadric and let f_1, f_2 be a basis for the vector space of all homogeneous cubics with projective variety affinely represented by A . Then f_1 and f_2 are relatively prime.*

This lemma yields that one can apply Bezout's theorem in order to compute a ninth intersection point v_9 of f_1 and f_2 . However, v_9 might not be different from v_1, \dots, v_8 (i.e. the intersection multiplicity might be greater than 1). But, again, generically this will not be the case as the following lemma shows (see [19]).

Lemma 2.4. *Let f_1, f_2 be two homogeneous polynomials in n variables (with $n \geq 2$) of degree d and generic coefficients. The discriminant $\Delta(f_1, f_2)$ vanishes if and only if $f_1(x) = f_2(x) = 0$ has a singular solution. The set of polynomials for which this is the case is a hypersurface.*

In Section 4 we will investigate the special case of polynomials $p \in P_{3,6} \setminus \Sigma_{3,6}$ with exactly seven zeros. In this context we use the following lemma (see [23]).

Lemma 2.5. *Suppose A is a set of seven distinct points in \mathbb{R}^3 , no four on a line and no seven on a quadric with basis f_1, f_2, f_3 for the vector space of homogeneous cubics with projective variety affinely represented by A . Then f_1, f_2, f_3 have no common zeros outside of A .*

2.2. Blekherman's results

In [4], Blekherman was able to fully characterize the extreme rays of the dual SOS cones $\Sigma_{3,6}^*$ and $\Sigma_{4,4}^*$, basically via the Cayley-Bacharach relation. We recall his main result, concentrating on ternary sextics.

Theorem 2.6 (Blekherman [4]). *Suppose that a ternary sextic p is nonnegative but not a sum of squares. Then there exist two real cubics $q_1, q_2 \in H_{3,3}$ intersecting transversely in 9 projective points $\gamma_1, \dots, \gamma_9$ which yield a certificate for $p \in P_{3,6} \setminus \Sigma_{3,6}$. More precisely, let v_1, \dots, v_9 be affine representatives of $\gamma_1, \dots, \gamma_9$. Then there exists a linear functional $l : H_{3,6} \rightarrow \mathbb{R}$ given by*

$$l(f) = \sum_{i=1}^9 a_i f(v_i)$$

for some $a_i \in \mathbb{C}$ such that $l(p) < 0$ and $l(\Sigma_{3,6}) \geq 0$. Furthermore, at most two of the points γ_i are complex.

This theorem works the same way for $p \in P_{4,4} \setminus \Sigma_{4,4}$ (see [4]).

Recall that for every $l \in \Sigma_{3,6}^*$ there is a corresponding quadratic for Q_l defined by $Q_l : H_{3,3} \rightarrow \mathbb{R}, f \mapsto l(f^2)$ (see e.g. [4, 16]). One defines the rank of a linear functional $l \in \Sigma_{3,6}^*$ by $\text{rank}(l) := \text{rank}(Q_l)$. In [4] it is shown that every extreme ray of $\Sigma_{3,6}^*$, which does not correspond to point evaluation (i.e., a rank 1 form), is given by a rank 7 form which comes from a 9-point evaluation.

Theorem 2.7 (Blekherman [4]). *Suppose l spans an extreme ray of $\Sigma_{3,6}^*$, which does not correspond to a point evaluation. Let W_l be the kernel of the corresponding quadratic form Q_l and suppose $q_1, q_2 \in W_l$ intersect transversely in 9 real projective points $\gamma_1, \dots, \gamma_9$ with affine representatives v_1, \dots, v_9 such that the unique Cayley-Bacharach relation is given by*

$$u_1 f(v_1) + \dots + u_9 f(v_9) = 0 \text{ for } f \in H_{3,3}.$$

Then Q_l can be uniquely written as

$$Q_l(f) = a_1 f(v_1)^2 + \dots + a_9 f(v_9)^2$$

with exactly one single negative coefficient a_k and the rest of the a_i being strictly positive and a_k is given by

$$a_k = \frac{-u_k^2}{\frac{u_1^2}{a_1} + \dots + \frac{u_9^2}{a_9} - \frac{u_k^2}{a_k}}. \quad (2.5)$$

Furthermore, any such form is extreme in $\Sigma_{3,6}^*$.

Again, an analogue version for $p \in P_{4,4} \setminus \Sigma_{4,4}$ holds (see [4]).

Suppose $p \in P_{3,6} \setminus \Sigma_{3,6}$ and we want to construct a separating extreme ray l for p using the upper theorem. Therefore, we need to find two coprime ternary cubics q_1, q_2 intersecting in 9 points. But, q_1, q_2 need to be contained in the kernel W_l of the quadratic form Q_l corresponding to l . Hence, one already needs to know l in advance to determine q_1, q_2 .

This problem can be avoided by choosing a 9-point configuration $A = \{v_1, \dots, v_9\}$ coming from an intersection of two real ternary cubics q_1, q_2 . So, a separating extreme ray l is obtained by finding an appropriate $a = (a_1, \dots, a_9)$ satisfying (2.5) with respect to A such that $l_a(p) < 0$. The question whether such an a exists is unclear a priori though it can be answered by quantifier elimination methods (see, e.g., [1, 8]). But, to the best of our knowledge, no methods are known to compute an appropriate a in a symbolic, exact way efficiently.

However, one can solve this problem numerically. Let $p \in P_{3,6} \setminus \Sigma_{3,6}$ be a ternary sextic and $r \in \text{int}(\Sigma_{3,6})$ (e.g., $r = x^6 + y^6 + z^6$ or $r = (x^2 + y^2 + z^2)^3$). Consider the following semidefinite optimization problem.

$$\min_{\lambda \in \mathbb{R}} \lambda \quad \text{such that} \quad p + \lambda r \in \Sigma_{3,6}.$$

For λ minimal the polynomial $p + \lambda r$ is strictly positive and lies on the boundary of $\Sigma_{3,6}$. Hence, $p + \lambda r$ is a sum of exactly three squares $s_1^2 + s_2^2 + s_3^2$ (see [4]).

The polynomials s_1, s_2, s_3 have no common zeros and an appropriate linear combination of two of these polynomials can be used as q_1 and q_2 in Blekherman's theorem (see [4, Corollary 2.3]). Of course, the computation of the corresponding nine intersection points will be difficult and not exact, too. Furthermore, getting "nice" values (e.g., a rational minimal λ) depends also highly on the choice of the polynomial $r \in \text{int}(\Sigma_{3,6})$. It is not clear how to choose r in dependence of p .

In the case $p \in P_{4,4} \setminus \Sigma_{4,4}$ this approach works the same way. For λ minimal the polynomial $p + \lambda r$ is a sum of exactly four squares $p + \lambda r = s_1^2 + s_2^2 + s_3^2 + s_4^2$. Three of these four s_j have a common zero (see [4]).

3. A certificate for boundary polynomials not to be SOS

Our approach to construct a separating extreme ray for a given boundary polynomial $p \in \partial P_{3,6} \setminus \Sigma_{3,6}$ is to investigate certain point sets $A := \{v_1, \dots, v_9\}$ containing the variety

$\mathcal{V}(p)$, satisfying the genericity condition (2.2) and for which we can certify that there are coprime polynomials $q_1, q_2 \in H_{3,3}$ with $\mathcal{V}(q_1) \cap \mathcal{V}(q_2) = A$.

Note that, if we talk about zeros of homogeneous polynomials in this and the following sections, then we always consider their affine representatives with slight abuse of notation.

The easiest case is when p has at least eight zeros v_1, \dots, v_8 (satisfying (2.2)). Lemma 2.3 provides the existence of coprime q_1, q_2 vanishing on v_1, \dots, v_8 and thus a ninth point v_9 is given by Bezout's Theorem. For a generic set of zeros v_1, \dots, v_8 the corresponding coprime polynomials q_1, q_2 have generic coefficients and hence, due to Lemma 2.4, we have $v_9 \notin \{v_1, \dots, v_8\}$ generically. Thus, $A := \{v_1, \dots, v_9\}$ satisfies (2.2) generically. This yields a certificate l immediately since for any choice of a_1, \dots, a_8 we obtain an $a_9 < 0$ by (2.5) and hence

$$l(p) = \sum_{j=1}^9 a_j p(v_j) = a_9 p(v_9) < 0.$$

In the following we generalize this idea to any number of zeros between one and eight. We choose the zeros v_1, \dots, v_k of a polynomial $p \in \partial P_{3,6} \setminus \Sigma_{3,6}$ as a subset of the nine intersection points $A := \{v_1, \dots, v_9\}$ of two coprime ternary cubics. We provide a symbolic method based on genericity conditions, which yields a separating extreme ray if one finds a $(9 - k)$ -point configuration satisfying some quadratic relation. Specifically, the following theorem holds.

Theorem 3.1. *Let $p \in \partial P_{3,6}$ such that p is not SOS. Let $A := \{v_1, \dots, v_9\} \subset \mathbb{R}^3$ such that the genericity condition (2.2) holds and $\mathcal{V}(p) = \{v_1, \dots, v_k\}$ with $1 \leq k \leq 8$ (i.e.: the projectivization of A is the intersection of two coprime polynomials $q_1, q_2 \in H_{3,3}$). Then one can compute a certificate $l_a : H_{3,6} \rightarrow \mathbb{R}, f \mapsto \sum_{j=1}^9 a_j f(v_j)$, $a := (a_1, \dots, a_9) \in \mathbb{R}^9$ with respect to A for p to be not SOS, if the following inequality holds*

$$(u_{k+1}^2 + \dots + u_8^2)(p(v_{k+1}) + \dots + p(v_8)) < u_9^2 p(v_9). \quad (3.1)$$

Here the u_j are given by the unique Cayley-Bacharach relation on A and l_a is an extreme ray of $\Sigma_{3,6}^*$.

Note that the Cayley-Bacharach coefficients u_j can be computed by solving a system of linear equations (see exemplarily in Section 5). Additionally, all u_j are rational, if every point in A is rational. Note furthermore that, for an arbitrary p , it is not clear whether an A with $\mathcal{V}(p) \subset A$ satisfying (3.1) does always exist. We discuss certain special cases in the two following sections.

Proof. Let $p \in \partial P_{3,6}$ with $\mathcal{V}(p) = \{v_1, \dots, v_k\}$, $1 \leq k \leq 8$ such that the genericity condition (2.2) holds for $\mathcal{V}(p)$. We choose points v_{k+1}, \dots, v_8 such that (2.2) is still satisfied. We obtain v_9 as the intersection of two relatively prime, cubic polynomials spanning the vector space of all ternary cubics vanishing on v_1, \dots, v_8 (see Lemma 2.3). Notice that, generically, we obtain $v_9 \notin \{v_1, \dots, v_8\}$ due to Lemma 2.4 and v_9 has to be real since v_1, \dots, v_9

is the intersection of two real polynomials (see, e.g., [23]). Let u_1, \dots, u_9 be the unique Cayley-Bacharach coefficients for v_1, \dots, v_9 in the sense of (2.1). Since $v_1, \dots, v_9 \in \mathbb{R}^3$ we have $u_1, \dots, u_9 \in \mathbb{R}$ (see [4, Lemma 4.1]).

By Theorem 2.7 every vector $a := (a_1, \dots, a_9) \in \mathbb{R}^9$ satisfying (2.5) with $a_1, \dots, a_8 > 0$, $a_9 < 0$ yields an extreme ray $l_a : H_{3,6} \rightarrow \mathbb{R}, f \mapsto \sum_{j=1}^9 a_j f(v_j)$ of the dual SOS cone $\Sigma_{3,6}^*$. l_a is the dual of a separating hyperplane for p if $l_a(p) < 0$, i.e., since $\mathcal{V}(p) = \{v_1, \dots, v_k\}$, if $a_{k+1}p(v_{k+1}) + \dots + a_9p(v_9) < 0$. By (2.5) this is equivalent to

$$\begin{aligned} a_{k+1}p(v_{k+1}) + \dots - \frac{u_9^2}{\frac{u_1^2}{a_1} + \dots + \frac{u_8^2}{a_8}} p(v_9) &< 0 \\ \Leftrightarrow (a_{k+1}p(v_{k+1}) + \dots + a_8p(v_8)) \cdot \left(\frac{u_1^2}{a_1} + \dots + \frac{u_8^2}{a_8} \right) &< u_9^2 p(v_9). \end{aligned}$$

Let $\lambda_{a_1, \dots, a_k} := \sum_{j=1}^k \frac{u_j^2}{a_j} \cdot (a_{k+1}p(v_{k+1}) + \dots + a_8p(v_8)) > 0$. Thus, $l_a(p) < 0$, if

$$\lambda_{a_1, \dots, a_k} + \sum_{j=k+1}^8 p(v_j) \left(u_j^2 + \sum_{i \in \{k+1, \dots, 8\} \setminus \{j\}} \frac{a_j u_i^2}{a_i} \right) < u_9^2 p(v_9).$$

We choose $a_{k+1} := 1, \dots, a_8 := 1$ and obtain

$$\lambda_{a_1, \dots, a_k} + (u_{k+1}^2 + \dots + u_8^2)(p(v_{k+1}) + \dots + p(v_8)) < u_9^2 p(v_9).$$

Since $\lim_{a_1, \dots, a_k \rightarrow \infty} \lambda_{a_1, \dots, a_k} \searrow 0$ the relaxation (3.1) yields an extreme ray l_a on A separating p from the SOS cone $\Sigma_{3,6}$.

If on the other hand any SOS polynomial $g \in \Sigma_{3,6}$ with $\mathcal{V}(g) = \mathcal{V}(p)$ would satisfy (3.1), then it follows from the upper construction that $l_a(g) < 0$ for a_1, \dots, a_k sufficiently large, $a_{k+1}, \dots, a_8 = 1$ and a_9 given by (2.5). This is a contradiction to Blekherman's Theorems 2.6 and 2.7. Thus, (3.1) is indeed a certificate for p to be not SOS. \square

Note that it is also easy to show that condition (3.1) is never satisfied for SOS polynomials $g \in \Sigma_{3,6}$ with $\mathcal{V}(g) = \mathcal{V}(p)$ in order to prove extremality of l_a by direct calculation and without usage of Blekherman's Theorems. This follows already from $g = \sum_{j=1}^r h_j^2$ and the Cayley-Bacharach relations.

In order to prove an analogon of the above theorem for $P_{4,4} \setminus \Sigma_{4,4}$, we need to show that Lemma 2.3 also holds for a seven point set $A := \{v_1, \dots, v_7\} \subset \mathbb{R}^4$. Generically, the vector space of all quadrics vanishing on A has dimension three (see [12]).

Lemma 3.2. *Suppose $A := \{v_1, \dots, v_7\}$ is a set of seven distinct points in \mathbb{R}^4 , no four on a plane such that q_1, q_2, q_3 is a basis for the vector space of all homogeneous quadrics with projective variety affinely represented by A . Then q_1, q_2, q_3 are relatively prime.*

Proof. Suppose q_1, q_2, q_3 have a common factor g . Then $q_j = g \cdot q'_j$ for $j \in \{1, 2, 3\}$ and g, q'_j have to be linear in $\mathbb{R}[x_1, \dots, x_4]$. Due to the genericity condition at most three zeros (w.l.o.g. v_1, v_2, v_3) are located on $\mathcal{V}(g)$ since otherwise there would exist at least five points are contained in a plane. Hence, $\mathcal{V}(q'_1), \mathcal{V}(q'_2)$ and $\mathcal{V}(q'_3)$ share four points, which is a contradiction since for each j all points in $\mathcal{V}(q'_j)$ are contained in a line. \square

Theorem 3.3. *Let $p \in \partial P_{4,4}$ such that p is not SOS. Let $A := \{v_1, \dots, v_8\} \subset \mathbb{R}^4$ such that the genericity condition (2.4) holds and $\mathcal{V}(p) = \{v_1, \dots, v_k\}$ with $1 \leq k \leq 7$ (i.e.: the projectivization of A is the intersection of three coprime polynomials $q_1, q_2, q_3 \in H_{4,2}$). Then there exists a certificate $l_a : H_{4,4} \rightarrow \mathbb{R}, f \mapsto \sum_{j=1}^8 a_j f(v_j)$, $a := (a_1, \dots, a_8) \in \mathbb{R}^8$ with respect to A for p to be not SOS, if the following inequality holds*

$$(u_{k+1}^2 + \dots + u_7^2)(p(v_{k+1}) + \dots + p(v_7)) < u_8^2 p(v_8). \quad (3.2)$$

Here the u_j are given by the unique Cayley-Bacharach relation on A and l_a is an extreme ray of $\Sigma_{4,4}^*$.

The proof works the same way as for Theorem 3.1 with the obvious modifications.

In fact, the proof of Theorem 3.1 already shows one possible way how to choose $a = (a_1, \dots, a_9) \in \mathbb{R}^9$ to obtain a separating extreme ray l_a .

Corollary 3.4. *For $p \in \partial P_{3,6} \setminus \Sigma_{3,6}$ and $A = \{v_1, \dots, v_9\} \supset \mathcal{V}(p)$ with (3.1) satisfied, one valid certificate is given by $a_1 = \dots = a_k = N \in \mathbb{R}$ (for N sufficiently large), $a_{k+1} = \dots = a_8 = 1$ and a_9 given by (2.5). For $p \in \partial P_{4,4} \setminus \Sigma_{4,4}$ and $A = \{v_1, \dots, v_8\} \supset \mathcal{V}(p)$ with (3.2) satisfied, one valid certificate is given by $a_1 = \dots = a_k = N \in \mathbb{R}$ (for N sufficiently large), $a_{k+1} = \dots = a_7 = 1$ and a_8 given by the analogon of (2.5) for $\Sigma_{4,4}^*$ (see [4]).*

In particular, l_a is a rational certificate, i.e., every a_j is rational, if every point $v_j \in A$ is rational.

\square

If one is interested in computing rational certificates, then, from an application view-point, there is the following problem. Suppose, we have a rational variety $\mathcal{V}(p) = \{v_1, \dots, v_k\}$ and we choose $v_{k+1}, \dots, v_8 \in \mathbb{Q}^3$ (resp. $v_{k+1}, \dots, v_7 \in \mathbb{Q}^4$) such that the genericity condition (2.2) (resp. (2.4)) holds (which is always possible). Then it is not clear a priori that the ninth intersection point $v_9 \in \mathbb{R}^3$ (resp. eighth intersection point $v_8 \in \mathbb{R}^4$), given by Bezout, is rational, too.

By results in [20] and [21], for $p \in P_{3,6} \setminus \Sigma_{3,6}$ (resp. $p \in P_{4,4} \setminus \Sigma_{4,4}$), the ninth (resp. eighth) intersection point can always be computed exactly. In particular, it can be deduced that this last point will always be rational whenever the remaining points are rational and hence, whenever $\mathcal{V}(p)$ is rational.

Corollary 3.5. *Let $p \in \partial P_{3,6} \setminus \Sigma_{3,6}$ with $\mathcal{V}(p) = \{v_1, \dots, v_k\} \subset \mathbb{Q}^3$ and $\{v_{k+1}, \dots, v_8\} \subset \mathbb{Q}^3$ such that (2.2) holds. Then there is a rational certificate l_a on $A = \{v_1, \dots, v_9\}$ with v_9 given by Bezout, whenever (3.1) holds.*

□

Obviously, an analogous result also holds in the case $(n, 2d) = (4, 4)$.

Note that in our Theorems 3.1 and 3.3 we only consider real points v_1, \dots, v_9 , whereas in Blekherman's Theorem 2.6 (at most) one pair of complex conjugated points are allowed. However, in [4], Blekherman states the following conjecture

Conjecture 3.6. *Every extreme ray $l \in \Sigma_{3,6}^*$, which is not a point evaluation, is given by two ternary cubics intersecting in only real points. Analogously, for $l \in \Sigma_{4,4}^*$.*

Based on our results, we formulate a slightly modified conjecture here.

Conjecture 3.7. *For $p \in P_{3,6} \setminus \Sigma_{3,6}$ there exist $v_1, \dots, v_9 \in \mathbb{R}^3$ yielding a separating extreme ray for p in the sense of Theorem 2.6. Analogously, for $p \in P_{4,4} \setminus \Sigma_{4,4}$.*

Clearly, Conjecture 3.6 implies Conjecture 3.7, since if every extreme ray is real representable, every nonnegative polynomial that is not SOS can be separated by a real intersection. It is unclear whether the two conjectures are indeed equivalent, however, we strongly suppose it.

4. The seven point case

Let $p \in P_{3,6} \setminus \Sigma_{3,6}$ and assume we are interested in finding a separating extreme ray l_a in $\Sigma_{3,6}^*$ for p . That means we need to find a generic 9-point set A being the intersection of two coprime polynomials $q_1, q_2 \in H_{3,3}$ such that the conditions in Theorem 2.6 hold. If p is located on the boundary of $P_{3,6}$, i.e., $\mathcal{V}(p) = \{v_1, \dots, v_k\} \neq \emptyset$, then Theorem 3.1 and Corollary 3.4 yield a certificate l_a whenever one can fill up $\mathcal{V}(p)$ with points v_{k+1}, \dots, v_9 such that condition (3.1) is satisfied.

However, it is not obvious a priori if and how points v_{k+1}, \dots, v_9 can be chosen such that the sufficient condition (3.1) of Theorem 3.1 holds (note that v_9 is always given by Bezout in this approach). It turns out that for $k = 7$, i.e., the easiest non-trivial case, the problem of choosing an appropriate v_8 is easy, since almost every point v_8 yields a point v_9 such that (3.1) is satisfied. Note that for $p \in \partial P_{3,6} \setminus \Sigma_{3,6}$ the variety $\mathcal{V}(p)$ always satisfies the genericity condition (2.2) (see [23]).

One reason why this case is of special interest is that $k = 7$ with v_1, \dots, v_7 satisfying the genericity condition (2.2) is the smallest number of zeros of a nonnegative polynomial such that the dimensional difference between exposed faces of $P_{3,6}$ and $\Sigma_{3,6}$ given by the vanishing of polynomials on these zeros is strictly positive (see [7]). In particular, by results in [7], this implies that for every generic configuration of at least seven points, one can always construct nonnegative polynomials that are not sums of squares vanishing at these points.

Theorem 4.1. *Let $p \in \partial P_{3,6} \setminus \Sigma_{3,6}$. Let $A := \{v_1, \dots, v_9\} \subset \mathbb{R}^3$ such that the genericity condition (2.2) holds and $\mathcal{V}(p) = \{v_1, \dots, v_7\}$. Then there exists a certificate $l_a : H_{3,6} \rightarrow \mathbb{R}, f \mapsto \sum_{j=1}^9 a_j f(v_j)$, $a_j \in \mathbb{R}$ with respect to A for p to be not SOS if*

$$u_8^2 p(v_8) \neq u_9^2 p(v_9). \quad (4.1)$$

Furthermore, l_a is an extreme ray of $\Sigma_{3,6}^*$.

Theorem 4.1 holds analogously in the case of $p \in P_{4,4} \setminus \Sigma_{4,4}$ and $k = 6$ with the obvious modifications. We omit to formulate the result for this case separately.

Proof. We choose v_8 such that the genericity condition (2.2) still holds for $\{v_1, \dots, v_8\}$ and obtain a real $v_9 \notin \{v_1, \dots, v_8\}$ such that (2.2) and Cayley-Bacharach hold for A generically (see proof of Theorem 3.1). Since all conditions of Theorem 3.1 are satisfied, there is a certificate l if $u_8^2 p(v_8) < u_9^2 p(v_9)$. But since $\{v_1, \dots, v_7, v_9\}$ also yields A with Bezout's theorem, this condition holds w.l.o.g. as long as $u_8^2 p(v_8) \neq u_9^2 p(v_9)$. \square

A nice consequence of this theorem is that it immediately verifies Conjecture 3.7 for special instances.

Corollary 4.2. *Conjecture 3.7 holds for $p \in \partial P_{3,6} \setminus \Sigma_{3,6}$ with $\#\mathcal{V}(p) \geq 7$ and $p \in \partial P_{4,4} \setminus \Sigma_{4,4}$ with $\#\mathcal{V}(p) \geq 6$. Furthermore, Conjecture 3.7 also holds for every exposed extremal form in $P_{3,6} \setminus \Sigma_{3,6}$.*

Proof. The first part immediately follows from Theorem 4.1. Furthermore, in [6], it is shown that every exposed extremal form in $P_{3,6} \setminus \Sigma_{3,6}$ has exactly ten zeros. By choosing seven of them, the results follow. \square

Let $p \in P_{3,6}$ such that p is not SOS with $\mathcal{V}(p) = \{v_1, \dots, v_k\} \subset \mathbb{R}^3$. Assume that we want to test for a large amount of point sets $\{v_{k+1}, \dots, v_8\}$ in \mathbb{R}^3 if (3.1) can be satisfied for each point set (under the condition that (2.2) holds for v_1, \dots, v_8). We have to do two steps. Firstly, we have to compute v_9 by calculating two coprime polynomials q_1, q_2 vanishing on $\{v_1, \dots, v_8\}$ via solving a system of linear equations on the coefficients of a ternary cubic and then calculate their Gröbner basis. If we are in the case $(n, 2d) = (3, 6)$, then this can also be done by using the formula given in [21]. Secondly, we have to compute the Cayley-Bacharach relation on $A := \{v_1, \dots, v_9\}$ via solving a system of linear equations again.

In the seven point case, i.e. if we want to check (4.1) for a large amount of point sets, we can avoid calculating the Cayley-Bacharach relation for each $\{v_8, v_9\}$ if we are not interested in computing the separating hyperplane given by l_a explicitly. The trick is to use Reznick's Lemma 2.5. Let $V(\mathcal{V}(p)) := \{h \in H_{3,3} : \mathcal{V}(p) \subseteq \mathcal{V}(h)\}$ be the vector space of all real ternary cubics vanishing on $\mathcal{V}(p)$. By Lemma 2.5 there is a basis $\{h_1, h_2, h_3\}$ of $V(\mathcal{V}(p))$ such that $\mathcal{V}(h_1) \cap \mathcal{V}(h_2) \cap \mathcal{V}(h_3) = \mathcal{V}(p)$ and h_1, h_2, h_3 do not depend on the choice of v_8 and v_9 .

Corollary 4.3. *Let p and A be as in Theorem 4.1. Let $h \in V(\mathcal{V}(p))$ with $\mathcal{V}(h) = \mathcal{V}(p)$. Then there exists a certificate $l_a : H_{3,6} \rightarrow \mathbb{R}, f \mapsto \sum_{j=1}^9 a_j f(v_j)$, $a_j \in \mathbb{R}$ with respect to A for p to be not SOS if*

$$h^2(v_9)p(v_8) \neq h^2(v_8)p(v_9).$$

Furthermore, l_a is an extreme ray of $\Sigma_{3,6}^*$.

Proof. By Lemma 2.5 there is a $h \in V(\mathcal{V}(p))$ with $\mathcal{V}(h) = \mathcal{V}(p)$. Since A satisfies (2.2) and $h \in H_{3,3}$ the Cayley-Bacharach relation (2.1) holds for h . Since $\mathcal{V}(h) = \mathcal{V}(p) = \{v_1, \dots, v_7\}$ this means $u_8 h(v_8) + u_9 h(v_9) = 0$ and thus $u_8^2 h^2(v_8) = u_9^2 h^2(v_9)$. Hence,

$$u_8^2 p(v_8) \neq u_9^2 p(v_9) \Leftrightarrow h^2(v_9)p(v_8) \neq h^2(v_8)p(v_9).$$

The assertion follows with Theorem 4.1. □

If we want to compute a specific l_a , then the approach of Corollary 4.3 does not suffice, since, by (2.5), we need to know the coefficients u_j of the Cayley-Bacharach relation (2.1) on A to compute the scalar a_9 for l_a .

5. An application: the Motzkin polynomial

In this section we demonstrate applications of our method. It turns out that finding a separating extreme ray becomes more difficult for polynomials in $p \in \partial P_{3,6} \setminus \Sigma_{3,6}$ with six or less zeros. The condition (3.1) provides more degrees of freedom and, in particular, the left hand side of this inequality has more than one summand. This fact yields that the set of point configurations $A := \{v_1, \dots, v_9\} \subset \mathbb{R}^3$ with $\mathcal{V}(p) \subset A$ which do not satisfy (3.1) will not be a lower dimensional subset in general – in contrast to the seven point case (independently from which point corresponds to the negative entry a_9 in an extreme ray). The same difficulties arise for $p \in P_{4,4} \setminus \Sigma_{4,4}$ with five or less zeros.

As an example, we investigate the Motzkin polynomial

$$m(x, y, z) = x^4 y^2 + x^2 y^4 - 3x^2 y^2 z^2 + z^6.$$

The Motzkin polynomial has six zeros

$$\begin{aligned} v_1 &:= (1, 0, 0), & v_2 &:= (0, 1, 0), & v_3 &:= (1, 1, 1), \\ v_4 &:= (-1, 1, 1), & v_5 &:= (1, -1, 1), & v_6 &:= (1, 1, -1). \end{aligned}$$

As a first instance we choose $v_7 := (0, 4, 1)$ and $v_8 := (4, 0, 1)$. These eight points satisfy the genericity condition (2.2). This can be checked by looking at the (3×3) -minors of the (8×3) -matrix given by the coordinates (x, y, z) of the points v_1, \dots, v_8 and looking

at the (6×6) -minors of the (8×6) -matrix given by the coordinates $(x^2, y^2, z^2, xy, xz, yz)$ of the points v_1, \dots, v_8 . Hence we can compute two coprime ternary cubics

$$\begin{aligned} q_1 &= -16z^3 + 15x^2z + y^2z + 56y^2x - 56z^2x \\ q_2 &= -4z^3 - x^2y + 4x^2z + 15y^2x - 15z^2x + z^2y \end{aligned}$$

vanishing on v_1, \dots, v_8 by solving the system of linear equations

$$h(v_1) = 0, \dots, h(v_8) = 0$$

on the coefficients of h . Here,

$$h := b_1x^3 + b_2y^3 + b_3z^3 + b_4x^2y + b_5x^2z + b_6y^2z + b_7y^2x + b_8z^2x + b_9z^2y + b_{10}xyz$$

We compute the Gröbner basis

$$\{7z - 26z^2 - 15z^3 + 26z^4 + 8z^5, -2z + 8z^2 + 2z^3 - 105y - 8z^4 + 105z^2y, \\ 8z^4 - 422z^3 + 1575y^2 - 1583z^2 + 422z, x - 1\}$$

of q_1, q_2 and $x - 1$ with lexicographic ordering. We obtain $v_9 = (1, 1, -7/2)$ and compute the Cayley-Bacharach coefficients u_j by solving the system of linear equations

$$u_1h(v_1) + \dots + u_9h(v_9) = 0$$

in u_1, \dots, u_9 . The solution is (up to scalar multiplication)

$$u = \left(-64, -64, -\frac{40}{9}, -4, -4, \frac{24}{5}, 1, 1, \frac{118098}{5} \right)^T$$

We have

$$\begin{aligned} (u_7^2 + u_8^2) \cdot (m(v_7) + m(v_8)) &= 4, \\ u_9^2 m(v_9) &= 228. \end{aligned}$$

Hence, the condition (3.1) of Theorem 3.1 is satisfied and we find a separating hyperplane for m on A . According to Corollary 3.4, we choose $a_1, \dots, a_6 := 100$ and $a_7, a_8 := 1$. By (2.5) we obtain

$$a_9 = \frac{-u_9^2}{\frac{u_1^2}{a_1} + \dots + \frac{u_8^2}{a_8}} = \frac{-14121476824050}{2143157}.$$

We check the correctness of our result by

$$\begin{aligned} l_a(m) &= a_1m(v_1) + \dots + a_9m(v_9) \\ &= m(0, 4, 1) + m(4, 0, 1) - \frac{14121476824050}{2143157} \cdot m\left(1, 1, -\frac{7}{2}\right) \\ &= -\frac{1484936}{2143157} < 0. \end{aligned}$$

Thus, by Blekherman's Theorem 2.7, l_a is a (rational) extreme ray of $\Sigma_{3,6}^*$ separating the Motzkin polynomial m from the SOS cone.

In contrast to the seven point case, not every generic point configuration yields a separating certificate. For example, with the same approach it is easy to show that the instance $v_7 := (2/7, 2/3, 1)$ and $v_8 := (2/3, 2/7, 1)$ does not satisfy the condition (3.1).

We show that for a symmetric choice of v_7, v_8 , i.e., $v_7 = (q, s, 1), v_8 = (s, q, 1)$ with $q, s \in \mathbb{R}$, the set

$$S := \{(q, s) \in \mathbb{R}^2 : (u_7^2 + u_8^2) \cdot (m(v_7) + m(v_8)) < u_9^2 m(v_9)\}$$

yielding a 9-point configuration, which satisfies (3.1) is full dimensional with some nice geometric structure (see Figure (5)).

With the formula in [21] we obtain the ninth Cayley-Bacharach point

$$v_9 = \frac{1}{n(q, s)} \cdot \begin{pmatrix} q^3 s + 2q^2 s^2 - q^2 + s^3 q - 2sq - s^2 \\ q^3 s + 2q^2 s^2 - q^2 + s^3 q - 2sq - s^2 \\ q^3 + q^2 s - 2q + s^2 q - 2s + s^3 \end{pmatrix},$$

where

$$n(q, s) = 12 \cdot (q^3 s^2 - 2q^3 s + q^3 + s^3 q^2 - 2q^2 s^2 + q^2 s - 3q + 4sq - 2s^3 q + s^2 q + 2 - 3s + s^3)$$

vanishing at $q = -2 - s, q = 1, s = 1$. Furthermore we obtain a non-generic point set for $q = s, q = -s, q = -1, s = -1$ and $q = 2 - s$. We compute the Cayley-Bacharach coefficients in dependence of q, s and obtain

$$\begin{aligned} (u_7^2 + u_8^2) \cdot (m(v_7) + m(v_8)) &= \frac{64(1 + s^4 q^2 + q^4 s^2 - 3q^2 s^2)}{(q - s)^4} \\ u_9^2 m(v_9) &= \frac{16(2q^4 s^2 + q^4 + 4s^3 q^3 - 4q^3 s + 2s^4 q^2 - 6q^2 s^2 - 2q^2 - 4s^3 q + 4sq + s^4 - 2s^2 + 4)}{(q - s)^4}. \end{aligned}$$

Note that the numerator of $(u_7^2 + u_8^2) \cdot (m(v_7) + m(v_8))$ is exactly the dehomogenized Motzkin polynomial in s, q for $z = 1$. We set

$$\begin{aligned} K(q, s)/L(q, s) &:= (u_7^2 + u_8^2) \cdot (m(v_7) + m(v_8)) - u_9^2 m(v_9) \\ &= 16(2q^2 s^2 - q^2 + 2sq - s^2 + 2)/(q - s)^2. \end{aligned}$$

Since $q \neq s$ by assumption we just need to investigate $K(q, s)$. Thus, by (3.1) we have $(q, s) \in S$ if and only if $K(q, s) < 0$ and $q \notin \{\pm s, \pm 1, \pm 2 - s\}$. Equivalently, $S = \emptyset$ if and only if $K(q, s)$ is PSD and $q \notin \{\pm s, \pm 1, \pm 2 - s\}$. Since $K(q, s)$ is a bivariate polynomial of degree 4, $K(q, s)$ is PSD if only if it is SOS. It can be checked easily that this is not the case.

We provide a plot of the set $S := \{(q, s) : K(q, s) < 0\} \setminus \{(q, s) : q = 2 - s \text{ or } q = -2 - s\}$ in Figure 1 on Page 15 (note that the other non-generic cases are not part of S although they are relevant for the computation). Obviously, this set is symmetric in $q = s$ and $q = -s$, semi-algebraic and for every q there is an s such that $(q, s) \in S$.




Figure 1: The set $S := \{(q, s) : K(q, s) < 0\} \setminus \{(q, s) : q = 2 - s \text{ or } q = -2 - s\}$ is given by the red (dark) area without the diagonal blue (dark) lines.

Due to the rich geometric structure of S , it would be of interest to investigate the geometric structure of the set of appropriate point configurations satisfying (3.1) for general nonnegative polynomials with k zeros.

In contrast, we briefly demonstrate the numerical method for finding a 9-point certificate given in Section 2.2 and the corresponding problems. Let $r = (x^2 + y^2 + z^2)^3 \in \text{int}(\Sigma_{3,6})$. Then, consider the following semidefinite optimization problem

$$\min_{\lambda \in \mathbb{R}} \lambda \quad \text{such that} \quad m + \lambda(x^2 + y^2 + z^2)^3 \in \Sigma_{3,6}.$$

The optimal λ is given numerically by $\lambda \approx 0.004596411406567$ and the corresponding sum of squares decomposition of $m + \lambda r \approx s_1^2 + s_2^2 + s_3^2$ is given by

$$m + \lambda(x^2 + y^2 + z^2)^3 \approx (-0.858558xz^2 + 0.941354xy^2 + 0.067796x^3)^2 + (-0.858558yz^2 + 0.067796y^3 + 0.941354x^2y)^2 + (-1.002295z^3 + 0.360838y^2z + 0.360838x^2z)^2.$$

Now, one has to choose an appropriate linear combination of two of the polynomials s_i to obtain the nine intersection points. But it seems unclear how to do this. E.g., since the coefficients of the s_i are given numerically, computing the Gröbner basis of two of the s_i and, say, $x - 1$ cannot be expected to work properly.

Finally, we remark that our method allows to generate strictly positive polynomials that are not sums of squares including a certificate without optimization, if (3.1) is satisfied for a polynomial $p \in \partial P_{3,6} \setminus \Sigma_{3,6}$ (resp. for $(n, 2d) = (4, 4)$). Let l_a be a separating extreme

ray for p and define $n := p + \lambda \cdot r$ with $\lambda \in \mathbb{R}_{>0}$ and $r \in \text{int}(\Sigma_{3,6})$ (resp. $r \in \text{int}(\Sigma_{4,4})$). Then n is strictly positive and evaluating n on l_a yields

$$l_a(n) = l_a(p) + \lambda \cdot l_a(r)$$

with $l_a(p) < 0$ and $l_a(r) > 0$. Hence, we can immediately solve for λ such that $l_a(n) < 0$.

Acknowledgements

We would like to thank Grigoriy Blekherman for his helpful advices. Furthermore, we would like to thank Cordian Riener and Thorsten Theobald for their support during the development of this article. We thank the anonymous referees for their helpful suggestions.

The second author was supported by DFG grant TH 1333/2-1.

References

- [1] Basu, S., Pollack, R., Roy, M.-F., 2006. Algorithms in real algebraic geometry, 2nd Edition. Vol. 10 of Algorithms and Computation in Mathematics. Springer-Verlag, Berlin.
- [2] Blekherman, G., 2004. Convexity properties of the cone of nonnegative polynomials. *Discrete Comput. Geom.* 32 (3), 345–371.
- [3] Blekherman, G., 2006. There are significantly more nonnegative polynomials than sums of squares. *Israel J. Math.* 153, 355–380.
- [4] Blekherman, G., 2012. Nonnegative polynomials and sums of squares. *J. Amer. Math. Soc.* 25 (3), 617–635.
- [5] Blekherman, G., 2012. Positive gorenstein ideals. To appear in Proceedings of the AMS, Preprint, [arXiv:1305.0642](https://arxiv.org/abs/1305.0642).
- [6] Blekherman, G., Hauenstein, J., Ottem, J., Ranestad, K., Sturmfels, B., 2012. Algebraic boundaries of Hilbert’s SOS cones. *Compos. Math.* 148 (6), 1717–1735.
- [7] Blekherman, G., Iliman, S., Kubitze, M., 2013. Dimensional differences between faces of the cone of nonnegative polynomials and sums of squares. Preprint, [arXiv:1305.0642](https://arxiv.org/abs/1305.0642).
- [8] Bochnak, J., Coste, M., Roy, M.-F., 1998. Real algebraic geometry. Vol. 36 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*. Springer-Verlag, Berlin.
- [9] Bosse, H., 2007. Positive Polynomials that are not Sums of Squares. Preprint, oai.cwi.nl/oai/asset/12303/12303D.pdf.

- [10] Choi, M., Lam, T., Reznick, B., 1980. Real zeros of positive semidefinite forms. I. *Math. Z.* 171 (1), 1–26.
- [11] Delzell, C., Prestel, A., 2001. *Positive polynomials*. Springer Monographs in Mathematics. Springer-Verlag, Berlin.
- [12] Eisenbud, D., 2005. *The geometry of syzygies*. Vol. 229 of Graduate Texts in Mathematics. Springer-Verlag, New York.
- [13] Eisenbud, D., Green, M., Harris, J., 1996. Cayley-Bacharach theorems and conjectures. *Bull. Amer. Math. Soc. (N.S.)* 33 (3), 295–324.
- [14] Hilbert, D., 1888. Ueber die Darstellung definiten Formen als Summe von Formenquadraten. *Math. Ann.* 32 (3), 342–350.
- [15] Lasserre, J., 2000/01. Global optimization with polynomials and the problem of moments. *SIAM J. Optim.* 11 (3), 796–817.
- [16] Laurent, M., 2009. Sums of squares, moment matrices and optimization over polynomials. In: *Emerging applications of algebraic geometry*. Vol. 149 of IMA Vol. Math. Appl. Springer, New York, pp. 157–270.
- [17] Marshall, M., 2008. *Positive polynomials and sums of squares*. Vol. 146 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI.
- [18] Motzkin, T., 1983. *Selected papers*. Contemporary Mathematicians. Birkhäuser Boston, Mass.
- [19] Nie, J., 2012. Discriminants and nonnegative polynomials. *J. Symbolic Comput.* 47 (2), 167–191.
- [20] Plaumann, D., Sturmfels, B., Vinzant, C., 2011. Quartic curves and their bitangents. *J. Symbolic Comput.* 46 (6), 712–733.
- [21] Ren, Q., 2011. A formula for the Cayley–Bacharach Theorem, preprint, <http://math.berkeley.edu/~qingchun/>.
- [22] Reznick, B., 2000. Some concrete aspects of Hilbert’s 17th Problem. In: *Real algebraic geometry and ordered structures* (Baton Rouge, LA, 1996). Vol. 253 of *Contemp. Math.* Amer. Math. Soc., Providence, RI, pp. 251–272.
- [23] Reznick, B., 2007. On Hilbert’s Construction of positive Polynomials, preprint, [arXiv: 0707.2156](https://arxiv.org/abs/0707.2156).

Appendix A. An example for the seven point case

As an example of our method we investigate the polynomial

$$p := x^2y^2(x-y)^2 + y^2z^2(y-z)^2 + z^2x^2(z-x)^2 + xyz(x-y)(y-z)(z-x),$$

given by Reznick in [23]. It has the seven zeros

$$\begin{aligned} v_1 &:= (1, 0, 0), & v_2 &:= (0, 1, 0), & v_3 &:= (0, 0, 1), & v_4 &:= (1, 1, 0) \\ v_5 &:= (1, 0, 1), & v_6 &:= (0, 1, 1), & v_7 &:= (1, 1, 1). \end{aligned}$$

We choose $v_8 := (-2, 5, -1)$ which satisfies (2.2). We obtain a ninth intersection point $v_9 = (3, 10, 1)$ by the formula in [21] and compute the Cayley-Bacharach coefficients. According to Corollary 3.4 we choose $a_1, \dots, a_7 := 10^9$ and $a_8 := 1$. By (2.5) we obtain

$$\begin{aligned} a_9 &= \frac{-u_9^2}{\frac{u_1^2}{a_1} + \dots + \frac{u_8^2}{a_8}} = \frac{-10^9}{84^2 + (-1260)^2 + (-36)^2 + (-90)^2 + 63^2 + 35^2 + (-60)^2 + 9 \cdot 10^9} \\ &= -\frac{500000000}{4500806423}. \end{aligned}$$

We check correctness of our result by

$$\begin{aligned} l_a(p) &= a_1p(v_1) + \dots + a_9p(v_9) = p(-2, 5, -1) - \frac{500000000}{4500806423} \cdot p(3, 10, 1) \\ &= -\frac{471957277321}{5626008028750} < 0. \end{aligned}$$