



**HAL**  
open science

## Extension from Precoloured Sets of Edges

Katherine Edwards, Jan van den Heuvel, Ross J. Kang, Jean-Sébastien Sereni

► **To cite this version:**

Katherine Edwards, Jan van den Heuvel, Ross J. Kang, Jean-Sébastien Sereni. Extension from Precoloured Sets of Edges. [Research Report] Loria & Inria Grand Est. 2014. hal-01024843v1

**HAL Id: hal-01024843**

**<https://hal.science/hal-01024843v1>**

Submitted on 16 Jul 2014 (v1), last revised 9 Dec 2019 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Extension from Precoloured Sets of Edges

Katherine Edwards\*      Jan van den Heuvel†      Ross J. Kang‡  
Jean-Sébastien Sereni§

July 16, 2014

## Abstract

We consider precolouring extension problems for proper edge-colourings of graphs and multigraphs, in an attempt to prove stronger versions of Vizing’s and Shannon’s bounds on the chromatic index of (multi)graphs in terms of their maximum degree  $\Delta$ . We are especially interested in the following question: when is it possible to extend a precoloured matching to a colouring of all edges of a (multi)graph? For simple graphs, we conjecture that it is guaranteed as long as the predefined set of available colours has cardinality  $\Delta + 1$ . This turns out to be related to the notorious list colouring conjecture and other classic notions of choosability.

## 1 Introduction

Let  $G = (V, E)$  be a (multi)graph and let  $\mathcal{K} = [K] = \{1, \dots, K\}$  be a palette of available colours. (In this paper, a *multigraph* can have multiple edges, but no loops; while a *graph* is always simple.) We consider the following question: given a subset  $S \subseteq E$  of edges and a proper colouring of elements of  $S$  (i.e., adjacent edges must receive distinct colours) using only colours from  $\mathcal{K}$ , is there a proper colouring of all edges of  $G$  (again using only colours from  $\mathcal{K}$ ) in concordance with the given colouring on  $S$ ? We may consider the set  $S$  as a set of *precoloured edges*, while the full colouring, if it exists, may be considered as *extending the precolouring*. Clearly, if the set  $S$  forms a matching in  $G$ , then the precolouring of  $S$  can be arbitrary from  $\mathcal{K}$ .

An early appearance of a problem regarding precolouring extension of edge-colourings can be found in Marcotte and Seymour [22]. Nevertheless, since then its counterpart for vertex-colourings has been more comprehensively studied. We hope to provoke interest in edge-precolouring extension and in particular introduce the following conjecture.

### Conjecture 1.

Let  $G$  be a multigraph with maximum degree  $\Delta(G)$  and maximum multiplicity  $\mu(G)$ . Using

---

\* Department of Computer Science, Princeton University, Princeton, NJ, USA; [ke@princeton.edu](mailto:ke@princeton.edu).

† Department of Mathematics, London School of Economics, London, UK; [j.van-den-heuvel@lse.ac.uk](mailto:j.van-den-heuvel@lse.ac.uk).

‡ Applied Stochastics, IMAPP, Radboud University, Nijmegen, the Netherlands; [ross.kang@gmail.com](mailto:ross.kang@gmail.com). This author is supported by a NWO Veni grant.

§ CNRS (LORIA), Vandœuvre-lès-Nancy, France; [sereni@kam.mff.cuni.cz](mailto:sereni@kam.mff.cuni.cz). This author’s work was partially supported by the French *Agence Nationale de la Recherche* under reference ANR 10 JCJC 0204 01.




Figure 1: A representative of a class of bipartite planar graphs  $G$ , with a non-extendable precoloured (distance-2) matching using palette  $[\Delta(G)] = [\chi'(G)]$ . Dashed lines indicate edges precoloured with colour 1.

the palette  $\mathcal{K} = [\Delta(G) + \mu(G)]$ , any precoloured matching can be extended to a proper edge-colouring of all of  $G$ .

If true, Conjecture 1 would extend Vizing's theorem [33], which is independently due to Gupta, cf. [15]. It strengthens a conjecture of Albertson and Moore [3, Conj. 1], which deals with graphs and imposes a stronger minimum distance condition between precoloured edges. A weaker form of Conjecture 1 was proved by Berge and Fournier [6, Cor. 2] — they showed that extension is guaranteed, provided that all edges of the matching have been precoloured with the same colour.

Note that even for trees, Conjecture 1 is false if we replace the palette  $\mathcal{K}$  by  $[\Delta(G)]$  or by  $[\chi'(G)]$  (where  $\chi'(G)$  is, as usual, the chromatic index of  $G$ ): consider a star with all leaves subdivided exactly once; see Figure 1.

The main contribution of this paper is to provide evidence in support of Conjecture 1 by confirming it in some important special cases, in particular, for bipartite multigraphs, subcubic multigraphs, and planar graphs of large enough maximum degree. We discuss this further in Subsection 1.1; however, first allow us to place the conjecture in context by giving some preliminary observations.

The obvious relationship between edge-precolouring and its vertex counterpart — in which we can see edge-precolouring extension of  $G$  as vertex-precolouring extension in its line graph  $L(G)$  — yields immediate implications. For us, the distance between two edges in  $G$  is their corresponding distance in  $L(G)$ , i.e., the number of *vertices* contained in a shortest path in  $G$  between any of their end-vertices. A *distance- $t$  matching* is a set of edges having pairwise distance greater than  $t$ . (This means that a matching is a distance-1 matching, while an induced matching is a distance-2 matching. Any set of edges is a distance-0 matching.) We point out the following consequence of a result of Albertson [1, Thm. 4] (see Subsection 1.2) and Vizing's theorem. This consequence is in two senses a weaker statement than that of Conjecture 1.

**Proposition 2.**

Let  $G$  be a multigraph with maximum degree  $\Delta(G)$  and maximum multiplicity  $\mu(G)$ . Using the palette  $\mathcal{K} = [\Delta(G) + \mu(G) + 1]$ , any precoloured distance-3 matching can be extended to a proper edge-colouring of all of  $G$ .

Evidently, we believe that in edge-precolouring the distance requirement ought to be not as strong as it is for vertex-precolouring extension. Naturally, one might wonder if a strong enough distance requirement on the precoloured set yields an improved palette bound, i.e., with palette  $[\Delta(G)]$  or  $[\chi'(G)]$ . This fails however even for bipartite graphs; see Figure 2.




Figure 2: A representative of a class of bipartite graphs  $G$ , with a non-extendable matching consisting of two edges that can be made arbitrarily distant, using the palette  $[\Delta(G)] = [\chi'(G)]$ . Dashed lines indicate edges precoloured with colour 1. (There is a general construction for  $\Delta(G)$  even: within each diamond-shaped block replace each of the two independent sets of size three with one of size  $\frac{1}{2}\Delta(G)$  and replace the central independent set of size five with one of size  $\Delta(G) - 1$ ; next increase the distance between the precoloured edges by replacing the two blocks by a chain of arbitrarily many blocks.)

By the following easy observation, Conjecture 1 is also related to list edge-colouring, and therefore to the *List Colouring Conjecture* (LCC) which states that  $\text{ch}'(G) = \chi'(G)$  for any multigraph  $G$  (where  $\text{ch}'(G)$  is, as usual, the list chromatic index of  $G$ ). For a non-precoloured edge, we define its *precoloured degree* as the number of adjacent precoloured edges.

**Proposition 3.**

Let  $G$  be a multigraph with list chromatic index  $\text{ch}'(G)$ . For a positive integer  $k$ , take the palette as  $\mathcal{K} = [\text{ch}'(G) + k]$ . If  $G$  is properly precoloured so that the precoloured degree of any non-precoloured edge is at most  $k$ , then the precolouring can be extended to a proper edge-colouring of all of  $G$ .

So, if we assume that the LCC holds, then the following weak form of Conjecture 1 holds as well: any precoloured matching can be extended to a proper edge-colouring of all of  $G$  with the palette  $\mathcal{K} = [\Delta(G) + \mu(G) + 2]$ ; moreover, if the precoloured matching is induced, then we may use  $\mathcal{K} = [\Delta(G) + \mu(G) + 1]$ .

Due to the remarkable work of Kahn [19, 20, 21] on edge-colourings and list edge-colourings of (multi)graphs, not only does an asymptotic form of Conjecture 1 hold, but so does a precolouring extension of an asymptotic form of the Goldberg–Seymour Conjecture (which we review in Subsection 1.2). Kahn’s theorem and Proposition 3 together imply the following.

**Proposition 4.**

For any  $\varepsilon > 0$ , there exists a constant  $C_\varepsilon$  such that the following holds. For any multigraph  $G$  with  $\chi'(G) \geq C_\varepsilon$ , any precoloured matching using the palette  $\mathcal{K} = [(1+\varepsilon)\chi'(G)]$  can be extended to a proper edge-colouring of all of  $G$ .

If we replace  $\chi'(G)$  in the statement above by  $\Delta(G) + \mu(G)$  or by the Goldberg–Seymour bound, then the statement remains valid, either due to Vizing’s theorem or due to another theorem of Kahn.

One of our motivations for the formulation and study of Conjecture 1 comes from the close connections with vertex-precolouring and with the LCC.

## 1.1 Main Results

Although it appears that the LCC and our conjecture are independent statements, we have obtained several results corresponding to specific areas of success in list edge-colouring. For clarity, in this subsection we mostly present our main results restricted to precoloured matchings, but more general statements hold. In summary, we have confirmed Conjecture 1 for bipartite multigraphs, subcubic multigraphs, and planar graphs of large enough maximum degree; we have also obtained a precolouring extension variant of Shannon's theorem.

Our first result is an edge-precolouring extension of Kőnig's theorem that any bipartite multigraph  $G$  is  $\Delta(G)$ -edge-colourable, whereas the subsequent result is an edge-precolouring analogue of Shannon's theorem that any multigraph  $G$  is  $\lfloor \frac{3}{2}\Delta(G) \rfloor$ -edge-colourable.

### Theorem 5.

*Let  $G$  be a bipartite multigraph with maximum degree  $\Delta(G)$ . With the palette  $\mathcal{K} = [\Delta(G) + 1]$ , any precoloured matching can be extended to a proper edge-colouring of all of  $G$ .*

As we saw in Figures 1 and 2, the palette size in Theorem 5 is sharp.

### Theorem 6.

*Let  $G$  be a multigraph with maximum degree  $\Delta(G)$ . With the palette  $\mathcal{K} = [\lfloor \frac{3}{2}\Delta(G) + \frac{1}{2} \rfloor]$ , any precoloured matching can be extended to a proper edge-colouring of all of  $G$ .*

Due to the Shannon multigraphs, this last statement is sharp if  $\Delta(G)$  is even, and within 1 of being sharp if  $\Delta(G)$  is odd. Theorems 5 and 6 are proved in Section 2 using powerful list colouring tools developed by Borodin, Kostochka and Woodall [10].

The following theorem concerns multigraphs that are subcubic, i.e. of maximum degree at most 3. Note that Theorem 7 improves upon Theorem 6 for  $\Delta(G) = 3$ .

### Theorem 7.

*Let  $G$  be a subcubic multigraph. With the palette  $\mathcal{K} = [4]$ , any precoloured matching can be extended to a proper edge-colouring of all of  $G$ .*

A form of Theorem 7, for subcubic graphs and with a distance condition on the precoloured matching, was observed by Albertson and Moore [3]. Although the LCC remains open for subcubic graphs, Juvan, Mohar and Škrekovski [17] have made a significant attempt. (They showed that for any subcubic graph  $G$ , if lists of 3 colours are given to the edges of a subgraph  $H$  with  $\Delta(H) \leq 2$  and lists of 4 colours to the other edges, then  $G$  has a proper edge-colouring using colours from those lists.)

Theorem 7 is a direct corollary of the following theorem, which may be of interest in its own right. Its proof uses a degree-choosability condition and can be found in Section 3. For a vertex, we define its *precoloured degree* as the number of incident precoloured edges.

### Theorem 8.

*Let  $G$  be a connected multigraph. Choose a non-negative integer  $k$  so that  $\Delta(L(G)) \leq \Delta(G) + k$ , and take the palette as  $\mathcal{K} = [\Delta(G) + k]$ . If  $G$  is properly precoloured so that the precoloured degree of any vertex is at most  $k$ , then the precolouring can be extended to a proper edge-colouring of all of  $G$ , except in the following cases:*

- (a)  $k = 0$  and  $G$  is a simple odd cycle;
- (b)  $G$  is a triangle with edges of multiplicity  $m_1, m_2, m_3$  and  $k = \min\{m_1, m_2, m_3\} - 1$ .

	Palette $\mathcal{K}$	Distance $t$	Max. degree $\Delta$	Reference
1.	$[\Delta + 4]$	1	all $\Delta$	Thm. 6 ( $\Delta \leq 7$ ) and Prop. 3 with [7] ( $\Delta \geq 8$ )
2.	$[\Delta + 3]$	1	$\Delta \leq 5$ ; $\Delta \geq 8$	Thm. 6; Prop. 3 with [7]
3.	$[\Delta + 2]$	1	$\Delta \geq 12$	Prop. 3 with [10]
4.	$[\Delta + 2]$	2	$\Delta \leq 4$ ; $\Delta \geq 8$	[18]; Prop. 3 with [7]
5.	$[\Delta + 2]$	3	all $\Delta$	Prop. 2
6.	$[\Delta + 1]$	1	$\Delta \leq 3$ ; $\Delta \geq 17$	Thm. 7; Thm. 9
7.	$[\Delta + 1]$	2	$\Delta \geq 12$	Prop. 3 with [10]
8.	$[\Delta]$	3	$\Delta \geq 20$	Thm. 10

Table 1: Summary of edge-precolouring extension results for planar graphs with maximum degree  $\Delta$ , when a distance- $t$  matching  $M$  is precoloured using the palette  $\mathcal{K}$ . See Section 4 for further details how these results can be obtained.

Note that, when restricted to precoloured matchings, this theorem produces weak or limited bounds for larger maximum degree. On the other hand, if we replace every precoloured edge in the example of Figure 1 by a precoloured multi-edge of multiplicity  $k$  (or  $k + 1$ ) and a precolouring from  $[k]$  (or  $[k + 1]$ ), we see that the palette bound (or precoloured degree condition) is best possible.

In the case where  $k = \Delta(L(G)) - \Delta(G)$  in Theorem 8, the number of colours used is equal to the maximum degree of the line graph. In that sense the theorem can be considered as a precolouring extension of Brooks's theorem restricted to line graphs. It is relevant to mention that vertex-precolouring extension versions of Brooks's theorem [2, 5, 25] require, among other conditions, a large minimum distance between the precoloured vertices.

The class of planar graphs could be of particular interest. There is a prominent line of work on (list) edge-colouring for this class, which we discuss further in Subsection 1.2 and Section 4. Our main contributions to this area are the following results, the second one of which can be viewed as a strengthening of an old result of Vizing [34], provided the graph's maximum degree is large enough.

**Theorem 9.**

*Let  $G$  be a planar graph with maximum degree  $\Delta(G) \geq 17$ . Using the palette  $\mathcal{K} = [\Delta(G) + 1]$ , any precoloured matching can be extended to a proper edge-colouring of all of  $G$ .*

**Theorem 10.**

*Let  $G$  be a planar graph with maximum degree  $\Delta(G) \geq 20$ . Using the palette  $\mathcal{K} = [\Delta(G)]$ , any precoloured distance-3 matching can be extended to a proper edge-colouring of all of  $G$ .*

Due to the trees exhibited in Figure 1, the palette size in Theorem 9 cannot be reduced, while the minimum distance condition in Theorem 10 cannot be weakened. In Section 4, we give some more results on when it is possible for a precoloured matching in a planar graph to be extended. A summary of the results is given in Table 1.

Some basic knowledge of edge-colouring is a prerequisite to the consideration of edge-precolouring extension problems — we provide some related background in the next subsection. To our frustration, many of the major methods for colouring edges (such as Kempe

chains, Vizing fans, Kierstead paths, Tashkinov trees) seem to be rendered useless by pre-coloured edges. Though Conjecture 1 may at first seem as if it should be an “easy extension” of Vizing’s theorem, it might well be very difficult to confirm (if true at all). We are keen to learn of related edge-precolouring results independent of current list colouring methodology.

## 1.2 Further Background

Edge-colouring is a classic area of graph theory. We give a quick overview of some of the most relevant history for our study. The reader is referred to the recent book by Stiebitz, Scheide, Toft and Favrholt [30] for detailed references and fuller insights. The lower bound  $\chi'(G) \geq \Delta(G)$  is obviously true for any multigraph  $G$ . Close to a century ago, König proved that all bipartite multigraphs meet this lower bound with equality. Shannon [29] in 1949 proved that  $\chi'(G) \leq \lfloor \frac{3}{2}\Delta(G) \rfloor$  for any multigraph  $G$ . Somewhat later, Gupta (as mentioned in [15]) and, independently, Vizing [33] proved that  $\chi'(G) \leq \Delta(G) + \mu(G)$  for any multigraph  $G$ , so  $\chi'(G) \in \{\Delta(G), \Delta(G) + 1\}$  if  $G$  is simple. Both the Shannon bound and the Gupta–Vizing bound are tight in general due to the Shannon multigraphs, which are triangles whose multi-edges have balanced multiplicities. (Note however that the latter bound can be improved for specific choices of  $\Delta(G)$  and  $\mu(G)$  as described in the work of Scheide and Stiebitz [27].)

A notable conjecture on edge-colouring arose in the 1970s, on both sides of the iron curtain. The Goldberg–Seymour Conjecture, due independently to Goldberg [14] and Seymour [28], asserts that  $\chi'(G) \in \{\Delta(G), \Delta(G) + 1, \lceil \rho(G) \rceil\}$  for any multigraph  $G$ , where

$$\rho(G) = \max \left\{ \frac{2|E(G[T])|}{|T| - 1} : T \subseteq V, |T| \geq 3, |T| \text{ odd} \right\}.$$

The parameter  $\rho(G)$  is a lower bound on  $\chi'(G)$  based on the maximum ratio between the number of edges in  $H$  and the number of edges in a maximum matching of  $H$ , taken over induced subgraphs  $H$  of  $G$ . This conjecture remains open and is regarded as one of the most important problems in chromatic graph theory. Perhaps the most outstanding progress on this problem is due to Kahn, who established an asymptotic form [20].

The list variant of edge-colouring can be traced as far back as list colouring itself. The concept of list colouring was devised independently by Vizing [35] and Erdős, Rubin and Taylor [11], with the iron curtain playing its customary role here too. The List Colouring Conjecture (LCC) was already formulated by Vizing as early as 1975 and was independently reformulated several times, a brief historical account of which is given in, e.g., Häggkvist and Janssen [16]. For more on the LCC, particularly with respect to the probabilistic method, consult the monograph of Molloy and Reed [24]. The results on the LCC most relevant to our investigations also happen to be two of the most striking, both from the mid-1990s. First, Galvin [13] used a beautiful short argument to prove Dinitz’s Conjecture (which concerned the extension of partially completed Latin squares), thereby confirming the LCC for bipartite multigraphs. Not long after Galvin’s work, Kahn applied powerful probabilistic methods, with inspiration from extremal combinatorics and statistical physics, to asymptotically affirm the LCC [20, 21]. For more background on Kahn’s proof, related methods, and improvements, consult [16, 23, 24].

Inspiration for this class of problems may also be taken from list vertex-colouring. For instance, we utilise a degree-choosability criterion due independently to Borodin [8] and Erdős,

Rubin and Taylor [11]. See for example a survey of Alon [4] for an excellent (if older) survey on list colouring in somewhat more generality. We should mention that part of the motivation for studying list colouring was to use it to attack other, less constrained colouring problems. The connection has gone back in the other direction as well, as precolouring extension demonstrates.

Activity in the area of precolouring extension increased dramatically as a result of the startling proof by Thomassen of planar 5-choosability [31]; a key ingredient in that proof was a particular type of precolouring extension from some pair of adjacent vertices, according to a specific planar embedding. A little bit later, Thomassen asked about precolouring extension for planar graphs under a more general setup [32]. Eliding the planarity condition, Albertson [1] quickly answered Thomassen's question and proved more: in any  $k$ -colourable graph, for any set of vertices with pairwise minimum distance at least 4, any precolouring of that set from the palette  $[k + 1]$  can be extended to a proper colouring of the entire graph. (This implies Proposition 2 above.)

Since Albertson's seminal work, a large body of research has developed around precolouring extension. But this research has focused almost exclusively on extension of vertex-colourings. One of the few papers we are aware of that deals with edge-precolouring extension is by Marcotte and Seymour [22], in which a different type of necessary condition for extension is studied — curiously, this paper predates the above mentioned activity in vertex-precolouring.

For planar graphs, there has been significant interest in both edge-colouring and list edge-colouring. It is known that planar graphs  $G$  with  $\Delta(G) \geq 7$  satisfy  $\chi'(G) = \Delta(G)$ . This was proved in 1965 by Vizing [34] in the case  $\Delta(G) \geq 8$ , and much later by Sanders and Zhao [26] for  $\Delta(G) = 7$ . We remark that Theorem 10 strengthens this for  $\Delta(G)$  somewhat larger. Vizing conjectured that the same can be said for planar graphs  $G$  with  $\Delta(G) = 6$ , but this long-standing question remains open. Vizing also noted that not every planar graph  $G$  with  $\Delta(G) \in \{4, 5\}$  is  $\Delta(G)$ -edge-colourable. Regarding list edge-colouring, Borodin, Kostochka and Woodall [10] proved the LCC for planar graphs with maximum degree at least 12, i.e., they proved that such graphs have list chromatic index equal to their maximum degree. The LCC remains open for planar graphs with smaller maximum degree, though it is known that if  $\Delta(G) \leq 4$ , or  $\Delta(G) \geq 8$ , then  $\text{ch}'(G) \leq \Delta(G) + 1$ . This is due to Juvan, Mohar and Škrekovski [18], and to Bonamy [7] and Borodin [9], respectively. As noted above, it is not true that planar graphs  $G$  with  $\Delta(G) \in \{4, 5\}$  are always  $\Delta(G)$ -edge-choosable.

## 2 Extensions of König's and Shannon's Theorems

Theorem 11 below implies Theorem 5, and hence verifies Conjecture 1 for bipartite multigraphs. Theorem 12 implies Theorem 6. Recall that, for a vertex, we define its precoloured degree as the number of incident precoloured edges.

### **Theorem 11.**

*Let  $G$  be a bipartite multigraph and  $k \geq 1$ . Take the palette as  $\mathcal{K} = [\Delta(G) + k]$ . If  $G$  is properly precoloured so that the precoloured degree of any vertex is at most  $k$ , then this precolouring can be extended to a proper edge-colouring of all of  $G$ .*

**Theorem 12.**

Let  $G$  be a multigraph and  $k \geq 1$ . Take the palette as  $\mathcal{K} = \lfloor \lfloor \frac{3}{2}\Delta(G) + \frac{1}{2}k \rfloor \rfloor$ . If  $G$  is properly precoloured so that the precoloured degree of any vertex is at most  $k$ , then this precolouring can be extended to a proper edge-colouring of all of  $G$ .

The two results are corollary to two theorems of Borodin, Kostochka and Woodall [10]. Let  $G$  be a multigraph and  $f: E(G) \rightarrow \mathbb{Z}^+$ . We say  $G$  is *f-edge-choosable* if, for any assignment of lists where every edge  $e$  receives a list of size at least  $f(e)$ , there is a proper edge-colouring of  $G$  using colours from the lists.

**Theorem 13** (Borodin, Kostochka & Woodall [10]).

Let  $G$  be a bipartite multigraph, and set  $f(uv) = \max\{d(u), d(v)\}$  for each edge  $uv \in E(G)$ . Then  $G$  is *f-edge-choosable*.

**Theorem 14** (Borodin, Kostochka & Woodall [10]).

Let  $G$  be a multigraph, and set  $f(uv) = \max\{d(u), d(v)\} + \lfloor \frac{1}{2} \min\{d(u), d(v)\} \rfloor$  for each edge  $uv \in E(G)$ . Then  $G$  is *f-edge-choosable*.

Note that Theorem 13 is a strengthening of Galvin's theorem; while Theorem 14 is a list colouring version of Shannon's theorem (and in fact follows from Theorem 13).

*Proof of Theorems 11 and 12.* For a vertex  $v$  in a properly precoloured multigraph  $G$ , let  $k(v)$  be the number of precoloured edges incident with  $v$ . Let  $G'$  be obtained from  $G$  by deleting all precoloured edges. To each edge  $e \in E(G')$ , assign a list  $\ell(e)$  consisting of those colours in  $\mathcal{K}$  that do not appear on precoloured edges adjacent to  $e$  in  $G$ .

Next consider any uncoloured edge  $e = uv$ , and assume that  $d_G(u) - k(u) = d_{G'}(u) \leq d_{G'}(v) = d_G(v) - k(v)$ .

In the bipartite case, since  $\Delta(G) \geq d_G(v)$  and  $k \geq k(u)$ , we infer that

$$\begin{aligned} |\ell(e)| &\geq (\Delta(G) + k) - k(u) - k(v) = (\Delta(G) - k(v)) + (k - k(u)) \\ &\geq d_{G'}(v) = \max\{d_{G'}(u), d_{G'}(v)\}. \end{aligned}$$

As this holds for every edge  $e = uv$  in  $G'$ , Theorem 13 guarantees a colouring of the edges of  $G'$  from their lists. This colouring is an extension of the precolouring of  $G$ , completing the proof of Theorem 11.

In the general case, again because  $\Delta(G) \geq \max\{d_G(u), d_G(v)\}$  and  $k \geq k(u)$ , we infer that

$$\begin{aligned} |\ell(e)| &\geq \lfloor \frac{3}{2}\Delta(G) + \frac{1}{2}k \rfloor - k(u) - k(v) = (\Delta(G) - k(v)) + \lfloor \frac{1}{2}\Delta(G) + \frac{1}{2}k - k(u) \rfloor \\ &\geq d_{G'}(v) + \lfloor \frac{1}{2}(\Delta(G) - k(u)) \rfloor \geq d_{G'}(v) + \lfloor \frac{1}{2}d_{G'}(u) \rfloor \\ &= \max\{d_{G'}(u), d_{G'}(v)\} + \lfloor \frac{1}{2} \min\{d_{G'}(u), d_{G'}(v)\} \rfloor. \end{aligned}$$

Analogously to the previous paragraph, we use Theorem 14 to complete the proof of Theorem 12.  $\square$

### 3 An Approach using Gallai Trees

In this section, we use a slight refinement of a result due independently to Borodin [8] and Erdős, Rubin and Taylor [11]. This is a list version of an older result of Gallai [12] on colour-critical graphs, and it follows for instance from Theorem 4.2 in [32]. A connected graph all of whose blocks are either complete graphs or odd cycles is called a *Gallai tree*.

**Theorem 15** (Borodin [8], Erdős, Rubin & Taylor [11]).

*Given a connected graph  $G = (V, E)$ , let  $\ell(v)$ , for  $v \in V$ , be an assignment of lists where each vertex  $v$  receives at least  $d(v)$  colours. Then there is a proper colouring of  $G$  using colours from the lists, unless  $G$  is a Gallai tree and  $|\ell(v)| = d(v)$  for all  $v$ .*

With this we prove Theorem 8, which implies Theorem 7.

*Proof of Theorem 8.* Assume to the contrary that the connected multigraph  $G$  and the non-negative integer  $k$  satisfy  $\Delta(L(G)) \leq \Delta(G) + k$ , but that, using the palette  $\mathcal{K} = [\Delta(G) + k]$ , there is a proper edge-precolouring of  $G$  of the required type that does not extend to a proper edge-colouring of  $G$ . For a vertex  $v$ , let  $K(v) \subseteq \mathcal{K}$  be the set of colours appearing on the precoloured edges incident with  $v$ , and set  $k(v) = |K(v)|$ .

Let  $G'$  be obtained from  $G$  by deleting all precoloured edges. To each edge  $e = uv$  in  $G'$ , we assign a list  $\ell(e)$  containing those colours in  $\mathcal{K}$  not appearing on precoloured edges adjacent to  $e$  in  $G$ . For any edge  $e = uv$  in  $G'$  we obtain, using that  $\Delta(G) + k \geq \Delta(L(G)) \geq d_{L(G)}(e)$ ,

$$\begin{aligned} |\ell(e)| &= |\mathcal{K}| - |K(u) \cup K(v)| = (\Delta(G) + k) - |K(u) \cup K(v)| \\ &\geq d_{L(G)}(e) - |K(u) \cup K(v)| = d_{L(G')}(e). \end{aligned} \tag{1}$$

Since there is no extension of the precolouring of  $L(G)$  to a full colouring of  $L(G)$ , it follows that  $L(G')$  is not edge-choosable with the lists  $\ell(e)$ , for  $e \in E(G')$ . In particular, there is a component  $C'$  of  $G'$  such that  $L(C')$  is not edge-choosable with the lists  $\ell(e)$ , for  $e \in E(C')$ . By Theorem 15,  $L(C')$  must be a Gallai tree such that  $|\ell(e)| = d_{L(C')}(e)$  for every  $e$ . This also means that we must have equality in all inequalities used to derive (1); in particular:

$$\forall e \in E(C'), \quad d_{L(G)}(e) = \Delta(L(G)) = \Delta(G) + k; \tag{2a}$$

$$\forall e \in E(C'), \quad |\ell(e)| = d_{L(C')}(e) = d_{L(G)}(e) - |K(u) \cup K(v)|. \tag{2b}$$

Now note that  $d_{C'}(v) + k(v) = d_G(v) \leq \Delta(G)$  for every vertex  $v$ . So, analogously to (1) above, we infer that for each edge  $e = uv$  in  $C'$  the order of  $\ell(e)$  is at least the degree in  $C'$  of each of its end-vertices:

$$\begin{aligned} |\ell(e)| &= (\Delta(G) + k) - |K(u) \cup K(v)| = (\Delta(G) + k) - k(u) - k(v) + |K(u) \cap K(v)| \\ &\geq d_{C'}(v) + (k - k(u)) + |K(u) \cap K(v)| \geq d_{C'}(v). \end{aligned} \tag{3}$$

We require the following statements.

**Claim 1.** *Every vertex in  $C'$  has at least two neighbours.*

*Proof.* Suppose to the contrary that the vertex  $u$  has the vertex  $v$  as its unique neighbour. Then for the edge  $e = uv$  we have  $d_{L(C')}(e) = d_{C'}(v) - 1$  (this holds even if  $uv$  is a multi-edge). But since  $|\ell(e)| = d_{L(C')}(e)$ , this gives  $|\ell(e)| < d_{C'}(v)$ , contradicting (3).  $\square$

**Claim 2.** *If  $C'$  is a simple odd cycle, then  $k = 0$  and  $G = C'$ .*

*Proof.* Suppose that  $C'$  is a simple odd cycle. If  $e = uv$  is an edge in  $C'$ , then  $|\ell(e)| = d_{L(C')}(e) = d_{C'}(v) = 2$ . From this we can assume, by permuting the colours, that  $\ell(e) = \{1, 2\}$  for every  $e \in L(C')$ . (Indeed, the only way to assign lists of length 2 to the edges of an odd cycle in such a way that there is no proper colouring of the cycle using colours from the lists is by making all lists identical.) There must also be equality everywhere in (3). Combining that with (2b) means in particular that for every edge  $e = uv$  we have  $K(u) \cup K(v) = \{3, 4, \dots, \Delta(G) + k\}$  and  $K(u) \cap K(v) = \emptyset$ . By an easy parity argument, we can see that this is only possible if all the sets  $K(u)$ , for  $u \in V(G)$ , are empty. This means that  $k = 0$ . Since  $G$  is connected, if there are no precoloured edges, then  $G$  can have only one component, which must be  $C'$ .  $\square$

We continue by considering the case that  $C'$  is not an odd cycle. Since line graphs are claw-free, it follows that odd cycle blocks of length at least five are impossible in  $L(C')$ . We deduce that all blocks of  $L(C')$  are cliques. The only way that a leaf block  $B$  of  $L(C')$  could be part of a nontrivial block structure is if it corresponds to a set of edges in  $C'$  that are all incident with a unique vertex, with one of the edges corresponding to the cut-vertex of  $B$ . This is ruled out by Claim 1. We conclude that  $L(C')$  must itself be a clique. In turn, the only way that a line graph  $L(C')$  of a multigraph is a clique is if  $C'$  is a star or a triangle, with possibly multiple edges. The first option is ruled out by Claim 1, so  $C'$  must be a triangle, possibly with multi-edges.

Let  $u, v, w$  be the vertices in  $C'$  and set  $m = |E(C')|$ . Then for all  $e \in E(C')$  we have  $|\ell(e)| = d_{L(C')}(e) = m - 1$ . It is easy to check that with lists of this size, the only way that  $C'$  is not edge-choosable is if all the lists are the same. This also means that the sets  $K(u) \cup K(v)$ ,  $K(u) \cup K(w)$  and  $K(v) \cup K(w)$  are the same.

Let  $A(u)$  be the set of colours that appear on precoloured edges which are incident with  $u$ , but not with  $v$  or  $w$ ; define  $A(v)$  and  $A(w)$  analogously. (In other words, these are colours on the edges that connect  $C'$  to the rest of the graph  $G$ .) Let  $D$  be the set of colours that appear on precoloured edges with end-vertices contained in  $\{u, v, w\}$ . From (2a) and (2b) we deduce that  $|\ell(e)| = d_{L(C')}(e)$  for every edge  $e$  in  $C'$ , which, applied to an edge between  $u$  and  $v$ , implies that  $A(u) \cap A(v) = \emptyset$ ,  $A(u) \cap D = \emptyset$  and  $A(v) \cap D = \emptyset$ . By symmetry,  $A(v) \cap A(w) = \emptyset$ ,  $A(u) \cap A(w) = \emptyset$  and  $A(w) \cap D = \emptyset$ .

Now recall that all edges in  $C'$  must have the same list. Consequently, the disjointness of the sets  $A(u)$ ,  $A(v)$  and  $A(w)$  implies that these three sets are empty. Thus we find that there are no precoloured edges between any of  $u, v, w$  and the rest of the graph. Since  $G$  is connected, it follows that  $V(G) = \{u, v, w\}$ . Let  $m(uv), m(uw), m(vw)$  be the multiplicities of the edges of  $G$ . Then  $\Delta(L(G)) = m(uv) + m(uw) + m(vw) - 1$ , while  $\Delta(G) = m(uv) + m(uw) + m(vw) - \min\{m(uv), m(uw), m(vw)\}$ . Since  $\Delta(L(G)) = \Delta(G) + k$ , we have shown that part (b) of the statement of the theorem holds, completing the proof.  $\square$

## 4 Planar Graphs

In this section, for brevity we usually write  $\Delta$  for  $\Delta(G)$ .

In the next subsection we prove Conjecture 1 for planar graphs of large enough maximum degree (at least 17), which is the assertion of Theorem 9. As mentioned earlier, the LCC is known to hold for planar graphs with maximum degree at least 12. This is yet another result of Borodin, Kostochka and Woodall [10]: they indeed show that  $\text{ch}'(G) \leq \Delta$  for such graphs  $G$ . Combining this with Proposition 3 gives the bounds in lines 3 and 7 of Table 1.

Borodin [9] showed  $\text{ch}'(G) \leq \Delta + 1$  for planar graphs  $G$  of maximum degree  $\Delta \geq 9$ . Recently, Bonamy [7] extended this last statement to the case  $\Delta = 8$ . Combining this result with Proposition 3 implies that for planar graphs with maximum degree  $\Delta \geq 8$  a precoloured matching can be extended to a proper colouring of the entire graph with the palette  $[\Delta + 3]$ , while a precoloured distance-2 matching can be extended with the palette  $[\Delta + 2]$ .

For smaller values of  $\Delta$ , we can use Theorems 6 and 7, and the result of Juvan, Mohar and Škrekovski [18] that  $\text{ch}'(G) \leq \Delta(G) + 1$  for a planar graph  $G$  with  $\Delta(G) \leq 4$ , to achieve several of the bounds in Table 1. In particular, it follows that  $\Delta + 4$  colours suffice for any planar graph with maximum degree  $\Delta$ .

The final proof we present is of Theorem 10. As discussed in Subsection 1.2, Vizing conjectured [34] that any planar graph with maximum degree  $\Delta \geq 6$  has a  $\Delta$ -edge-colouring. The examples in Figure 1 show that this statement is false if we allow an adversarial precolouring of a distance-2 matching. But does it remain true with the adversarial precolouring of any distance-3 matching? We prove that this is indeed the case if  $\Delta \geq 20$ . We expect that this lower bound on  $\Delta$  can be reduced, though, as noted before, certainly not below 6.

The proofs of Theorems 9 and 10 can be found in the next two subsections. They use a common framework, terminology and notation, which we outline now.

Whenever considering a planar graph  $G$ , we fix a drawing of  $G$  in the plane. (So we really should talk about a *plane graph*.) Because of this fixed embedding we can talk about the *faces* of the graph. If  $G$  is connected, then the boundary of any face  $f$  forms a closed walk  $W_f$ .

We adopt the following notation to classify vertices of a graph  $G$  according to degree and incidence with vertices of degree 1. Let  $V_i$  be the set of vertices of degree  $i$ . Also, identify by  $T_i \subseteq V_i$  those vertices of degree  $i$  that are adjacent to a vertex of degree 1, and set  $U_i = V_i \setminus T_i$ . Write  $T = \cup_{i \geq 1} T_i$  and  $U = V(G) \setminus T$ . We also adopt the shorthand notation  $V_{[i,j]}$ ,  $U_{[i,j]}$  and  $T_{[i,j]}$  to mean, respectively, the sets of vertices in  $V$ ,  $U$  and  $T$  with degrees between  $i$  and  $j$  inclusively.

#### 4.1 Proof of Theorem 9

The statement of Theorem 9 is true for graphs with maximum degree 17 and exactly 17 edges. We use induction on  $E(G)$ , and proceed with the induction step. We may easily assume that  $G$  is connected with at least 18 vertices, since  $\Delta \geq 17$ . Let  $M$  be a precoloured matching.

We first observe that

$$\text{if } uv \in E(G) \setminus M, \text{ then } d(u) + d(v) \geq \Delta + 3. \quad (4)$$

Indeed, suppose that the inequality does not hold for some edge  $uv \notin M$ . Then, by induction,  $M$  can be extended to a colouring of  $G - uv$  so that at most  $\Delta$  colours are used on the edges adjacent to  $uv$ , and so we can easily extend the colouring further to  $uv$ . It follows from this

observation that  $G$  has no vertices of degree 2, that every vertex with degree 1 is incident with an edge of  $M$  and that any vertex has at most one neighbour of degree 1. We will use these facts often without reference in the remainder of the proof.

For a face  $f$ , let  $V^-(f) = V(f) \setminus V_1$ , and denote by  $W_f^-$  the sequence of vertices on the boundary walk  $W_f$  after removing vertices from  $V_1$ . For a vertex  $v$ , let  $v_1, v_2, \dots, v_{d(v)}$  be the neighbours of  $v$ , listed in clockwise order according to the drawing of  $G$ . Write  $f_i$  for the face incident with  $v$  lying between the edges  $vv_i$  and  $vv_{i+1}$  (taking addition modulo  $d(v)$ ).

If  $v \in T$  has a (unique) neighbour in  $V_1$ , then we always choose  $v_1$  to be this neighbour. In that case we have  $f_{d(v)} = f_1$ ; we denote that face by  $f_1$  again. Note that it is possible for other faces to be the same as well (if  $v$  is a cut-vertex), but we will not identify those multiple names of the same face. So, if  $v \in U$ , then the faces around  $v$  in consecutive order are  $f_1, f_2, \dots, f_{d(v)}$ ; while, if  $v \in T$ , then the faces around  $v$  are  $f_1, f_2, \dots, f_{d(v)-1}$ .

**Claim 3.**  $|V_\Delta| > |V_3|$ .

*Proof.* Consider the set  $F$  of edges in  $E(G) \setminus M$  with one end-vertex in  $V_3$  and the other in  $V_\Delta$ . The subgraph with vertex set  $V_3 \cup V_\Delta$  and edge set  $F$  is bipartite; we claim it is acyclic. For suppose there exists an (even) cycle  $C \in F$ . By induction, we can extend the precolouring of  $E(G) \setminus M$  to  $G - C$ . But then we can further extend this colouring to the edges of  $C$ , since each edge of  $C$  is adjacent to only  $\Delta - 1$  coloured edges, and even cycles are 2-edge-choosable.

Since each vertex in  $V_3$  is incident with at least two edges in  $F$ , we have  $|V_\Delta| + |V_3| > |F| \geq 2|V_3|$ . The claim follows.  $\square$

We use a discharging argument to continue the proof of the theorem. First, let us assign to each vertex  $v$  a charge

$$\alpha 1: \quad \alpha(v) = 3d(v) - 6,$$

and to each face  $f$  a charge

$$\alpha 2: \quad \alpha(f) = -6.$$

For each vertex  $v$  we define  $\beta(v)$  as follows.

$$\beta 1: \quad \text{If } v \in V_\Delta, \text{ then } \beta(v) = -2.$$

$$\beta 2: \quad \text{If } v \in V_3, \text{ then } \beta(v) = 2.$$

$$\beta 3: \quad \text{In all other cases, } \beta(v) = 0.$$

For each vertex  $v$  and edge  $e = vu$ , we define  $\gamma_e(v)$  and  $\gamma_e(u)$  as follows.

$$\gamma 1: \quad \text{If } v \in V_1, \text{ then } \gamma_e(v) = -\gamma_e(u) = 3.$$

$$\gamma 2: \quad \text{If } v, u \notin V_1, \text{ then } \gamma_e(v) = \gamma_e(u) = 0.$$

Finally, for each face  $f$  and vertex  $v \in W_f^-$  we define  $\delta_f(v)$  and  $\delta_v(f)$  as follows.

$$\delta 1: \quad \text{If } v \in T_3, \text{ then } \delta_v(f) = -\delta_f(v) = 1.$$

$$\delta 2: \quad \text{If } v \in U_3, \text{ then } \delta_v(f) = -\delta_f(v) = \frac{5}{3}.$$

$$\delta 3: \quad \text{If } v \in T \text{ and } 4 \leq d(v) \leq \Delta - 2, \text{ then } \delta_v(f) = -\delta_f(v) = 3 - \frac{6}{d(v) - 1}.$$

$$\delta 4: \quad \text{If } v \in U \text{ and } 4 \leq d(v) \leq \Delta - 2, \text{ then } \delta_v(f) = -\delta_f(v) = 3 - \frac{6}{d(v)}.$$

- $\delta 5$ : If  $d(v) \geq \Delta - 1$ ,  $|V^-(f)| = 3$ , and both neighbours of  $v$  in  $V^-(f)$  are vertices in  $U_{[3,8]}$  that are joined by an edge in  $M$ , then  $\delta_v(f) = -\delta_f(v) = 3$ .
- $\delta 6$ : If  $d(v) \geq \Delta - 1$ ,  $|V^-(f)| = 3$ , and  $v$  has a neighbour in  $V^-(f) \cap (T_{[3,6]} \cup U_{[3,5]})$ , then  $\delta_v(f) = -\delta_f(v) = \frac{5}{2}$ .
- $\delta 7$ : If  $d(v) \geq \Delta - 1$ ,  $|V^-(f)| = 3$ , and none of  $\delta 6$  and  $\delta 7$  applies, then  $\delta_v(f) = -\delta_f(v) = 2$ .
- $\delta 8$ : If  $d(v) \geq \Delta - 1$ ,  $|V^-(f)| \geq 4$ , and  $v$  has a neighbour in  $V^-(f) \cap T_{[3,6]}$ , then  $\delta_v(f) = -\delta_f(v) = 2$ .
- $\delta 9$ : If  $d(v) \geq \Delta - 1$ ,  $|V^-(f)| \geq 4$ , and  $\delta 9$  does not apply, then  $\delta_v(f) = -\delta_f(v) = \frac{3}{2}$ .

For a vertex  $v$ , write  $\gamma(v)$  for the sum of  $\gamma_e(v)$  over all edges  $e$  that have  $v$  as an end-vertex. For a vertex  $v$  of degree 1 we set  $\delta(v) = 0$ . For any other vertices, write  $\delta(v)$  for the sum over the faces  $f$  around  $v$  of  $\delta_f(v)$ . Similarly, for a face  $f$ , write  $\delta(f)$  for the sum over the vertices  $v$  on the reduced walk  $W_f^-$  around  $f$  of the values of  $\delta_v(f)$ .

By the definitions of  $\gamma$  and  $\delta$ ,

$$\sum_v \gamma(v) + \sum_v \delta(v) + \sum_f \delta(f) = 0.$$

It follows from Claim 3 that

$$\sum_v \beta(v) < 0.$$

Finally, from Euler's Formula for simple plane graphs, we obtain

$$\sum_v \alpha(v) + \sum_f \alpha(f) < 0.$$

Thus, in order to reach a contradiction, it is enough to show that for every vertex  $v$ :

$$\alpha(v) + \beta(v) + \delta(v) + \gamma(v) \geq 0, \quad (5)$$

and that for every face  $f$ :

$$\alpha(f) + \delta(f) \geq 0. \quad (6)$$

Let  $f$  be a face. As  $G$  is simple,  $|V^-(f)| \geq 3$ . Since  $\alpha(f) = -6$ , to establish (6) it is enough to show that  $\delta(f) \geq 6$ . Let  $v$  be the vertex in  $V^-(f)$  for which  $\delta_v(f)$  is minimum. If  $\delta_v(f) \cdot |V^-(f)| \geq 6$ , then (6) clearly holds, and so we only need to deal with cases  $\delta 1 - \delta 4$ . Also, if  $v \in T_{[7, \Delta-2]} \cup U_{[6, \Delta-2]}$ , then  $\delta 3$  and  $\delta 4$  give  $\delta_v(f) \geq 2$ , and hence again (6) is verified.

If  $v \in T_{[3,4]}$ , then  $\delta_v(f) \geq 1$  by  $\delta 1$  or  $\delta 3$  and, by (4), the neighbours  $u$  and  $w$  of  $v$  in  $V^-(f)$  have degree at least  $\Delta - 1$ . If  $|V^-(f)| = 3$ , then  $\delta 6$  applies to both  $u$  and  $w$ , so  $\delta_u(f) = \delta_w(f) = \frac{5}{2}$ . If  $|V^-(f)| \geq 4$ , then  $\delta 8$  applies, so  $\delta_u(f) = \delta_w(f) = 2$ , while a fourth vertex  $z$  in  $V^-(f)$  satisfies  $\delta_z(f) \geq 1$ . So (6) always follows.

If  $v \in T_{[5,6]}$ , then  $\delta_v(f) \geq \frac{3}{2}$ , so we may assume that  $|V^-(f)| = 3$  (as  $\delta_v(f) \leq \delta_u(f)$  whenever  $u \in V^-(f)$ ). Moreover,  $v$  has neighbours  $u, w$  in  $V^-(f)$  with degree at least  $\Delta - 3 \geq 9$ . If  $d(u) \geq \Delta - 1$ , then  $\delta 6$  gives  $\delta_u(f) = \frac{5}{2}$ . If  $d(u) \in \{\Delta - 3, \Delta - 2\}$ , then  $\delta 3$  and  $\delta 4$  give  $\delta_u(f) \geq \frac{9}{4}$ , as  $\Delta \geq 5$ . Since similar bounds hold for  $\delta_w(f)$ , we deduce that (6) holds.

We are left with the case where  $v \in U_{[3,5]}$ . By  $\delta 2$  and  $\delta 4$  we find that  $\delta_v(f) \geq \frac{3}{2}$ , and hence we again only have to consider the case where  $|V^-(f)| = 3$ . Rules  $\delta 3 - \delta 6$  ensure that any other vertex  $u$  in  $V^-(f)$  with  $d(u) \geq 9$  satisfies  $\delta_u(f) \geq \frac{9}{4}$ . So we may suppose that there is a vertex  $u \in V^-(f)$  with  $d(u) \leq 8$ . Since  $|V^-(f)| = 3$ , we must in fact have  $uv \in E(G)$ .

Moreover, as  $\Delta \geq 12$ , we know by (4) that the edge  $uv$  belongs to the matching  $M$ . This means that  $u \in U_{[3,8]}$ . Let  $w$  be the third vertex in  $V^-(f)$ . If  $v \in U_{[3,4]}$ , then  $d(w) \geq \Delta - 1$  by (4), and so  $\delta_w(f) = 3$  by  $\delta 5$ , confirming (6). As the final case, assume that  $v \in U_5$  and  $\delta_u(f) \geq \delta_v(f) = \frac{9}{5}$ . Since also  $d(w) \geq \Delta - 2 \geq 11$ , one of  $\delta 3$ – $\delta 6$  applies, yielding that  $\delta_w(f) \geq \frac{12}{5}$ . So again  $\delta(f) \geq 6$ , confirming (6) for all faces.

Now let  $v$  be a vertex. Recall the convention that if  $v \in T$ , then the two consecutive faces incident with  $v$  neighbouring the neighbour of degree 1 are counted as one face, while all other faces are counted separately.

If  $d(v) = 1$ , then  $\alpha(v) = -3$  and  $\gamma(v) = 3$ . Since  $\beta(v) = \delta(v) = 0$ , we immediately obtain (5).

Recall that  $G$  has no vertices of degree 2. If  $d(v) = 3$ , then  $\alpha(v) = 3$ , while  $\beta(v) = 2$  by  $\beta 2$ . If  $v \in T_3$ , then  $\gamma(v) = -3$  and  $\delta 1$  implies that  $\delta(v) = -2$ . If  $v \in U_3$ , then  $\gamma(v) = 0$  and  $\delta 2$  implies that  $\delta(v) = -5$ . This confirms (5) if  $d(v) = 3$ .

Next suppose that  $4 \leq d(v) \leq \Delta - 2$ . Recall that  $\alpha(v) = 3d(v) - 6$ , and observe that  $\beta(v) = 0$ . If  $v \in T$ , then  $\gamma(v) = -3$  by  $\gamma 1$ . By  $\delta 3$  we have  $\delta(v) = (d(v) - 1) \cdot \left(-3 - \frac{6}{d(v) - 1}\right) = 9 - 3d(v)$ . Similarly, if  $v \in U$ , then  $\gamma(v) = 0$ , and  $\delta 4$  implies that  $\delta(v) = 6 - 3d(v)$ . This proves (5) for those vertices  $v$ .

Now suppose that  $d(v) \geq \Delta - 1$ . As a next step towards proving (5), we consider the average value of  $\delta_f(v)$  over the faces incident with  $v$ .

**Claim 4.** *For any two consecutive faces  $f_i$  and  $f_{i+1}$  incident with  $v$  it holds that  $\delta_{f_i}(v) + \delta_{f_{i+1}}(v) \geq -5$  (where the addition is modulo  $d(v)$  if  $v \in U$ , and modulo  $d(v) - 1$  if  $v \in T$ ).*

*Proof.* We may assume that one of  $\delta_{f_i}(v)$  or  $\delta_{f_{i+1}}(v)$  is smaller than  $-\frac{5}{2}$ , otherwise the stated inequality holds. Suppose first that  $\delta_{f_i}(v) < -\frac{5}{2}$ . By the definitions of  $\delta 5$ – $\delta 9$ , it follows that  $\delta 5$  was applied to  $f_i$ , and consequently  $\delta_{f_i}(v) = -3$ . Hence  $v_i$  and  $v_{i+1}$  both belong to  $U$ , have degree at most 8, and there is an edge from  $M$  between them. Since  $M$  is a matching, it cannot also be the case that  $\delta_{f_{i+1}}(v) = -3$ . If  $\delta_{f_{i+1}}(v) = -\frac{5}{2}$ , then  $\delta 6$  was applied to the face  $f_{i+1}$ . But then the vertex  $v_{i+2}$  has degree at most 6 and there is an edge  $v_{i+1}v_{i+2}$ , which contradicts (4). So  $\delta_{f_{i+1}}(v) \geq -2$  and the statement follows.

By symmetry, the same arguments apply if  $\delta_{f_{i+1}}(v) < -\frac{5}{2}$ . □

If we add up the values of  $\delta_{f_i}(v) + \delta_{f_{i+1}}(v)$  for  $i \in \{1, 2, \dots, d(v)\}$  if  $v \in U$ , or for  $i \in \{1, 2, \dots, d(v) - 1\}$  if  $v \in T$ , then Claim 4 yields that

$$\delta(v) = \frac{1}{2} \sum_i (\delta_{f_i}(v) + \delta_{f_{i+1}}(v)) \geq \begin{cases} -\frac{5}{2}d(v) + \frac{5}{2}, & \text{if } v \in T; \\ -\frac{5}{2}d(v), & \text{if } v \in U. \end{cases} \quad (7)$$

First suppose that  $d(v) = \Delta - 1$ . Then  $\alpha(v) = 3\Delta - 9$  and  $\beta(v) = 0$ . If  $v \in T$ , then  $\gamma(v) = -3$  and (7) gives  $\delta(v) \geq -\frac{5}{2}(\Delta - 1) + \frac{5}{2}$ . Since  $\Delta \geq 14$ , inequality (5) follows. If  $v \in U$ , then  $\gamma(v) = 0$  and  $\delta(v) \geq -\frac{5}{2}(\Delta - 1)$ . The hypothesis that  $\Delta \geq 13$  guarantees that (5) is valid again.

Finally, suppose that  $d(v) = \Delta$ . Now  $\alpha(v) = 3\Delta - 6$  and  $\beta(v) = -2$ . If  $v \in T$ , then  $\gamma(v) = -3$  and  $\delta(v) \geq -\frac{5}{2}\Delta + \frac{5}{2}$ . We see that (5) holds, as  $\Delta \geq 17$ . If  $v \in U$ , then  $\gamma(v) = 0$  and  $\delta(v) \geq -\frac{5}{2}\Delta$ . So (5) is verified, provided that  $\Delta \geq 16$ .

This confirms (5) for all vertices and completes the proof of the theorem. □

## 4.2 Proof of Theorem 10

Recall the notation and terminology introduced in the introduction of this section. The statement of Theorem 10 is true for graphs with maximum degree 20 and exactly 20 edges. We use induction on  $E(G)$ , and proceed with the induction step. We may easily assume that  $G$  is connected with at least 21 vertices, since  $\Delta \geq 20$ . Let  $M$  be a precoloured distance-3 matching.

We first observe that

$$\text{if } uv \in E(G) \setminus M, \text{ then } d(u) + d(v) \geq \Delta + 2. \quad (8)$$

Indeed, suppose that the inequality does not hold for some  $uv \notin M$ . Then  $M$  can be extended to a colouring of  $G - uv$  so that at most  $\Delta - 1$  colours are used on the edges adjacent to  $uv$ , and so we can easily extend the colouring further to  $uv$ . From (8) it follows that every vertex with degree 1 is incident with an edge in  $M$  and that if  $v$  has degree 2 and  $uv \notin M$ , then  $d(u) = \Delta$ . In particular, if a vertex  $v$  with degree greater than 1 has a neighbour in  $T_2$ , then  $d(v) = \Delta$ . Moreover, since edges in  $M$  are at distance at least 4 in  $G$ , a vertex can have at most one neighbour in  $V_1 \cup T_2$ .

Let  $V'_2$  be the set of vertices of degree 2 that are not incident with an edge of  $M$ . For a face  $f$ , let  $V^-(f) = V(f) \setminus (V_1 \cup T_2)$ , and let  $W_f^-$  be the sequence of vertices on the boundary walk  $W_f$  after removing vertices from  $V_1 \cup T_2$ . For a vertex  $v$ , let  $v_1, v_2, \dots, v_{d(v)}$  be the neighbours of  $v$ , listed in clockwise order according to the drawing of  $G$ . Write  $f_i$  for the face incident with  $v$  lying between the edges  $vv_i$  and  $vv_{i+1}$  (taking addition modulo  $d(v)$ ).

If a vertex  $v$  has a (unique) neighbour in  $V_1 \cup T_2$ , then we always choose  $v_1$  to be this neighbour. In that case  $f_{d(v)} = f_1$ , and that face is called  $f_1$  again. Note that it is possible for other faces to be the same as well (if  $v$  is a cut-vertex), but we will not identify those multiple names of the same face.

**Claim 5.**  $|V_\Delta| > |V'_2|$ .

*Proof.* Consider the set  $F$  of edges in  $E(G) \setminus M$  with one end-vertex in  $V'_2$  and the other in  $V_\Delta$ . The subgraph with vertex set  $V'_2 \cup V_\Delta$  and edge set  $F$  is bipartite; we claim it is acyclic. For suppose there exists an (even) cycle  $C \in F$ . By induction, we can extend the precolouring of  $E(G) \setminus M$  to  $G - C$ . But then we can further extend this colouring to the edges of  $C$ , since each one sees only  $\Delta - 2$  coloured edges, and even cycles are 2-edge-choosable.

Since each vertex in  $V'_2$  is incident with at least two edges in  $F$ , we have  $|V_\Delta| + |V'_2| > |F| \geq 2|V'_2|$ . The claim follows.  $\square$

We use a discharging argument to complete the proof. First, let us assign to each vertex  $v$  a charge

$$\alpha 1: \quad \alpha(v) = 3d(v) - 6,$$

and to each face  $f$  a charge

$$\alpha 2: \quad \alpha(f) = -6.$$

For each vertex  $v$  we define  $\beta(v)$  as follows.

$$\beta 1: \quad \text{If } v \in V_\Delta, \text{ then } \beta(v) = -2.$$

$$\beta 2: \quad \text{If } v \in V'_2, \text{ then } \beta(v) = 2.$$

$\beta 3$ : In all other cases,  $\beta(v) = 0$ .

For each vertex  $v$  and edge  $e = vu$ , we define  $\gamma_e(v)$  and  $\gamma_e(u)$  as follows.

$\gamma 1$ : If  $v \in V_1$ , then  $\gamma_e(v) = -\gamma_e(u) = 3$ .

$\gamma 2$ : If  $v \in T_2$  and  $u \in V_\Delta$ , then  $\gamma_e(v) = -\gamma_e(u) = 3$ .

$\gamma 3$ : If  $v \in U_2 \setminus V'_2$  and  $u \in V_\Delta$ , then  $\gamma_e(v) = -\gamma_e(u) = 2$ .

$\gamma 4$ : In all other cases,  $\gamma_e(v) = 0$ .

Finally, for each face  $f$  and vertex  $v \in W_f^-$  we define  $\delta_f(v)$  and  $\delta_v(f)$  as follows.

$\delta 1$ : If  $v \in U_2$ , then  $\delta_v(f) = -\delta_f(v) = 1$ .

$\delta 2$ : If  $v \in T$  and  $3 \leq d(v) \leq \Delta - 4$ , then  $\delta_v(f) = -\delta_f(v) = 3 - \frac{6}{d(v) - 1}$ .

$\delta 3$ : If  $v \in U$  and  $3 \leq d(v) \leq \Delta - 4$ , then  $\delta_v(f) = -\delta_f(v) = 3 - \frac{6}{d(v)}$ .

$\delta 4$ : If  $d(v) \geq \Delta - 3$ ,  $|V^-(f)| = 3$ , and both neighbours of  $v$  in  $V^-(f)$  are joined by an edge in  $M$ , then  $\delta_v(f) = -\delta_f(v) = 4$ .

$\delta 5$ : If  $d(v) \geq \Delta - 3$  and  $v$  has a neighbour in  $V^-(f) \cap T_3$ , then  $\delta_v(f) = -\delta_f(v) = 3$ .

$\delta 6$ : If  $d(v) \geq \Delta - 3$  and none of  $\delta 4$  and  $\delta 5$  applies, then  $\delta_v(f) = -\delta_f(v) = \frac{5}{2}$ .

For a vertex  $v$  and face  $f$ , let  $\gamma(v)$ ,  $\delta(v)$  and  $\delta(f)$  be defined as in the proof of Theorem 9. By definition we have  $\sum_v \gamma(v) + \sum_v \delta(v) + \sum_f \delta(f) = 0$ . It follows from Claim 5 that  $\sum_v \beta(v) < 0$ . From Euler's Formula we obtain  $\sum_v \alpha(v) + \sum_f \alpha(f) < 0$ .

Thus, in order to reach a contradiction, it is enough to show that for every vertex  $v$ :

$$\alpha(v) + \beta(v) + \delta(v) + \gamma(v) \geq 0, \quad (9)$$

and that for every face  $f$ :

$$\alpha(f) + \delta(f) \geq 0. \quad (10)$$

Let  $f$  be a face. As  $G$  is simple,  $|V^-(f)| \geq 3$ . Since  $\alpha(f) = -6$ , it follows that (10) is verified if we can show that  $\delta(f) \geq 6$ . Let  $v$  be the vertex in  $V^-(f)$  for which  $\delta_v(f)$  is minimum. If  $\delta_v(f) \cdot |V^-(f)| \geq 6$ , then (10) clearly holds. So, by checking  $\delta 1 - \delta 6$ , we see we only have to consider the case where  $v \in T_{[3,6]} \cup U_{[2,5]}$ . (Recall that vertices from  $V_1 \cup T_2$  do not appear in  $W_f^-$ .)

If  $v \in U_2$ , then let  $u$  and  $w$  be the neighbours of  $v$ . Consider first the case where both  $u$  and  $w$  have degree  $\Delta$ . Then they both belong to  $V^-(f)$ , so (10) follows, since  $\delta_v(f) = 1$  and  $\delta_u(f) \geq \frac{5}{2}$ ,  $\delta_w(f) \geq \frac{5}{2}$  by  $\delta 4 - \delta 6$ . Suppose now that  $u$  has degree less than  $\Delta$ , which implies by (8) that  $uv \in M$  and, consequently,  $vw \notin M$ . In particular,  $w \in V^-(f)$  and  $w$  has degree  $\Delta$ . Note also that necessarily  $u \in V^-(f)$ . If  $|V^-(f)| = 3$ , then  $\delta_w(f) = 4$  by  $\delta 4$ . As  $\delta_u(f) \geq \delta_v(f) = 1$ , it follows that (10) holds. If  $|V^-(f)| \geq 4$ , then  $u$  has a neighbour  $u'$  in  $V^-(f) \setminus \{v, w\}$ . We assert that  $\delta_u(f) + \delta_{u'}(f) \geq \frac{5}{2}$ . Indeed, this holds if either of  $u$  and  $u'$  has degree at least  $\Delta - 3$ , by  $\delta 5$  and  $\delta 6$ . Otherwise,  $\delta 2$  or  $\delta 3$  applies to both  $u$  and  $u'$ . So  $\delta_u(f) + \delta_{u'}(f) \geq 6 - \frac{6}{d(u) - 1} - \frac{6}{d(u') - 1}$ . Since  $d(u) \geq 3$ ,  $d(u') \geq 3$  and  $d(u') \geq \Delta + 2 - d(u)$ , it is easy to see that the expression  $6 - \frac{6}{d(u) - 1} - \frac{6}{d(u') - 1}$  is minimised

if  $d(u) = d(u') = \frac{1}{2}(\Delta + 2)$ . We infer that  $\delta_u(f) + \delta_{u'}(f) \geq 6 - 2 \cdot \frac{6}{\frac{1}{2}\Delta} \geq \frac{5}{2}$ , because  $\Delta \geq 7$ .

The assertion follows. As a result, we deduce that (10) holds, since  $\delta_w(f) \geq \frac{5}{2}$  by  $\delta 5$  and  $\delta 6$ .

If  $v \in T_3$ , then  $\delta_v(f) = 0$ , but  $v$  has two neighbours in  $V^-(f)$  that have degree at least  $\Delta - 1$  each. Equation (10) then follows from  $\delta 5$ .

For the remaining cases we always have  $\delta_v(f) \geq 1$ . Rules  $\delta 2 - \delta 6$  ensure that any vertex  $u \in V^-(f)$  with  $d(u) \geq 13$  satisfies  $\delta_u(f) \geq \frac{5}{2}$ ; hence there can be at most one such vertex and, in particular, a neighbour  $u$  of  $v$  in  $V^-(f)$  must have degree at most 12. As  $v$  itself has degree at most 6, by (8) we have  $uv \in M$ , which also implies that  $u, v \in U$ . Hence in particular  $v \in U_{[3,5]}$ . Let  $w$  be the neighbour of  $v$  in  $V^-(f) \setminus \{u\}$ . Since  $v \in U_{[3,5]}$  and  $vw \notin M$ , it necessarily holds that  $d(w) \geq \Delta - 3$ . If  $|V^-(f)| = 3$ , then (10) holds by  $\delta 4$  since  $\delta_u(f) \geq \delta_v(f) \geq 1$ . If  $|V^-(f)| \geq 4$ , then  $u$  has a neighbour  $u'$  in  $V^-(f) \setminus \{v, w\}$ , which has degree at least  $\Delta + 2 - d(u) \geq 10$ . Consequently,  $\delta_{u'}(f) \geq \frac{5}{2}$  by  $\delta 3$ ,  $\delta 5$  or  $\delta 6$ . We deduce that (10) holds, as  $\delta_w(f) \geq \frac{5}{2}$  by  $\delta 5$  or  $\delta 6$ .

This confirms (10) for all faces.

Now let  $v$  be a vertex. Recall that  $\alpha(v) = 3d(v) - 6$ . Furthermore, if  $v$  has a neighbour in  $V_1 \cup T_2$ , then the two consecutive faces incident with that neighbour are counted as one face; all other faces are counted separately. Finally, as noted earlier, a vertex can have at most one neighbour in  $V_1 \cup T_2$ .

If  $d(v) = 1$ , then  $\alpha(v) = -3$  and  $\gamma(v) = 3$ . Since  $\beta(v) = \delta(v) = 0$ , we immediately obtain (9).

If  $d(v) = 2$ , then  $\alpha(v) = 0$ . If  $v \in T_2$ , then both  $\gamma 1$  and  $\gamma 2$  apply; hence  $\gamma(v) = 0$ . Again one can check that  $\beta(v) = \delta(v) = 0$ , confirming (9). Otherwise  $v \in U_2$ , and  $\delta 1$  implies that  $\delta(v) \geq -2$ , as  $v$  is incident with at most two faces. If  $v \in V_2'$  as well, then  $\beta 2$  yields that  $\beta(v) = 2$  and  $\gamma(v) = 0$ . If  $v \notin V_2'$ , then  $\gamma(v) = 2$  while  $\beta(v) = 0$ . In either case (9) follows.

Next suppose that  $3 \leq d(v) \leq \Delta - 4$ . Observe that  $\beta(v) = 0$ . If  $v \in T$ , then  $\gamma(v) = -3$  by  $\gamma 1$ . Since  $v$  has a neighbour with degree one, we know that  $v$  is incident with  $d(v) - 1$  regions, and so  $\delta 2$  yields that  $\delta(v) = (d(v) - 1) \cdot \left(-3 - \frac{6}{d(v) - 1}\right) = 9 - 3d(v)$ . Similarly, if  $v \in U$ , then  $\gamma(v) = 0$ , and  $\delta 4$  yields that  $\delta(v) = 6 - 3d(v)$ . This proves (9) for those vertices  $v$ .

Suppose now that  $d(v) \in \{\Delta - 3, \Delta - 2, \Delta - 1\}$ . Then  $\beta(v) = 0$ . If  $v \in T$ , then  $\gamma(v) = -3$  by  $\gamma 1$ . Since  $M$  is distance-3, none of  $\delta 4$  and  $\delta 5$  applies to  $v$ , and  $v$  is incident with  $d(v) - 1$  faces. From  $\delta 6$  we deduce that  $\delta(v) = -\frac{5}{2}(d(v) - 1)$ . Since  $d(v) \geq \Delta - 3 \geq 13$ , it follows that  $3d(v) - 6 - 3 - \frac{5}{2}(d(v) - 1) = \frac{1}{2}d(v) - \frac{13}{2} \geq 0$ , and hence (9) is satisfied again. Next assume that  $v \in U$ , and so  $\gamma(v) = 0$ . The fact that  $M$  is distance-3 ensures that  $\delta 6$  applies to at least  $d(v) - 1$  faces incident with  $v$ . This implies that  $\delta(v) \geq -(4 + \frac{5}{2}(d(v) - 1))$ . Combined with the assumption that  $\Delta \geq 15$ , this is always enough to satisfy (9).

Finally, suppose that  $d(v) = \Delta$ . In this case  $\beta(v) = -2$ . If  $v \in T$ , then the distance condition on  $M$  ensures that  $\gamma(v) = -3$  and  $\delta(v) = -\frac{5}{2}(\Delta - 1)$ . Since  $\Delta \geq 17$  this confirms (9).

So we are left with the case where  $v \in U$ . Since  $M$  is a distance-3 matching, at most one of  $\gamma 2$ ,  $\gamma 3$ ,  $\delta 4$  or  $\delta 5$  applies. Moreover, if  $\gamma 2$  does apply, then  $\gamma(v) = -3$ , the vertex  $v$  is incident with  $\Delta - 1$  faces and for all those faces  $f$  we have  $\delta_f(v) = -\frac{5}{2}$ . If  $\gamma 3$  does apply, then  $\gamma(v) = -2$ , the vertex  $v$  is incident with  $\Delta$  faces and for all those faces  $f$  we have  $\delta_f(v) = -\frac{5}{2}$ . If  $\delta 4$  or  $\delta 5$  does apply, then  $\gamma(v) = 0$ , the vertex  $v$  is incident with  $\Delta$  faces, for all those faces  $f$  but one we have  $\delta_f(v) = -\frac{5}{2}$ , while for the final face  $f$  we have  $\delta_f(v) \in \{-4, -3\}$ .

Finally, if none of  $\gamma_2$ ,  $\gamma_3$ ,  $\delta_4$  or  $\delta_5$  applies, then  $\gamma(v) = 0$ , the vertex  $v$  is incident with  $\Delta$  faces and for all those faces  $f$  we have  $\delta_f(v) = -\frac{5}{2}$ . Using that  $\Delta \geq 20$ , we can check that (9) is satisfied in all cases.

This confirms (9) for all vertices and completes the proof of the theorem.  $\square$

## 5 A Final Conjecture

Suppose that we would go to any means to get an extension form of Vizing's theorem, say, by weakening the precolouring condition. We still let  $\mathcal{K} = [K]$  be a palette of available colours. Given a subset  $S \subseteq E$  of edges and an arbitrary (i.e., not necessarily proper) colouring of elements of  $S$  using only colours from  $\mathcal{K}$ , is there a proper colouring of all edges of  $G$  (using colours from  $\mathcal{K}$ ) that *disagrees* with the given colouring on every edge of  $S$ ? We may consider the coloured set  $S$  as a set of *forbidden* (coloured) edges, while the full colouring, if it can be produced, is called an *avoidance* of the forbidden edges. The following is a significant weakening of Conjecture 1, and is not implied by the LCC, nevertheless its proof eludes us.

### Conjecture 16.

Let  $G$  be a multigraph with maximum degree  $\Delta(G)$  and maximum multiplicity  $\mu(G)$ . Using the palette  $\mathcal{K} = [\Delta(G) + \mu(G)]$ , any forbidden matching can be avoided by a proper edge-colouring of all of  $G$ .

## References

- [1] M. O. Albertson. You can't paint yourself into a corner. *J. Combin. Theory Ser. B*, 73(2):189–194, 1998.
- [2] M. O. Albertson, A. V. Kostochka, and D. B. West. Precolouring extensions of Brooks' theorem. *SIAM J. Discrete Math.*, 18(3):542–553, 2004/05.
- [3] M. O. Albertson and E. H. Moore. Extending graph colorings using no extra colors. *Discrete Math.*, 234(1-3):125–132, 2001.
- [4] N. Alon. Restricted colorings of graphs. In *Surveys in Combinatorics, 1993 (Keele)*, volume 187 of *London Math. Soc. Lecture Note Ser.*, pages 1–33. Cambridge Univ. Press, Cambridge, 1993.
- [5] M. Axenovich. A note on graph coloring extensions and list-colorings. *Electron. J. Combin.*, 10, 2003. 5 pp.
- [6] C. Berge and J.-C. Fournier. A short proof for a generalization of Vizing's theorem. *J. Graph Theory*, 15(3):333–336, 1991.
- [7] M. Bonamy. Planar graphs with  $\Delta \geq 8$  are  $(\Delta + 1)$ -edge-choosable. arXiv:1303.4025 [cs.DM], 2013.
- [8] O. V. Borodin. Criterion of chromaticity of a degree prescription (in Russian). In *Abstracts of IV All-Union Conf. on Theoretical Cybernetics (Novosibirsk)*, pages 127–128, 1977.

- [9] O. V. Borodin. A generalization of Kotzig's theorem and prescribed edge coloring of planar graphs. *Math. Notes*, 48(5-6):1186–1190, 1990.
- [10] O. V. Borodin, A. V. Kostochka, and D. R. Woodall. List edge and list total colourings of multigraphs. *J. Combin. Theory Ser. B*, 71(2):184–204, 1997.
- [11] P. Erdős, A. L. Rubin, and H. Taylor. Choosability in graphs. In *Proceedings of the West Coast Conference on Combinatorics, Graph Theory and Computing (Humboldt State Univ., Arcata, Calif., 1979)*, volume XXVI of *Congress. Numer.*, pages 125–157. Utilitas Math., Winnipeg, Man., 1980.
- [12] T. Gallai. Kritische Graphen. I. *Magyar Tud. Akad. Mat. Kutató Int. Közl.*, 8:165–192, 1963.
- [13] F. Galvin. The list chromatic index of a bipartite multigraph. *J. Combin. Theory Ser. B*, 63(1):153–158, 1995.
- [14] M. K. Gol'dberg. Multigraphs with a chromatic index that is nearly maximal (in Russian). *Diskret. Analiz*, 23:3–7, 72, 1973.
- [15] R. P. Gupta. On decompositions of a multi-graph into spanning subgraphs. *Bull. Amer. Math. Soc.*, 80:500–502, 1974.
- [16] R. Häggkvist and J. Janssen. New bounds on the list-chromatic index of the complete graph and other simple graphs. *Combin. Probab. Comput.*, 6(3):295–313, 1997.
- [17] M. Juvan, B. Mohar, and R. Škrekovski. On list edge-colorings of subcubic graphs. *Discrete Math.*, 187(1-3):137–149, 1998.
- [18] M. Juvan, B. Mohar, and R. Škrekovski. Graphs of degree 4 are 5-edge-choosable. *J. Graph Theory*, 32(3):250–264, 1999.
- [19] J. Kahn. Asymptotically good list-colorings. *J. Combin. Theory Ser. A*, 73(1):1–59, 1996.
- [20] J. Kahn. Asymptotics of the chromatic index for multigraphs. *J. Combin. Theory Ser. B*, 68(2):233–254, 1996.
- [21] J. Kahn. Asymptotics of the list-chromatic index for multigraphs. *Random Structures Algorithms*, 17(2):117–156, 2000.
- [22] O. Marcotte and P. Seymour. Extending an edge-coloring. *J. Graph Theory*, 14(5):565–573, 1990.
- [23] M. Molloy and B. Reed. Near-optimal list colorings. *Random Structures Algorithms*, 17(3-4):376–402, 2000.
- [24] M. Molloy and B. Reed. *Graph Colouring and the Probabilistic Method*, volume 23 of *Algorithms and Combinatorics*. Springer-Verlag, Berlin, 2002.

- [25] T. Rackham. A note on  $K_{\Delta+1}^-$ -free precolouring with  $\Delta$  colours. *Electron. J. Combin.*, 16(1), 2009. 8 pp.
- [26] D. P. Sanders and Y. Zhao. Planar graphs of maximum degree seven are class I. *J. Combin. Theory Ser. B*, 83(2):201–212, 2001.
- [27] D. Scheide and M. Stiebitz. On Vizing’s bound for the chromatic index of a multigraph. *Discrete Math.*, 309(15):4920–4925, 2009.
- [28] P. D. Seymour. On multicolourings of cubic graphs, and conjectures of Fulkerson and Tutte. *Proc. London Math. Soc. (3)*, 38(3):423–460, 1979.
- [29] C. E. Shannon. A theorem on coloring the lines of a network. *J. Math. Physics*, 28:148–151, 1949.
- [30] M. Stiebitz, D. Scheide, B. Toft, and L. M. Favrholdt. *Graph Edge Coloring. Vizing’s Theorem and Goldberg’s Conjecture*. Wiley Series in Discrete Mathematics and Optimization. John Wiley & Sons Inc., Hoboken, NJ, 2012.
- [31] C. Thomassen. Every planar graph is 5-choosable. *J. Combin. Theory Ser. B*, 62(1):180–181, 1994.
- [32] C. Thomassen. Color-critical graphs on a fixed surface. *J. Combin. Theory Ser. B*, 70(1):67–100, 1997.
- [33] V. G. Vizing. On an estimate of the chromatic class of a  $p$ -graph (in Russian). *Diskret. Analiz*, 3:25–30, 1964.
- [34] V. G. Vizing. Critical graphs with given chromatic class (in Russian). *Diskret. Analiz*, 5:9–17, 1965.
- [35] V. G. Vizing. Coloring the vertices of a graph in prescribed colors (in Russian). *Diskret. Analiz*, 29:3–10, 101, 1976.