



## A. Stern's analysis of the nodal sets of some families of spherical harmonics revisited

Pierre Bérard, Bernard Helffer

### ► To cite this version:

| Pierre Bérard, Bernard Helffer. A. Stern's analysis of the nodal sets of some families of spherical harmonics revisited. 2014. hal-01026363v1

HAL Id: hal-01026363

<https://hal.science/hal-01026363v1>

Preprint submitted on 21 Jul 2014 (v1), last revised 2 Jul 2015 (v4)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A. Stern's analysis of the nodal sets of some families of spherical harmonics revisited

P. Bérard

Institut Fourier, Université de Grenoble and CNRS, B.P.74,  
F 38402 Saint Martin d'Hères Cedex, France.

and

B. Helffer

Laboratoire de Mathématiques, Univ. Paris-Sud 11 and CNRS,  
F 91405 Orsay Cedex, France, and  
Laboratoire Jean Leray, Université de Nantes.

July 21, 2014

## Abstract

In this paper, we revisit the analyses of Antonie Stern (1925) and Hans Lewy (1977) devoted to the construction of spherical harmonics with two or three nodal domains. This is a natural continuation of our critical reading of A. Stern's results for Dirichlet eigenfunctions in the square [2].

Keywords: Nodal lines, Nodal domains, Courant theorem.

MSC 2010: 35B05, 35P20, 58J50.

## 1 Introduction

Let  $D$  be a bounded domain in  $\mathbb{R}^n$ . Let  $\Delta$  be the non-positive Laplacian with Dirichlet or Neumann boundary conditions. We arrange the eigenvalues  $(\lambda_j)_{j \in \mathbb{N}^*}$  of  $-\Delta$  in increasing order,

$$\lambda_1 < \lambda_2 \leq \lambda_3 \leq \dots .$$

Courant's 1923 celebrated nodal domain theorem [4], [5, p. 452] states that an eigenfunction associated with the  $n$ -th eigenvalue  $\lambda_n$ , has at most  $n$  nodal domains. On the other hand, an eigenfunction associated with  $\lambda_n$ , has a least two nodal domains when  $n \geq 2$ . The question remained of an eventual lower bound for the number of nodal domains of an  $n$ -th eigenfunction, as in the Sturm-Liouville theory.

For 2-dimensional domains with Dirichlet boundary conditions, using the Faber-Krahn inequality in an essential way, Åke Pleijel [16, 1956] proved that the number of nodal domains of an  $n$ -th eigenfunction is asymptotically less\* than  $0.7n$ . As a corollary, one can conclude that Courant's theorem is sharp for finitely many eigenvalues only. Pleijel's result was later generalized by other authors [15, 1], including an analysis for the Neumann boundary condition [17, 2009] (dimension 2, real analytic boundary)<sup>†</sup>. More recently, starting from 2009, there has been a renewed interest for Courant's theorem in the context of minimal partitions, and the investigation of the cases in which Courant's theorem is sharp [8, 9], see Section 5. These developments motivated [2] and motivate the present paper.

Let us now go back to the 1920's. Antonie Stern's 1925 thesis [18], written under the supervision of Richard Courant, contains the following three results.

**Theorem 1.1** *Let  $D$  be the unit square in  $\mathbb{R}^2$ , and  $\Delta$  the non-positive Laplacian with Dirichlet boundary conditions. Then, for any integer  $m$ , there exists an eigenfunction  $u$  of  $-\Delta$ , associated with the eigenvalue  $(4m^2+1)\pi^2$ , whose nodal set inside the square consists of a single simple closed curve. As a consequence,  $u$  has exactly two nodal domains.*

**Theorem 1.2** *Let  $\mathbb{S}^2$  be the unit sphere in  $\mathbb{R}^3$ , and  $\Delta$  the non-positive spherical Laplacian. For any odd integer  $\ell$ , there exists a spherical harmonic, of degree  $\ell$ , whose nodal set consists of a single simple closed curve. As a consequence,  $u$  has exactly two nodal domains.*

**Theorem 1.3** *Let  $\mathbb{S}^2$  be the unit sphere in  $\mathbb{R}^3$ , and  $\Delta$  the non-positive spherical Laplacian. For any even integer  $\ell \geq 2$ , there exists a spherical harmonic, of degree  $\ell$ , whose nodal set consists of two disjoint simple closed curves. As a consequence,  $u$  has exactly three nodal domains.*

Theorem 1.1 is stated without proof in [5, p. 455], with a reference to Stern's thesis [18], and illustrated by two figures taken from [18]. Theorems 1.2 and 1.3 do not seem to be mentioned in [5]. On the other hand, Stern's results on spherical harmonics appear in the 1977 paper [11] by Hans Lewy (Theorems 1 and 2), without any reference to A. Stern.

In [19], we provide extracts from Stern's thesis, with annotations and highlighting of the main assertions and ideas.

Stern's thesis is rather discursive. The main results are not stated as propositions or theorems. They appear in the course of the thesis, for example in [19, tags E1, K1, K2]:

[E1] ... es läßt sich beispielweise leicht zeigen, daß auf der Kugel bei jedem Eigenwert die Gebietszahlen 2 oder 3 auftreten, und daß bei Ordnung nach wachsenden Eigenwerten auch beim Quadrat die Gebietszahl 2 immer wieder vorkommt.

---

\*More precisely  $4\pi/\lambda(Disk_1)$  where  $\lambda(Disk_1)$  is the lowest eigenvalue of the Dirichlet Laplacian in the disk of area 1.

<sup>†</sup>The case of the Neumann problem for the square was already considered in Pleijel's article.

[K1] Zunächst wollen wir zeigen, daß es zu jedem Eigenwert Eigenfunktionen gibt, deren Nulllinien die Kugelfläche nur in zwei oder drei Gebiete teilen. .... Die Gebietszahl zwei tritt somit bei allen Eigenwerten  $\lambda_n = (2r+1)(2r+2)$   $r = 1, 2, \dots$  auf ;

[K2] ebenso wollen wir jetzt zeigen, daß die Gebietszahl drei bei allen Eigenwerten  $\lambda_n = 2r(2r+1)$ ,  $r = 1, 2, \dots$  immer wieder vorkommt.

Stern's proofs are far from being complete, but she provides nice geometric arguments [19, tags I1-I3], and figures.

[I1] Legen wir die beiden Knotenliniensysteme übereinander und schraffieren wir die Gebiete, in denen beide Funktionen gleiches Vorzeichen haben, so kann die Knotenlinie der Kugelfunktion

$$P_{2r+1}^{2r+1}(\cos \vartheta) \cos(2r+1)\varphi + \mu P_{2r+1}(\cos \vartheta), \quad \mu > 0$$

nur in der nichtschraffierten Gebieten verlaufen

[I3] und zwar für hinreichend kleine  $\mu$  in beliebiger Nachbarschaft der Knotenlinien von  $P_{2r+1}^{2r+1}(\cos \vartheta) \cos(2r+1)\varphi$ , d. h. der  $2r+1$  Meridiane, da sich bei stetiger Änderung von  $\mu$  das Knotenliniensystem stetig ändert ....

[I2] Da die Knotenlinie ferner durch die  $2(2r+1)^2$  Schnittpunkte der Nulllinien der beiden obenstehenden Kugelfunktionen gehen muß ...

Let us now explain Stern's main ideas. In both cases, A. Stern starts from an eigenfunction  $u$  whose nodal set can be completely described. In the case of the square, this function  $u$  is chosen to be  $\sin(2m\pi x) \sin(\pi y) + \sin(\pi x) \sin(2m\pi y)$ . In the case of the sphere, it is chosen to be the restriction to the sphere of the homogeneous harmonic polynomial  $\Im(x+iy)^\ell$ . A. Stern then perturbs the eigenfunction  $u$  by some eigenfunction  $v$  (in the same eigenspace), looking at the family  $u^\mu = u + \mu v$  for  $\mu$  small. The function  $v$  is chosen to be  $\sin(\pi x) \sin(2m\pi y)$  for the square, and a spherical harmonic whose nodal sets mainly contains latitude circles (parallels) in the case of the sphere.

The main observation made by A. Stern is that for  $\mu > 0$ , the nodal set  $N(u^\mu)$  satisfies

$$\mathcal{N} \subset N(u^\mu) \subset \mathcal{N} \cup \{uv < 0\},$$

where  $\mathcal{N} = N(u) \cap N(v)$  is the set of zeros common to  $u$  and  $v$ . In the case of the square, the connected components of the set  $\{uv \neq 0\}$  are small squares, whose vertices belong to  $\mathcal{N}$ . In the case of the sphere, they are square-like domains with vertices in  $\mathcal{N}$ , and triangle-like domains one of whose vertices is the north or south pole, and some others belong to  $\mathcal{N}$ . In both cases, the domains form a kind of black/white checkerboard (the connected open sets on which  $uv$  is positive/negative) on the unit square or on the sphere. The above inclusions say that the nodal set  $N(u^\mu)$  contains  $\mathcal{N}$ , and has to avoid the black squares, see [19, tags I1, I2]. A. Stern concludes by saying that the nodal set  $N(u^\mu)$  deforms continuously, and remains close to the nodal set  $N(u)$  when  $\mu$  is small, [19, tag I3].

In [2], we gave a complete geometric proof of Theorem 1.1. In this paper, we give a complete geometric proof of Theorems 1.2 and 1.3. In each case, our proofs use Stern's geometric ideas, together with an analysis of the possible local nodal patterns in the square-like and triangle-like domains. For this latter analysis,

- (i) we study *critical zeros* (points in the nodal set which are also critical points), and we show in particular that  $u^\mu$  does not have any critical zero when  $\mu \neq 0$  is small enough (Lemmas 3.1 and 4.2);
- (ii) we use a separation lemma (Lemmas 3.2 and 4.3) to exclude certain local nodal patterns;
- (iii) we use an energy argument to show that a connected component of the set  $\{uv \neq 0\}$  cannot contain a simple closed nodal curve of  $u^\mu$ , see Properties 3.3 (iii) and Properties 4.4 (vii);
- (iv) we also use classical properties of nodal sets of eigenfunctions, as summarized in [2, Section 5.2].

In her thesis, A.Stern refers to the continuity of the nodal set under a small perturbation [19, tag I3], but does not give any detail. In our proofs, the continuity argument is replaced by the separation lemmas.

In his 1977 paper [11], Hans Lewy's uses the harmonic homogeneous polynomials rather than the expression of spherical harmonics in terms of Legendre functions. He starts off from the spherical harmonic  $W(x, y, z) = \Im(x + iy)^\ell$ . When  $\ell$  is odd (Theorem 1), he perturbs  $W$  with a spherical harmonic which is positive at  $(0, 0, 1)$ . When  $\ell$  is even (Theorem 2), he perturbs  $W$  with a spherical harmonic of the form  $xyq(x, y, z)$ , where  $q(0, 0, 1) > 0$ . Lewy provides complete perturbation arguments, in particular near the singular points of the nodal set of  $\Im(x + iy)^\ell$ . He also indicates (Introduction) that a spherical harmonic of even degree has at least three nodal domains.

Finally, let us point out that our method yields sharp quantitative results. We indeed show that there exists some positive  $\mu_c$  such that for  $0 < \mu < \mu_c$  the nodal set of  $u^\mu$  is a regular 1-dimensional submanifold of the sphere, while the nodal set of  $u^{\mu_c}$  has self-intersections.

The paper is organized as follows. In Section 2 we recall some properties of spherical harmonics and Legendre functions. In Sections 3 and 4, we give detailed geometric proofs of Stern's first and second theorems for the sphere, with quantitative statements, Propositions 3.4 and 4.5. In Section 5, we recall the state of the art on the question of Courant sharpness for the sphere. In the appendix, we provide some numerical computations of nodal sets of spherical harmonics.

## 2 Preliminaries

### 2.1 Spherical harmonics

Denote by  $\mathbb{S}^2 := \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1\}$  the round 2-sphere. Given an integer  $\ell \in \mathbb{N}$ , we call  $\mathcal{H}_\ell$  the vector space of spherical harmonics of degree  $\ell$ , *i.e.* the restriction to the sphere of the harmonic homogeneous polynomials in 3 variables in  $\mathbb{R}^3$ . This is the eigenspace of  $-\Delta$  on  $\mathbb{S}^2$ , associated with the eigenvalue  $\ell(\ell+1)$ . It has dimension  $2\ell+1$ . Given a spherical harmonic  $h(\xi, \eta, \zeta)$  of degree  $\ell$ , with  $(\xi, \eta, \zeta) \in \mathbb{S}^2$ , one can recover the harmonic homogeneous polynomial  $H$  it comes from as follows. Let  $r = (x^2 + y^2 + z^2)^{1/2}$ . Then,

$$H(x, y, z) = r^\ell h(xr^{-1}, yr^{-1}, zr^{-1}). \quad (2.1)$$

For simplicity, we shall henceforth identify the spherical harmonic  $h$  and the polynomial  $H$ .

The space  $\mathcal{H}_0$  is 1-dimensional, associated with the eigenvalue 0. The space  $\mathcal{H}_1$  has dimension 3. It is associated with the eigenvalue 2, and is generated by the coordinate functions  $x, y$  and  $z$  which have exactly two nodal domains. The space  $\mathcal{H}_2$  has dimension 5. It is associated with the eigenvalue 6, and is generated by the polynomials  $yz, xz, xy, 2z^2 - x^2 - y^2$  and  $x^2 - z^2$ . It is easy to check that for  $\mu > 0$ , small enough, the spherical harmonic  $xy + \mu(2z^2 - x^2 - y^2)$  has exactly three nodal domains: the nodal set of the spherical harmonic consists of two simple closed curves given by the intersection of the sphere with a right cylinder over a hyperbola in the  $\{x, y\}$ -plane. Following A. Stern, we shall later on consider a perturbation of the degree  $\ell$  spherical harmonic  $\Im(x + iy)^\ell$ .

We denote the north and south poles of  $\mathbb{S}^2$  respectively by  $p_+ = (0, 0, 1)$  and  $p_- = (0, 0, -1)$ .

By abuse of language, we shall call *spherical coordinates* on the sphere  $\mathbb{S}^2$ , the map

$$\begin{cases} E : [0, \pi] \times \mathbb{R} \rightarrow \mathbb{S}^2, \\ E(\vartheta, \varphi) = (\sin \vartheta \cos \varphi, \sin \vartheta \sin \varphi, \cos \vartheta), \end{cases} \quad (2.2)$$

where  $\vartheta$  is the *co-latitude*, and  $\varphi$  the *longitude*.

The map  $E$  is a diffeomorphism from  $(0, \pi) \times (\varphi_0, 2\pi + \varphi_0)$  onto  $\mathbb{S}^2 \setminus M_{\varphi_0}$ , where  $M_{\varphi_0} = E([0, \pi], \varphi_0)$  is the meridian from  $p_+$  to  $p_-$  with longitude  $\varphi_0$ . To cover  $\mathbb{S}^2 \setminus \{p_\pm\}$ , we will work in  $(0, \pi) \times \mathbb{R}_{2\pi}$ , *i.e.* modulo  $2\pi$  in the  $\varphi$  variable ( $\mathbb{R}_{2\pi} = \mathbb{R}/(2\pi\mathbb{Z})$ ).

The map  $E$  can be viewed as the polar coordinates in the exponential map  $\exp_{p_+}$ , which sends the disk  $D(0, \pi)$ , with center 0 and radius  $\pi$  in  $T_{p_+}\mathbb{S}^2$ , onto  $\mathbb{S}^2 \setminus \{p_-\}$  diffeomorphically. The variable  $\vartheta$  is the distance to the north pole.

In the spherical coordinates, the antipodal map is given by  $(\vartheta, \varphi) \rightarrow (\pi - \vartheta, \pi + \varphi)$ .

In the sequel, we will illustrate the proofs by figures representing the nodal patterns viewed through the exponential map, *i.e.* in the disk  $D(0, \pi)$ .

Using the spherical coordinates, the spherical harmonics can be described in terms of Legendre functions and polynomials. In the next section, we fix some notation, and recall useful properties of Legendre functions and polynomials.

## 2.2 Legendre functions and polynomials

The  $(2\ell + 1)$ -dimensional vector space  $\mathcal{H}_\ell$  of spherical harmonics of degree  $\ell$  admits the basis,

$$P_\ell(\cos \vartheta), \quad P_\ell^m(\cos \vartheta) \cos(m\varphi), \quad P_\ell^m(\cos \vartheta) \sin(m\varphi), \quad (2.3)$$

where  $m$  is an integer  $1 \leq m \leq \ell$ ,  $P_\ell$  the Legendre polynomial of degree  $\ell$ , and  $P_\ell^m$  the Legendre functions. We use the notation and the normalizations of [13].

**Properties 2.1** For  $0 \leq m \leq \ell$ , the Legendre function  $P_\ell^m$  satisfies the differential equation,

$$((1 - t^2)P'(t))' + \left( \ell(\ell + 1) - \frac{m^2}{1 - t^2} \right) P(t) = 0. \quad (2.4)$$

When  $m$  is 0,  $P_\ell^m = P_\ell$ . Furthermore, the following properties hold.

(i) *Identities.*

$$P_\ell(t) = \frac{1}{2^\ell \ell!} \left( \frac{d}{dt} \right)^\ell (t^2 - 1)^\ell, \quad (2.5)$$

$$P_\ell^m(t) = (-1)^m (1 - t^2)^{m/2} \left( \frac{d}{dt} \right)^m P_\ell(t), \quad (2.6)$$

and

$$(1 - t^2) P'_\ell(t) = \ell P_{\ell-1}(t) - \ell t P_\ell(t). \quad (2.7)$$

In particular,  $P_\ell^\ell(\cos \vartheta) = C_\ell \sin^\ell(\vartheta)$ , where  $C_\ell$  is a constant.

(ii) *The polynomial  $P_\ell$  has degree  $\ell$ , the same parity as the integer  $\ell$ , and satisfies*

$$P_\ell(1) = 1, \quad P_\ell(-1) = (-1)^\ell, \quad (2.8)$$

and

$$\sup_{[-1,1]} |P_\ell(t)| = 1. \quad (2.9)$$

(iii) *The polynomial  $P_\ell(t)$  has  $\ell$  simple roots  $t_j = t_j(\ell)$  ( $j = 1, \dots, \ell$ ) in the interval  $(-1, 1)$ , enumerated in decreasing order. We write these roots as  $t_j(\ell) = \cos(\vartheta_j(\ell))$ , with*

$$0 < \vartheta_1(\ell) < \vartheta_2(\ell) < \dots < \vartheta_{\ell-1}(\ell) < \vartheta_\ell(\ell) < \pi. \quad (2.10)$$

*The derivative  $P'_\ell(t)$  has  $(\ell - 1)$  simple roots which we write as  $\cos(\vartheta'_j(\ell))$ . They satisfy*

$$0 < \vartheta_1(\ell) < \vartheta'_1(\ell) < \vartheta_2(\ell) < \dots < \vartheta_{\ell-1}(\ell) < \vartheta'_{\ell-1}(\ell) < \vartheta_\ell(\ell) < \pi. \quad (2.11)$$

*Note that the values  $\vartheta_j(\ell)$  and  $\vartheta'_j(\ell)$  are symmetrical with respect to  $\frac{\pi}{2}$ . As a consequence, for  $\ell$  odd,  $P_\ell(0) = 0$  and  $P'_\ell(0) \neq 0$ , and for  $\ell$  even,  $P_\ell(0) \neq 0$  and  $P'_\ell(0) = 0$ .*

(iv) The polynomials  $P_\ell$  and  $P_{\ell-1}$  have no common zero. More precisely, the zeros of  $P_\ell$  and  $P_{\ell-1}$  are intertwined:

$$0 < \vartheta_1(\ell) < \vartheta_1(\ell-1) < \vartheta_2(\ell) < \cdots < \vartheta_{\ell-1}(\ell) < \vartheta_{\ell-1}(\ell-1) < \vartheta_\ell(\ell) < \pi. \quad (2.12)$$

(v) For the zeros of  $P_\ell$ , one has the inequalities,

$$\frac{2j-1}{2\ell+1}\pi < \vartheta_j(\ell) < \frac{2j}{2\ell+1}\pi, \text{ for } j = 1, \dots, \ell. \quad (2.13)$$

(vi) Call  $p_j(\ell)$ ,  $1 \leq j \leq [\frac{\ell}{2}]$ , the local maxima of  $|P_\ell(t)|$ , when  $t$  decreases from 1 to 0. Then,

$$0 < p_{[\frac{\ell}{2}]}(\ell) < \cdots < p_2(\ell) < p_1(\ell) < 1. \quad (2.14)$$

Here  $[\cdot]$  denotes the integer part.

(vii) For the derivative of the Legendre polynomial  $P_\ell(t)$ , one has the inequality,

$$|P'_\ell(t)| \leq \frac{\ell(\ell+1)}{2} \text{ for } -1 \leq t \leq 1, \quad (2.15)$$

where the equality is achieved for  $\ell = 0, 1$  and when  $\ell \geq 2$ , for  $t = \pm 1$ .

For these properties, we refer to [13, Chapters IV and V] and to [20]. In particular, Properties (v)-(vii) can be found in [20], resp. under Theorems 6.21.2, 7.3.1, and Inequality (7.33.8).

**Remarks.** (i) Using (2.7), one can prove that for  $1 \leq j \leq \ell-1$ ,

$$\vartheta_j(\ell) < \vartheta_j(\ell-1) < \vartheta'_j(\ell). \quad (2.16)$$

(ii) One can also relate the asymptotic behavior of  $\vartheta_1(\ell)$  as  $\ell \rightarrow +\infty$ , to the first zero  $\mathbf{j}$  of the zero-th Bessel function  $J_0$ , [20, Theorem 8.1.2],

$$\vartheta_1(\ell) \sim \mathbf{j}/\ell. \quad (2.17)$$

(iii) One can relate the asymptotic behavior of  $p_1(\ell)$  as  $\ell \rightarrow +\infty$ , to the first zero  $\mathbf{j}_1$  of the Bessel function  $J_1 = -J'_0$ ,

$$p_1(\ell) \sim -J_0(\mathbf{j}_1). \quad (2.18)$$

### 3 Stern's first theorem: odd case

The purpose of this section is to prove Theorem 1.2. As a matter of fact, we shall prove a more quantitative result, Proposition 3.3, which implies the theorem. We use Stern's ideas sketched in the introduction.

Fix an integer  $\ell \in \mathbb{N}$ , without any parity assumption for the time being.

### 3.1 Notation

Up to scaling, there is a unique spherical harmonic  $Z_\ell$ , of degree  $\ell$ , which is invariant under the rotations about the  $z$ -axis. Viewed in the spherical coordinates, this zonal spherical harmonic is given by  $P_\ell(\cos \vartheta)$ . Let  $\vartheta_1(\ell) < \vartheta_2(\ell) < \dots < \vartheta_\ell(\ell)$  be the zeros of the function  $\vartheta \rightarrow P_\ell(\cos \vartheta)$  in the interval  $(0, \pi)$ , see Properties 2.1 (iii). The nodal set of the spherical harmonic  $Z_\ell$ , denoted  $N(Z_\ell)$ , consists of precisely  $\ell$  latitude circles (parallels),

$$L_i := \{(\vartheta, \varphi) \mid \vartheta = \vartheta_i(\ell)\}, \quad 1 \leq i \leq \ell.$$

They determine sectors on the sphere,

$$\mathcal{L}_i := \{(\vartheta, \varphi) \mid \vartheta_i(\ell) < \vartheta < \vartheta_{i+1}(\ell)\}, \quad 0 \leq i \leq \ell,$$

where  $\vartheta_0(\ell) = 0$  and  $\vartheta_{\ell+1}(\ell) = \pi$ . In the sector  $\mathcal{L}_i$ , the function  $Z_\ell$  has the sign of  $(-1)^i$ .

Call  $W_\ell$  the spherical harmonic of degree  $\ell$  obtained by restricting the harmonic homogeneous polynomial  $\Im(x+iy)^\ell$  to the sphere. Viewed in spherical coordinates, this spherical harmonic is given by  $\sin^\ell(\vartheta) \sin(\ell\varphi)$ . Its nodal set  $N(W_\ell)$  consists of  $\ell$  great circles of  $\mathbb{S}^2$ , *i.e.* of  $2\ell$  meridians,

$$M_j = \{(\vartheta, \varphi) \mid \varphi = j\frac{\pi}{\ell}\}, \quad 0 \leq j \leq 2\ell - 1.$$

They determine sectors on the sphere,

$$\mathcal{M}_j = \{(\vartheta, \varphi) \mid j\frac{\pi}{\ell} < \varphi < (j+1)\frac{\pi}{\ell}\} \quad 0 \leq j \leq 2\ell - 1.$$

In the sector  $\mathcal{M}_j$  the function  $W_\ell$  has the sign of  $(-1)^j$ .

Note that these meridians meet at the north and south poles  $p_\pm$  which are the only singular points of the nodal set  $N(W_\ell)$ .

The intersection

$$\mathcal{N} = N(Z_\ell) \cap N(W_\ell)$$

is the finite set of zeros common to  $Z_\ell$  and  $W_\ell$ .

We call  $q_{i,j}$  the intersection point of  $L_i$  with  $M_j$ ,  $1 \leq i \leq \ell$ ,  $0 \leq j \leq 2\ell - 1$ , so that

$$\mathcal{N} = \{q_{i,j}, \quad 1 \leq i \leq \ell, \quad 0 \leq j \leq 2\ell - 1\}.$$

For  $0 \leq i \leq \ell$  and  $0 \leq j \leq 2\ell - 1$ , we introduce the sets  $\mathcal{Q}_{i,j} = \mathcal{L}_i \cap \mathcal{M}_j$ , which are the connected components of the open set  $\{Z_\ell W_\ell \neq 0\}$ . In  $\mathcal{Q}_{i,j}$  the sign of the function  $Z_\ell W_\ell$  is  $(-1)^{i+j}$ .

The sets  $\mathcal{Q}_{i,j}$  form a *checkerboard* over the sphere, and following the idea of A. Stern, they can be colored according to the sign of the function  $Z_\ell W_\ell$ .

Finally, for  $0 \leq j \leq 2\ell - 1$ , we introduce the meridian  $B_j$ ,

$$B_j = \{(\vartheta, \varphi) \mid \varphi = (j + \frac{1}{2})\frac{\pi}{\ell}\}, \quad (3.1)$$

which bisects the sector  $\mathcal{M}_j$ .

Figure 3.1 displays the latitude circles and the meridians viewed through the exponential map at the north pole  $p_+$ , in the cases  $\ell = 3$  and  $\ell = 4$ . The common zeros of  $W_\ell$  and  $Z_\ell$  are the big dots. The coloring white/grey illustrates the sign of  $W_\ell Z_\ell$  (see below). The outer circle is mapped to the south pole by the exponential map.




Figure 3.1: Checkerboards in the cases  $\ell = 3$  and  $\ell = 4$ .

### 3.2 The family $H^{\mu,\ell}$

Following Stern [18], we consider the one-parameter family of spherical harmonics,

$$H^{\mu,\ell} = W_\ell + \mu Z_\ell, \quad (3.2)$$

which may be written in spherical coordinates as

$$h^{\mu,\ell}(\vartheta, \varphi) = \sin^\ell(\vartheta) \sin(\ell\varphi) + \mu P_\ell(\cos \vartheta). \quad (3.3)$$

Note that

$$h^{-\mu,\ell}(\vartheta, \varphi) = -h^{\mu,\ell}(\vartheta, \varphi + \frac{\pi}{\ell}), \quad (3.4)$$

for  $(\vartheta, \varphi) \in (0, \pi) \times \mathbb{R}_{2\pi}$ . It follows that we can restrict to the case  $\mu > 0$ . We shall do so for the remaining part of Section 3.

#### 3.2.1 Critical zeros of $H^{\mu,\ell}$

We call *critical zero* of a function a point which is both a zero and a critical point.

According to Properties 2.1 (ii),  $H^{\mu,\ell}(p_+) = \mu$ , and  $H^{\mu,\ell}(p_-) = (-1)^\ell \mu$ , and hence the north and south poles do not belong to the nodal set  $N(H^{\mu,\ell})$  when  $\mu \neq 0$ . As a consequence, for  $\mu > 0$ , the critical zeros of  $H^{\mu,\ell}$  are located in  $\mathbb{S}^2 \setminus \{p_\pm\}$ , and we can look for them in the spherical coordinates  $(\vartheta, \varphi) \in (0, \pi) \times \mathbb{R}_{2\pi}$ .

For  $\mu > 0$ , the point  $(\vartheta, \varphi)$  corresponds to a critical zero of  $H^{\mu,\ell}$ , if and only if

$$\begin{aligned} h^{\mu,\ell}(\vartheta, \varphi) &= 0, \\ \partial_\vartheta h^{\mu,\ell}(\vartheta, \varphi) &= 0, \\ \partial_\varphi h^{\mu,\ell}(\vartheta, \varphi) &= 0. \end{aligned} \tag{3.5}$$

This is equivalent for  $(\vartheta, \varphi)$  to satisfy the relations,

$$\begin{aligned} \cos(\ell\varphi) &= 0 \quad i.e. \quad \sin(\ell\varphi) = \pm 1, \\ \pm \sin^\ell(\vartheta) + \mu P_\ell(\cos \vartheta) &= 0, \\ \pm \ell \cos \vartheta \sin^{\ell-1}(\vartheta) - \mu \sin \vartheta P'_\ell(\cos \vartheta) &= 0. \end{aligned} \tag{3.6}$$

We plug (2.7) into the third line in the above system, and notice that the second line implies that  $\vartheta \neq 0, \pi$ . It follows that, for  $\mu \neq 0$ , (3.6) is equivalent to

$$\begin{aligned} \cos(\ell\varphi) &= 0 \quad i.e. \quad \sin(\ell\varphi) = \pm 1, \\ \frac{1}{\mu} &= \mp \frac{P_\ell(\cos \vartheta)}{\sin^\ell(\vartheta)}, \\ P_{\ell-1}(\cos \vartheta) &= 0. \end{aligned} \tag{3.7}$$

By Properties 2.1 (iii)-(iv), the last equation in (3.7) has exactly  $(\ell - 1)$  simple roots in  $[0, \pi]$ . We denote them by,  $\vartheta_1(\ell - 1) < \dots < \vartheta_{\ell-1}(\ell - 1)$ . They are symmetrical with respect to  $\frac{\pi}{2}$  due to the parity of  $P_{\ell-1}$ .

It follows that, for  $\mu > 0$ , the only possible critical zeros of the spherical harmonic  $H^{\mu,\ell}$  are given in spherical coordinates by the points  $(\vartheta_i(\ell - 1), (j + \frac{1}{2})\frac{\pi}{\ell})$  for  $1 \leq i \leq \ell - 1$  and  $0 \leq j \leq 2\ell - 1$ . These points can only occur as critical zeros for finitely many values of  $\mu$ , given by the second equation in (3.7). Away from these values of  $\mu$ , the spherical harmonic  $H^{\mu,\ell}$  has no critical zero.

Since we restrict to  $\mu > 0$ , the *critical* values of  $\mu$  are given by

$$\mu_i(\ell) = \frac{\sin^\ell(\vartheta_i(\ell - 1))}{|P_\ell(\cos \vartheta_i(\ell - 1))|}, \tag{3.8}$$

for  $1 \leq i \leq \ell - 1$ .

They are well-defined because the denominators do not vanish, since the zeros of  $P_\ell$  and  $P_{\ell-1}$  are intertwined, see Properties 2.1 (iv). For the value  $\mu_i$ , the spherical harmonic  $H^{\mu_i,\ell}$  has finitely many critical zeros which are well determined by equations (3.7). Note that the values  $\mu_i(\ell)$  are positive.

Taking the parity of the Legendre polynomials into account, it suffices to consider the values  $\mu_i(\ell)$  for  $1 \leq i \leq [\frac{\ell}{2}]$ , where  $[\frac{\ell}{2}]$  denotes the integer part of  $\frac{\ell}{2}$ . Let us summarize the preceding discussion in the following lemma.

**Lemma 3.1** Assume  $\mu > 0$ , and define  $\mu_c(\ell) > 0$  to be the infimum

$$\mu_c(\ell) = \inf_{1 \leq i \leq [\frac{\ell}{2}]} \mu_i(\ell), \quad (3.9)$$

where the positive values  $\mu_i(\ell)$  are given by (3.8).

The spherical harmonic

$$H^{\mu,\ell} = W_\ell + \mu Z_\ell$$

does not vanish at the north and south poles. Except for the values  $\{\mu_i\}_{i=1}^{[\frac{\ell}{2}]}$ ,  $H^{\mu,\ell}$  has no critical zero. In particular, for  $0 < \mu < \mu_c(\ell)$ , the function  $H^{\mu,\ell}$  has no critical zero, its nodal set is a 1-dimensional submanifold of the sphere, and hence consists of finitely many disjoint regular simple closed curves.

Note. The last assertion in the lemma follows from the fact that self-intersections in the nodal set of an eigenfunction correspond to critical zeros, see [2], Section 5.2.

**Remark.** One can easily estimate  $\mu_c(\ell)$  from below. Using Properties 2.1 (ii) and (vi), one finds that

$$\mu_i(\ell) > \frac{\sin^\ell(\vartheta_i(\ell-1))}{p_i(\ell)} > \frac{\sin^\ell(\vartheta_1(\ell-1))}{p_1(\ell)}.$$

Hence,

$$\mu_c(\ell) \geq \frac{\sin^\ell(\vartheta_1(\ell-1))}{p_1(\ell)}. \quad (3.10)$$

Note that one also has,

$$\mu_c(\ell) \geq \min \left( \mu_1(\ell), \frac{\sin^\ell(\vartheta_2(\ell))}{p_2(\ell)} \right). \quad (3.11)$$

One can also obtain the asymptotics of  $\mu_1(\ell)$  as  $\ell$  tends to infinity. Recall that  $\frac{\pi}{2\ell+1} \leq \vartheta_1(\ell) \leq \frac{2\pi}{2\ell+1}$ , and use Hilb's formula [20, Theorems 8.21.6],

$$P_\ell(\cos \vartheta) = \left( \frac{\vartheta}{\sin \vartheta} \right)^{\frac{1}{2}} J_0 \left( (\ell + \frac{1}{2})\vartheta \right) + R(\vartheta),$$

with

$$R(\vartheta) = \mathcal{O}(\vartheta^2), \quad \text{if } |\vartheta| \leq C/\ell.$$

It follows that

$$\vartheta_1(\ell) = \frac{\mathbf{j}}{\ell + \frac{1}{2}} + \mathcal{O}(1/\ell^3),$$

where  $\mathbf{j}$  is the least positive zero of the Bessel function  $J_0$ .

Compute

$$P_\ell(\cos \vartheta_1(\ell-1)) = \left( 1 + \mathcal{O}\left(\frac{1}{\ell^2}\right) \right) J_0 \left( (\ell + \frac{1}{2}) \frac{\mathbf{j}}{\ell - \frac{1}{2}} \right) + \mathcal{O}\left(\frac{1}{\ell^2}\right) = \mathbf{j} J'_0(\mathbf{j}) / (\ell - \frac{1}{2}) + \mathcal{O}\left(\frac{1}{\ell^2}\right).$$

Recall that

$$\mu_1(\ell) = \sin^\ell(\vartheta_1(\ell - 1)) / |P_\ell(\cos \vartheta_1(\ell - 1))|.$$

Observe that

$$\sin(\vartheta_1(\ell - 1)) = \frac{\mathbf{j}}{\ell - \frac{1}{2}} + \mathcal{O}(1/\ell^3).$$

Taking the power  $\ell$ , the remainder term does not change the main term of the asymptotics,

$$\sin^\ell(\vartheta_1(\ell - 1)) \sim \left( \frac{\mathbf{j}}{\ell - \frac{1}{2}} \right)^\ell.$$

Finally, we obtain

$$\mu_1(\ell) \sim \left( \frac{\mathbf{j}}{\ell - \frac{1}{2}} \right)^{\ell-1} \frac{1}{|J'_0(\mathbf{j})|}.$$

It turns out that the second term in the right hand side of (3.11) is asymptotically bigger than the first one. It follows that the preceding formula holds with  $\mu_1(\ell)$  replaced by  $\mu_c(\ell)$  as well.

### 3.2.2 A separation lemma for $N(H^{\mu,\ell})$

For  $0 \leq j \leq 2\ell - 1$ , we look at the function  $H^{\mu,\ell}$  restricted to the meridian  $B_j$ . Let

$$b^{\mu,\ell,j} = H^{\mu,\ell}|_{B_j},$$

i.e.

$$b^{\mu,\ell,j}(\vartheta) = (-1)^j \sin^\ell(\vartheta) + \mu P_\ell(\cos \vartheta).$$

Recall the notation for  $\vartheta_1(\ell)$  and  $p_1(\ell)$ , see Properties 2.1, (iii) and (vi).

**Lemma 3.2** *The functions  $b^{\mu,\ell,j}$  satisfy the following properties.*

- (i) *For  $0 < \mu < \mu_c(\ell)$ , the function  $b^{\mu,\ell,j}$  does not vanish in the interval  $[\vartheta_1(\ell), \pi - \vartheta_1(\ell)]$ .*
- (ii) *When  $\ell$  and  $j$  are even,  $b^{\mu,\ell,j}(\vartheta) > 0$  in  $[0, \vartheta_1(\ell)] \cup [\pi - \vartheta_1(\ell), \pi]$ .*
- (iii) *When  $\ell$  is even and  $j$  odd, the function  $b^{\mu,\ell,j}(\vartheta)$  vanishes exactly once in each interval  $(0, \vartheta_1(\ell))$  and  $(\pi - \vartheta_1(\ell), \pi)$ .*
- (iv) *When  $\ell$  is odd and  $j$  even, the function  $b^{\mu,\ell,j}(\vartheta)$  is positive in  $[0, \vartheta_1(\ell)]$  and vanishes exactly once in  $(\pi - \vartheta_1(\ell), \pi)$ .*
- (v) *When  $\ell$  and  $j$  are odd, the function  $b^{\mu,\ell,j}(\vartheta)$  vanishes exactly once in  $(0, \vartheta_1(\ell))$  and is negative in  $[\pi - \vartheta_1(\ell), \pi]$ .*

The previous items describe the possible intersections of the nodal set  $N(H^{\mu,\ell})$  with the meridian  $B_j$  which bisects the sector  $\mathcal{M}_j$ .

**Proof.** (i) The function  $b^{\mu,\ell,j}$  vanishes in the given interval if and only if

$$\frac{P_\ell(\cos \vartheta)}{\sin^\ell(\vartheta)} = \frac{(-1)^{j+1}}{\mu}.$$

Call  $\beta_1(\vartheta)$  the function in the left hand side. Using (2.7), we obtain its derivative,  $\beta'_1(\vartheta) = -\frac{\ell P_\ell(\cos \vartheta)}{\sin^{\ell+1}(\vartheta)}$ . It follows that the local extrema of the function  $\beta_1$  are the numbers,

$$b^{\mu,\ell,j}(\vartheta_i(\ell-1)) = (-1)^j \sin^\ell(\vartheta_i(\ell-1)) \left[ 1 + (-1)^j \mu \frac{P_\ell(\cos \vartheta_i(\ell-1))}{\sin^\ell(\vartheta_i(\ell-1))} \right].$$

Using (3.8), the assertion follows. To prove assertions (ii)-(v), look at the signs of the functions  $P_\ell(\cos \vartheta)$  and  $P'_\ell(\cos \vartheta)$  in the intervals  $(0, \vartheta_1(\ell))$  and  $(\pi - \vartheta_1(\ell), \pi)$  in order to determine the signs of  $b^{\mu,\ell,j}(\vartheta)$  and  $\partial_\vartheta b^{\mu,\ell,j}(\vartheta)$ . We leave the details to the reader.  $\square$

### 3.2.3 General properties of $N(H^{\mu,\ell})$

We now state simple general properties of the nodal set of the spherical harmonic  $H^{\mu,\ell} = W_\ell + \mu Z_\ell$ . We use the notation of Subsection 3.1.

**Properties 3.3** For  $\mu > 0$ , the nodal sets of the spherical harmonics  $H^{\mu,\ell}$  share the following properties.

(i) The nodal set of  $H^{\mu,\ell}$  satisfies

$$\mathcal{N} \subset N(H^{\mu,\ell}) \subset \mathcal{N} \cup \{Z_\ell W_\ell < 0\}.$$

This means that a point in the nodal set of  $H^{\mu,\ell}$  is either one of the points in  $\mathcal{N}$ , or a point in some open domain  $\mathcal{Q}_{i,j}$ , with  $(-1)^{i+j} = -1$ .

- (ii) The nodal set of  $H^{\mu,\ell}$  near each point  $q_{i,j} \in \mathcal{N}$  consists of a single regular arc which is transversal to the latitude circle  $L_i$  and to the meridian  $M_j$ . In other words, an arc in the nodal set inside some domain  $\mathcal{Q}_{i,j}$ , with  $(-1)^{i+j} = -1$ , can only enter/exit  $\mathcal{Q}_{i,j}$  through a point in  $\mathcal{N}$  (a vertex), and cannot cross the boundary of  $\mathcal{Q}_{i,j}$  elsewhere.
- (iii) For  $0 < \mu < \mu_c(\ell)$  (defined in Lemma 3.1), no connected component of the nodal set  $N(H^{\mu,\ell})$  can be entirely contained in some  $\mathcal{Q}_{i,j}$ .

**Proof.** Property (i) is clear. Property (ii) follows from the fact that both  $\partial_\vartheta h^{\mu,\ell}$  and  $\partial_\varphi h^{\mu,\ell}$  do not vanish at the points  $(\vartheta_i, j \frac{\pi}{\ell})$ . Indeed,  $\mu > 0$ , and  $P_\ell$  and  $P'_\ell$  have no common zero according to Properties 2.1 (iv). For the proof of Property (iii), we observe that the assumption on  $\mu$  implies that the nodal set  $N(H^{\mu,\ell})$  is a 1-dimensional submanifold of the sphere, *i.e.* a finite collection of disjoint regular simple closed curves. Assume that one

such closed curve is entirely contained in some domain  $\mathcal{Q}_{i,j}$ . This domain would contain some nodal domain of  $H^{\mu,\ell}$ , and hence its first Dirichlet eigenvalue  $\lambda$  would be strictly less than  $\ell(\ell+1)$ , the eigenvalue associated with  $H^{\mu,\ell}$ . On the other-hand, since  $\mathcal{Q}_{i,j}$  is contained in one of the nodal domains of  $Z_\ell$  (or of  $W_\ell$ ), its first eigenvalue satisfies  $\lambda > \ell(\ell+1)$ . This leads to a contradiction.  $\square$

### 3.2.4 Local nodal patterns for $H^{\mu,\ell}$

Assume that  $0 < \mu < \mu_c(\ell)$ .

Figure 3.2 is drawn in the case  $\ell = 4$ , but aims at illustrating the possible local nodal patterns in the general case. We look at square-like domains  $\mathcal{Q}_{i,j}$  which stay away from the poles, and can be visited by the nodal set. This means that  $1 \leq i \leq \ell-1$ ,  $0 \leq j \leq 2\ell-1$  and that  $(-1)^{i+j} = -1$ , corresponding to the white domains on the checkerboard, see Properties 3.3 (i).




Figure 3.2: Local nodal patterns away from the poles.

The local nodal pattern at the vertices is shown in the domain labelled (A). According to Lemma 3.1, and our assumption on  $\mu$ , the nodal set  $N(H^{\mu,\ell})$  consists of finitely many disjoint simple closed regular curves. According to Properties 3.3 (ii), it follows from Jordan's separation theorem that any such nodal curve can only enter a domain  $\mathcal{Q}_{i,j}$  at a vertex, and exit at another one. Taking into account Properties 3.3 (iii), this leaves exactly three possibilities for the nodal pattern in a domain  $\mathcal{Q}_{i,j}$ , illustrated in the domains

labelled (B), (C) and (D). According to the separation Lemma 3.2, both (C) and (D) are impossible. Notice that case (D) could also be discarded by the fact that the nodal curves do not intersect (absence of critical zeros). Finally, the only possible local nodal pattern in the square-like domain  $\mathcal{Q}_{i,j}$  is the one shown in (B): the nodal curves “follow” the meridians.

Remark. In Stern’s thesis this conclusion follows from the claim that the nodal set depends continuously on  $\mu$  and that  $\mu$  is small enough.

Figure 3.3 is drawn in the case  $\ell = 3$ , but aims at illustrating the possible local nodal patterns in the general case. We look at triangle-like domains  $\mathcal{Q}_{i,j}$  one of whose vertices is at the north or south pole and can be visited by the nodal set. This means that  $i = 0$  or  $\ell$ ,  $0 \leq j \leq 2\ell - 1$  and that  $(-1)^{i+j} = -1$ , see Properties 3.3 (i).




Figure 3.3: Local nodal patterns near the poles.

The same arguments as above show that there is only one possible nodal pattern.

### 3.2.5 A. Stern’s first theorem for the sphere

We can now state the following quantitative version of A. Stern’s first theorem, see Theorem 1.2. Recall the notations  $\vartheta_1(\ell)$  and  $p_1(\ell)$ , see Properties 2.1, (iii) and (vi).

**Proposition 3.4** *Assume that  $0 < \mu < \mu_c(\ell)$ .*

- (i) *When  $\ell$  is odd, the nodal set  $N(H^{\mu,\ell})$  is a unique regular simple closed curve and hence, the eigenfunction  $H^{\mu,\ell}$  has exactly two nodal domains.*

- (ii) When  $\ell$  is even, the nodal set  $N(H^{\mu,\ell})$  is the union of  $\ell$  regular disjoint simple closed curves and hence, the eigenfunction  $H^{\mu,\ell}$  has exactly  $(\ell + 1)$  nodal domains.

**Proof.** According to the remark following Lemma 3.1, under the assumption on  $\mu$ , the nodal set of  $H^{\mu,\ell}$  is a regular 1-dimensional submanifold. Since the eigenfunction  $H^{\mu,\ell}$  does not vanish at the north and south poles, we can work in the exponential map at the north pole. For the proofs below, keep in mind Section 3.2.4.

*Proof of Proposition 3.4, Assertion (ii).* The integer  $\ell$  is assumed to be even. The proof is illustrated by Figure 3.4 which shows parts of the nodal patterns of  $Z_\ell$  (latitude circles) and  $W_\ell$  (meridians) viewed in the exponential map centered at the north pole  $p_+$ . The south pole corresponds to the outer circle (the boundary of the maximal domain in which the exponential map is a diffeomorphism). The meridian  $M_j$  is labelled  $[j]$ ; the meridian  $B_j$  is labelled  $\{j\}$ .




Figure 3.4: Case  $\ell$  even,  $j$  odd.

Call  $B'_j$  the intersection

$$B'_j = B_j \cap \{\vartheta_1(\ell) \leq \vartheta \leq \pi - \vartheta_1(\ell)\}.$$

We now use Lemma 3.2.

- (a) When  $j$  is even, the function  $b^{\mu,\ell,j}(\vartheta)$  is positive in  $[0, \vartheta_1(\ell)] \cup [\pi - \vartheta_1(\ell), \pi]$ , and hence, by Lemma 3.2,  $N(H^{\mu,\ell}) \cap B_j = \emptyset$ . When  $j$  is odd, the function  $b^{\mu,\ell,j}(\vartheta)$  has exactly one zero in each of the intervals  $(0, \vartheta_1(\ell))$  and  $(\pi - \vartheta_1(\ell), \pi)$ . It follows that  $N(H^{\mu,\ell}) \cap B_j$  consists of exactly two points, one in  $\mathcal{Q}_{0,j}$  and one in  $\mathcal{Q}_{\ell,j}$ .

(b) Choose  $j = 2k + 1$ , odd (see Figure 3.4). Note that there are exactly  $\ell$  such values of  $j$  between 0 and  $2\ell - 1$ . In  $\mathcal{Q}_{0,j}$  the nodal set  $N(H^{\mu,\ell})$  can only consist of a curve from the point  $q_{1,j}$  to the point  $q_{1,j+1}$  (see Subsection 3.1 for the notation, and use Properties 3.3), intersecting  $B_j$  at exactly one point. This curve is part of a connected component (a simple closed curve)  $\gamma_k \subset N(H^{\mu,\ell})$ . We now follow the curve  $\gamma_k$ , starting from  $q_{1,j}$  in the direction of  $q_{1,j+1}$ . According to the preceding point (a),  $\gamma_k$  can meet neither  $B_{j+1}$ , nor  $B'_j$ . Therefore, according to Paragraph 3.2.4, the curve has to go through the points  $q_{1,j+1}, q_{2,j+1}, \dots, q_{\ell,j+1}$  passing alternatively inside  $\mathcal{M}_{j+1}$  or  $\mathcal{M}_j$ . Since  $\ell$  is even, at  $q_{\ell,j+1}$ , the curve enters  $\mathcal{Q}_{\ell,j}$ , crosses  $B_j$  (at a single point), and exits  $\mathcal{Q}_{\ell,j}$  at  $q_{\ell,j}$ . Since it can cross neither  $B_{j-1}$ , nor  $B'_{j-1}$ , the curve  $\gamma_k$  has to go back to  $q_{1,j}$ , through the points  $q_{\ell-1,j}, \dots, q_{2,j}$ , alternatively inside  $\mathcal{M}_j$  or  $\mathcal{M}_{j-1}$ . This means that the simple closed curve  $\gamma_k$  goes through all the points in  $\mathcal{N} \cap \mathcal{M}_j$ , with  $j = 2k + 1$ .

(c) In this way, we obtain  $\ell$  simple closed curves  $\gamma_1, \dots, \gamma_\ell$  which are connected components of  $N(H^{\mu,\ell})$ , with the curve  $\gamma_k$  (where  $j = 2k + 1$ ) contained in the sector bounded by the meridians  $B_{j-1}$  and  $B_{j+1}$  and containing  $B_j$ . Furthermore, these  $\ell$  curves visit all the points  $q_{i,j} \in \mathcal{N}$ . It follows from Properties 3.3 (iii) that there can be no other components, and that

$$N(H^{\mu,\ell}) = \bigcup_{k=1}^{\ell} \gamma_k.$$

This finishes the proof of Proposition 3.4, Assertion (ii).  $\square$

*Proof of Proposition 3.4, Assertion (i).* The integer  $\ell$  is now assumed to be odd. The proof is illustrated by Figure 3.5 which shows parts of the nodal patterns of  $Z_\ell$  (latitude circles) and  $W_\ell$  (meridians) viewed in the exponential map centered at the north pole  $p_+$ . The south pole corresponds to the outer circle (the boundary of the domain in which the exponential map is a diffeomorphism). The meridian  $M_j$  is labelled  $[j]$ ; the meridian  $B_j$  is labelled  $\{j\}$ .

As in the previous proof, call  $B'_j$  the intersection

$$B'_j = B_j \cap \{\vartheta_1(\ell) \leq \vartheta \leq \pi - \vartheta_1(\ell)\}.$$

(a) When  $j$  is even, the function  $b^{\mu,\ell,j}(\vartheta)$  is positive in  $[0, \vartheta_1]$ , and admits exactly one zero in the interval  $(\pi - \vartheta_1, \pi)$ . Hence  $N(H^{\mu,\ell}) \cap B_j$  contains exactly one point located in  $\mathcal{Q}_{\ell,j}$ . When  $j$  is odd, the function  $b^{\mu,\ell,j}(\vartheta)$  has exactly one zero in the interval  $(0, \vartheta_1)$  and is negative in  $[\pi - \vartheta_1, \pi]$ . By Lemma 3.2, it follows that  $N(H^{\mu,\ell}) \cap B_j$  consists of exactly one point located in  $\mathcal{Q}_{0,j}$ .

(b) Choose  $j = 1$ . In  $\mathcal{Q}_{0,1}$  the nodal set  $N(H^{\mu,\ell})$  can only consist of a curve from the point  $q_{1,1}$  to the point  $q_{1,2}$  (see Subsection 3.1 for the notation and use Properties 3.3), intersecting  $B_1$  at exactly one point. This curve is part of a connected component (a simple closed curve)  $\gamma \subset N(H^{\mu,\ell})$ . We now follow the curve  $\gamma$ , starting from  $q_{1,1}$  in the direction of  $q_{1,2}$ . According to the preceding point (a),  $\gamma$  can meet neither  $B'_2$ , nor  $B_1$ , so that it has to go through the points  $q_{1,2}, q_{2,2}, \dots, q_{\ell,2}$  passing alternatively inside  $\mathcal{M}_2$  or  $\mathcal{M}_1$ . Because  $\ell$  is odd, at  $q_{\ell,2}$ , the curve exits  $\mathcal{Q}_{\ell-1,1}$ , enters  $\mathcal{Q}_{\ell,2}$ , crosses  $B_2$  (at a single point), and exits  $\mathcal{Q}_{\ell,2}$  at  $q_{\ell,3}$  into  $\mathcal{M}_3$ . Since it can cross neither  $B'_3$ , nor  $B'_2$ , the curve  $\gamma$  has to go to  $q_{1,3}$ , through the points  $q_{\ell-1,3}, \dots, q_{2,3}$ , alternatively inside  $\mathcal{M}_2$  or  $\mathcal{M}_3$ . The




Figure 3.5: Case  $\ell$  odd.

curve  $\gamma$  therefore goes from  $q_{1,1}$  to  $q_{1,3}$  where we can start again with the same argument as before. Iterating  $\ell$  times the argument, the curve  $\gamma$  gets back to its initial point  $q_{1,1}$ .

(c) In this way, we obtain a simple closed curve  $\gamma$  in  $N(H^{\mu,\ell})$ , which crosses all the meridians  $B_j$  once, and which visits all the points  $q_{i,j} \in \mathcal{N}$ . It follows from Properties 3.3 (iii) that there can be no other component and that

$$N(H^{\mu,\ell}) = \gamma.$$

This finishes the proof of Proposition 3.4, Assertion (i).  $\square$

It is easy to follow the above proofs on Figure 3.6 which shows the nodal set of  $H^{\mu,\ell}$ , in the exponential map at  $p_+$ , for  $\mu > 0$  small enough and for  $\ell = 3$  (left) and  $\ell = 4$  (right).

## 4 Stern's second theorem: even case

The purpose of this section is to prove Theorem 1.3. As a matter of fact, we shall give a more quantitative result, Proposition 4.5, which implies the theorem. As in Section 3, we follow the ideas of A. Stern sketched in the introduction.

Fix an integer

$$\ell = 2r \geq 2,$$




Figure 3.6: Cases  $\ell = 3$  and  $\ell = 4$ .

as well as an angle  $\alpha$  defined by

$$\alpha = \frac{\epsilon\pi}{2r}, \text{ with } 0 < \epsilon < \frac{1}{2}. \quad (4.1)$$

## 4.1 Notation

As in the first example, we consider the spherical harmonic of degree  $\ell = 2r$

$$W(x, y, z) = \Im(x + iy)^{2r}, \quad (4.2)$$

whose expression in spherical coordinates  $(\vartheta, \varphi) \in (0, \pi) \times \mathbb{R}_{2\pi}$  is given by

$$w(\vartheta, \varphi) = \sin^{2r}(\vartheta) \sin(2r\varphi). \quad (4.3)$$

The perturbation of  $W$  is chosen to be the spherical harmonic  $V_\alpha$ , of degree  $2r$ , whose expression in spherical coordinates is given by

$$v_\alpha(\vartheta, \varphi) = P_{2r}^1(\cos \vartheta) \sin(\varphi - \alpha). \quad (4.4)$$

According to Properties 2.1 (i), we have  $P_{2r}^1(t) = -(1 - t^2)^{1/2} \frac{d}{dt} P_{2r}(t)$ , so that

$$v_\alpha(\vartheta, \varphi) = -\sin \vartheta P'_{2r}(\cos \vartheta) \sin(\varphi - \alpha). \quad (4.5)$$

According to (2.1), the corresponding harmonic homogeneous polynomial of degree  $2r$  in  $\mathbb{R}^3$  is given by the formula

$$V_\alpha(x, y, z) = (\sin \alpha x - \cos \alpha y) \sum_{j=0}^{r-1} a_j z^{2r-2j-1} (x^2 + y^2 + z^2)^j, \quad (4.6)$$

where the  $a_j$ 's are the coefficients of the polynomial  $P'_{2r}$ ,

$$P'_{2r}(t) = \sum_{j=0}^{r-1} a_j t^{2r-2j-1}.$$

The nodal set of the spherical harmonic  $W$  consists of the  $2\ell = 4r$  meridians  $M_j$ ,  $0 \leq j \leq 4r - 1$ , defined as in Subsection 3.1,

$$N(W) = \bigcup_{j=0}^{4r-1} M_j ,$$

with the corresponding open sectors  $\mathcal{M}_j$  on the sphere.

These meridians meet at the north and south poles which are the only critical zeros of  $W$ ,  $W(p_\pm) = 0$ , and  $d_{p_\pm} W = 0$  (the differential of the function  $W$  at the poles).

The nodal set of the spherical harmonic  $V_\alpha$  consists of  $(2r - 1)$  latitude circles  $L'_i$  ( $1 \leq i \leq 2r - 1$ ), and two meridians  $M'_0$  and  $M'_1$ ,

$$N(V_\alpha) = \bigcup_{j=0}^{2r-1} L'_i \bigcup M'_0 \bigcup M'_1 . \quad (4.7)$$

The latitude circles,

$$L'_i = \{(\vartheta, \varphi) \mid \vartheta = \vartheta'_i(2r)\} , \quad 1 \leq i \leq 2r - 1 , \quad (4.8)$$

are associated with the  $(2r - 1)$  zeros,  $0 < \vartheta'_1(2r) < \dots < \vartheta'_{2r-1}(2r) < \pi$ , of the function  $P'_{2r}(\cos \vartheta)$ , see Properties 2.1 (iii), and we let  $\vartheta'_0(2r) = 0$  and  $\vartheta'_{2r}(2r) = \pi$ . They determine sectors

$$\mathcal{L}'_i = \{(\vartheta, \varphi) \mid \vartheta'_i(2r) < \vartheta < \vartheta'_{i+1}(2r)\} , \quad 0 \leq i \leq 2r - 1 . \quad (4.9)$$

The meridians  $M'_k$  are given by

$$M'_0 = \{(\vartheta, \varphi) \mid \varphi = \alpha\} \text{ and } M'_1 = \{(\vartheta, \varphi) \mid \varphi = \alpha + \pi\} . \quad (4.10)$$

They determine sectors

$$\mathcal{M}'_0 = \{(\vartheta, \varphi) \mid \alpha < \varphi < \alpha + \pi\} \text{ and } \mathcal{M}'_1 = \{(\vartheta, \varphi) \mid \alpha + \pi < \varphi < 2\pi + \alpha\} . \quad (4.11)$$

Figure 4.1 shows the nodal sets  $N(W)$  and  $N(V_\alpha)$ , in the case  $\ell = 2r = 4$ . They are viewed in the exponential map  $\exp_{p_+}$ , *i.e.* in the disk  $D(0, \pi)$ , whose boundary corresponds to the cut-locus of  $p_+$  *i.e.*  $p_-$ . In the figures, the meridians  $M_j$  are labelled  $[j]$ ; the latitude circles  $L'_i$  are labelled  $\{i'\}$ , and the meridians  $M'_k$  are labelled  $[k']$ .

As in the first example, the set  $\mathcal{N} = N(W) \cap N(V_\alpha)$  of common zeros to the spherical harmonics  $W$  and  $V_\alpha$ , plays a special role. We have

$$\mathcal{N} = \{p_+, p_-\} \bigcup \{q_{i,j} \mid 1 \leq i \leq 2r - 1, 0 \leq j \leq 4r - 1\} , \quad (4.12)$$




Figure 4.1: Nodal sets  $N(W)$  and  $N(V_\alpha)$ , when  $\ell = 2r = 4$ .

where  $q_{i,j}$  is the intersection point of the latitude circle  $L'_i$  with the meridian  $M_j$ . Figure 4.2 shows the set  $\mathcal{N}$  in the exponential map. The points in  $\mathcal{N}$  appear as the big dots : the intersection points of the latitude circles  $L'_i$ , with the meridians  $M_j$ , and the poles. Note that the south pole is represented by two small dots, one on the meridian  $M'_0$ , one on the meridian  $M'_1$ . Note that there are no other dots on these meridians, see Properties 4.4 (iii).

We also introduce the connected components of the set  $\{W V_\alpha \neq 0\}$ ,

$$\mathcal{Q}_{i,j,k} = \mathcal{L}'_i \cap \mathcal{M}_j \cap \mathcal{M}'_k, \quad 0 \leq i \leq 2r-1, \quad 0 \leq j \leq 4r-1, \quad k = 0, 1. \quad (4.13)$$

Note that

$$\operatorname{sgn}(W V_\alpha) = (-1)^{i+j+k+1} \text{ on } \mathcal{Q}_{i,j,k}. \quad (4.14)$$




Figure 4.2: The set  $\mathcal{N}$  in the case  $\ell = 2r = 4$ .

## 4.2 The family $H^\mu$

### 4.2.1 Definition

Following the ideas of Stern [18], we consider the one-parameter family of spherical harmonics of degree  $\ell = 2r$ ,

$$H^\mu(x, y, z) = W(x, y, z) - \mu V_\alpha(x, y, z), \quad (4.15)$$

whose expression in spherical coordinates is given by

$$\begin{aligned} h^\mu(\vartheta, \varphi) &= w(\vartheta, \varphi) - \mu v_\alpha(\vartheta, \varphi), \\ &= \sin^{2r}(\vartheta) \sin(2r\varphi) + \mu \sin \vartheta P'_{2r}(\cos \vartheta) \sin(\varphi - \alpha). \end{aligned} \quad (4.16)$$

Note that

$$h^{-\mu}(\vartheta, \varphi) = h^\mu(\vartheta, \varphi + \pi). \quad (4.17)$$

It follows that it suffices to consider the case  $\mu > 0$ . We shall therefore assume that  $\mu > 0$  for the remainder of Section 4.

### 4.2.2 Critical zeros

We now investigate the critical zeros of  $H^\mu$ . The spherical harmonic  $W$  vanishes at order at least 3 at the poles, while the nodal set of  $V_\alpha$  is a piece of great circle at each pole. It follows that the north and south poles are not critical points of  $N(H^\mu)$ , see also Properties 4.4 (i). We can therefore look for critical zeros in the spherical coordinates, *i.e.* look for critical zeros of  $h^\mu$  in  $(0, \pi) \times \mathbb{R}_{2\pi}$ .

The point  $(\vartheta, \varphi) \in (0, \pi) \times \mathbb{R}_{2\pi}$  is a critical zero of  $h^\mu$  if and only if,

$$\begin{aligned} h^\mu(\vartheta, \varphi) &= 0, \\ \partial_\vartheta h^\mu(\vartheta, \varphi) &= 0, \\ \partial_\varphi h^\mu(\vartheta, \varphi) &= 0. \end{aligned} \tag{4.18}$$

Using the second order differential equation (2.4) satisfied by the Legendre polynomial  $P_{2r}$ , we find that

$$\begin{aligned} \partial_\vartheta h^\mu(\vartheta, \varphi) &= 2r \cos \vartheta \sin^{2r-1}(\vartheta) \sin(2r\varphi) \\ &\quad + \mu \sin(\varphi - \alpha) [2r(2r+1)P_{2r}(\cos \vartheta) - \cos \vartheta P'_{2r}(\cos \vartheta)]. \end{aligned}$$

It follows that the point  $(\vartheta, \varphi) \in (0, \pi) \times \mathbb{R}_{2\pi}$  is a critical zero of  $h^\mu$  if and only if,

$$\begin{aligned} \sin^{2r-1}(\vartheta) \sin(2r\varphi) + \mu P'_{2r}(\cos \vartheta) \sin(\varphi - \alpha) &= 0, \\ 2r \cos \vartheta \sin^{2r-1}(\vartheta) \sin(2r\varphi) \\ + \mu \sin(\varphi - \alpha) [2r(2r+1)P_{2r}(\cos \vartheta) - \cos \vartheta P'_{2r}(\cos \vartheta)] &= 0, \\ 2r \sin^{2r-1}(\vartheta) \cos(2r\varphi) + \mu P'_{2r}(\cos \vartheta) \cos(\varphi - \alpha) &= 0. \end{aligned} \tag{4.19}$$

The pair of the first and third equations in (4.19) is equivalent to the pair of the first and third equations in (4.20) below. Plugging the first equation in (4.19) into the second one, using the fact that  $\mu > 0$  and that  $\vartheta \in (0, \pi)$ , we get the second equation in (4.20). It follows that the point  $(\vartheta, \varphi) \in (0, \pi) \times \mathbb{R}_{2\pi}$  is a critical zero of  $h^\mu$  if and only if,

$$\begin{aligned} \mu P'_{2r}(\cos \vartheta) + \sin^{2r-1}(\vartheta) [2r \cos(2r\varphi) \cos(\varphi - \alpha) + \sin(2r\varphi) \sin(\varphi - \alpha)] &= 0, \\ \sin(\varphi - \alpha) [2r P_{2r}(\cos \vartheta) - \cos \vartheta P'_{2r}(\cos \vartheta)] &= 0, \\ 2r \cos(2r\varphi) \sin(\varphi - \alpha) - \sin(2r\varphi) \cos(\varphi - \alpha) &= 0. \end{aligned} \tag{4.20}$$

**Property 4.1** *Assume that  $\mu > 0$ . Then, the product  $\sin(2r\varphi) \sin(\varphi - \alpha)$  does not vanish at the critical zeros of  $H^\mu$ .*

This property follows immediately from the third equation in (4.20).

Finally, it follows that the point  $(\vartheta, \varphi) \in (0, \pi) \times \mathbb{R}_{2\pi}$  is a critical zero of  $h^\mu$  if and only if,

$$\begin{aligned} \mu P'_{2r}(\cos \vartheta) + \sin^{2r-1}(\vartheta) [2r \cos(2r\varphi) \cos(\varphi - \alpha) + \sin(2r\varphi) \sin(\varphi - \alpha)] &= 0, \\ 2r P_{2r}(\cos \vartheta) - \cos \vartheta P'_{2r}(\cos \vartheta) &= 0, \\ 2r \cos(2r\varphi) \sin(\varphi - \alpha) - \sin(2r\varphi) \cos(\varphi - \alpha) &= 0. \end{aligned} \tag{4.21}$$

We first analyze the second equation in (4.21). Define  $Q(t) := 2rP_{2r}(t) - tP'_{2r}(t)$ . This is an even polynomial of degree less than or equal to  $(2r - 2)$ . For parity reasons the roots of the polynomials  $Q$ ,  $P_{2r}$  and  $P'_{2r}$  are symmetric with respect to 0, and it suffices to look at  $t \geq 0$ . According to Properties 2.1 (iii), the positive roots  $t_i$  of  $P_{2r}$ , and  $t'_i$  of  $P'_{2r}$  satisfy

$$0 = t'_r < t_r < t'_{r-1} < t_{r-1} < \cdots < t_2 < t'_1 < t_1 < 1.$$

The following equalities are easy to check,

$$\begin{aligned} \operatorname{sgn}(P_{2r}(t'_i)) &= (-1)^i \text{ and } \operatorname{sgn}(P'_{2r}(t_i)) = (-1)^{i-1}, \\ \operatorname{sgn}(Q(t_i)) &= (-1)^i \text{ and } \operatorname{sgn}(Q(t'_{i-1})) = (-1)^{i-1}. \end{aligned}$$

It follows that  $Q$  vanishes at least once in each interval  $(t_{i+1}, t'_i)$  for  $1 \leq i \leq r - 1$ . Since  $Q$  has at most  $(r - 1)$  non-negative zeros, we can conclude that  $Q$  has exactly  $(r - 1)$  zeros in  $(0, 1)$ , and more precisely one zero, which we denote by  $\cos \omega_i$ , in each interval  $(t_{i+1}, t'_i)$ , so that  $\omega_i \in (\vartheta'_i(2r), \vartheta_{i+1}(2r))$ , and

$$0 < \vartheta_1(2r) < \vartheta'_1(2r) < \omega_1 < \vartheta_2(2r) < \cdots < \vartheta'_{r-1}(2r) < \omega_{r-1} < \vartheta_r(2r) < \vartheta'_r(2r) = \frac{\pi}{2}.$$

Note that the inequalities are strict, *i.e.* that  $P_{2r}(\cos \omega_i) \neq 0$  and  $P'_{2r}(\cos \omega_i) \neq 0$ , and that the zeros  $\omega_i$  depend on  $\ell = 2r$ .

We now analyze the third equation in (4.21). Define the function,

$$f(\varphi) = 2r \cos(2r\varphi) \sin(\varphi - \alpha) - \sin(2r\varphi) \cos(\varphi - \alpha). \quad (4.22)$$

The function  $f$  satisfies  $f(\pi + \varphi) + f(\varphi) = 0$ , and  $f'(\varphi) = -(4r^2 - 1) \sin(2r\varphi) \sin(\varphi - \alpha)$ . An easy analysis in  $[0, \pi]$  (using the choice of  $\alpha$ ) shows that  $f$  does not vanish in  $[0, \frac{\pi}{2r}]$ , and has exactly one zero in each interval  $[j \frac{\pi}{2r}, (j + 1) \frac{\pi}{2r}]$  for  $1 \leq j \leq 2r - 1$ . It follows that  $f$  has exactly  $(4r - 2)$  zeros in  $[0, 2\pi]$ ,  $0 < \varphi_1 < \varphi_2 < \cdots < \varphi_{4r-2} < 2\pi$ , and that  $\varphi_j \in (j \frac{\pi}{2r}, (j + 1) \frac{\pi}{2r})$ .

The only possible critical zeroes of  $H^\mu$  are given in spherical coordinates by the points  $(\omega_i, \varphi_j)$  and  $(\pi - \omega_i, \varphi_j)$ , for  $1 \leq i \leq r - 1$  and  $1 \leq j \leq 4r - 2$ . These points can only occur as critical zeros for finitely many values of  $\mu$  given by the first equation in (4.21). Since we work with  $\mu > 0$ , these critical values of  $\mu$  are given (see also the first line in (4.19)) by

$$\mu_{i,j}(\alpha) = \left| \frac{\sin^{2r-1}(\omega_i) \sin(2r\varphi_j)}{P'_{2r}(\cos \omega_i) \sin(\varphi_j - \alpha)} \right|, \quad (4.23)$$

for  $1 \leq i \leq r - 1$ ,  $1 \leq j \leq 2r - 1$ , and can be numerically computed.

Let us summarize the preceding analysis.

**Lemma 4.2** *For  $\mu > 0$ , the spherical harmonic  $H^\mu$  has no critical zero except for finitely many values of  $\mu$  which are given by (4.23), and for which  $H^\mu$  has finitely many critical zeros. Define the number  $\mu_c(\alpha, 2r)$  to be*

$$\mu_c(\alpha, 2r) = \inf \mu_{i,j}(\alpha), \quad (4.24)$$

*where the infimum is taken over  $1 \leq i \leq r - 1$ , and  $1 \leq j \leq 2r - 1$ . Then, for  $0 < \mu < \mu_c(\alpha, 2r)$ , the function  $H^\mu$  has no critical zero, so that its nodal set  $N(H^\mu)$  consists of finitely many disjoint simple closed curves.*

**Remark.** One can bound  $\mu_c(\alpha, 2r)$  from below using the inequalities satisfied by the  $\omega_i$ , and Properties 2.1 (vii).

### 4.2.3 A separation lemma for $N(H^\mu)$

For  $1 \leq j \leq 4r - 2$ , call  $C_j$  the meridian,

$$C_j = \{(\vartheta, \varphi) \mid \varphi = \varphi_j\}, \quad (4.25)$$

where  $\varphi_j$  are the zeros of the function  $f$  defined in (4.22). We now look at the restriction of the spherical harmonic  $H^\mu$  to the meridian  $C_j$ . Recall from Properties 2.1 (iii), that  $\cos \vartheta'_1(2r)$  is the largest zero of the function  $P'_{2r}(\cos \vartheta)$  in  $[0, \pi]$ .

**Lemma 4.3** Define the functions,

$$\begin{aligned} b^{\mu,j}(\vartheta) &= h^\mu(\vartheta, \varphi_j), \\ &= \sin(2r\varphi_j) \sin^{2r}(\vartheta) + \mu \sin \vartheta P'_{2r}(\cos \vartheta) \sin(\varphi_j - \alpha). \end{aligned} \quad (4.26)$$

Assume that  $2r + 1 \leq j \leq 4r - 1$ .

- (i) For  $0 < \mu < \mu_c(\alpha, 2r)$ , the functions  $b^{\mu,j}(\vartheta)$  do not vanish in the interval  $[\vartheta'_1(2r), \pi - \vartheta'_1(2r)]$ .
- (ii) When  $j$  is odd, the function  $b^{\mu,j}(\vartheta)$  vanishes exactly once in the interval  $(0, \vartheta'_1(2r))$  and does not vanish in the interval  $[\pi - \vartheta'_1(2r), \pi]$ .
- (iii) When  $j$  is even, the function  $b^{\mu,j}(\vartheta)$  vanishes exactly once in the interval  $(\pi - \vartheta'_1(2r), \pi)$  and does not vanish in the interval  $[0, \vartheta'_1(2r)]$ .

The above assertions determine the possible intersections of the nodal set  $N(H^\mu)$  with the meridian  $C_j$ .

**Proof.** Notice that the assumptions on  $j$  and  $\alpha$  imply that  $\sin(\varphi_j - \alpha) < 0$ , and that  $(-1)^j \sin(2r\varphi_j) > 0$ . The function  $b^{\mu,j}$  vanishes at the points such that

$$\frac{\sin \vartheta P'_{2r}(\cos \vartheta)}{\sin^{2r}(\vartheta)} = -\frac{\sin(2r\varphi_j)}{\mu \sin(\varphi_j - \alpha)}.$$

Call  $\beta_2(\vartheta)$  the function in the left-hand side. Using the differential equation satisfied by  $P_{2r}$ , one finds that

$$\beta'_2(\vartheta) = (2r+1) \frac{2r P_{2r}(\cos \vartheta) - \cos \vartheta P'_{2r}(\cos \vartheta)}{\sin^{2r}(\vartheta)}.$$

The local extrema of  $\beta_2$  in the interval  $[\vartheta'_1(2r), \pi - \vartheta'_1(2r)]$  are achieved at the zeros  $\omega_i \in (\vartheta'_i(2r), \vartheta'_{i+1}(2r))$  of the second equation in (4.21), for  $1 \leq i \leq 2r - 2$ . We have at these points,

$$h^\mu(\omega_i, \varphi_j) = \sin^{2r}(2r\varphi_j) \sin(\omega_i) \left[ 1 + \mu \frac{P'_{2r}(\cos \omega_i) \sin(\varphi_j - \alpha)}{\sin(2r\varphi_j) \sin^{2r-1}(\omega_i)} \right].$$

The first assertion follows from (4.23).

We now look at what happens in the intervals  $(0, \vartheta'_1(2r))$  and  $(\pi - \vartheta'_1(2r), \pi)$ .

Write  $b^{\mu,j}(\vartheta) = (-1)^j \sin \vartheta f_j(\vartheta)$ . The derivative  $f'_j(\vartheta)$  is given by

$$f'_j(\vartheta) = (2r-1)(-1)^j \sin(2r\varphi_j) \cos \vartheta \sin^{2r-2}(\vartheta) - (-1)^j \mu \sin \vartheta P''_{2r}(\cos \vartheta) \sin(\varphi_j - \alpha).$$

Recall that  $P_{2r}$  and  $P''_{2r}$  are even functions and that  $P'_{2r}$  is odd. The largest zeros of these functions in  $[-1, 1]$  satisfy, with an obvious notation,

$$t_2 < t''_1 < t'_1 < t_1 < 1.$$

Looking at the signs of these functions in the various intervals between  $t_2$  and 1, and using the parity to look at what happens near  $-1$ , we can make the following observations.

- Case  $j$  odd. For  $\vartheta \in (0, \vartheta'_1)$ ,  $f_j(\vartheta) > 0$ . On the other-hand,  $f_j(\pi - \vartheta'_1) > 0$  and  $f_j(\pi) < 0$ , while  $f'_j(\vartheta) < 0$  in  $(\pi - \vartheta'_1, \pi)$ .
- Case  $j$  even. For  $\vartheta \in (\pi - \vartheta'_1, \pi)$ ,  $f_j(\vartheta) > 0$ . On the other-hand,  $f_j(0) < 0$  and  $f_j(\vartheta'_1) > 0$ , while  $f'_j > 0$  in  $(0, \vartheta'_1)$ .

The second and third assertion follows.  $\square$

#### 4.2.4 General properties of $N(H^\mu)$

**Properties 4.4** For  $\mu > 0$ , the nodal sets  $N(H^\mu)$  share the following properties.

- (i) The north and south poles are zeroes of order 1 of  $H^\mu$ ,  $H^\mu(p_\pm) = 0$ , and  $d_{p_\pm} H^\mu \neq 0$ . In particular, near each pole, the nodal set  $N(H^\mu)$  consists of a single arc, tangent to the great circle  $M'_0 \cup M'_1$ .
- (ii) The nodal set  $N(H^\mu)$  satisfies,

$$\mathcal{N} \subset N(H^\mu) \subset \mathcal{N} \bigcup \{W V_\alpha > 0\}. \quad (4.27)$$

- (iii) Since  $\alpha = \frac{\epsilon\pi}{2r}$ , with  $0 < \epsilon < \frac{1}{2}$ , the nodal set  $N(H^\mu)$  meets the great circle  $M'_0 \cup M'_1$  at the poles tangentially, and nowhere else.
- (iv) The connected components of  $N(H^\mu)$  are contained in either the closed hemisphere  $\overline{\mathcal{M}'_0}$  or in  $\overline{\mathcal{M}'_1}$ .
- (v) The points in  $\mathcal{N}$  (common zeros to the spherical harmonics  $W$  and  $V_\alpha$ ) are not critical zeros of  $H^\mu$ . At the points  $q_{i,j} \in \mathcal{N}$ ,  $1 \leq i \leq 2r-1$ ,  $0 \leq j \leq 4j-1$ , the nodal set  $N(H^\mu)$  consists of a single arc which is transversal to the latitude circles  $L'_i$  and to the meridians  $M_j$ .
- (vi) For  $0 < \mu < \mu_{a,\alpha}$  (defined in (4.24)), no closed component of the nodal set  $N(H^\mu)$  can be entirely contained in some domain  $\mathcal{Q}_{i,j,k}$ .

**Proof.** Assertion (i) follows from the fact that near the poles,  $N(V_\alpha)$  is a piece of great circle, while  $W$  vanishes at order at least 3. Assertion (ii) is clear (this is the checkerboard property introduced by A. Stern as we recalled in the introduction). Assertion (iii) is clear because the great circle  $M'_0 \cup M'_1$  only meets the nodal set  $N(W)$  at the poles. Assertion (iv) follows from Assertion (ii), the choice of  $\alpha$  and the parity of  $\ell = 2r$ . We can indeed look at a neighborhood of the north pole (the pattern near the south pole is the image of the pattern at the north pole under the antipodal map). The nodal curve at  $p_+$  must visit the domains  $\mathcal{Q}_{0,0,1}$  and  $\mathcal{Q}_{0,2r,1}$ , both in  $\mathcal{M}'_1$ , and cannot visit the domains  $\mathcal{Q}_{0,0,0}$  and  $\mathcal{Q}_{0,2r,0}$ . On the other-hand, as we already pointed out, the nodal set cannot meet the great circle  $M'_0 \cup M'_1$ . Assertion (v) follows by checking that the partial derivatives  $\partial_\theta h^\mu$  and  $\partial_\varphi h^\mu$  do not vanish at the points  $(\vartheta'_i, \frac{j\pi}{2r})$ . Assertion (vi) follows by using the same energy argument as in Properties 3.3.  $\square$

Figure 4.3 illustrates the proof of Properties 4.4. The checkerboard appears in white/grey (allowed/forbidden domains).




Figure 4.3: The checkerboard for  $\ell = 2r = 4$  and  $\mu > 0$ .

#### 4.2.5 Local nodal patterns for $H^\mu$

The arguments to determine the local nodal patterns for  $H^\mu$  are the same as in Paragraph 3.2.4, with an extra case. Namely, at each pole, the nodal set  $N(H^\mu)$  is a single arc tangent to the great circle  $M'_0 \cup M'_1$ , going through two triangle-like domains, one of whose

vertices is the pole. One of the two remaining vertices does not belong to  $\mathcal{N}$ , the other does belong to  $\mathcal{N}$  so that the local nodal pattern is well determined. See Figures 3.2, 3.3 and 4.3.

#### 4.2.6 A. Stern's second theorem

We can now state the following improved version of Stern's second theorem, Theorem 1.3. Recall the definition of  $\mu_c(\alpha, 2r)$  given in (4.24).

**Proposition 4.5** *For  $\alpha$  satisfying (4.1) and  $0 < \mu < \mu_c(\alpha, 2r)$ ,*

- (i) *the spherical harmonic  $H^\mu$ , of degree  $2r$ , introduced in (4.15), has no critical zero,*
- (ii) *the nodal set  $N(H^\mu)$  of  $H^\mu$  has exactly two connected components, i.e. consists of exactly two simple closed curves which do not intersect.*

*In particular, for  $0 < \mu < \mu_c(\alpha, 2r)$ , the spherical harmonic  $H^\mu$  has exactly three nodal domains.*

#### Proof of Proposition 4.5

Note that  $H^\mu$  is even, so that it is invariant under the antipodal map, and so is its nodal set  $N(H^\mu)$ . We have already seen, Properties 4.4, that a connected component of  $N(H^\mu)$  is contained in either  $\overline{\mathcal{M}'_0}$  or  $\overline{\mathcal{M}'_1}$ . Furthermore, there is one connected component, call it  $\gamma$ , which is contained in  $\overline{\mathcal{M}'_1}$ , and which is tangent to the great circle  $M'_0 \cup M'_1$  at the north pole  $p_+$ . Similarly, there is another connected component which is contained in  $\overline{\mathcal{M}'_0}$ , and which is tangent to  $M'_0 \cup M'_1$  at the south pole  $p_-$ . The second can be deduced from  $\gamma$  by applying the antipodal map.

It follows that it suffices to look at the part of the nodal set  $N(H^\mu)$  which is contained in  $\overline{\mathcal{M}'_1}$ . For this reason, we only have to consider the meridians  $C_j$  for  $2r+1 \leq j \leq 4r-1$ . The connected component  $\gamma$  is a simple closed curve. Start from the north pole, tangentially to  $M'_0$ , inside the domain  $\mathcal{Q}_{0,0,1}$ . The only possibility for  $\gamma$  is to exit  $\mathcal{Q}_{0,0,1}$  through the point  $q_{1,0,1}$ . Using the separation lemma, Lemma 4.2, and the analysis of local nodal patterns, we see that  $\gamma$  has to wind around  $M_0$ , inside the white domains, until it reaches the last point  $q_{2r-1,0,1}$ , at which it has to enter the white domain  $\mathcal{Q}_{2r-1,4r-1,1}$ , cross the meridian  $B_{4r-1}$ , exit through the point  $q_{2r-1,4r-1,1}$  and wind along  $M_{4r-1}$  until it reaches the domain  $\mathcal{Q}_{0,4r-2,1}$ , etc. The situation is similar to the one we encountered in the proof of Proposition 3.4 (i). Indeed, the important point in this proof was that the number  $\ell$  of latitude circle  $L_i$  was odd. In the present case we have  $\ell = 2r$ , but the number of latitude circles  $L'_i$  is  $2r-1$ , an odd integer. The proof of Section 3 applies mutatis mutandis, and the conclusion is that  $\gamma$  goes back to the north pole after going up and down  $r$  times, visiting all the points in  $\mathcal{N} \cap \overline{\mathcal{M}'_1}$ . Using Properties 4.4, it follows that  $N(H^\mu) \cap \overline{\mathcal{M}'_1}$  has exactly one connected component  $\gamma$ . Using the antipodal map, this means that  $N(H^\mu)$  has exactly two connected components.  $\square$

Figure 4.4 shows the nodal pattern of  $H^\mu$  in the exponential map, with one component tangent to the great circle  $M'_0 \cup M'_1$  at the north pole, the other at the south pole.




Figure 4.4: The nodal set  $N(H^\mu)$ .

## 5 Courant sharp property and open questions for minimal partitions for the sphere.

Leydold's thesis is devoted to this question (see [12] and more recently [6]). We reproduce below some synthesis essentially extracted from [9]. Given a spherical harmonic  $u$ , let  $\mu(u)$  denote the number of nodal domains of  $u$  (this notation should not induce confusion with the parameter  $\mu$ ).

- Courant's theorem for the sphere says that for any  $u_\ell \in \mathcal{H}_\ell$ ,

$$\mu(u_\ell) \leq \ell^2 + 1.$$

- Pleijel's Theorem extends to bounded domains in  $\mathbb{R}^n$ , and more generally to compact  $n$ -manifolds with boundary, with a constant  $\gamma(n) < 1$  replacing the constant  $\gamma(2) = 4/\mathbf{j}^2$  in the right-hand side of (5.1) (Peetre [15], Bérard-Meyer [1]). It is also interesting to note that this constant is independent of the geometry. In particular Pleijel's theorem is true in the case of the sphere. For any sequence of eigenfunctions  $u_\ell \in \mathcal{H}_\ell$

$$\lim_{\ell \rightarrow +\infty} \sup \frac{\mu(u_\ell)}{\ell(\ell-1)} \leq \frac{4}{j^2}. \quad (5.1)$$

- The conjecture of Leydold on the maximal cardinal of nodal sets is the following.

### Conjecture 5.1

$$\max_{u \in \mathcal{H}_\ell} \mu(u) = \begin{cases} \frac{1}{2}(\ell+1)^2 & \text{if } \ell \text{ is odd} \\ \frac{1}{2}\ell(\ell+2) & \text{if } \ell \text{ is even} \end{cases}$$

The theorem is proved in [12] for  $\ell \leq 6$ . Note that the example treated in the appendix for  $\ell = 3$  (middle subfigure in Fig. A.3) shows the optimality in this case. The conjecture corresponds to the idea that the maximum is obtained for product functions. The conjecture implies that the unique Courant sharp situation corresponds to the second eigenvalue. This last statement is true as a consequence of a theorem by Karpushkin [10].

### Theorem 5.2

$$\max_{u \in \mathcal{H}_\ell} \mu(u) = \begin{cases} \ell(\ell-2)+5 & \text{if } \ell \text{ is odd} \\ \ell(\ell-2)+4 & \text{if } \ell \text{ is even} \end{cases}$$

For Pleijel's theorem, Conjecture 5.1 gives the conjecture that

**Conjecture 5.3** *For any sequence of eigenfunctions  $u_\ell \in \mathcal{H}_\ell$ , we have*

$$\limsup_{\ell \rightarrow +\infty} \frac{\mu(u_\ell)}{\ell(\ell-1)} \leq \frac{1}{2}.$$

It is easy to see that this cannot be improved.

- Spectral minimal partitions are for example defined in [8]. Motivated by a conjecture in harmonic analysis popularized by Bishop [3] (who refers to [7]), the authors of [8] have proved in [9] that up to rotation the minimal 3-partition is the so-called  $Y$ -partition ( $\{0 < \phi < \frac{2\pi}{3}\}$ ,  $\{\frac{2\pi}{3} < \phi < \frac{4\pi}{3}\}$ , and  $\{\frac{4\pi}{3} < \phi < 2\pi\}$ ). There is a conjecture that the four faces of a spherical tetrahedron determine a minimal 4-partition on  $\mathbb{S}^2$ . What we get from the previous item and the general theory of [8] (nodal minimal partitions should correspond to a Courant sharp situation) is that minimal partitions cannot be nodal.
- With a different point of view, let us mention the contributions of [14] on random spherical harmonics.

## A Some simulations with Maple

In this appendix, we provide some pictures issued from numerical computations with Maple. The nodal sets are viewed in the exponential map at the north pole. The outer circle, at distance  $\pi$ , is the cut-locus of  $p_+$  and corresponds to the south pole.

Figure A.1 illustrates Proposition 3.4 in the cases  $\ell = 3$  (left) and  $\ell = 4$  (right).

The figures in the top line show the checkerboard associated with  $Z_\ell$  and  $W_\ell$ . The figures in the bottom line show the nodal set of  $H^{\mu,\ell}$  for  $\mu$  small enough.

Figure A.2 illustrates Proposition 3.4 in the cases  $\ell = 5$  (left) and  $\ell = 6$  (right).




Figure A.1: Example 1, with  $\ell = 3$  and  $\ell = 4$ .




Figure A.2: Example 1, with  $\ell = 5$  and  $\ell = 6$ .




Figure A.3: Appearance and disappearance of critical zeroes.

Figure A.3 illustrates the occurrence of critical zeros in Stern's Example 1, with  $\ell = 3$ . The corresponding Legendre polynomial is  $P_3(t) = \frac{1}{2}t(5t^2 - 3)$ . The polynomial  $P_2(t) = \frac{1}{2}(3t^2 - 1)$  has two roots  $\pm \frac{1}{\sqrt{3}}$ . According to Section 3.2.1, there are twelve possible critical zeros, given in spherical coordinates by the points  $\left(\arccos(\pm \frac{1}{\sqrt{3}}), j\frac{\pi}{6}\right)$  with  $j \in \{1, 3, 5, 7, 9, 11\}$ , and exactly two critical values of the parameter,  $\mu = \pm \sqrt{2}$ . For  $\mu > 0$ , there is exactly one critical value  $\mu = \sqrt{2}$ , which is associated with six critical zeros. Figure A.3 shows the nodal set  $N(H^{\mu,3})$  for  $\mu < \sqrt{2}$  (left), for  $\mu = \sqrt{2}$  (center) and for  $\mu > \sqrt{2}$  (right).




Figure A.4: Stern's Example 2 in the cases  $\ell = 4$  and  $\ell = 6$ .

Figure A.4 illustrates Proposition 4.4. The figures in the top line show the nodal sets of  $W$ ,  $V_\alpha$ , and  $H^\mu$  for  $\mu$  small. The great circle  $M'_0 \cup M'_1$  divides the sphere into two closed hemispheres. Each one contains a simple closed nodal curve tangent to the great circle at one of the poles. As usual, the south pole is represented by the outer circle (dotted line), the cut-locus of 0 in the tangent space at  $p_+$ . The figures in the bottom line show the nodal sets of  $H^\mu$  alone.

## References

- [1] P. Bérard and D. Meyer. Inégalités isopérimétriques et applications. Annales scientifiques de l’Ecole Normale Supérieure, Sér. 4, 15 (3), p. 513-541 (1982).
- [2] P. Bérard and B. Helffer. Dirichlet eigenfunctions of the square membrane: Courant’s property, and A. Stern’s and Å. Pleijel’s analyses. Preprint 2014, arXiv:14026054.
- [3] C.J. Bishop. Some questions concerning harmonic measure. Dahlberg, B. (ed.) et al., Partial Differential equations with minimal smoothness and applications. IMA Vol. Math. Appl. 42 (1992), p. 89-97.
- [4] R. Courant. Ein allgemeiner Satz zur Theorie der Eigenfunktionen selbstadjungierter Differentialausdrücke. Nachr. Ges. Göttingen (1923), p. 81-84.
- [5] R. Courant and D. Hilbert. Methods of Mathematical Physics. Volume 1. John Wiley & Sons, 1989.
- [6] A. Eremenko, D. Jakobson, and N. Nadirashvili. On nodal sets and nodal domains on  $\mathbb{S}^2$ . Annales Institut Fourier 57 (7) (2007), p. 2345-2360.
- [7] S. Friedland and W.K. Hayman. Eigenvalue Inequalities for the Dirichlet problem on spheres and the growth of subharmonic functions. Comment. Math. Helvetici 51 (1976), p. 133-161.
- [8] B. Helffer, T. Hoffmann-Ostenhof and S. Terracini. Nodal domains and spectral minimal partitions. Annales Institut Henri Poincaré (Section Analyse non linéaire) (2009).
- [9] B. Helffer, T. Hoffmann-Ostenhof and S. Terracini. On spectral minimal partitions: the case of the sphere. Around the research of Vladimir Maz’ya. III, p. 153-178, Int. Math. Ser. (N.Y.), 13 Springer 2010.
- [10] V.N. Karpushkin. Topology of the zeros of eigenfunctions. Funktional Anal. i Prilozhen 23:3 (1989), p. 59-60.
- [11] H. Lewy. On the minimum number of domains in which the nodal lines of spherical harmonics divide the sphere. Comm. Partial Differential Equations 2 (1977), no. 12, p. 1233-1244.
- [12] J. Leydold. On the number of nodal domains of spherical harmonics. Topology 35, p. 301-321, 1996.
- [13] W. Magnus, F. Oberhettinger and R.P. Soni. Formulas and Theorems for the Special Functions of Mathematical Physics. Third Edition. Berlin: Springer-Verlag, 1966.
- [14] F. Nazarov and M. Sodin. On the number of nodal domains of random spherical harmonics. Amer. J. Math. 131 (2009), p. 1337-1357.
- [15] J. Peetre. A generalization of Courant nodal theorem. Math. Scandinavica 5, p. 15-20 (1957).

- [16] Å. Pleijel. Remarks on Courant's nodal theorem. *Comm. Pure. Appl. Math.* **9**, p. 543–550 (1956) .
- [17] I. Polterovich. Pleijel's nodal domain theorem. *Proc. Amer. Math. Soc.* 137 (2009), p. 1021-1024.
- [18] A. Stern. Bemerkungen über asymptotisches Verhalten von Eigenwerten und Eigenfunctionen. Inaugural-Dissertation zur Erlangung der Doktorwürde der Hohen Mathematisch-Naturwissenschaftlichen Fakultät der Georg August-Universität zu Göttingen. Druck der Dieterichschen Univertisäts-Buchdruckerei (W. Fr. Kaestner). Göttingen, 1925
- [19] A. Stern. Bemerkungen über asymptotisches Verhalten von Eigenwerten und Eigenfunctionen. Diss. Göttingen 1925. Extracts and annotations by P. Bérard and B. Helffer. [Available here](#).
- [20] G. Szegö. Orthogonal Polynomials. Fourth edition. AMS colloquium publications, Vol. XXIII, Amer. Math. Soc. Providence, R.I. (1975).