



HAL
open science

Flows and functional inequalities for fractional operators

Jean Dolbeault, An Zhang

► **To cite this version:**

Jean Dolbeault, An Zhang. Flows and functional inequalities for fractional operators. *Applicable Analysis*, 2017, 96 (9), pp.1547-1560. hal-01404580

HAL Id: hal-01404580

<https://hal.science/hal-01404580>

Submitted on 28 Nov 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Flows and functional inequalities for fractional operators

Jean Dolbeault^{*,a} and An Zhang^{*,b}

**CEREMADE (CNRS UMR n° 7534), PSL research university, Université Paris-Dauphine, Place de Lattre de Tassigny, 75775 Paris 16, France*

November 28, 2016

This paper collects results concerning global rates and large time asymptotics of a fractional fast diffusion on the Euclidean space, which is deeply related with a family of fractional Gagliardo-Nirenberg-Sobolev inequalities. Generically, self-similar solutions are not optimal for the Gagliardo-Nirenberg-Sobolev inequalities, in strong contrast with usual standard fast diffusion equations based on non-fractional operators. Various aspects of the stability of the self-similar solutions and of the entropy methods like *carré du champ* and *Rényi entropy powers* methods are investigated and raise a number of open problems.

Keywords: fractional Gagliardo-Nirenberg-Sobolev inequality; fractional Sobolev inequality; fractional fast diffusion equation; self-similar solutions; asymptotic behavior; intermediate asymptotics; rate of convergence; entropy methods; *carré du champ*; Rényi entropy powers; entropy – entropy production inequality; self-similar variables; linearization

AMS Subject Classifications: Primary: 26A33, 35R11; Secondary: 35A23, 35B33, 35B40, 35K65.

1. Introduction

Recently many papers have been devoted to the extension of results involving second order partial differential operators to fractional operators. More specifically, fractional diffusion equations have been studied from the point of view of existence, comparison and regularity of the solutions, not only in the linear case but also in the framework of nonlinear models of porous medium or fast diffusion type. Concerning modeling issues, fractional diffusions are usually motivated by microscopic jump processes. One can refer to [1, 2] for an extended review of models arising from various areas of physics. We will not go in this direction. Relying on known theoretical results like the ones of [3–7], our approach aims at the description of fundamental qualitative properties of such equations in relation with closely associated, basic functional inequalities.

Standard fast diffusion or porous medium equations have simple features which arise from the homogeneity of the nonlinear term or from the fact that the diffusion operator is of second-order. These features explain the special role of self-similar solutions, known as Barenblatt-Pattle, in the large time regimes: see [8] for an historical presentation. Also remarkable is the fact discovered in [9] that the Barenblatt-Pattle solutions are optimal for some Gagliardo-Nirenberg-Sobolev inequalities, which are essential to measure the asymptotic stability of the self-

^aJean Dolbeault, Email: dolbeaul@ceremade.dauphine.fr

^bCorresponding author: An Zhang, Email: zhang@ceremade.dauphine.fr

similar solutions using relative entropy methods. Entropy functionals are indeed deeply connected with fast diffusion equations, as these equations can be seen as gradient flows of the entropies with respect to the Wasserstein's distance (see [10]).

In the context of fractional fast diffusion or porous medium equations, it is therefore very natural to question the role of self-similar solutions in terms of large time asymptotics and related functional inequalities. However, fractional order derivatives and nonlinearities do not combine well and this raises a number of difficulties. Self-similar solutions which generalize the Barenblatt-Pattle solutions to fractional diffusion have been studied in connection with large time asymptotics in [4, 5] and [11–13]. There is a certain flexibility in the generalization of the standard nonlinear diffusion equations: this has been investigated in [14] and [15].

In the present paper, we shall specifically rely on the fractional fast diffusion equation

$$\frac{\partial u}{\partial t} = \nabla \cdot \left(\sqrt{u} \nabla (-\Delta)^{-s} u^{m-\frac{1}{2}} \right)$$

with $m < 1$, which seems particularly well adapted to entropy – entropy production inequalities as we shall see below. Such an equation has already been considered in [15, Equation (MG)] from the point of view of self-similarity. Our purpose is to clarify the interplay of this equation with fractional Gagliardo-Nirenberg-Sobolev inequalities (see Section 2.3), and observe that, generically, optimal functions for these inequalities differ from self-similar solutions that are supposed to govern the large time behavior of the solutions of the evolution equation (Proposition 2.3). Beyond some results about, *e.g.*, the existence of optimal solutions for the interpolation inequalities (Proposition 2.1), we raise a number of open questions concerning the large time asymptotics in Section 3 and the applicability of the Bakry-Emery, or *carré du champ*, method in Section 4.

2. Preliminaries: a fractional interpolation inequality and a fractional fast diffusion flow

2.1. The fractional Sobolev inequality

According to [16], for any $\alpha \in (0, d)$, with $q = \frac{2d}{d-\alpha}$, the fractional Sobolev inequality in \mathbb{R}^d can be written as

$$\|w\|_{\dot{H}^{\frac{\alpha}{2}}(\mathbb{R}^d)}^2 \geq S_{d,\alpha} \left(\int_{\mathbb{R}^d} |w|^q dx \right)^{\frac{2}{q}} \quad \forall w \in \dot{H}^{\frac{\alpha}{2}}(\mathbb{R}^d) \quad (1)$$

where $\dot{H}^{\frac{\alpha}{2}}(\mathbb{R}^d)$ is the space of all tempered distributions w such that

$$\hat{w} \in L^1_{\text{loc}}(\mathbb{R}^d) \quad \text{and} \quad \|w\|_{\dot{H}^{\frac{\alpha}{2}}(\mathbb{R}^d)}^2 := \int_{\mathbb{R}^d} |\xi|^\alpha |\hat{w}|^2 d\xi < \infty.$$

Here \hat{w} denotes the Fourier transform of w and the optimal constant is given by

$$S_{d,\alpha} = 2^\alpha \pi^{\frac{\alpha}{2}} \frac{\Gamma(\frac{d+\alpha}{2})}{\Gamma(\frac{d-\alpha}{2})} \left(\frac{\Gamma(\frac{d}{2})}{\Gamma(d)} \right)^{\alpha/d}.$$

Up to translations, dilations and multiplications by a nonzero constant, the optimal function is

$$w_\star(x) = (1 + |x|^2)^{-\frac{d-\alpha}{2}} \quad \forall x \in \mathbb{R}^d.$$

It is easy to check that w_\star solves

$$(-\Delta)^{\frac{\alpha}{2}} w_\star = C_{d,\alpha} w_\star^{\frac{d+\alpha}{d-\alpha}} \quad \text{where} \quad C_{d,\alpha} = 2^\alpha \frac{\Gamma(\frac{d+\alpha}{2})}{\Gamma(\frac{d-\alpha}{2})}. \quad (2)$$

With the notation $\mathcal{D}_\alpha := \nabla(-\Delta)^{\frac{\alpha-1}{2}}$, we get that $(-\Delta)^{\frac{\alpha}{2}} = \mathcal{D}_{\alpha/2}^* \mathcal{D}_{\alpha/2}$ and (1) can then be written as

$$\|w\|_{\dot{H}^{\frac{\alpha}{2}}(\mathbb{R}^d)}^2 = \|\mathcal{D}_{\alpha/2} w\|_{L^2(\mathbb{R}^d)}^2 \geq S_{d,\alpha} \|w\|_{L^q(\mathbb{R}^d)}^2 \quad \forall w \in \dot{H}^{\frac{\alpha}{2}}(\mathbb{R}^d).$$

The case $\alpha = 2$ was established in [17, 18], but one may refer to [19] in case $d = 3$ and even to [20] for some early considerations on optimal functions. In the fractional case $\alpha \neq 2$, the dual form of the optimal inequality, *i.e.*, the Hardy-Littlewood-Sobolev inequality, was established in [21], while considerations on the case $\alpha = 1$ in connection with trace issues can be found in [22]. Later work include [23] and [24] among many other related papers. The issue of the symmetry of the optimal functions, up to translations, was considered in [21] and established in [25] using the moving planes method.

2.2. A fractional Gagliardo-Nirenberg-Sobolev inequality

With $q = \frac{2d}{d-\alpha}$, a simple Hölder interpolation shows that

$$\|w\|_{L^{2p}(\mathbb{R}^d)} \leq \|w\|_{L^{p+1}(\mathbb{R}^d)}^{1-\vartheta} \|w\|_{L^q(\mathbb{R}^d)}^\vartheta \quad \text{with} \quad \vartheta = \frac{d}{p} \frac{p-1}{d+\alpha-p(d-\alpha)}.$$

Together with Sobolev's inequality, this provides an interpolation inequality of Gagliardo-Nirenberg-Sobolev type. More precisely, we have the following statement.

PROPOSITION 2.1 *Assume that $\alpha \in (0, d)$ and $p \in (1, q]$ with $q = \frac{2d}{d-\alpha}$. With the above notations, the following Gagliardo-Nirenberg-Sobolev inequality*

$$\|w\|_{L^{2p}(\mathbb{R}^d)} \leq C_{\text{GNS}} \|w\|_{\dot{H}^{\frac{\alpha}{2}}(\mathbb{R}^d)}^\vartheta \|w\|_{L^{p+1}(\mathbb{R}^d)}^{1-\vartheta} \quad \forall w \in \mathcal{D}(\mathbb{R}^d) \quad (3)$$

holds with an optimal constant $C_{\text{GNS}} \leq S_{d,\alpha}^{\vartheta/2}$. Moreover, there exists a positive optimal function in $\dot{H}^{\frac{\alpha}{2}}(\mathbb{R}^d) \cap L^{p+1}(\mathbb{R}^d)$ which solves the Euler-Lagrange equation

$$(-\Delta)^{\frac{\alpha}{2}} w + w^p - w^{2p-1} = 0, \quad x \in \mathbb{R}^d. \quad (4)$$

In the above statement $\mathcal{D}(\mathbb{R}^d)$ denotes the space of smooth functions with compact support, but it is simple to extend it by density to the space $\{w \in \dot{H}^{\frac{\alpha}{2}}(\mathbb{R}^d) : \int_{\mathbb{R}^d} w^{p+1} dx < \infty\}$. See [26, Theorem 2.1] for details. When $p = \frac{q}{2} = \frac{d}{d-\alpha}$, then $\vartheta = 1$ and $C_{\text{GNS}} = S_{d,\alpha}$. The symmetry of the optimal functions is not considered here and the interested reader is invited to refer for instance to [27] for a related problem.

Sketch of the proof. This result is out of the scope of the present paper and the proof is relatively standard, so let us give only a simple sketch. A reader interested in further details is invited to refer to [28] and [29] for further details in similar cases. The above result can actually be seen as a special case of those dealt with in [26]. The convergence of sequences in the critical case has been analyzed in details in [30]. We can therefore assume that $p \in (1, q)$.

An optimal function for (3) can be obtained by considering

$$\mathcal{I}_M := \inf \left\{ \|w\|_{\dot{H}^{\frac{\alpha}{2}}(\mathbb{R}^d)}^2 + \|w\|_{L^{p+1}(\mathbb{R}^d)}^{p+1} : \int_{\mathbb{R}^d} w^{2p} dx = M \right\}.$$

An optimisation on $\lambda > 0$ of $w_\lambda = \lambda^{\frac{d}{2p}} w(\lambda \cdot)$ shows that the minimization of \mathcal{I}_M is equivalent to the characterization of the optimality case in (3), and also that

$$\mathcal{I}_M = M^\gamma \mathcal{I}_1 \quad \text{with} \quad \gamma = \frac{d + \alpha - p(d - \alpha)}{d - p(d - 2\alpha)} < 1.$$

With these preliminary observations, a concentration-compactness analysis can be done as follows. Let us consider a sequence $(w_n)_{n \in \mathbb{N}}$ of nonnegative function such that $\int_{\mathbb{R}^d} w_n^{2p} dx = M$ for any $n \in \mathbb{N}$ and

$$\lim_{n \rightarrow \infty} \|w_n\|_{\dot{H}^{\frac{\alpha}{2}}(\mathbb{R}^d)}^2 + \|w_n\|_{L^{p+1}(\mathbb{R}^d)}^{p+1} = \mathcal{I}_M.$$

1) The following result has been stated in [31, Lemma I.1, p. 231].

LEMMA 2.2 *Let $R > 0$ and $p \in [1, 2^*/2)$. If $(w_n)_n$ is bounded in $H^1(\mathbb{R}^d)$ and if*

$$\limsup_{n \rightarrow \infty} \int_{B_R(x)} |w_n|^{2p} dy = 0$$

for any $x \in \mathbb{R}^d$, then $\lim_{n \rightarrow \infty} \|w_n\|_{L^r(\mathbb{R}^d)} = 0$ for any $r \in [2, 2^)$.*

Here $2^* = \infty$ if $d = 1$ or 2 , and $2^* = 2d/(d-2)$ if $d \geq 3$. By an analogue of this result in $\dot{H}^{\frac{\alpha}{2}}(\mathbb{R}^d)$, with $2^*/2$ replaced by q , *vanishing* cannot occur, which means that, after eventually replacing w_n by $w_n(\cdot + y_n)$ for some unbounded sequence $(y_n)_{n \in \mathbb{N}}$ of points in \mathbb{R}^d , we have

$$\lim_{R \rightarrow \infty} \lim_{n \rightarrow \infty} \int_{B_R(0)} |w_n|^{2p} dy > 0.$$

2) Since $\gamma < 1$, it is straightforward to check that

$$\mathcal{I}_{M_1+M_2} \leq \mathcal{I}_{M_1} + \mathcal{I}_{M_2},$$

which prevents *dichotomy*. As a consequence, for any $\varepsilon > 0$, there exists some $R > 0$, large enough, such that

$$\lim_{n \rightarrow \infty} \int_{B_R(0)} |w_n|^{2p} dy \geq M - \varepsilon.$$

In other words, this means that the sequence $(w_n^{2p})_{n \in \mathbb{N}}$ is tightly relatively compact in the sense of measures.

3) *Concentration* is forbidden by Sobolev's inequality (1). Altogether, this proves that $(w_n)_{n \in \mathbb{N}}$ is weakly relatively compact in $L^{2p}(\mathbb{R}^d)$ (see for instance [32, Lemma 2.2]) and the conclusion holds by lower semi-continuity.

Let us take the logarithm of both sides in (3). An optimal function is a minimizer of the functional

$$w \mapsto \vartheta \log \|w\|_{\dot{H}^{\frac{\alpha}{2}}(\mathbb{R}^d)} + (1 - \vartheta) \log \|w\|_{L^{p+1}(\mathbb{R}^d)} - \log \|w\|_{L^{2p}(\mathbb{R}^d)} .$$

A variation shows that

$$\frac{\vartheta (-\Delta)^{\frac{\alpha}{2}} w}{\|w\|_{\dot{H}^{\frac{\alpha}{2}}(\mathbb{R}^d)}^{\frac{\alpha}{2}}} + \frac{(1 - \vartheta) w^p}{\|w\|_{L^{p+1}(\mathbb{R}^d)}^p} - \frac{w^{2p-1}}{\|w\|_{L^{2p}(\mathbb{R}^d)}^{2p-1}} = 0 .$$

Using the homogeneity in (3) and a multiplication by a constant, the coefficients can be adjusted so that w solves (4). ■

2.3. A fractional fast diffusion equation

Let us consider the case

$$\alpha = 2(1 - s) .$$

Any nonnegative solution u of

$$\frac{\partial u}{\partial t} = \nabla \cdot (\sqrt{u} \nabla (-\Delta)^{-s} w) \quad \text{with} \quad w = u^{m-\frac{1}{2}} \quad (5)$$

which is smooth enough and has good decay properties as $|x| \rightarrow \infty$ is such that

$$E' = \frac{2m(1-m)}{2m-1} I , \quad (6)$$

where the *entropy* E and the *Fisher information* I are defined respectively by

$$E[u] := \int_{\mathbb{R}^d} u^m dx \quad \text{and} \quad I[u] := \int_{\mathbb{R}^d} \left| \nabla^{(1-s)} u^{m-\frac{1}{2}} \right|^2 dx ,$$

and $\nabla^{(1-s)}$ is defined by

$$\nabla^{(1-s)} w := \nabla (-\Delta)^{-s/2} w .$$

Above we consider $E[u(t, \cdot)]$ as a function of t if u solves (5) and E' denotes the t -derivative of $E[u(t, \cdot)]$.

Because the right hand side in (5) is in divergence form, we know that

$$M := \int_{\mathbb{R}^d} u dx$$

is independent of $t \geq 0$ for any solution u which is smooth enough, and with enough

decay as $|x| \rightarrow +\infty$. Let us assume that

$$m \geq m_1 := \frac{1}{2} + \frac{1}{q} = 1 - \frac{\alpha}{2d}.$$

If $m = m_1$ and $v_\star := w_\star^q$, then $v_\star(x) = (1 + |x|^2)^{-d}$ satisfies

$$v_\star^{m_1 - \frac{1}{2}} = w_\star \quad \text{and} \quad (-\Delta)^{1-s} \left(v_\star^{m_1 - \frac{1}{2}} \right) = C_{d,\alpha} v_\star^{\frac{d+\alpha}{2d}} = C_{d,\alpha} (1 + |x|^2)^{s-1} \sqrt{v_\star}$$

according to (2).

Assume now that $m \in [m_1, 1)$. If we introduce a *generalized Rényi entropy power functional* F as in [33, 34], defined by

$$F := E^\sigma$$

with

$$\sigma := \frac{2}{\vartheta} \frac{1 - \vartheta}{p + 1} + 1 = \frac{m - m_c}{1 - m}, \quad p = \frac{1}{2m - 1} \quad \text{and} \quad m_c := 1 - \frac{\alpha}{d},$$

then by applying Proposition 2.1 to $w = u^{m - \frac{1}{2}}$ we obtain that

$$F' \geq \kappa$$

using (6), where the constant κ depends only on d, s (or α), m and M , and can be computed as

$$\kappa := \frac{2m(1-m)}{2m-1} \sigma \left(\frac{M}{C_{\text{GNS}}^{2p}} \right)^{\frac{1}{p\vartheta}} = \frac{2m}{2m-1} (m - m_c) \left(\frac{M}{C_{\text{GNS}}^{2p}} \right)^{\frac{d+\alpha-p(d-\alpha)}{d(p-1)}}.$$

2.4. Self-similar solutions

Let us introduce the time-dependent change of variables

$$u(t, x) = \frac{1}{R^d} v \left(\log R, \frac{x}{R} \right) \tag{7}$$

where $R = R(t)$ is given by the ordinary differential equation

$$R^{\mu-1} \frac{dR}{dt} = 1, \quad R(0) = 1, \tag{8}$$

and the parameter μ is defined by

$$\mu := d(m - m_c) \quad \text{with} \quad m_c = 1 - \frac{\alpha}{d}.$$

If m is in the range $[m_1, 1)$, it is straightforward to check that μ is positive and the scale R has the explicit expression

$$R(t) = (1 + \mu t)^{1/\mu} \quad \forall t \geq 0.$$

If u solves (5), it is then easy to check that v is a solution of

$$\frac{\partial v}{\partial t} = \nabla \cdot \left[\sqrt{v} \left(\nabla(-\Delta)^{-s} v^{m-\frac{1}{2}} + x \sqrt{v} \right) \right]. \quad (9)$$

This change of variable is classical in case of non-fractional operators but has also been considered in the case of the fractional Laplacian, for instance in [11, 12]. A detailed analysis of self-similar profiles is available in [15]. In the fast diffusion case, for an equation related with (5), we refer to [35], where the role of Barenblatt type solutions for regularization effects has been clarified.

PROPOSITION 2.3 *Let $p = \frac{1}{2m-1}$. Under the assumptions of Proposition 2.1, if w is an optimal function for (3), then $v = w^{2p}$ is not a bounded, positive, stationary solution of (9) such that $\mathbb{I}[v]$ is finite, unless $s = 0$, $\alpha = 2$ or $m = \frac{d+1}{d+2(1-s)}$.*

In other words, we claim that (4) does not provide stationary solutions of (9) as it is the case in the non-fractional case, except in the special case corresponding to $m = \frac{d+1}{d+2(1-s)}$, which was observed in [14, 15].

Proof. We argue by contradiction and consider a solution of

$$\nabla \cdot \left[\sqrt{v} \left(\nabla(-\Delta)^{-s} v^{m-\frac{1}{2}} + x \sqrt{v} \right) \right] = 0.$$

We deduce that $w = v^{m-\frac{1}{2}}$ solves

$$0 = \nabla(-\Delta)^{-s} v^{m-\frac{1}{2}} + x \sqrt{v} = \nabla(-\Delta)^{-s} w + x w^p$$

where $p = 1/(2m-1)$ and, as a consequence,

$$-(-\Delta)^{1-s} w + \nabla \cdot (x w^p) = 0.$$

With $\alpha = 2(1-s)$, if w simultaneously solves (4) up to a multiplication by a constant, a scaling and a translation, this means that $w^p = v$ is such that

$$\nabla \cdot (x w^p) = a w^{2p-1} - b w^p$$

for some positive constants a and b that can be adjusted. With

$$a = \frac{p}{p-1} \quad \text{and} \quad b = \frac{p}{p-1} - d,$$

we find that

$$w(x) = (1 + |x|^2)^{\frac{1}{1-p}} \quad \forall x \in \mathbb{R}^d.$$

The result follows from [14, Theorem 3.1]. ■

The case $m = m_1$ is almost explicit. In order to illustrate Proposition 2.3, let us give some details. With the notations of Section 2.1, if we choose

$$v_\star^{m_1-\frac{1}{2}} = w_\star \quad \iff \quad v_\star = w_\star^{\frac{2d}{d-\alpha}} \quad \iff \quad v_\star(x) = (1 + |x|^2)^{-d} \quad \forall x \in \mathbb{R}^d,$$

we obtain that

$$\begin{aligned}\nabla(-\Delta)^{-1}(-\Delta)^{1-s}v_\star^{m_1-\frac{1}{2}}+x\sqrt{v_\star}&=\nabla(-\Delta)^{-1}C_{d,\alpha}w_\star^{\frac{d+\alpha}{d-\alpha}}+x\sqrt{v_\star} \\ &=\nabla(-\Delta)^{-1}\left(C_{d,\alpha}v_\star^{\frac{d+\alpha}{2d}}-C_{d,2}v_\star^{\frac{d+2}{2d}}\right)\neq 0\end{aligned}$$

unless $\alpha = 2$.

For later purpose, let us define the self-similar profile \mathfrak{B} as the unique radial solution of

$$\nabla(-\Delta)^{-s}\mathfrak{B}^{m-\frac{1}{2}}+x\sqrt{\mathfrak{B}}=0$$

such that $\int_{\mathbb{R}^d}\mathfrak{B}dx=M$. We recall that these solutions are the so-called *Barenblatt profiles* when $s=0$ and refer to [36] for more details. It is straightforward to check that

$$u_\star(t,x)=\frac{1}{R^d}\mathfrak{B}\left(\log R,\frac{x}{R}\right)$$

is a self-similar solution of (5) if $R=R(t)$ is given by (8).

3. Global rates of convergence and formal large time asymptotics

The *generalized Rényi entropy power functional* defined as in Section 2.3 by

$$F[u]=\left(\int_{\mathbb{R}^d}u^m dx\right)^\sigma$$

is such that

$$F[u(t,\cdot)]\geq F[u_0]+\kappa t \quad \forall t\geq 0$$

if u solves (5) with initial datum u_0 on the one hand, and a direct computation shows that the self-similar solution u_\star satisfies the relation

$$F[u_\star(t,\cdot)]=F[\mathfrak{B}](1+\mu t) \quad \forall t\geq 0.$$

COROLLARY 3.1 *With the above notations and under the assumptions of Proposition 2.3, we have the estimate*

$$\kappa<\kappa_\star:=\mu F[\mathfrak{B}].$$

Proof. The inequality $\kappa\leq\kappa_\star$ is a straightforward consequence of the above computations, with $u_0=\mathfrak{B}$. The strict inequality follows from Proposition 2.3. \blacksquare

Now let us investigate the large time asymptotics at formal level. Inspired by [8, 35], we expect that for any $m\in[m_1,1)$,

$$u(t,x)\sim u_\star(t,x) \quad \text{as } t\rightarrow+\infty$$

as in [37, Theorem 1.4], and thus

$$F[u(t, \cdot)] = \kappa_\star t (1 + o(1)) = F[\mathfrak{B}] (1 + \mu t) (1 + o(1)).$$

Using the self-similar change variables (7)-(8), the equivalence $u \sim u_\star$ amounts to

$$v(t, \cdot) \rightarrow \mathfrak{B} \quad \text{as } t \rightarrow +\infty.$$

So far this question is open. Next let us investigate at formal level the question of the attraction of the solutions by the self-similar solutions.

If u solves (5) and u_\star is the self-similar solution with same mass, we define the relative entropy by

$$\mathcal{E}[u] = \frac{1}{m-1} \int_{\mathbb{R}^d} (u^m - u_\star^m - m u_\star^{m-1} (u - u_\star)) dx.$$

A straightforward computation shows that

$$\begin{aligned} \frac{d}{dt} \mathcal{E}[u(t, \cdot)] &= \frac{m}{m-1} \int_{\mathbb{R}^d} (u^{m-1} - u_\star^{m-1}) \frac{\partial u}{\partial t} dx - m \int_{\mathbb{R}^d} u_\star^{m-2} (u - u_\star) \frac{\partial u_\star}{\partial t} dx \\ &= -\frac{m}{m-1} \int_{\mathbb{R}^d} \nabla (u^{m-1} - u_\star^{m-1}) \cdot \left(\sqrt{u} \nabla (-\Delta)^{-s} u^{m-\frac{1}{2}} \right) dx \\ &\quad + m \int_{\mathbb{R}^d} \nabla (u_\star^{m-2} (u - u_\star)) \cdot \left(\sqrt{u_\star} \nabla (-\Delta)^{-s} u_\star^{m-\frac{1}{2}} \right) dx. \end{aligned}$$

At formal level, we investigate the limit as $\varepsilon \rightarrow 0$ of

$$u_\varepsilon = u_\star \left(1 + \varepsilon u_\star^{\frac{1}{2}-m} f \right).$$

First, let us notice that

$$\begin{aligned} \sqrt{u_\varepsilon} &= \sqrt{u_\star} + \frac{1}{2} \varepsilon u_\star^{1-m} f + o(\varepsilon), \quad u_\varepsilon^{m-\frac{1}{2}} = u_\star^{m-\frac{1}{2}} + (m - \frac{1}{2}) \varepsilon f + o(\varepsilon), \\ u_\varepsilon^{m-1} &= u_\star^{m-1} + (m-1) \varepsilon \frac{f}{\sqrt{u_\star}} + \frac{1}{2} (m-1)(m-2) \varepsilon \frac{f^2}{u_\star^m} + o(\varepsilon^2). \end{aligned}$$

As a consequence, we find that

$$\mathcal{E}[u] = \frac{m}{2} \varepsilon^2 \int_{\mathbb{R}^d} u_\star^{1-m} f^2 dx + o(\varepsilon^2)$$

and

$$\frac{d}{dt} \mathcal{E}[u(t, \cdot)] = -\frac{m}{2} \varepsilon^2 \mathcal{Q}_{u_\star}[f] + o(\varepsilon^2)$$

where the quadratic form \mathcal{Q}_{u_\star} is defined by

$$\begin{aligned} \mathcal{Q}_{u_\star}[f] := & \int_{\mathbb{R}^d} \nabla\left(\frac{f}{\sqrt{u_\star}}\right) \cdot \left((2m-1) \sqrt{u_\star} \nabla(-\Delta)^{-s} f + u_\star^{1-m} f \nabla(-\Delta)^{-s} u_\star^{m-\frac{1}{2}} \right) dx \\ & + (m-2) \int_{\mathbb{R}^d} \nabla\left(\frac{f^2}{u_\star^m}\right) \cdot \left(\sqrt{u_\star} \nabla(-\Delta)^{-s} u_\star^{m-\frac{1}{2}} \right) dx. \end{aligned}$$

If $s = 0$, we observe that

$$\frac{1}{2m-1} \mathcal{Q}_{u_\star}[f] = \int_{\mathbb{R}^d} u_\star \left| \nabla\left(\frac{f}{\sqrt{u_\star}}\right) \right|^2 dx - \frac{1-m}{2} \int_{\mathbb{R}^d} \left((2-m) \frac{|\nabla u_\star|^2}{u_\star^2} - \frac{\Delta u_\star}{u_\star} \right) f^2 dx.$$

To characterize the asymptotic stability of u_\star , the point is to notice that $\mathcal{Q}_{u_\star}[f]$ controls $\int_{\mathbb{R}^d} u_\star^{1-m} f^2 dx$.

PROPOSITION 3.2 *With the previous notations, if $s = 0$ and $m_1 \leq m < 1$, there exists a positive constant C which depends only on m and d such that*

$$\mathcal{Q}_{u_\star}[f] \geq \frac{2}{1-m} \frac{C}{R^\mu} \int_{\mathbb{R}^d} u_\star^{1-m} f^2 dx$$

for any smooth function f such that $\int_{\mathbb{R}^d} u_\star^{\frac{3}{2}-m} f dx = 0$.

We recall that $R = R(t)$ is given by (8), and u_\star depends on t , so that the above inequality holds for any $t \geq 0$. When $s = 0$, $\mu = dm - (d-2)$ and $R^\mu(t) = 1 + \mu t$ for any $t \geq 0$. By density, it is easy to extend the above inequality to any function $f \in L^\infty(dt, L^2(\mathbb{R}^d, u_\star^{1-m} dx))$ such that $\mathcal{Q}_{u_\star}[f]$ is finite for any $t \geq 0$.

Proof. The result follows from the change of variables (7) and [38, Corollary 1]: the above inequality can be rewritten in the form of a spectral gap inequality with a constant which is independent of t . Also see [36] for earlier spectral gap computations. \blacksquare

In self-similar variables, Proposition 3.2 shows that \mathfrak{B} is linearly stable, and this is enough to prove that u_\star is asymptotically linearly stable when $s = 0$. Actually one also knows that \mathfrak{B} is globally stable when $s = 0$, with explicit global rates, which is a far deeper issue. Proposition 3.2 immediately raises two open questions:

- (i) Is u_\star asymptotically linearly stable for any $s \in (0, n)$?
- (ii) Can we prove that u_\star is, under the appropriate mass normalization, a global attractor and deduce that the asymptotic rate of convergence of any solution towards u_\star is determined by the spectral gap Λ associated with the inequality

$$\mathcal{Q}_{\mathfrak{B}}[g] \geq \Lambda \int_{\mathbb{R}^d} \mathfrak{B}^{1-m} g^2 dx$$

for any admissible function g such that $\int_{\mathbb{R}^d} \mathfrak{B}^{\frac{3}{2}-m} g dx = 0$?

4. On the Bakry-Emery method applied to the fractional fast diffusion equation and generalized Rényi entropy powers

In this section we do a computation based on the generalized Rényi entropy powers in case $s = 0$ and emphasize the points which differ when $s > 0$. So far such a computation has been done using the so-called *pressure* variable u^{m-1} as, for instance, in [33, 34]. Here we base our computations on

$$f := u^{m-\frac{1}{2}}.$$

With this simple change of variables, we obtain that

$$\frac{\partial f}{\partial t} = u^{m-1} \left[\left(m - \frac{1}{2}\right) \Delta f + \frac{1}{2} \frac{|\nabla f|^2}{f} \right]$$

and compute the time derivative of the Fisher information

$$I[u] = \int_{\mathbb{R}^d} |\nabla f|^2 dx$$

as

$$I' = -2 \int_{\mathbb{R}^d} (\Delta f) \frac{\partial f}{\partial t} dx = -(2m-1) \int_{\mathbb{R}^d} u^{m-1} (\Delta f)^2 dx - \int_{\mathbb{R}^d} u^{m-1} \Delta f \frac{|\nabla f|^2}{f} dx.$$

The computation of the r.h.s. relies on two identities. Here we assume that integrations by parts can be taken without any special precautions. A detailed discussion of this issue can be found in [39].

- *First identity:* we use an integration by parts to obtain

$$\begin{aligned} & - \int_{\mathbb{R}^d} u^{m-1} \Delta f \frac{|\nabla f|^2}{f} dx \\ &= \int_{\mathbb{R}^d} \nabla f \cdot \nabla \left(u^{m-1} \frac{|\nabla f|^2}{f} \right) dx \\ &= \int_{\mathbb{R}^d} (\nabla u^{m-1}) \cdot \nabla f \frac{|\nabla f|^2}{f} dx + 2 \int_{\mathbb{R}^d} u^{m-1} \text{H}f : \frac{\nabla f \otimes \nabla f}{f} dx \\ & \quad - \int_{\mathbb{R}^d} u^{m-1} \frac{|\nabla f|^4}{f^2} dx \end{aligned}$$

where $\text{H}f$ denotes the Hessian matrix of f . After integrating by parts again, we obtain

$$- \int_{\mathbb{R}^d} u^{m-1} \Delta f \frac{|\nabla f|^2}{f} dx = 2 \int_{\mathbb{R}^d} u^{m-1} \text{H}f : \frac{\nabla f \otimes \nabla f}{f} dx - \frac{1}{2m-1} \int_{\mathbb{R}^d} u^{m-1} \frac{|\nabla f|^4}{f^2} dx \quad (10)$$

using

$$\nabla (u^{m-1}) = \frac{2(m-1)}{2m-1} u^{m-1} \frac{\nabla f}{f}. \quad (11)$$

• *Second identity:* using again (11), we find that

$$\begin{aligned} - (2m-1) \int_{\mathbb{R}^d} u^{m-1} (\Delta f)^2 dx &= (2m-1) \int_{\mathbb{R}^d} \nabla f \cdot \nabla (u^{m-1} \Delta f) dx \\ &= 2(m-1) \int_{\mathbb{R}^d} u^{m-1} \Delta f \frac{|\nabla f|^2}{f} dx + (2m-1) \int_{\mathbb{R}^d} u^{m-1} \nabla f \cdot \nabla \Delta f dx. \end{aligned}$$

Next we use the identity $\nabla f \cdot \nabla \Delta f = \frac{1}{2} \Delta(|\nabla f|^2) - \|\mathbf{H}f\|^2$ and obtain

$$\begin{aligned} - (2m-1) \int_{\mathbb{R}^d} u^{m-1} (\Delta f)^2 dx &= 2(m-1) \int_{\mathbb{R}^d} u^{m-1} \Delta f \frac{|\nabla f|^2}{f} dx - (2m-1) \int_{\mathbb{R}^d} u^{m-1} \|\mathbf{H}f\|^2 dx \\ &\quad + (m-\frac{1}{2}) \int_{\mathbb{R}^d} u^{m-1} \Delta(|\nabla f|^2) dx \end{aligned}$$

where, using twice (11), we obtain that

$$\begin{aligned} \int_{\mathbb{R}^d} u^{m-1} \Delta(|\nabla f|^2) dx &= \frac{2(m-1)}{2m-1} \int_{\mathbb{R}^d} |\nabla f|^2 \nabla \cdot \left(u^{m-1} \frac{\nabla f}{f} \right) dx \\ &= \frac{2(m-1)}{2m-1} \int_{\mathbb{R}^d} u^{m-1} \Delta f \frac{|\nabla f|^2}{f} dx + \frac{2(m-1)}{2m-1} \left[\frac{2(m-1)}{2m-1} - 1 \right] \int_{\mathbb{R}^d} u^{m-1} \frac{|\nabla f|^4}{f^2} dx \\ &= \frac{2(m-1)}{2m-1} \int_{\mathbb{R}^d} u^{m-1} \Delta f \frac{|\nabla f|^2}{f} dx - \frac{2(m-1)}{(2m-1)^2} \int_{\mathbb{R}^d} u^{m-1} \frac{|\nabla f|^4}{f^2} dx. \end{aligned}$$

Notice that the above computation will cause trouble for a generalization to the case of a function f which is defined as a non-local expression of $u^{m-\frac{1}{2}}$. Hence

$$\begin{aligned} - (2m-1) \int_{\mathbb{R}^d} u^{m-1} (\Delta f)^2 dx &= 3(m-1) \int_{\mathbb{R}^d} u^{m-1} \Delta f \frac{|\nabla f|^2}{f} dx - (2m-1) \int_{\mathbb{R}^d} u^{m-1} \|\mathbf{H}f\|^2 dx \\ &\quad - \frac{m-1}{2m-1} \int_{\mathbb{R}^d} u^{m-1} \frac{|\nabla f|^4}{f^2} dx. \end{aligned}$$

We can evaluate $\int_{\mathbb{R}^d} u^{m-1} \Delta f \frac{|\nabla f|^2}{f} dx$ using (10) and get that

$$\begin{aligned} - (2m-1) \int_{\mathbb{R}^d} u^{m-1} (\Delta f)^2 dx &= -3(m-1) \left(2 \int_{\mathbb{R}^d} u^{m-1} \mathbf{H}f : \frac{\nabla f \otimes \nabla f}{f} dx - \frac{1}{2m-1} \int_{\mathbb{R}^d} u^{m-1} \frac{|\nabla f|^4}{f^2} dx \right) \\ &\quad - (2m-1) \int_{\mathbb{R}^d} u^{m-1} \|\mathbf{H}f\|^2 dx - \frac{m-1}{2m-1} \int_{\mathbb{R}^d} u^{m-1} \frac{|\nabla f|^4}{f^2} dx. \end{aligned}$$

This provides the second identity

$$\begin{aligned}
& - (2m - 1) \int_{\mathbb{R}^d} u^{m-1} (\Delta f)^2 dx \\
& = - (2m - 1) \int_{\mathbb{R}^d} u^{m-1} \|\mathbf{H}f\|^2 dx - 6(m - 1) \int_{\mathbb{R}^d} u^{m-1} \mathbf{H}f : \frac{\nabla f \otimes \nabla f}{f} dx \\
& \quad + \frac{2(m-1)}{2m-1} \int_{\mathbb{R}^d} u^{m-1} \frac{|\nabla f|^4}{f^2} dx. \quad (12)
\end{aligned}$$

Collecting terms of our two identities (10) and (12), we have found that

$$\begin{aligned}
I' & = - (2m - 1) \int_{\mathbb{R}^d} u^{m-1} \|\mathbf{H}f\|^2 dx - 2(3m - 4) \int_{\mathbb{R}^d} u^{m-1} \mathbf{H}f : \frac{\nabla f \otimes \nabla f}{f} dx \\
& \quad - \frac{3-2m}{2m-1} \int_{\mathbb{R}^d} u^{m-1} \frac{|\nabla f|^4}{f^2} dx,
\end{aligned}$$

that we can also write as

$$I' = - (2m - 1) \left(1 - \frac{m_1}{m}\right) \int_{\mathbb{R}^d} u^{m-1} \left| \Delta f - \frac{1}{2m-1} \frac{|\nabla f|^2}{f} \right|^2 dx - \mathcal{R}$$

with

$$\begin{aligned}
\mathcal{R} & := (2m - 1) \frac{m_1}{m} \int_{\mathbb{R}^d} u^{m-1} \|\mathbf{H}f\|^2 dx \\
& \quad - 2 \left(3(1 - m_1) + \frac{m_1}{m}\right) \int_{\mathbb{R}^d} u^{m-1} \mathbf{H}f : \frac{\nabla f \otimes \nabla f}{f} dx \\
& \quad + \frac{2m(1-m_1)+m_1}{m(2m-1)} \int_{\mathbb{R}^d} u^{m-1} \frac{|\nabla f|^4}{f^2} dx.
\end{aligned}$$

Recalling that $m_1 = (d - 1)/d$, the reader is invited to check that

$$\mathcal{R} = \frac{2m-1}{m} \int_{\mathbb{R}^d} u^{m-1} \left\| \mathbf{L}f - \frac{1}{2m-1} \mathbf{M}f \right\|^2 dx$$

using the notations

$$\mathbf{L}f := \mathbf{H}f - \frac{1}{d} \Delta f \text{Id} \quad \text{and} \quad \mathbf{M}f := \frac{\nabla f \otimes \nabla f}{f} - \frac{1}{d} \frac{|\nabla f|^2}{f} \text{Id},$$

so that

$$\mathbf{H}f : \frac{\nabla f \otimes \nabla f}{f} = \mathbf{H}f : \mathbf{M}f + \frac{1}{d} \Delta f \frac{|\nabla f|^2}{f} = \mathbf{L}f : \mathbf{M}f + \frac{1}{d} \Delta f \frac{|\nabla f|^2}{f}$$

and

$$\|\mathbf{L}f\|^2 = \|\mathbf{H}f\|^2 - \frac{1}{d} (\Delta f)^2 \quad \text{and} \quad \|\mathbf{M}f\|^2 = \left(1 - \frac{1}{d}\right) \frac{|\nabla f|^4}{f^2}.$$

Finally we introduce the Rényi entropy power F defined by

$$F := E^\sigma \quad \text{with} \quad \sigma := \frac{2}{d} \frac{1}{1-m} - 1.$$

If $1 - \frac{1}{d} = m_1 \leq m < 1$, proving that, as a function of t , F is concave amounts to proving that

$$G := \frac{2m-1}{2m(1-m)} E^{2-\sigma} F'' = (\sigma - 1) \frac{2m(1-m)}{2m-1} |I|^2 + E|I'$$

is nonpositive, because of (6). We observe that

$$\begin{aligned} \int_{\mathbb{R}^d} u^{m-1} \left| \Delta f - \frac{1}{2m-1} \frac{|\nabla f|^2}{f} + \frac{2m}{2m-1} \frac{1}{E} \sqrt{u} \right|^2 dx \\ = \int_{\mathbb{R}^d} u^{m-1} \left| \Delta f - \frac{1}{2m-1} \frac{|\nabla f|^2}{f} \right|^2 dx - \left(\frac{2m}{2m-1} \right)^2 \frac{|I|^2}{E} \end{aligned}$$

and, as a consequence,

$$\begin{aligned} G = - (2m-1) \left(1 - \frac{m_1}{m} \right) E \int_{\mathbb{R}^d} u^{m-1} \left| \Delta f - \frac{1}{2m-1} \frac{|\nabla f|^2}{f} + \frac{2m}{2m-1} \frac{1}{E} \sqrt{u} \right|^2 dx \\ - \frac{2m-1}{m} E \int_{\mathbb{R}^d} u^{m-1} \left\| \mathbb{L}f - \frac{1}{2m-1} \mathbb{M}f \right\|^2 dx. \end{aligned}$$

Summarizing, when $s = 0$ and by assuming that there are no boundary terms in the integrations by parts, we establishes that F is a concave increasing function, and shows that

$$F' \geq \kappa = \kappa_\star$$

with the notations of Sections 2.3 and Corollary 3.1. According to [39], this is equivalent to the proof of Inequality (3). The main difference with [39] is that the computations are done in terms of $f = u^{m-\frac{1}{2}}$ instead of u^m .

When $s \neq 0$, it would be desirable to do a similar computation with

$$f = (-\Delta)^{-s/2} u^{m-\frac{1}{2}}$$

but there are several obstructions. First of all, according to Proposition 2.3, we cannot expect that $\kappa = \kappa_\star$. In other words, the self-similar solution u_\star is not optimal for Inequality (3). We can hope that monotonicity of F' holds for some non optimal constant as a consequence of the Bakry-Emery method applied to the fractional fast diffusion equation and generalized Rényi entropy powers: proving it is still open. Technically another obstruction arises from the computations as we have no identity equivalent to (11). As in [11, 12] in the porous medium case, one could expect that a Stroock-Varadhan identity [40–42] allows to give a proof of the optimality with non-optimal constants, and also characterize the stability of the self-similar solutions as $t \rightarrow +\infty$, but this is so far also an open question.

Acknowledgements. This work is supported by a public grant overseen by the French National Research Agency (ANR) as part of the ‘‘Investissements d’Avenir’’ program (A.Z.,

reference: ANR-10-LABX-0098, LabEx SMP) and by the projects *STAB* (J.D., A.Z.) and *Kibord* (J.D.) of the French National Research Agency (ANR). A.Z. thanks the ERC Advanced Grant BLOWDISOL (Blow-up, dispersion and solitons; PI: Frank Merle) # 291214 for support. The authors thank Maria J. Esteban for fruitful discussions and suggestions.

© 2016 by the authors. This paper may be reproduced, in its entirety, for non-commercial purposes.

References

- [1] Bouchaud JP, Georges A. Anomalous diffusion in disordered media: statistical mechanisms, models and physical applications. *Physics reports*. 1990;195(4):127–293; Available from: <http://www.sciencedirect.com/science/article/pii/037015739090099N>.
- [2] Metzler R, Klafter J. The random walk’s guide to anomalous diffusion: a fractional dynamics approach. *Physics reports*. 2000;339(1):1–77; Available from: <http://www.sciencedirect.com/science/article/pii/S0370157300000703>.
- [3] Caffarelli L, Soria F, Vázquez JL. Regularity of solutions of the fractional porous medium flow. *J Eur Math Soc (JEMS)*. 2013;15(5):1701–1746; Available from: <https://dx.doi.org/10.4171/JEMS/401>.
- [4] Caffarelli L, Vazquez JL. Nonlinear porous medium flow with fractional potential pressure. *Arch Ration Mech Anal*. 2011;202(2):537–565; Available from: <https://dx.doi.org/10.1007/s00205-011-0420-4>.
- [5] Caffarelli LA, Vázquez JL. Asymptotic behaviour of a porous medium equation with fractional diffusion. *Discrete Contin Dyn Syst*. 2011;29(4):1393–1404; Available from: <http://aimsciences.org/journals/pdfs.jsp?paperID=5715&mode=full>.
- [6] de Pablo A, Quirós F, Rodríguez A, Vázquez JL. A general fractional porous medium equation. *Comm Pure Appl Math*. 2012;65(9):1242–1284; Available from: <https://dx.doi.org/10.1002/cpa.21408>.
- [7] de Pablo A, Quirós F, Rodríguez A, Vázquez JL. Classical solutions for a logarithmic fractional diffusion equation. *J Math Pures Appl (9)*. 2014;101(6):901–924; Available from: <https://dx.doi.org/10.1016/j.matpur.2013.10.009>.
- [8] Vázquez JL. Barenblatt solutions and asymptotic behaviour for a nonlinear fractional heat equation of porous medium type. *J Eur Math Soc (JEMS)*. 2014;16(4):769–803; Available from: <https://dx.doi.org/10.4171/JEMS/446>.
- [9] Del Pino M, Dolbeault J. Best constants for Gagliardo-Nirenberg inequalities and applications to nonlinear diffusions. *J Math Pures Appl (9)*. 2002;81(9):847–875; Available from: [https://dx.doi.org/10.1016/S0021-7824\(02\)01266-7](https://dx.doi.org/10.1016/S0021-7824(02)01266-7).
- [10] Otto F. The geometry of dissipative evolution equations: the porous medium equation. *Comm Partial Differential Equations*. 2001;26(1-2):101–174; Available from: <https://dx.doi.org/10.1081/PDE-100002243>.
- [11] Biler P, Imbert C, Karch G. Barenblatt profiles for a nonlocal porous medium equation. *C R Math Acad Sci Paris*. 2011;349(11-12):641–645; Available from: <http://dx.doi.org/10.1016/j.crma.2011.06.003>.
- [12] Biler P, Imbert C, Karch G. The nonlocal porous medium equation: Barenblatt profiles and other weak solutions. *Arch Ration Mech Anal*. 2015;215(2):497–529; Available from: <https://dx.doi.org/10.1007/s00205-014-0786-1>.
- [13] Carrillo JA, Huang Y, Santos MC, Vázquez JL. Exponential convergence towards stationary states for the 1D porous medium equation with fractional pressure. *J Differential Equations*. 2015;258(3):736–763; Available from: <https://dx.doi.org/10.1016/j.jde.2014.10.003>.
- [14] Huang Y. Explicit Barenblatt profiles for fractional porous medium equations. *Bull Lond Math Soc*. 2014;46(4):857–869; Available from: <https://dx.doi.org/10.1112/blms/bdu045>.
- [15] Stan D, del Teso F, Vázquez JL. Transformations of self-similar solutions for porous

- medium equations of fractional type. *Nonlinear Anal.* 2015;119:62–73; Available from: <http://dx.doi.org/10.1016/j.na.2014.08.009>.
- [16] Chen S, Frank RL, Weth T. Remainder terms in the fractional Sobolev inequality. *Indiana Univ Math J.* 2013;62(4):1381–1397; Available from: <http://dx.doi.org/10.1512/iumj.2013.62.5065>.
- [17] Aubin T. Problèmes isopérimétriques et espaces de Sobolev. *C R Acad Sci Paris Sér A-B.* 1975;280(5):Aii, A279–A281.
- [18] Talenti G. Best constant in Sobolev inequality. *Ann Mat Pura Appl* (4). 1976;110:353–372; Available from: <https://link.springer.com/article/10.1007%2FBF02418013>.
- [19] Rosen G. Minimum value for c in the Sobolev inequality $\|\phi^3\| \leq c\|\nabla\phi\|^3$. *SIAM J Appl Math.* 1971;21:30–32; Available from: <https://epubs.siam.org/doi/abs/10.1137/0121004>.
- [20] Bliss G. An integral inequality. *J London Math Soc.* 1930;1(1):40; Available from: <doi:10.1112/jlms/s1-5.1.40>.
- [21] Lieb EH. Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities. *Ann of Math* (2). 1983;118(2):349–374; Available from: <https://dx.doi.org/10.2307/2007032>.
- [22] Escobar JF. Sharp constant in a Sobolev trace inequality. *Indiana Univ Math J.* 1988;37(3):687–698; Available from: <https://dx.doi.org/10.1512/iumj.1988.37.37033>.
- [23] Swanson CA. The best Sobolev constant. *Appl Anal.* 1992;47(4):227–239; Available from: <https://dx.doi.org/10.1080/00036819208840142>.
- [24] Chang SYA, González MdM. Fractional Laplacian in conformal geometry. *Adv Math.* 2011;226(2):1410–1432; Available from: <http://dx.doi.org/10.1016/j.aim.2010.07.016>.
- [25] Chen W, Li C, Ou B. Classification of solutions for an integral equation. *Comm Pure Appl Math.* 2006;59(3):330–343; Available from: <http://dx.doi.org/10.1002/cpa.20116>.
- [26] Hajaiej H, Yu X, Zhai Z. Fractional Gagliardo-Nirenberg and Hardy inequalities under Lorentz norms. *J Math Anal Appl.* 2012;396(2):569–577; Available from: <http://dx.doi.org/10.1016/j.jmaa.2012.06.054>.
- [27] Hajaiej H. Symmetry of minimizers of some fractional problems. *Applicable Analysis.* 2015;94(4):694–700; Available from: <http://dx.doi.org/10.1080/00036811.2014.898273>.
- [28] Trabelsi S. Optimal functions in fractional Gagliardo-Nirenberg-Sobolev inequalities. 2016; in preparation.
- [29] Hajaiej H. Existence of minimizers of functionals involving the fractional gradient in the absence of compactness, symmetry and monotonicity. *Journal of Mathematical Analysis and Applications.* 2013;399(1):17 – 26; Available from: <http://www.sciencedirect.com/science/article/pii/S0022247X1200755X>.
- [30] Gérard P. Description du défaut de compacité de l’injection de Sobolev. *ESAIM Control Optim Calc Var.* 1998;3:213–233 (electronic); Available from: <http://dx.doi.org/10.1051/cocv:1998107>.
- [31] Lions PL. The concentration-compactness principle in the calculus of variations. The locally compact case. II. *Ann Inst H Poincaré Anal Non Linéaire.* 1984;1(4):223–283; Available from: http://www.numdam.org/item?id=AIHPC_1984__1_4_223_0.
- [32] Autuori G, Pucci P. Elliptic problems involving the fractional laplacian in \mathbb{R}^n . *Journal of Differential Equations.* 2013 10;255(8):2340–2362; Available from: <http://www.sciencedirect.com/science/article/pii/S0022039613002532>.
- [33] Savaré G, Toscani G. The concavity of Rényi entropy power. *IEEE Trans Inform Theory.* 2014;60(5):2687–2693; Available from: <http://dx.doi.org/10.1109/TIT.2014.2309341>.
- [34] Dolbeault J, Toscani G. Nonlinear diffusions: Extremal properties of Barenblatt profiles, best matching and delays. *Nonlinear Analysis: Theory, Methods & Applications.* 2016 6;138:31–43; Available from:

- <http://www.sciencedirect.com/science/article/pii/S0362546X15003880>.
- [35] Vázquez JL, Volzone B. Optimal estimates for fractional fast diffusion equations. *J Math Pures Appl* (9). 2015;103(2):535–556; Available from: <http://dx.doi.org/10.1016/j.matpur.2014.07.002>.
 - [36] Blanchet A, Bonforte M, Dolbeault J, Grillo G, Vázquez JL. Asymptotics of the fast diffusion equation via entropy estimates. *Archive for Rational Mechanics and Analysis*. 2009 02;191(2):347–385; Available from: <http://dx.doi.org/10.1007/s00205-008-0155-z>.
 - [37] de Pablo A, Quirós F, Rodríguez A. Nonlocal filtration equations with rough kernels. *Nonlinear Anal*. 2016;137:402–425; Available from: <https://dx.doi.org/10.1016/j.na.2016.01.026>.
 - [38] Dolbeault J, Toscani G. Fast diffusion equations: matching large time asymptotics by relative entropy methods. *Kinetic and Related Models*. 2011;4(3):701–716; Available from: <http://dx.doi.org/10.3934/krm.2011.4.701>.
 - [39] Dolbeault J, Esteban MJ, Loss M. Interpolation inequalities, nonlinear flows, boundary terms, optimality and linearization. 2016 Nov; hal: hal-01395771 and ArXiv: 1611.04321; Available from: <https://hal.archives-ouvertes.fr/hal-01395771>.
 - [40] Liskevich VA, Semenov YA. Some problems on Markov semigroups. In: *Schrödinger operators, Markov semigroups, wavelet analysis, operator algebras*. Vol. 11 of *Math. Top.*; Akademie Verlag, Berlin; 1996. p. 163–217; Available from: <http://onlinelibrary.wiley.com/doi/10.1002/cpa.3160250603/>.
 - [41] Stroock D, Varadhan SRS. On degenerate elliptic-parabolic operators of second order and their associated diffusions. *Comm Pure Appl Math*. 1972;25:651–713; Available from: <http://onlinelibrary.wiley.com/doi/10.1002/cpa.3160250603>.
 - [42] Stroock DW, Varadhan SRS. On the support of diffusion processes with applications to the strong maximum principle. In: *Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability* (Univ. California, Berkeley, Calif., 1970/1971), Vol. III: Probability theory; June-July 1971. Univ. California Press, Berkeley, Calif.; 1972. p. 333–359.