



HAL
open science

Discrete self-similar and ergodic Markov chains

Laurent Miclo, Pierre Patie, Rohan Sarkar

► **To cite this version:**

Laurent Miclo, Pierre Patie, Rohan Sarkar. Discrete self-similar and ergodic Markov chains. *Annals of Probability*, In press. hal-03777200

HAL Id: hal-03777200

<https://hal.science/hal-03777200>

Submitted on 14 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DISCRETE SELF-SIMILAR AND ERGODIC MARKOV CHAINS

LAURENT MICLO[†], PIERRE PATIE, AND ROHAN SARKAR

ABSTRACT. The first aim of this paper is to introduce a class of Markov chains on \mathbb{Z}_+ which are discrete self-similar in the sense that their semigroups satisfy an invariance property expressed in terms of a discrete random dilation operator. After showing that this latter property requires the chains to be upward skip-free, we first establish a gateway relation, a concept introduced in [26], between the semigroup of such chains and the one of spectrally negative self-similar Markov processes on \mathbb{R}_+ . As a by-product, we prove that each of these Markov chains, after an appropriate scaling, converge in the Skorohod metric, to the associated self-similar Markov process. By a linear perturbation of the generator of these Markov chains, we obtain a class of ergodic Markov chains, which are non-reversible. By means of intertwining and interweaving relations, where the latter was recently introduced in [27], we derive several deep analytical properties of such ergodic chains including the description of the spectrum, the spectral expansion of their semigroups, the study of their convergence to equilibrium in the ϕ -entropy sense as well as their hypercontractivity property.

Keywords: Discrete self-similarity, Markov chains, generalized Meixner polynomials, intertwining, non-reversible, spectral theory, ergodicity constants, convergence to equilibrium, hypercontractivity.

2010 Mathematical Subject Classification: 41A60, 47G20, 33C45, 47D07, 37A30, 60J27, 60J60

CONTENTS

1. Introduction	2
2. Main Results	5
3. Examples	15
4. Proof of the Main Results	18

[†] Funding from the grant ANR-17-EURE-0010 is acknowledged.

The authors are very grateful to the referees for the careful reading of the manuscript and their positive and constructive comments.

1. INTRODUCTION

Self-similar processes are ubiquitous in the theory of Markov processes and they have been studied intensively over the last three decades, both from theoretical and applied perspectives. A Markov process on \mathbb{R}_+ is called self-similar of index 1 if for all $\alpha > 0$, one has the following commutation type relation

$$(1.1) \quad Q_t d_\alpha = d_\alpha Q_{\alpha t}$$

where $Q = (Q_t)_{t \geq 0}$ is the semigroup associated with the process and $d_\alpha f(x) = f(\alpha x)$ is the dilation operator, that satisfies the semigroup property $d_\alpha d_\beta = d_{\alpha\beta}$ for all $\alpha, \beta > 0$. Motivated by limit theorems, Lamperti [24] obtained a complete characterization of these processes.

In this paper, we first aim at introducing continuous-time Markov processes with state space the set of all nonnegative integers that also enjoy a scaling type property. Naturally, one cannot expect (1.1) to hold in this setting, because the set of integers is not stable by the dilation operators as defined above. However, in [26], the authors introduced the following signed Binomial kernel defined by

$$\mathbb{D}_\alpha f(n) = \sum_{k=0}^n \binom{n}{k} \alpha^k (1-\alpha)^{n-k} f(k)$$

which resembles the dilation operator through the multiplicative semigroup property $\mathbb{D}_{\alpha\beta} = \mathbb{D}_\alpha \mathbb{D}_\beta$ for all $\alpha, \beta > 0$, which will be proved in Proposition 4.1 below. Furthermore, they showed that the linear birth-death Markov chain, see Remark 2.3 below for definition, satisfies the following commutation type relation

$$Q_t \mathbb{D}_\alpha = \mathbb{D}_\alpha Q_{\alpha t}$$

where Q is the associated semigroup. Motivated from this result, we introduce a class of continuous-time Markov chains on \mathbb{Z}_+ that satisfy the scaling property as above and are upward skip-free, that is, at any instant the Markov chains do not jump more than one step above and name them **discrete self-similar Markov chains**, see Definition 2.1. This class of Markov chains, to the best of our knowledge, have not been identified before. Moreover, we want to understand their connections with self-similar Markov processes. To this end, we resort to **intertwining relationship**

between Markov processes. More specifically, for two Markov semigroups P and Q , we say that they are intertwined if, for all $t \geq 0$,

$$P_t \Lambda = \Lambda Q_t$$

for some linear operator Λ . Note that when the underlying processes have different state spaces, one lattice and the other one continuous, we use the terminology gateway relation, coined in [26], to emphasize the unexpected two-sided connection between the two worlds. The term duality is also used in a fast growing and fascinating literature on this topic related to differential operators arising in statistical mechanics, see e.g. [3, 12, 18, 33, 19] and references therein. More generally, the concept of intertwining relation goes back to Dynkin [16] who used it to construct new Markov semigroups from a reference one. These ideas were extended by Rogers and Pitman in [32], leading to the characterization of Markov functions; that is, measurable maps that preserve the Markov property. With the help of the intertwining relationship, we prove the Feller property of the discrete self-similar Markov chains, see Theorem 2.6, and obtain the spectrally negative self-similar Markov processes as the scaling limit of these Markov chains, see Theorem 2.6(2). The use of intertwining relations to prove limit theorems is not new and, in fact, a general framework was built up by Borodin and Olshanski [10], where they apply it to construct a class of Markov chains on the Thoma cone. Unfortunately, their strategy is not applicable in our situation because their conditions are too stringent for us, namely the set of finitely supported functions are not invariant with respect to the discrete self-similar Markov semigroups. Nonetheless, still resorting to the intertwining relation, we are able to derive explicit formulas for the moments of these Markov chains and we identify their scaling limits by the method of moments. We emphasize that there are many instances of the appearance of positive self-similar Markov processes as the scaling limits of models, such as coalescence-fragmentation processes, see Bertoin [7], random planar maps, see Le Gall and Miermont [25]. We also mention the recent paper by Bertoin and Kortchemski [9] where the authors introduce a class of discrete-time Markov chains whose appropriate scaling limits are positive self-similar Markov processes. It appears that our work offers another class of Markov chains in the domain of attraction of such self-similar Markov processes, with the additional surprising feature that the connection between the two objects goes, thanks to the gateway relation, in both directions.

We proceed by introducing another class of ergodic Markov chains which are obtained by a linear first order perturbation of the generators of the discrete self-similar Markov chains. We name them **skip-free Laguerre** chains. The motivation behind this comes from the fact that their continuous analogue are the generalized Laguerre processes, studied in [29], which are

also constructed by perturbation of the generator of self-similar processes by a linear convection term, that is a first order differential operator with a linear coefficient. We show that they generate a class of Feller semigroups of ergodic Markov chains which intertwine with the class of the generalized Laguerre semigroups. Using this connection, we develop the spectral theory, including the spectrum and the eigenvalues expansions, in the Hilbert space ℓ^2 of nonnegative integers weighted with the invariant distributions \mathbf{n}_ϕ of the semigroups of these non-reversible chains. As by-product, and under some mild conditions, we prove compactness and also obtain a hypercoercivity estimate for the $\ell^2(\mathbf{n}_\phi)$ convergence to equilibrium, which is given explicitly as a perturbed spectral gap inequality. This part involves a deep theory of non-self-adjoint operators as developed in [29], see Section 4.11 for more details.

We continue our analysis of these skip-free Laguerre semigroups by investigating the entropy decay to equilibrium as well as the hypercontractivity property. For self-adjoint Markov semigroups, these two phenomena are equivalent to the (modified) log-Sobolev inequalities. Unfortunately, in our context, this relation fails due to the non-self-adjointness of the semigroups. However, resorting to the idea of interweaving relation, introduced recently in [27], we relate the skip-free Laguerre semigroups with the self-adjoint diffusion Laguerre semigroups and deduce, up to some universal random time, both the entropy decay and the hypercontractivity. Finally, showing that this random time is infinitely divisible, we develop a thorough analysis of the skip-free Laguerre semigroups subordinated with the associated subordinator, which generate a class of ergodic Markov chains with two-sided jumps, for which all the results described above are obtained explicitly.

The remaining part of the paper is organized as follows. Most of the frequently used notations are defined in Section 1.1 while Section 2 contains all the main results of the paper. We provide some examples in Section 3 and Section 4 is devoted to the proofs of the main results. Some aspects of spectral theory for non-self-adjoint operators have been reviewed in Subsection 4.11 and the results related to interweaving relations have been proved in Subsection 4.14.

1.1. Notations and Preliminaries. For any locally compact topological space E we write $\mathbf{C}(E)$, $\mathbf{C}_b(E)$, $\mathbf{C}_c(E)$ and $\mathbf{C}_0(E)$ to denote the class of continuous functions (the set of all functions when $E = \mathbb{Z}_+$), class of all bounded continuous functions, class of all compactly supported continuous functions and class of all continuous functions vanishing at infinity on E respectively. In addition, when $E = \mathbb{R}$ or \mathbb{R}_+ , we write $\mathbf{C}_b^\infty(E)$ to denote the class of all bounded smooth functions with bounded derivatives on E .

Next, for any nonnegative sigma-finite measure μ on \mathbb{R}_+ and $p \in [1, \infty]$, $\mathbf{L}^p(\mu)$ denotes the L^p space with weight μ . When $p = 2$, the corresponding Hilbert space is endowed with the inner product denoted by $\langle f, g \rangle_\mu = \int_{\mathbb{R}_+} f(x)\overline{g(x)}\mu(dx)$. When μ is the Lebesgue measure, we simply write $\mathbf{L}^2(\mu) = \mathbf{L}^2(\mathbb{R}_+)$ associated with the inner product $\langle \cdot, \cdot \rangle$. If the underlying space is the set of all integers \mathbb{Z}_+ , then for any nonnegative discrete measure \mathbf{m} on \mathbb{Z}_+ , we write $\ell^p(\mathbf{m})$ to denote the weighted ℓ^p space on \mathbb{Z}_+ and for $p = 2$, the inner product is written as $\langle \mathbf{f}, \mathbf{g} \rangle_{\mathbf{m}} = \sum_{n \in \mathbb{Z}_+} \mathbf{f}(n)\overline{\mathbf{g}(n)}\mathbf{m}(n)$. When \mathbf{m} is the counting measure, we use the notation $\ell^2(\mathbb{Z}_+) = \ell^2(\mathbf{m})$. For any measurable function $f \geq 0$ or $f \in \mathbf{L}^1(E, \mu)$, we write $\mu f = \int_E f d\mu$.

For any two Banach spaces $\mathbf{B}_1, \mathbf{B}_2$, $\mathcal{B}(\mathbf{B}_1, \mathbf{B}_2)$ denotes the set of all bounded linear operators defined from \mathbf{B}_1 to \mathbf{B}_2 . Finally, for any operator A (possibly unbounded) defined on some Banach space, $\mathbf{D}(A)$ denotes the domain of the operator and we represent the operator as $(A, \mathbf{D}(A))$ and in case of Hilbert spaces, we denote the adjoint of A by \hat{A} .

We denote the complex plane by \mathbb{C} and for any $z \in \mathbb{C}$, $\operatorname{Re}(z), \operatorname{Im}(z)$ denote the real and imaginary part of z respectively. Next, for any $S \subset \mathbb{R}$, we write $\mathbb{C}_S = \{z \in \mathbb{C}; \operatorname{Re}(z) \in S\}$. In particular, when $S = \mathbb{R}_+$ (resp. \mathbb{R}_-), we simply write \mathbb{C}_+ (resp. \mathbb{C}_-).

For two functions f, g defined on the real line, we use the following notation.

$$f \asymp g \text{ means that } \exists c > 0 \text{ such that, for all } x, c^{-1} \leq \frac{f(x)}{g(x)} \leq c$$

$$f \stackrel{a}{\sim} g \text{ means that } \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = 1 \text{ for some } a \in [0, \infty]$$

$$f(x) \stackrel{a}{=} O(g(x)) \text{ means that } \overline{\lim}_{x \rightarrow a} \left| \frac{f(x)}{g(x)} \right| < \infty$$

$$f(x) \stackrel{a}{=} o(g(x)) \text{ means that } \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = 0.$$

2. MAIN RESULTS

2.1. Discrete dilation and discrete self-similar Markov chains. We start by introducing a transformation on $\mathbf{C}(\mathbb{Z}_+)$, which we name the **discrete dilation operator**. For any $\alpha > 0$ and $\mathbf{f} \in \mathbf{C}(\mathbb{Z}_+)$, we define

$$(2.1) \quad \mathbb{D}_\alpha \mathbf{f}(n) = \sum_{r=0}^n \binom{n}{r} \alpha^r (1-\alpha)^{n-r} \mathbf{f}(r).$$

It should be noted that \mathbb{D}_α is well defined on $\mathbf{C}(\mathbb{Z}_+)$ for all $\alpha \geq 0$ and it is a Markov kernel when $\alpha \in [0, 1]$. When $\alpha > 1$, $\mathbb{D}_\alpha \mathbf{f}$ may not be bounded even

if f is bounded. For instance, taking $f(n) = (-1)^n$, for any $n \in \mathbb{Z}_+$, we have $|\mathbb{D}_\alpha f(n)| = (2\alpha - 1)^n$, which grows exponentially with respect to n . The operator \mathbb{D} shares the multiplicative semigroup property with the dilation operator, that is, for all $\alpha, \beta > 0$, we have $\mathbb{D}_{\alpha\beta} = \mathbb{D}_\alpha \mathbb{D}_\beta$, see Proposition 4.1 below. Next, we introduce the discrete self-similar Markov chains which are defined in terms of the operator \mathbb{D}_α .

Definition 2.1. We say that the semigroup $\mathbb{Q} = (\mathbb{Q}_t)_{t \geq 0}$ of a continuous-time Markov chain \mathbb{X} with state space \mathbb{Z}_+ is **discrete self-similar** if for all $t \geq 0, \alpha \in [0, 1]$, the following identity

$$(2.2) \quad \mathbb{Q}_t \mathbb{D}_\alpha = \mathbb{D}_\alpha \mathbb{Q}_{\alpha t}$$

holds on $\mathbf{C}_b(\mathbb{Z}_+)$.

In terms of the law of the Markov chain $\mathbb{X} = (\mathbb{X}(t, n), n \in \mathbb{Z}_+)_{t \geq 0}$, where $\mathbb{X}(t, n)$ means that it is issued from n , the discrete self-similarity can be interpreted by the following identity in distribution, for any $\alpha \in [0, 1]$, $t \geq 0$ and $n \in \mathbb{Z}_+$,

$$(2.3) \quad \mathbb{B}(\mathbb{X}(t, n), \alpha) \stackrel{(d)}{=} \mathbb{X}(\alpha t, \mathbb{B}(n, \alpha))$$

where $\mathbb{B}(n, \alpha)$ is a Binomial random variable with parameter n and α , and $\mathbb{X}(t, \mathbb{B}(n, \alpha))$ is the chain at time t with initial law the one of $\mathbb{B}(n, \alpha)$.

Next, we consider the class of triplets (m, σ^2, Π) such that $m, \sigma^2 \geq 0$ and Π is a non-negative measure on \mathbb{R}_+ that satisfies

$$(2.4) \quad \int_0^\infty (y \wedge y^2) \Pi(dy) < \infty,$$

that is Π is a Lévy measure with a finite first moment away from 0. To each of these triplets, we associate the so-called Bernstein function defined as

$$(2.5) \quad \phi(u) = m + \sigma^2 u + \int_0^\infty (1 - e^{-uy}) \bar{\Pi}(y) dy$$

where $\bar{\Pi}(y) = \Pi(y, \infty)$ is the tail of the measure Π . Let \mathbf{B} denote the class of all functions of the form (2.5).

We are now ready to introduce a class of discrete operators on \mathbb{Z}_+ , which is the central object of this paper. For any $\phi \in \mathbf{B}$ associated with the triplet (m, σ^2, Π) and $f \in \mathbf{C}_c(\mathbb{Z}_+)$, we define

$$(2.6) \quad \mathbb{G}_\phi f(n) = \sigma^2 n (\partial_+ + \partial_-) f(n) + (m + \sigma^2) \partial_+ f(n) + \mathbb{G}_\Pi f(n)$$

where $\partial_{\pm}f(n) = f(n \pm 1) - f(n)$ for all $n \in \mathbb{Z}_+$ and

$$(2.7) \quad \mathbb{G}_{\Pi}f(n) = \frac{1}{n+1} \int_0^{\infty} [\mathbb{D}_{e^{-y}}f(n+1) - f(n+1) + y(n+1)\partial_+f(n)] \Pi(dy).$$

We are now ready to state our first main result.

Theorem 2.2. *The operator $(\mathbb{G}_{\phi}, \mathbf{C}_c(\mathbb{Z}_+))$ generates a Feller Markov chain on \mathbb{Z}_+ , denoted by $\mathbb{X}_{\phi} = (\mathbb{X}_{\phi}(t, n), n \in \mathbb{Z}_+)_{t \geq 0}$ which is self-similar, and $\mathbf{C}_c(\mathbb{Z}_+)$ serves as a core for \mathbb{G}_{ϕ} .*

This theorem is proved in Section 4.1.

Remark 2.3. When $\Pi \equiv 0$, \mathbb{X}_{ϕ} is the reversible linear birth-death chain with invariant measure

$$\frac{\Gamma(n+m+1)}{\Gamma(n+1)}, \quad n \in \mathbb{Z}_+.$$

For a detailed account on such Markov chains, we refer to [26].

Remark 2.4. In (2.2), we restrict $\alpha \in [0, 1]$ as \mathbb{D}_{α} is, in this case, a Markov kernel. However, since $\mathbb{D}_{\alpha}\mathbf{p}_k(n) = \alpha^k\mathbf{p}_k(n)$, for all $\alpha > 0$ and $k, n \in \mathbb{Z}_+$, where \mathbf{p}_k is defined in (4.22) below, Theorem 4.8, also below, yields that for all $\phi \in \mathbf{B}$ and $t \geq 0$, $\mathbb{Q}_t^{\phi}\mathbb{D}_{\alpha}\mathbf{p}_k(n) = \mathbb{D}_{\alpha}\mathbb{Q}_{\alpha t}^{\phi}\mathbf{p}_k(n)$, where \mathbb{Q}^{ϕ} is the discrete self-similar semigroup generated by \mathbb{G}_{ϕ} . This reveals that the discrete self-similarity property also holds in a more general framework than the one given in (2.2).

A continuous-time Markov chain is called upward skip-free if it does not jump more than one step above at any instant, that is, for any $n \in \mathbb{Z}_+$ and $l \geq n+2$, $\mathbb{G}(n, l) = 0$ where \mathbb{G} is the generator of the Markov chain. It can be easily shown that the discrete self-similar Markov chain \mathbb{X}_{ϕ} with generator \mathbb{G}_{ϕ} is upward skip-free, see (4.3) below. In the next theorem we show the converse claim that is any discrete self-similar Markov chains must be upward skip-free.

Theorem 2.5. *Let \mathbb{X} be any continuous-time discrete self-similar Markov chain on \mathbb{Z}_+ . Then \mathbb{X} is upward skip-free.*

This theorem is proved in Section 4.2.

2.2. Connections with self-similar Markov processes: gateway relation and scaling limit. Self-similar Markov processes on the positive real line are well studied as they appear as the weak limits of various Markov processes, see Lamperti [23]. When these processes are **spectrally negative**, that is, they do not have any positive jumps, and with 0 as an entrance-non-exit boundary, Lamperti [24] showed that they are in bijection with the subset of Bernstein functions \mathbf{B} defined in (2.5) and moreover, the generator of these processes are of the form

$$(2.8) \quad G_\phi f(x) = \sigma^2 x f''(x) + (m + \sigma^2) f'(x)$$

$$(2.9) \quad + \frac{1}{x} \int_0^\infty [d_{e^{-y}} f(x) - f(x) + y x f'(x)] \Pi(dy)$$

where ϕ is defined in terms of the triplet (m, σ^2, Π) , see (2.5) and $f \in \mathbf{C}_c^\infty(\mathbb{R}_+)$. The careful reader will have noticed that the operator \mathbb{G}_ϕ in (2.6) is the discrete analogue of the operator G_ϕ , revealing that the former is a natural approximation of the latter. However, we provide below a deeper connection between these class of Markov processes (operators) by establishing a gateway relation between their semigroups, a concept introduced in [27], meaning that the connection goes in both directions. As a by-product, we show that discrete self-similar Markov chains, after scaling appropriately, converge to the self-similar Markov processes in the Skorohod's J_1 -topology.

Theorem 2.6. (1) *Gateway relation.* For any $\phi \in \mathbf{B}$, let Q^ϕ and \mathbb{Q}^ϕ denote the Feller semigroups generated by G_ϕ and \mathbb{G}_ϕ respectively. Then, for any $f \in \mathbf{C}_0(\mathbb{Z}_+)$ and $t \geq 0$,

$$(2.10) \quad Q_t^\phi \Lambda f = \Lambda \mathbb{Q}_t^\phi f$$

where $\Lambda f(x) = \mathbb{E}[f(\text{Pois}(x))]$, $\text{Pois}(x)$ being a Poisson random variable with parameter $x > 0$.

(2) *Scaling limit.* For any $\phi \in \mathbf{B}$, let \mathbb{X}_ϕ (resp. $X_\phi = (X_\phi(t, x))_{t \geq 0}$) be the discrete self-similar Markov chain (resp. the positive self-similar Markov process issued from x), then, for all $x > 0$,

$$(2.11) \quad \left(\frac{1}{n} \mathbb{X}_\phi(nt, \lfloor nx \rfloor) \right)_{t \geq 0} \longrightarrow (X_\phi(t, x))_{t \geq 0}$$

in Skorohod's J_1 -topology.

Remark 2.7. As mentioned to us by an anonymous referee, the gateway relationship (2.10) has the following neat probabilistic interpretation, using the notation of item (2) above,

$$(2.12) \quad \mathbb{X}_\phi(t, \text{Pois}(x)) \stackrel{(d)}{=} \text{Pois}(X_\phi(t, x))$$

which is valid for any $t, x > 0$. Using the self-similarity property of X_ϕ , this identity yields, for any fixed $t, x > 0$ and large integer n (but not for $\lfloor nx \rfloor$), $\frac{1}{n}\mathbb{X}_\phi(t, \text{Pois}(nx)) \stackrel{(d)}{=} \frac{1}{n}\text{Pois}(X_\phi(t, nx)) \stackrel{(d)}{=} \frac{1}{n}\text{Pois}(nX_\phi(t, x)) \rightarrow X_\phi(t, x)$ in distribution. Moreover, the identity (2.12) boils down when x tends to 0 to

$$\mathbb{X}_\phi(t, 0) \stackrel{(d)}{=} \text{Pois}(X_\phi(t, 0))$$

where $X_\phi(t, 0)$ stands for the entrance law of X_ϕ which is known to exist as $m \geq 0$, see e.g. [29].

The intertwining relation in (1) is proved in Proposition 4.3(1) and the scaling limit in (2) is proved in Section 4.4.

2.3. Discrete Laguerre chains from discrete self-similar Markov chains. Let us now consider a perturbation of the discrete self-similar Markov chains, that is, we introduce a new family of discrete operators on $\mathbf{C}_c(\mathbb{Z}_+)$ defined by

$$(2.13) \quad \mathbb{L}_\phi \mathbf{f}(n) = \mathbb{G}_\phi \mathbf{f}(n) + n\partial_- \mathbf{f}(n)$$

where $\phi \in \mathbf{B}$ and \mathbb{G}_ϕ is defined in (2.6). Alternatively, the operator \mathbb{L}_ϕ can be represented, for any $\mathbf{f} \in \mathbf{C}_c(\mathbb{Z}_+)$, as follows

$$(2.14) \quad \mathbb{L}_\phi \mathbf{f}(n) = \sum_{l=0}^{n+1} \mathbb{L}_\phi(n, l) \mathbf{f}(l)$$

where

$$(2.15) \quad \mathbb{L}_\phi(n, l) = \begin{cases} \mathbb{G}_\phi(n, l) & \text{if } l \neq n, n-1 \\ \mathbb{G}_\phi(n, n-1) + n & \text{if } l = n-1 \\ \mathbb{G}_\phi(n, n) - n & \text{if } l = n \end{cases}$$

with $\mathbb{G}_\phi(n, l) = \mathbb{G}_\phi \delta_l(n)$ and $\delta_l(n) = \mathbb{1}_{\{l=n\}}$.

Theorem 2.8. (1) For any $\phi \in \mathbf{B}$, the operator $(\mathbb{L}_\phi, \mathbf{C}_c(\mathbb{Z}_+))$ generates a Feller Markov semigroup on $\mathbf{C}_0(\mathbb{Z}_+)$, which we denote by \mathbb{K}^ϕ .

(2) We have, for any $\mathbf{f} \in \mathbf{C}_0(\mathbb{Z}_+)$ and $t \geq 0$,

$$(2.16) \quad \mathbb{K}_t^\phi \mathbf{f} = \mathbb{Q}_{e^t-1}^\phi \mathbb{D}_{e^{-t}} \mathbf{f}.$$

(3) The semigroup \mathbb{K}^ϕ has a unique invariant distribution denoted by \mathbf{n}_ϕ and $\mathbf{n}_\phi(n) > 0$ for all $n \in \mathbb{Z}_+$. Moreover, \mathbf{n}_ϕ has moments of all orders and it is moment determinate.

(4) Finally, the semigroup \mathbb{K}^ϕ is self-adjoint in $\ell^2(\mathbf{n}_\phi)$ if and only if $\phi(u) = m + \sigma^2 u$ for some $m, \sigma^2 \geq 0$.

Remark 2.9. In Propositions 4.11 and 4.12, we provide additional properties, including several representations, of the invariant measure \mathbf{n}_ϕ .

We have omitted the proof of the item (1) since it can be obtained by following a line of reasoning similar to the proof of Theorem 2.2 from the claims given in Proposition 4.11. Item 2 is proved after this latter Proposition. The properties of the invariant distribution in item (3) are proved in Proposition 4.11(2) and Proposition 4.12. Item (4) is proved in Proposition 4.11(4).

We name the Markov semigroup \mathbb{K}^ϕ (resp. the Markov chain) the **skip-free Laguerre semigroup** (resp. **skip-free Laguerre chain**). This is motivated by the following observation. The operator \mathbb{L}_ϕ can be viewed as the discrete analogue of the generalized Laguerre operator on \mathbb{R}_+ , studied in [29], and defined by

$$(2.17) \quad \begin{aligned} L_\phi f(x) &= G_\phi f(x) - x f'(x) \\ &= \sigma^2 x f''(x) + (m + \sigma^2 - x) f'(x) \end{aligned}$$

$$(2.18) \quad + \frac{1}{x} \int_0^\infty (d_{e^{-y}} f(x) - f(x) + y x f'(x)) \Pi(dy)$$

where G_ϕ is defined in (2.8) and (σ, β, Π) is the characteristic triplet of ϕ .

We now aim to derive the spectral properties, convergence to the equilibrium and hypercontractivity phenomenon of \mathbb{K}^ϕ .

2.4. Spectral expansion and the spectrum of the skip-free Laguerre semigroups. Since the semigroup \mathbb{K}^ϕ has invariant distribution \mathbf{n}_ϕ , we can extend it on the Hilbert space $\ell^2(\mathbf{n}_\phi)$. If ϕ is as in (2.5), let σ_1 be defined as follows

$$(2.19) \quad \sigma_1 = \begin{cases} \sigma^2 & \text{if } \sigma^2 > 0 \\ 1 & \text{if } \sigma^2 = 0. \end{cases}$$

We now introduce a sequence of discrete (acting on \mathbb{Z}_+) polynomials defined, for $k, n \in \mathbb{Z}_+$, by

$$(2.20) \quad (1 + \sigma_1^{-1})^{-\frac{k}{2}} \sum_{r=0}^k (-1)^r \binom{k}{r} \frac{\mathbf{p}_r(n)}{W_\phi(r+1)}$$

where $W_\phi(k+1) = \prod_{r=1}^k \phi(r)$, $W_\phi(1) = 1$ and $\mathbf{p}_r(n) = \frac{\Gamma(n+1)}{\Gamma(n+1-r)}$. Since the invariant distribution \mathbf{n}_ϕ has finite moments of all order, see Theorem 2.8(3), it is plain that, for all $k \in \mathbb{Z}_+$, $\mathbf{P}_k^\phi \in \ell^2(\mathbf{n}_\phi)$. Next, for $k, n \in \mathbb{Z}_+$, we define

$$(2.21) \quad \mathbf{V}_k^\phi(n) = \frac{(1 + \sigma_1^{-1})^{\frac{k}{2}}}{\mathbf{n}_\phi(n)} \sum_{r=0}^{k \wedge n} (-1)^r \frac{(k+n-r)!}{(k-r)!(n-r)!r!} \mathbf{n}_\phi(k+n-r).$$

Theorem 2.10. (1) **Spectrum.** For any $\phi \in \mathbf{B}, t \geq 0$ and $k \in \mathbb{Z}_+$, $\mathbf{V}_k^\phi \in \ell^2(\mathbf{n}_\phi)$, and

$$\mathbb{K}_t^\phi \mathbf{P}_k^\phi = e^{-kt} \mathbf{P}_k^\phi, \quad \widehat{\mathbb{K}}_t^\phi \mathbf{V}_k^\phi = e^{-kt} \mathbf{V}_k^\phi$$

where $\widehat{\mathbb{K}}_t^\phi$ is the $\ell^2(\mathbf{n}_\phi)$ -adjoint of \mathbb{K}_t^ϕ . Hence, $\{e^{-kt}; k \in \mathbb{Z}_+\} \subseteq \text{Spec}_p(\mathbb{K}_t^\phi) \cap \text{Spec}_p(\widehat{\mathbb{K}}_t^\phi)$, where for an operator T , $\text{Spec}_p(T)$ denotes the point spectrum of T .

(2) **Biorthogonality.** $(\mathbf{P}_k^\phi)_{k \geq 0}$ and $(\mathbf{V}_k^\phi)_{k \geq 0}$ are biorthogonal sequences in $\ell^2(\mathbf{n}_\phi)$, that is, for all $k, l \in \mathbb{Z}_+$,

$$\left\langle \mathbf{P}_k^\phi, \mathbf{V}_l^\phi \right\rangle_{\mathbf{n}_\phi} = \mathbb{1}_{\{k=l\}}.$$

(3) **Spectral expansion.** If $\sigma^2 > 0$, then, for all $f \in \ell^2(\mathbf{n}_\phi)$ and $t > \frac{1}{2} \log(1 + \sigma^{-2})$,

$$(2.22) \quad \mathbb{K}_t^\phi f = \sum_{k=0}^{\infty} e^{-kt} \left\langle f, \mathbf{V}_k^\phi \right\rangle_{\mathbf{n}_\phi} \mathbf{P}_k^\phi.$$

(4) **Compactness.** If $\sigma^2 > 0$, then, for all $t > \frac{1}{2} \log(1 + \sigma^{-2})$, \mathbb{K}_t^ϕ is compact and, denoting by $\text{Spec}(\mathbb{K}_t^\phi)$ the spectrum of \mathbb{K}_t^ϕ , we have

$$\text{Spec}(\mathbb{K}_t^\phi) \setminus \{0\} = \text{Spec}_p(\mathbb{K}_t^\phi) = \{e^{-kt}; k \in \mathbb{Z}_+\}.$$

(5) **Transition probabilities.** If $(\mathbb{K}_t^\phi(\cdot, \cdot))_{t \geq 0}$ denotes the transition probabilities of the skip-free Laguerre chain and $\sigma^2 > 0$, then, for all $t > \frac{1}{2} \log(1 + \sigma^{-2})$ and $n, l \in \mathbb{Z}_+$, we have

$$\mathbb{K}_t^\phi(n, l) = \sum_{k=0}^{\infty} e^{-kt} \mathbf{P}_k^\phi(n) \mathbf{V}_k^\phi(l)_{\mathbf{n}_\phi(l)}$$

where the sum on the right-hand side of the above identity converges absolutely.

This theorem is proved in Section 4.12.

Remark 2.11. It should be noted that (1) in the above theorem is different from the result in the case of generalized Laguerre semigroups on \mathbb{R}_+ , their continuous analogue. Indeed, from [29, Theorem 1.22, 4(d)], $e^{-kt} \in \text{Spec}_p(\widehat{K}_t^\phi)$ only if $k \in \mathbb{Z}_\phi$ (see (4.57) for the definition of \mathbb{Z}_ϕ) and $e^{-kt} \in \text{Spec}_r(\widehat{K}_t^\phi)$ if $k \notin \mathbb{Z}_\phi$, where $\text{Spec}_r(\mathbb{K}_t^\phi)$ stands for the residual spectrum of \mathbb{K}_t^ϕ . However, for the discrete Laguerre semigroup \mathbb{K}^ϕ , $e^{-kt} \in \text{Spec}_p(\widehat{\mathbb{K}}_t^\phi)$ for all $k \in \mathbb{Z}_+$.

2.5. Convergence to equilibrium. In Theorem 2.8(3) we have seen that the non-self-adjoint skip-free Laguerre chains have an unique invariant distribution. In this section, we start by studying the rate of convergence to their invariant distributions via spectral gap inequality, which comes as a by-product of the spectral expansion obtained in the previous theorem. We proceed with explicit rate of convergence to equilibrium in the Φ -entropy sense, which is a consequence of a more subtle relation with the self-adjoint birth-death Laguerre chain, namely an interweaving relation discussed in Section 4.14. Before stating the result, let us introduce a few additional objects related to the Bernstein functions. For any $\phi \in \mathbf{B}$ let us define

$$(2.23) \quad d_\phi = \min\{u \geq 0; \phi(-u) = -\infty, \phi(-u) = 0\} \in [0, \infty].$$

If (m, σ^2, Π) is the triplet associated to ϕ , let us write

$$(2.24) \quad m_\phi = \lim_{u \rightarrow \infty} \frac{\phi(u) - \sigma^2 u}{\sigma^2} = \frac{m + \overline{\overline{\Pi}}(0)}{\sigma^2}$$

where $\overline{\overline{\Pi}}(0) = \int_0^\infty \Pi(y, \infty) dy$. The quantity m_ϕ is finite whenever $\sigma^2 > 0$ and $\overline{\overline{\Pi}}(0) \in [0, \infty)$.

Next, for an open interval $I \subseteq \mathbb{R}$, we say that a function $\Phi : I \rightarrow \mathbb{R}$ is **admissible** if

$$(2.25) \quad \Phi \in \mathbf{C}^4(I) \text{ with both } \Phi \text{ and } -\frac{1}{\Phi''} \text{ convex.}$$

Given an admissible function Φ , and a probability measure μ on \mathbb{R} , we write for any $f : \mathbb{R}_+ \rightarrow I$ with $f, \Phi(f) \in \mathbf{L}^1(\mu)$

$$(2.26) \quad \text{Ent}_\mu^\Phi(f) = \mu\Phi(f) - \Phi(\mu f)$$

for the so-called Φ -entropy of f . When $\Phi(x) = x^2, I = \mathbb{R}$, (2.26) is equal to $\text{Var}_\mu(f)$ and when $\Phi(x) = x \log x, I = \mathbb{R}_+$, (2.26) yields the Boltzmann entropy of f with respect to μ . From Jensen's inequality it is plain that the Φ -entropy is always nonnegative. We are now ready to state the following.

Theorem 2.12. *Let $\phi \in \mathbf{B}$ be associated with the triplet (m, σ^2, Π) such that $\sigma^2, d_\phi > 0$ and $\overline{\overline{\Pi}}(0) < \infty$. Then, the following holds.*

(1) **Hypo coercive estimate.** *For all $f \in \ell^2(\mathbf{n}_\phi)$ and $t \geq 0$, we have*

$$(2.27) \quad \left\| \mathbb{K}_t^\phi f - \mathbf{n}_\phi f \right\|_{\ell^2(\mathbf{n}_\phi)} \leq \sqrt{\frac{(m_\phi + 1)(1 + \sigma^2)}{\sigma^2(d_\phi + 1)}} e^{-t} \|f - \mathbf{n}_\phi f\|_{\ell^2(\mathbf{n}_\phi)}.$$

(2) **Entropy decay.** For all $\beta > m_\phi$, $t \geq 0$ and f such that $f, \Phi(f) \in \ell^1(\mathbf{n}_\phi)$, we have

$$(2.28) \quad \text{Ent}_{\mathbf{n}_\phi}^\Phi \left(\mathbb{K}_{t+\tau_\beta}^\phi f \right) \leq e^{-t} \text{Ent}_{\mathbf{n}_\phi}^\Phi (f)$$

where, we recall that $\mathbb{K}_{t+\tau_\beta}^\phi f(n) = \mathbb{E}[f(\mathbb{X}_\phi(t + \tau_\beta, n))]$ and τ_β is an infinitely divisible positive random variable whose Laplace transform is given by

$$(2.29) \quad \int_0^\infty e^{-us} \mathbb{P}(\tau_\beta \in ds) = e^{-\phi_\beta(u)}, \quad u > 0,$$

with $\phi_\beta(u) = u \log(1 + \sigma^{-2}) + \log\left(\frac{\Gamma(u+\beta+1)}{\Gamma(1+\beta)\Gamma(u+1)}\right)$.

Item (1) of the above theorem is proved in Section 4.13 and item (2) is proved in Section 4.16.

Remark 2.13. The estimate in (1) gives the hypocoercivity, in the sense of Villani [36], for the skip-free Laguerre semigroups. This notion continues to attract a lot of interests, especially in the area of kinetic Fokker–Planck equations; see e.g. Baudoin [6] and Dolbeault et al. [15] and the references therein. Unlike this literature, we are able to identify the hypocoercive constants, namely the exponential decay rate as the spectral gap and the constant in front of the exponential, which is greater than 1 as with $\sigma^2, d_\phi > 0$ we have $m_\phi > d_\phi$, is a measure of the deviation of the spectral projections from forming an orthogonal basis. Note that in general, the hypocoercive constants may be difficult to identify and may have little to do with the spectrum. Results in the spirit of (1) have already been obtained by Achleitner et al. [1], Patie and Savov [29] as well as in Patie and Vaidyanathan [31] where a general framework based on intertwining relation is developed.

2.6. Hypercontractivity. A Markov semigroup defined on the state space E with invariant distribution μ is said to be hypercontractive if there exists $\alpha > 0$ such that

$$\|P_t\|_{\mathbf{L}^2(E, \mu) \rightarrow \mathbf{L}^{p(\alpha t)}(E, \mu)} \leq 1$$

where $p(t) = 1 + e^t$ and

$$\|P_t\|_{\mathbf{L}^2(E, \mu) \rightarrow \mathbf{L}^{p(\alpha t)}(E, \mu)} = \sup_{f: \|f\|_{\mathbf{L}^2(E, \mu)} = 1} \|P_t f\|_{\mathbf{L}^{p(\alpha t)}(E, \mu)}.$$

It is readily seen that the hypercontractivity reflects the regularity of the semigroup. For self-adjoint Markov semigroups, hypercontractivity can be interpreted in terms of their (modified) log-Sobolev constants, see [5, Theorem 5.2.3] and references therein. Nonetheless, even for the self-adjoint

birth-death Laguerre chain, it is difficult to obtain a precise value of the (modified) log-Sobolev constant. Using the concept of interweaving, see Section 4.14, we circumvent this issue, and in fact, we are able to obtain the hypercontractivity estimates for (non self-adjoint) skip-free Laguerre semigroups up to a random warm-up time.

Theorem 2.14. *If $\sigma^2 > 0$ and $\overline{\Pi}(0) < \infty$, then, for all $\beta > m_\phi = \frac{m + \overline{\Pi}(0)}{\sigma^2}$ and $t \geq 0$,*

$$\left\| \mathbb{K}_{t+\tau_\beta}^\phi \right\|_{\ell^2(\mathbf{n}_\phi) \rightarrow \ell^{p(t)}(\mathbf{n}_\phi)} \leq 1$$

where τ_β is defined in (2.29).

This theorem is proved in Section 4.17.

2.7. Bochner subordination of skip-free Laguerre chains. In the previous two sections we have seen that Theorem 2 and Theorem 2.14 hold for skip-free Laguerre semigroups up to a random warm-up or delay time denoted by τ_β . However, applying a time-change on the skip-free Laguerre chains, we can obtain a new class of skip-free Markov chains for which the above theorems hold with a deterministic warm-up or delay time. In other words, we obtain a new class of Markov chains (with two sided-jumps of arbitrary size) for which the quantity τ_β can be replaced by a deterministic number. Since τ_β is an infinitely divisible random variable with $\phi_\beta \in \mathbf{B}$ as its Lévy-Khintchine exponent, one can consider the subordinator $(\tau_\beta(t), t \geq 0)$ such that $\tau_\beta(1) \stackrel{(d)}{=} \tau_\beta$. With an abuse of notation, we still denote this subordinator by τ_β . Now, let us consider the subordinated Laguerre semigroup defined by

$$(2.30) \quad \mathbb{K}_t^{\phi, \tau_\beta} = \int_0^\infty \mathbb{K}_s^\phi \mathbb{P}(\tau_\beta(t) \in ds).$$

Since $\lim_{t \rightarrow \infty} \tau_\beta(t) = \infty$ almost surely, the semigroup $\mathbb{K}^{\phi, \tau_\beta}$ has the same invariant measure \mathbf{n}_ϕ . Below, we provide the spectral expansion, the Φ -entropy convergence and the hypercontractivity property of $\mathbb{K}^{\phi, \tau_\beta}$.

Theorem 2.15. *Let $\phi \in \mathbf{B}$ be associated with the triplet (m, σ^2, Π) .*

(1) **Spectral Expansion.** *If $\sigma^2 > 0$ then for all $\beta > 0$, $\mathbf{f} \in \ell^2(\mathbf{n}_\phi)$ and $t > \frac{1}{2}$ we have*

$$\mathbb{K}_t^{\phi, \tau_\beta} \mathbf{f} = \sum_{k=0}^{\infty} e^{-t\phi_\beta(k)} \langle \mathbf{f}, \mathbf{V}_k^\phi \rangle_{\mathbf{n}_\phi} \mathbf{P}_k^\phi.$$

- (2) **Φ -entropy decay.** If $\sigma^2 > 0$, $\overline{\overline{\Pi}}(0) < \infty$ and $\beta > m_\phi$, then for all admissible (see (2.25) for definition) function Φ and \mathbf{f} such that $\mathbf{f}, \Phi(\mathbf{f}) \in \ell^1(\mathbf{n}_\phi)$, we have, for all $t \geq 0$,

$$\text{Ent}_{\mathbf{n}_\phi}^\Phi \left(\mathbb{K}_t^{\phi, \tau_\beta} \mathbf{f} \right) \leq e^{-\phi_\beta(1)(t-1)_+} \text{Ent}_{\mathbf{n}_\phi}^\Phi(\mathbf{f})$$

where $t_+ = \max(t, 0)$.

- (3) **Hypercontractivity.** If $\sigma^2 > 0$, $\overline{\overline{\Pi}}(0) < \infty$ and $\beta > m_\phi$, then, for all $t \geq 0$,

$$\left\| \mathbb{K}_{t+1}^{\phi, \tau_\beta} \right\|_{\ell^2(\mathbf{n}_\phi) \rightarrow \ell^{p(t)}(\mathbf{n}_\phi)} \leq 1$$

where $p(t) = 1 + e^t$.

3. EXAMPLES

3.1. The discrete Laguerre chains and the Meixner polynomials.

Let us consider the Bernstein function $\phi(u) = \sigma^2 u + m$ where $\sigma^2 > 0, m \geq 0$. If \mathbb{K}^ϕ is the skip-free Laguerre semigroup associated with ϕ , then the generator is given by

$$\mathbb{L}_\phi(n, l) \begin{cases} \sigma^2 n + m + \sigma^2 + 1 & \text{if } l = n + 1 \\ (1 + \sigma^2)n & \text{if } l = n - 1 \\ -(1 + 2\sigma^2)n - m - \sigma^2 - 1 & \text{if } l = n \\ 0 & \text{otherwise.} \end{cases}$$

From Proposition 4.12, the unique invariant distribution of \mathbb{L}_ϕ is given by

$$\mathbf{n}_\phi(n) = \frac{\Gamma(n + \frac{m}{\sigma^2} + 1)}{\Gamma(\frac{m}{\sigma^2} + 1) n!} 2^{-n - \frac{m}{\sigma^2} - 1}, \quad n \in \mathbb{Z}_+.$$

The semigroup \mathbb{K}^ϕ is self-adjoint in $\ell^2(\mathbf{n}_\phi)$ and it follows from Theorem 2.10 that the eigenfunctions \mathbf{P}_k^ϕ of \mathbb{K}_t^ϕ corresponding to its eigenvalue e^{-kt} form an orthogonal sequence in $\ell^2(\mathbf{n}_\phi)$. More specifically, writing $\beta = \frac{m}{\sigma^2}$, for all $k \in \mathbb{Z}_+$,

$$\begin{aligned} \mathbf{P}_k^\phi(n) &= (1 + \sigma^{-2})^{-\frac{k}{2}} \Gamma(\beta + 1) \sum_{r=0}^k (-\sigma)^{-2r} \binom{k}{r} \frac{\mathbf{p}_r(n)}{\Gamma(r + \beta + 1)} \\ &= (1 + \sigma^{-2})^{-\frac{k}{2}} {}_2F_1(-n, -k, \beta + 1; -\sigma^{-2}) \end{aligned}$$

where

$$(3.1) \quad {}_2F_1(a, b; c; x) = \frac{\Gamma(c)}{\Gamma(a)\Gamma(b)} \sum_{r=0}^{\infty} \frac{\Gamma(r+a)\Gamma(r+b)}{\Gamma(r+c)} \frac{x^r}{r!}.$$

From [21, Equation (7)], it follows that

$$\left\| \mathbf{P}_k^\phi \right\|_{\ell^2(\mathbf{n}_\phi)}^2 = \mathbf{c}_k(\beta)^{-1}$$

where for any $a > 0$, $\mathbf{c}_k(a) = \frac{\Gamma(a+k+1)}{\Gamma(a+1)\Gamma(k+1)}$. Finally, for all $\mathbf{f} \in \ell^2(\mathbf{n}_\phi)$ and $t > 0$, we have, in $\ell^2(\mathbf{n}_\phi)$,

$$\mathbb{K}_t^{\phi} \mathbf{f} = \sum_{k=0}^{\infty} \mathbf{c}_k \left(\frac{m}{\sigma^2} \right) e^{-kt} \left\langle \mathbf{f}, \mathbf{P}_k^\phi \right\rangle_{\mathbf{n}_\phi} \mathbf{P}_k^\phi.$$

3.2. The perturbed Laguerre skip-free chain. Consider the Bernstein function defined for $\mathbf{m} > 1$ by

$$\phi_{\mathbf{m}}(u) = \frac{(u + \mathbf{m} + 1)(u + \mathbf{m} - 1)}{u + \mathbf{m}} = \frac{\mathbf{m}^2 - 1}{\mathbf{m}} + u + \int_0^\infty (1 - e^{-uy}) e^{-my} dy.$$

Let $\mathbb{G}_{\phi_{\mathbf{m}}}$ be the generator of the discrete self-similar Markov semigroup associated with $\phi_{\mathbf{m}}$. Then, according to (2.6), $\sigma^2 = 1$, $m = \frac{\mathbf{m}^2 - 1}{\mathbf{m}}$ and $\Pi(dy) = \mathbf{m} e^{-my} dy$. So, the infinitesimal generator $\mathbb{G}_{\mathbf{m}}$ is given by

$$\mathbb{G}_{\phi_{\mathbf{m}}}(n, l) = \begin{cases} \frac{\mathbf{m}\Gamma(l+\mathbf{m})\Gamma(n-l+2)}{(n+1)\Gamma(n+\mathbf{m}+2)} & \text{if } l \in \llbracket 0, n-2 \rrbracket \\ \frac{2\mathbf{m}}{(n+1)(n+\mathbf{m})(n+\mathbf{m}+1)} + n & \text{if } l = n-1 \\ \mathbf{m} - \frac{1}{\mathbf{m}+n+1} & \text{if } l = n+1 \\ \frac{\mathbf{m}}{(n+\mathbf{m})(n+\mathbf{m}+1)} - \frac{1}{\mathbf{m}} & \text{if } l = n \\ 0 & \text{if } l > n+1. \end{cases}$$

Now, the corresponding skip-free Laguerre chain has the generator $\mathbb{L}_{\phi_{\mathbf{m}}}$ given by

$$\mathbb{L}_{\phi_{\mathbf{m}}}(n, l) = \begin{cases} \mathbb{G}_{\mathbf{m}}(n, l) & \text{if } l \neq n, n-1 \\ \mathbb{G}_{\mathbf{m}}(n, n-1) + n & \text{if } l = n-1 \\ \mathbb{G}_{\mathbf{m}}(n, n) - n & \text{if } l = n. \end{cases}$$

From Proposition 4.12, the unique invariant distribution of $\mathbb{L}^{\phi_{\mathbf{m}}}$ is given by

$$\mathbf{n}_{\phi_{\mathbf{m}}}(n) = \frac{(n + \mathbf{m} + 1)\Gamma(n + \mathbf{m})}{(\mathbf{m} + 1)\Gamma(\mathbf{m})n!} 2^{-(n+\mathbf{m}+1)}, \quad n \in \mathbb{Z}_+.$$

Let us compute the eigenfunctions and co-eigenfunctions of the semigroup $\mathbb{K}^{\phi_{\mathbf{m}}}$ generated by $\mathbb{L}^{\phi_{\mathbf{m}}}$. Denoting the eigenfunction (resp. the co-eigenfunction) of $\mathbb{K}_t^{\phi_{\mathbf{m}}}$ corresponding to the eigenvalue (resp. co-eigenvalue) e^{-kt} by $\mathbf{P}_k^{\phi_{\mathbf{m}}}$

(resp. $V_k^{\phi_m}$), we have

$$\begin{aligned}
P_k^{\phi_m}(n) &= 2^{-\frac{k}{2}} \sum_{r=0}^k (-1)^r \frac{\binom{k}{r}}{W_{\phi_m}(r+1)} p_r(n) \\
&= 2^{-\frac{k}{2}} [(\mathbf{m}+1) {}_2F_1(-k, -n, \mathbf{m}+1; -1) - {}_2F_1(-k, -n, \mathbf{m}+2; -1)], \\
V_k^{\phi_m}(n) &= \frac{2^{-\frac{k}{2}} \Gamma(n+k+\mathbf{m})}{\mathbf{n}_\phi(n) k! n!} ((n+\mathbf{m}+k) {}_2F_1(-k, -n, -n-k-\mathbf{m}; 2) \\
&\quad + {}_2F_1(-k, -n, -n-k-\mathbf{m}+1; 2))
\end{aligned}$$

where ${}_2F_1$ is the hypergeometric function defined in (3.1).

3.3. The Beta skip-free chain. We consider the Bernstein function ϕ_m corresponding to a compound Poisson process with exponential jumps which is defined, for $\mathbf{m} > 1$ and $u > 0$, by

$$\phi_m(u) = \frac{u}{\mathbf{m}(u+\mathbf{m})} = \int_0^\infty (1 - e^{-uy}) e^{-\mathbf{m}y} dy.$$

Therefore, according to (2.5), $\sigma^2 = 0$, $m = 0$, $\Pi(dy) = \mathbf{m}e^{-\mathbf{m}y} dy$ and $\phi_m(\infty) = \frac{1}{\mathbf{m}}$. If L_{ϕ_m} denotes the generator of the Laguerre semigroup corresponding to ϕ_m in continuous state space, we have for all $f \in C_c^\infty(\mathbb{R}_+)$,

$$L_{\phi_m} f(x) = -x f'(x) + \frac{\mathbf{m}}{x} \int_0^\infty (f(e^{-y}x) - f(x) + yx f'(x)) e^{-\mathbf{m}y} dy.$$

The Bernstein-gamma function associated with ϕ_m is

$$W_{\phi_m}(k+1) = \frac{\Gamma(\mathbf{m}+1)\Gamma(k+1)}{\mathbf{m}^k \Gamma(k+1+\mathbf{m})}, \quad k \in \mathbb{Z}_+.$$

From [29, Proposition 2.6(1)], the semigroup generated by L_{ϕ_m} admits an unique invariant measure ν_{ϕ_m} which is absolutely continuous with moment sequence $(W_{\phi_m}(k+1))_{k \geq 0}$ and given by

$$\nu_{\phi_m}(dx) = \mathbf{m}^2 (1 - \mathbf{m}x)^{\mathbf{m}-1} dx, \quad 0 < x < \frac{1}{\mathbf{m}}.$$

Now coming back to the corresponding skip-free Laguerre chain in the discrete state space, (4.39) implies that the unique invariant distribution of its

semigroup \mathbb{K}^{ϕ_m} is

$$\begin{aligned} \mathbf{n}_{\phi_m}(n) &= \frac{1}{n!} \sum_{r=0}^{\infty} W_{\phi_m}(n+r+1) \frac{(-1)^r}{r!} \\ &= \frac{1}{n!} \sum_{r=0}^{\infty} \frac{\Gamma(m+1)\Gamma(n+r+1)}{m^{n+r}\Gamma(n+r+m+1)} \frac{(-1)^r}{r!} \\ &= \frac{1}{m^n} \frac{\Gamma(m+1)}{\Gamma(n+m+1)} {}_1F_1\left(n, n+m; \frac{1}{m}\right) \end{aligned}$$

where ${}_1F_1$ is an hypergeometric function. Finally, from Proposition 4.17, the eigenfunction of $\mathbb{K}_t^{\phi_m}$ corresponding to e^{-kt} is given by

$$\mathbf{P}_k^{\phi_m}(n) = 2^{-\frac{k}{2}} \sum_{r=0}^k (-1)^r \binom{k}{r} \frac{\mathbf{p}_r(n)}{W_{\phi_m}(r+1)} = 2^{\frac{k}{2}} {}_3F_1(-k, -n, m+1; 1; -m)$$

where ${}_3F_1(a, b, c; d; x) = \frac{\Gamma(d)}{\Gamma(a)\Gamma(b)\Gamma(c)} \sum_{r=0}^{\infty} \frac{\Gamma(r+a)\Gamma(r+b)\Gamma(r+c)}{\Gamma(r+d)} \frac{x^r}{r!}$ and $\mathbf{V}_k^{\phi_m}$ is given by (2.21).

4. PROOF OF THE MAIN RESULTS

We begin this section with some useful facts about to the discrete dilation operator.

Proposition 4.1. (1) For all $\alpha > 0$ and $\mathbf{f} \in \mathbf{C}_b(\mathbb{Z}_+)$,

$$(4.1) \quad d_{\alpha}\Lambda\mathbf{f} = \Lambda\mathbb{D}_{\alpha}\mathbf{f}$$

where $d_{\alpha}f(x) = f(\alpha x)$ is the usual dilation operator on \mathbb{R}_+ and Λ is as in Theorem 2.6.

(2) For all $\mathbf{f} \in \mathbf{C}(\mathbb{Z}_+)$, and $\alpha, \beta > 0$, $\mathbb{D}_{\alpha\beta}\mathbf{f} = \mathbb{D}_{\alpha}\mathbb{D}_{\beta}\mathbf{f}$.

(3) $\mathbb{D}_1 = \text{Id}$ and for all $\alpha > 0$, $\mathbb{D}_{\alpha}^{-1} = \mathbb{D}_{1/\alpha}$.

(4) $(\mathbb{D}_{e^{-t}})_{t \geq 0}$ (resp. $(d_{e^{-t}})_{t \geq 0}$) form a semigroup on $\ell^2(\mathbb{Z}_+)$ (resp. on $\mathbf{L}^2(\mathbb{R}_+)$) with generator $\partial_-^n = n\partial_-$ (resp. $\partial^x = -x\frac{d}{dx}$) with $\partial^x\Lambda = \Lambda\partial_-^n$ on $\mathbf{C}_c(\mathbb{Z}_+)$.

Remark 4.2. There are several analogies between the two dilation operators which make our choice of the discrete one natural. Indeed, as its continuous analogue, the discrete dilation operator is a multiplicative semigroup, and, the generator of its associated additive semigroup is the discrete analogue of the continuous one, see item 4. However, unlike in the continuous case, \mathbb{D}_{α} does not have bounded inverse when $\alpha \in (0, 1)$.

Proof. The first item follows from [26, Proposition 1]. Next, we note that, for any $f \in \mathbf{C}_b(\mathbb{Z}_+)$ and $\alpha > 0$,

$$(4.2) \quad |\mathbb{D}_\alpha f(n)| \leq |(2\alpha - 1)|^n \|f\|_\infty.$$

Then, (4.2) implies that both $\Lambda \mathbb{D}_\alpha f$ and $d_\alpha \Lambda f$ are well defined. Now, (1) yields

$$\Lambda \mathbb{D}_{\alpha\beta} = d_{\alpha\beta} \Lambda = d_\alpha d_\beta \Lambda = d_\alpha \Lambda \mathbb{D}_\beta = \Lambda \mathbb{D}_\alpha \mathbb{D}_\beta.$$

Since $\Lambda : \mathbf{C}_b(\mathbb{Z}_+) \rightarrow \mathbf{C}_b(\mathbb{R}_+)$ is injective, see [26, Lemma 4(4)], item (2) follows. Item (3) is a direct consequence of item (2). For item (4), it is immediate from item (2) that $(\mathbb{D}_{e^{-t}})_{t \geq 0}$ is a translation semigroup on $\mathbf{C}_0(\mathbb{Z}_+)$. Moreover, from (4.1) we have that for all $t \geq 0$ and $f \in \mathbf{C}_0(\mathbb{Z}_+)$

$$d_{e^{-t}} \Lambda f = \Lambda \mathbb{D}_{e^{-t}} f.$$

Differentiating the above identity with respect to t when $f \in \mathbf{C}_c(\mathbb{Z}_+)$ and noting that $d_{e^{-t}} = e^{t\partial^x}$ one obtains that

$$\partial^x \Lambda f = \Lambda \frac{d}{dt} \mathbb{D}_{e^{-t}} f|_{t=0}$$

where we used that Λ is a bounded operator. However, from [26, Lemma 4.5], after observing that $\Lambda = \nabla^{-1}$, we have $\partial^x \Lambda f = \Lambda \partial^n f$ whenever $f \in \mathbf{C}_c(\mathbb{Z}_+)$. Since Λ is injective on $\mathbf{C}_c(\mathbb{Z}_+)$, we conclude that for all $f \in \mathbf{C}_c(\mathbb{Z}_+)$ one has

$$\frac{d}{dt} \mathbb{D}_{e^{-t}} f|_{t=0} = \partial^n f$$

which proves item (4). \square

4.1. Proof of Theorem 2.2. It is not difficult to see that the operator \mathbb{G}_Π can be simplified as follows. We can write $\mathbb{G}_\Pi f(n) = \sum_{l=0}^{n+1} \mathbb{G}(n, l) f(l)$ where

$$(4.3) \quad \mathbb{G}_\Pi(n, l) = \begin{cases} \int_0^\infty \frac{1}{n+1} \binom{n+1}{l} e^{-ly} (1 - e^{-y})^{n-l+1} \Pi(dy) & \text{if } l \in \llbracket 0, n-1 \rrbracket \\ \int_0^\infty \frac{1}{n+1} (e^{-(n+1)y} - 1 + (n+1)y) \Pi(dy) & \text{if } l = n+1 \\ 0 & \text{if } l > n+1 \end{cases}$$

and $\mathbb{G}_\Pi(n, n) = -\sum_{l \neq n} \mathbb{G}_\Pi(n, l)$.

To show that \mathbb{G}_ϕ is a Markov generator, we need to show that $\mathbb{G}_\phi(n, l) \geq 0$ for all $n \neq l$. From the expression of \mathbb{G}_ϕ in (2.6), it is enough to show that $\mathbb{G}_\Pi(n, l) \geq 0$ for all $l \neq n$, a fact which follows readily from (4.3). To get that \mathbb{G}_ϕ generates a Feller semigroup on $\mathbf{C}_0(\mathbb{Z}_+)$, we wish to combine Theorem 3.2 with Corollary 3.2 from [17, Chapter 8]. To this end, the following four conditions need to be checked

- (i) $\sup_{n \in \mathbb{Z}_+} \frac{|\mathbf{G}_\phi(n, n)|}{n+1} < \infty$
- (ii) $\lim_{n \rightarrow \infty} \mathbf{G}_\phi(n, l) = 0$ for all $l \in \mathbb{Z}_+$
- (iii) $\sup_{n \in \mathbb{Z}_+} \sum_{l \in \mathbb{Z}_+} \frac{n+1}{l+1} \mathbf{G}_\phi(n, l) < \infty$
- (iv) $\sup_{n \in \mathbb{Z}_+} \frac{1}{n+1} \sum_{l \in \mathbb{Z}_+} (l-n) \mathbf{G}_\phi(n, l) < \infty$.

First, we note that

$$\mathbf{G}_\phi(n, l) = \begin{cases} \sigma^2(n+1) + m + \mathbf{G}_\Pi(n, n+1) & \text{if } l = n+1 \\ \sigma^2 n + \mathbf{G}_\Pi(n, n-1) & \text{if } l = n-1 \\ -2\sigma^2 n - \sigma^2 - m + \mathbf{G}_\Pi(n, n) & \text{if } l = n \\ \mathbf{G}_\Pi(n, l) & \text{otherwise.} \end{cases}$$

It is plainly sufficient to check all four conditions above for \mathbf{G}_Π merely. From the definition of $\mathbf{G}_\Pi(n, n)$, we get that, for all $n \in \mathbb{Z}_+$,

$$\begin{aligned} \mathbf{G}_\Pi(n, n) &= - \int_0^\infty \frac{1}{n+1} \left(1 - \sum_{l=0}^{n-1} \binom{n+1}{l} e^{-ly} (1 - e^{-y})^{n-l+1} \right) \Pi(dy) \\ &\quad - \int_0^\infty \frac{1}{n+1} (1 - e^{-(n+1)y} + (n+1)y) \Pi(dy). \end{aligned}$$

Since for any $y > 0$, $\sum_{l=0}^{n+1} \binom{n+1}{l} e^{-ly} (1 - e^{-y})^{n+1-l} = 1$, the above expression reduces to

$$(4.4) \quad \mathbf{G}_\Pi(n, n) = \int_0^\infty (e^{-ny} - e^{-(n+1)y} - y) \Pi(dy).$$

Next, noting that

$$\begin{aligned} |e^{-ny} - e^{-(n+1)y} - y| &= \left| \int_n^{n+1} y(1 - e^{-ry}) dr \right| \\ &\leq (2n+1)y^2 \mathbb{1}_{\{y \leq 1\}} + y \mathbb{1}_{\{y > 1\}}, \end{aligned}$$

the integral in (4.4) is finite due to (2.4) and therefore,

$$\overline{\lim}_{n \rightarrow \infty} \frac{|\mathbf{G}_\phi(n, n)|}{n+1} \leq 2\sigma^2 + 2 \int_0^1 y^2 \Pi(dy) < \infty.$$

This verifies condition (i). Then, for any $l \in \mathbb{Z}_+$ and sufficiently large n ,

$$\mathbf{G}_\Pi(n, l) = \frac{1}{n+1} \int_0^\infty \binom{n+1}{l} e^{-ly} (1 - e^{-y})^{n-l+1} \Pi(dy).$$

When $l = 0$,

$$(4.5) \quad \mathbf{G}_\Pi(n, 0) = \frac{1}{n+1} \int_0^\infty (1 - e^{-y})^{n+1} \Pi(dy)$$

and clearly $n \mapsto \int_0^\infty (1 - e^{-y})^{n+1} \Pi(dy)$ is a decreasing sequence. Thus, $\lim_{n \rightarrow \infty} \mathbf{G}_\Pi(n, 0) = 0$. When $l \geq 1$, let us define, for all $n \in \mathbb{Z}_+$ with $n \geq l + 1$,

$$a_n = \int_0^1 e^{-ly} (1 - e^{-y})^{n-l+1} \Pi(dy),$$

$$b_n = \int_1^\infty e^{-ly} (1 - e^{-y})^{n-l+1} \Pi(dy).$$

We note that both a_n, b_n are well defined if $n \geq l + 1$. Since, for all $n \geq l + 1$, $a_{n+1} \leq (1 - e^{-1})a_n$, we have that $a_n \leq a_{l+1}(1 - e^{-1})^{n-l-1}$, and thus

$$\lim_{n \rightarrow \infty} \binom{n+1}{l} a_n = 0.$$

On the other hand, observing that, for any $y > 0$,

$$\lim_{n \rightarrow \infty} \binom{n+1}{l} e^{-ly} (1 - e^{-y})^{n-l+1} = 0$$

$$\sup_{n \geq 1} \binom{n+1}{l} e^{-ly} (1 - e^{-y})^{n+1-l} \leq 1,$$

a dominated convergence argument entails that

$$\lim_{n \rightarrow \infty} \binom{n+1}{l} b_n = 0$$

which verifies condition (ii). For condition (iii), we first observe that for any $y > 0$ and $l, n \in \mathbb{Z}_+$ with $l \leq n + 1$, the following identity

$$\begin{aligned} \frac{1}{n+2} \sum_{j=1}^{n+1} (1 - e^{-y})^j - \frac{y}{n+2} &= \sum_{l=0}^{n-1} \frac{1}{l+1} \binom{n+1}{l} e^{-ly} (1 - e^{-y})^{n-l+1} \\ &\quad + \frac{(e^{-(n+1)y} - 1 + (n+1)y)}{n+2} + (e^{-ny} - e^{-(n+1)y} - y) \end{aligned}$$

holds. As a result of the above identity and invoking (4.3) one gets

$$(4.6) \quad \sum_{l=0}^{n+1} \frac{n+1}{l+1} \mathbf{G}_\Pi(n, l) = \frac{1}{n+2} \int_0^\infty (1 - e^{-y} - y) + \frac{1}{n+2} \sum_{j=2}^{n+1} (1 - e^{-y})^j \Pi(dy).$$

Since for $y > 0$, $|1 - e^{-y} - y| \leq y \wedge \frac{y^2}{2}$, using (2.4), we get

$$\int_0^\infty |1 - e^{-y} - y| \Pi(dy) \leq \int_0^1 \frac{y^2}{2} \Pi(dy) + \int_1^\infty y \Pi(dy) < \infty$$

and, for all $2 \leq j \leq n + 1$,

$$\int_0^\infty (1 - e^{-y})^j \Pi(dy) \leq \int_0^1 y^2 \Pi(dy) + \int_1^\infty \Pi(dy) < \infty.$$

Therefore, condition (iii) is satisfied as well. Finally, the last condition follows since, plainly,

$$\lim_{n \rightarrow \infty} \frac{1}{n+1} \sum_{l=0}^{n+1} (l-n) \mathbb{G}_\Pi(n, l) = \lim_{n \rightarrow \infty} \frac{1}{n+1} \int_0^\infty (e^{-y} - 1 + y) \Pi(dy) = 0.$$

Therefore, \mathbb{G}_ϕ generates a Feller semigroup on $\mathbf{C}_0(\mathbb{Z}_+)$ with $\mathbf{C}_c(\mathbb{Z}_+)$ as its core.

Next, to prove the discrete self-similarity property of the generated semigroup above, we need the following.

Proposition 4.3. (1) *Let Q^ϕ and \mathbb{Q}^ϕ denote the Feller semigroups generated by G_ϕ and \mathbb{G}_ϕ respectively. Then, for any $f \in \mathbf{C}_0(\mathbb{Z}_+)$ and for all $t \geq 0$,*

$$(4.7) \quad Q_t^\phi \Lambda f = \Lambda Q_t^\phi f$$

where we recall that $\Lambda f(x) = \mathbb{E}[f(\text{Pois}(x))]$ with $\text{Pois}(x)$ a Poisson random variable with parameter $x > 0$.

(2) *The counting measure on \mathbb{Z}_+ , denoted by \mathfrak{m} , is an excessive measure for the semigroup Q^ϕ . Hence Q^ϕ can be extended uniquely to a strongly continuous contraction semigroup on $\ell^2(\mathbb{Z}_+)$, which we again denote by Q^ϕ .*

(3) *The operator Λ can be extended uniquely to an operator (also denoted by Λ) in $\mathcal{B}(\ell^2(\mathbb{Z}_+), \mathbf{L}^2(\mathbb{R}_+))$. Keeping the same notation for the extension of Q^ϕ on $\mathbf{L}^2(\mathbb{R}_+)$, we have, for all $f \in \ell^2(\mathbb{Z}_+)$ and $t \geq 0$,*

$$(4.8) \quad Q_t^\phi \Lambda f = \Lambda Q_t^\phi f.$$

Moreover, Λ is a **quasi-affinity**, that is, it is bounded, injective and has dense range.

We split its proof into several parts.

4.1.1. Proof of Proposition 4.3(1). First, let us write

$$(4.9) \quad \begin{aligned} G_\phi &= G_{m, \sigma^2} + G_\Pi = \sigma^2 x \frac{d^2}{dx^2} + (m + \sigma^2) \frac{d}{dx} + G_\Pi \\ \mathbb{G}_\phi &= \mathbb{G}_{m, \sigma^2} + \mathbb{G}_\Pi = (\sigma^2 n + m + \sigma^2) \partial_+ + n \partial_- + \mathbb{G}_\Pi \end{aligned}$$

where, for all $f \in \mathbf{C}_b^2(\mathbb{R}_+)$,

$$G_{\Pi}f(x) = \frac{1}{x} \int_0^{\infty} (f(xe^{-y}) - f(x) + yxf'(x))\Pi(dy).$$

Let $\mathbf{P}_{\mathfrak{e}}$ be the vector space of functions defined on \mathbb{R}_+ which are of the form $e^{-x}P(x)$, P being a polynomial. We define the linear operator $\nabla : \mathbf{P}_{\mathfrak{e}} \rightarrow \mathbf{C}_c(\mathbb{Z}_+)$ as follows

$$(4.10) \quad \nabla f(n) = \frac{d^n}{dx^n}(e^x f(x))(0).$$

Lemma 4.4. *For any $f \in \mathbf{P}_{\mathfrak{e}}$,*

$$\mathbf{G}_{\phi}\nabla f = \nabla G_{\phi}f.$$

Proof. From [26, Lemma 3], it is known that, for all $f \in \mathbf{P}_{\mathfrak{e}}$,

$$\mathbf{G}_{m,\sigma^2}\nabla f = \nabla G_{m,\sigma^2}f.$$

Thus, it suffices to prove this lemma replacing $G_{\phi}, \mathbf{G}_{\phi}$ by $G_{\Pi}, \mathbf{G}_{\Pi}$ respectively. For $y > 0$, let δ_y denote the Dirac measure at y . Taking $\Pi = \delta_y$ and writing G_{δ_y} and \mathbf{G}_{δ_y} simply as G_y, \mathbf{G}_y respectively, we get

$$(4.11) \quad G_y f(x) = \frac{1}{x}(f(xe^{-y}) - f(x) + yxf'(x))$$

and

$$(4.12) \quad \mathbf{G}_y(n, l) = \begin{cases} \frac{1}{n+1} \binom{n+1}{l} (1 - e^{-ly})(1 - e^{-y})^{n-l+1} & \text{if } l \in \llbracket 0, n-1 \rrbracket \\ \frac{1}{n+1} (e^{-(n+1)y} - 1 + (n+1)y) & \text{if } l = n+1 \\ 0 & \text{if } l > n+1 \end{cases}$$

with $\mathbf{G}_y(n, n)$ being such that $\sum_{l=0}^{n+1} \mathbf{G}_y(n, l) = 0$. Then, observing that $\mathbf{G}_{\Pi}(n, l) = \int_0^{\infty} \mathbf{G}_y(n, l)\Pi(dy)$ as well as $G_{\Pi}f(x) = \int_0^{\infty} G_y f(x)dy$ for all $f \in \mathbf{C}_b^2(\mathbb{R}_+)$. We claim that it suffices to show that, for all $y > 0$ and $f \in \mathbf{P}_{\mathfrak{e}}$,

$$(4.13) \quad \mathbf{G}_y\nabla f = \nabla G_y f.$$

Indeed, when $f \in \mathbf{P}_{\mathfrak{e}}$,

$$\begin{aligned} \mathbf{G}_{\Pi}\nabla f(n) &= \sum_{l=0}^{n+1} \mathbf{G}_{\Pi}(n, l)\nabla f(l) = \sum_{l=0}^{n+1} \nabla f(l) \int_0^{\infty} \mathbf{G}_y(n, l)\Pi(dy) \\ &= \int_0^{\infty} \mathbf{G}_y\nabla f(n)\Pi(dy). \end{aligned}$$

On the other hand,

$$\nabla G_{\Pi} f(n) = \frac{d^n}{dx^n} (e^x G_{\Pi} f(x))(0) = \frac{d^n}{dx^n} \left(e^x \int_0^{\infty} G_y f(x) \Pi(dy) \right) (0).$$

Since $\mathbf{P}_{\mathfrak{e}} \subset \mathbf{C}_b^{\infty}(\mathbb{R}_+)$, the above integration and differentiation can be interchanged, therefore yielding

$$\nabla G_{\Pi} f(n) = \int_0^{\infty} \nabla G_y f(n) \Pi(dy).$$

We now proceed to show (4.13). Since $\mathbf{P}_{\mathfrak{e}} = \text{Span}\{x \mapsto e^{-x} x^l; l \in \mathbb{Z}_+\}$, it suffices to prove (4.13) only for $f(x) = h_l(x) := e^{-x} x^l$. Now, for $l \geq 1$,

$$(4.14) \quad \nabla G_y h_l(n) = \frac{d^n}{dx^n} \left[e^{x(1-xe^{-y})} x^{l-1} e^{-yl} - x^{l-1} + y(lx^{l-1} - x^l) \right] (0).$$

When $l \in \llbracket 1, n-1 \rrbracket$, applying Leibniz rule we get

$$(4.15) \quad \begin{aligned} \nabla G_y h_l(n) &= e^{-ly} \sum_{m=0}^n \binom{n}{m} \frac{d^m}{dx^m} (x^{l-1})(0) \frac{d^{n-m}}{dx^{n-m}} \left(e^{x(1-e^{-y})} \right) (0) \\ &= \binom{n}{l-1} (l-1)! (1-e^{-y})^{n-l+1} \\ &= \frac{n!}{(n-l+1)!} e^{-ly} (1-e^{-y})^{n-l+1} = l! \mathbf{G}_y(n, l). \end{aligned}$$

Also, (4.14) entails

$$(4.16) \quad \begin{aligned} \nabla G_y h_n(n) &= n! (e^{-ny} (1-e^{-y}) - y) = n! \mathbf{G}_y(n, n) \\ \nabla G_y h_{n+1}(n) &= n! (e^{-(n+1)y} - n! + (n+1)y) = (n+1)! \mathbf{G}_y(n, n+1). \end{aligned}$$

Finally,

$$(4.17) \quad \nabla G_y h_0(n) = \frac{d^n}{dx^n} \left(\frac{1}{x} (e^{x(1-e^{-y})} - 1) \right) (0) = \frac{1}{n+1} (1-e^{-y})^{n+1} = \mathbf{G}_y(n, 0).$$

On the other hand, for all $l \in \mathbb{Z}_+$, $\mathbf{G}_y \nabla h_l(n) = l! \mathbf{G}_y(n, l)$. Therefore, combining (4.15), (4.16) and (4.17), we conclude that, for all $n, l \geq 0$,

$$\mathbf{G}_y \nabla h_l(n) = \nabla G_y h_l(n).$$

This completes the proof of the lemma. \square

The next lemma is a variant of [26, Lemma 4].

Lemma 4.5. $\nabla : \mathbf{P}_{\mathfrak{e}} \rightarrow \mathbf{C}_c(\mathbb{Z}_+)$ is bijective with inverse Λ such that $\Lambda f(x) = \mathbb{E}[f(\text{Pois}(x))]$ for all $f \in \mathbf{C}_c(\mathbb{Z}_+)$. Moreover, Λ extends to a bounded operator from $\mathbf{C}_0(\mathbb{Z}_+)$ to $\mathbf{C}_0(\mathbb{R}_+)$.

We also need the following useful result.

Lemma 4.6. *For all $\phi \in \mathbf{B}$, $\mathbf{P}_\epsilon \subset \mathbf{D}(G_\phi)$ where $\mathbf{D}(G_\phi)$ is the domain of the generator of the Feller semigroup Q^ϕ .*

Proof. Denoting the Dynkin characteristic operator of the semigroup Q^ϕ by G_ϕ^D , it follows from [24, Proposition 6.1] that $\mathbf{P}_\epsilon \subset \mathbf{D}(G_\phi^D)$. Also, for all $f \in \mathbf{P}_\epsilon$, $G_\phi^D f = G_\phi f$. In light of [?, Theorem 5.5, Chapter V.3], it suffices to show that for all $f \in \mathbf{P}_\epsilon$, $G_\phi f \in \mathbf{C}_0([0, \infty))$. Since any function $f \in \mathbf{P}_\epsilon$ is of the form $f(x) = e^{-x}P(x)$ for some polynomial P , clearly $G_{m, \sigma^2} f \in \mathbf{C}_0([0, \infty))$. Now, for any $f \in \mathbf{P}_\epsilon$, we have

$$\begin{aligned} G_\Pi f(x) &= \frac{1}{x} \int_0^\infty [f(xe^{-y}) - f(x) + yxf'(x)]\Pi(dy) \\ &= \frac{1}{x} \int_0^\infty [f(xe^{-y}) - f(x) - (e^{-y} - 1)xf'(x)]\Pi(dy) \\ &\quad + f'(x) \int_0^\infty (e^{-y} - 1 + y)\Pi(dy) \\ &= A_1(x) + A_2(x). \end{aligned}$$

Since $f \in \mathbf{P}_\epsilon$, it implies that $f' \in \mathbf{C}_0([0, \infty))$ and therefore $A_2(x) \rightarrow 0$ as $x \rightarrow \infty$. To deal with A_1 , by means of Taylor expansion of f up to order 2 we obtain that

$$(4.18) \quad \begin{aligned} &\frac{1}{x} \int_0^1 |f(xe^{-y}) - f(x) - (e^{-y} - 1)xf'(x)| \Pi(dy) \\ &\leq \frac{x}{2} \int_0^1 \sup_{t \in [e^{-y}x, x]} |f''(t)|(1 - e^{-y})^2 \Pi(dy). \end{aligned}$$

We note that any function $f \in \mathbf{P}_\epsilon$, f is either eventually increasing or decreasing, depending on the sign of the leading coefficient of the polynomial associated with the function. Without loss of generality, we assume that f is eventually decreasing. Since for all $y \in [0, 1]$, $e^{-y} \geq e^{-1}$, for all large values of x , we have $\sup_{t \in [e^{-y}x, x]} |f''(t)| \leq |f''(e^{-1}x)|$, as $f \in \mathbf{P}_\epsilon$ implies $f'' \in \mathbf{P}_\epsilon$. Thus, the right-hand side of (4.18) goes to 0 as $x \rightarrow \infty$. On the other hand,

$$\frac{1}{x} |f(xe^{-y}) - f(x) - (e^{-y} - 1)xf'(x)| \leq (1 - e^{-y}) \|f'\|_\infty \leq y \|f'\|_\infty.$$

Using the dominated convergence theorem, we obtain that

$$\lim_{x \rightarrow \infty} \frac{1}{x} \int_1^\infty [f(e^{-y}x) - f(x) + (e^{-y} - 1)xf'(x)]\Pi(dy) = 0.$$

This shows that $A_2(x) \rightarrow 0$ as $x \rightarrow \infty$ which completes the proof of the lemma. \square

Let $\mathbf{D}(\mathbf{G}_\phi)$ denote the domain of the Feller generator \mathbf{G}_ϕ . Now, coming back to the proofs of Proposition 4.3, Lemma 4.4, Lemma 4.5 and Lemma 4.6 imply that, for all $f \in \mathbf{C}_c(\mathbb{Z}_+)$,

$$(4.19) \quad \Lambda f \in \mathbf{P}_t \subset \mathbf{D}(G_\phi) \text{ and } G_\phi \Lambda f = \Lambda G_\phi f.$$

Since $\mathbf{C}_c(\mathbb{Z}_+)$ is a core for the generator \mathbf{G}_ϕ , for any $f \in \mathbf{D}(\mathbf{G}_\phi)$ there exists a sequence $\{f_n\} \subset \mathbf{C}_c(\mathbb{Z}_+)$ such that $\|f_n - f\|_\infty \rightarrow 0$ and $\|\mathbf{G}_\phi f_n - \mathbf{G}_\phi f\|_\infty \rightarrow 0$. Therefore, thanks to Lemma 4.5,

$$\|\Lambda f_n - \Lambda f\|_\infty \rightarrow 0, \quad \|\Lambda G_\phi f_n - \Lambda G_\phi f\|_\infty \rightarrow 0.$$

Thus, (4.19) entails that $G_\phi \Lambda f_n$ converges in $\mathbf{C}_0(\mathbb{R}_+)$ which implies that $\Lambda f \in \mathbf{D}(G_\phi)$ as $(G_\phi, \mathbf{D}(G_\phi))$ is a closed operator, and, for all $f \in \mathbf{D}(\mathbf{G}_\phi)$,

$$(4.20) \quad G_\phi \Lambda f = \lim_{n \rightarrow \infty} G_\phi \Lambda f_n = \lim_{n \rightarrow \infty} \Lambda G_\phi f_n = \Lambda G_\phi f.$$

Using Kolmogorov's forward and backward equations, we get, for all $f \in \mathbf{D}(\mathbf{G}_\phi)$, $t > 0$ and $s \in [0, t]$,

$$\begin{aligned} \frac{d}{ds} Q_s^\phi \Lambda Q_{t-s}^\phi f &= Q_s^\phi G_\phi \Lambda Q_{t-s}^\phi - Q_s^\phi \Lambda G_\phi Q_{t-s}^\phi f \\ &= Q_s^\phi [G_\phi \Lambda - \Lambda G_\phi] Q_{t-s}^\phi f \\ &= 0 \end{aligned}$$

which is due to (4.20) together with the fact that $Q_{t-s}^\phi f \in \mathbf{D}(\mathbf{G}_\phi)$. Integrating the above identity, we obtain, for all $f \in \mathbf{D}(\mathbf{G}_\phi)$,

$$Q_t^\phi \Lambda f = \Lambda Q_t^\phi f.$$

Finally, using the density of $\mathbf{D}(\mathbf{G}_\phi)$ and the boundedness of the operators Q^ϕ , Q_t^ϕ and Λ , (4.7) follows.

4.1.2. Proof of Proposition 4.3(2). It is plain that, for any $n \in \mathbb{Z}_+$,

$$\int_0^\infty e^{-x} \frac{x^n}{n!} dx = 1,$$

which implies that $\mu \Lambda = \mathfrak{m}$ where μ is the Lebesgue measure on \mathbb{R}_+ . Also, Λ being a positive operator, for any $f \in \mathbf{C}_0(\mathbb{Z}_+)$ with $f \geq 0$, we have

$$\mathfrak{m} f = \mu \Lambda f \geq \mu Q_t^\phi \Lambda f = \mu \Lambda Q_t^\phi f = \mathfrak{m} Q_t^\phi f$$

where the second inequality in the above line holds as μ is an excessive measure for Q^ϕ . This shows that \mathfrak{m} is an excessive measure for Q^ϕ .

4.1.3. **Proof of Proposition 4.3(3).** For any $f \in \mathbf{C}_c(\mathbb{Z}_+)$,

$$\begin{aligned} \|\Lambda f\|_{\mathbf{L}^2(\mathbb{R}_+)}^2 &= \int_0^\infty (\Lambda f(x))^2 dx = \int_0^\infty \left(\sum_{n=0}^\infty e^{-x} \frac{x^n}{n!} f(n) \right)^2 dx \\ &\leq \sum_{n=0}^\infty f(n)^2 \int_0^\infty e^{-x} \frac{x^n}{n!} dx \\ &= \sum_{n=0}^\infty f(n)^2 = \|f\|_{\ell^2(\mathbb{Z}_+)}^2. \end{aligned}$$

Using the density of $\mathbf{C}_c(\mathbb{Z}_+)$ in $\ell^2(\mathbb{Z}_+)$, Λ extends uniquely to a bounded operator from $\ell^2(\mathbb{Z}_+)$ to $\mathbf{L}^2(\mathbb{R}_+)$. Finally, for any $f \in \mathbf{C}_0(\mathbb{Z}_+) \cap \ell^2(\mathbb{Z}_+)$, $\Lambda f \in \mathbf{C}_0(\mathbb{R}_+) \cap \mathbf{L}^2(\mathbb{R}_+)$. Thus, for all $f \in \mathbf{C}_0(\mathbb{Z}_+) \cap \ell^2(\mathbb{Z}_+)$, item (1) ensures that

$$Q_t^\phi \Lambda f = \Lambda Q_t^\phi f.$$

Again, using the density of $\mathbf{C}_c(\mathbb{Z}_+)$ in $\ell^2(\mathbb{Z}_+)$, (4.8) follows. Now, it remains to show that Λ is a quasi-affinity. Boundedness of Λ follows from item (2), and one easily checks that, for all $f \in \ell^2(\mathbb{Z}_+)$,

$$\Lambda f(x) = \sum_{n=0}^\infty e^{-x} \frac{x^n}{n!} f(n) \quad \text{a.e.}$$

Therefore, $\ker(\Lambda) = \{0\}$, which proves the injectivity. The density of $\text{Range}(\Lambda)$ follows by observing that $\widehat{\Lambda} : \mathbf{L}^2(\mathbb{R}_+) \rightarrow \ell^2(\mathbb{Z}_+)$, the adjoint of Λ , takes the following form

$$\widehat{\Lambda} f(n) = \frac{1}{n!} \int_0^\infty f(x) e^{-x} x^n dx = \mathbb{E}[f(\text{Gamma}(n+1))]$$

where $\text{Gamma}(n+1)$ is a gamma random variable with $n+1$ as the scale parameter and 1 as the rate parameter. Approximating the $\mathbf{L}^2(\mathbb{R}_+)$ functions by compactly supported continuous functions, it can be shown that $\widehat{\Lambda}$ is an injective operator, which proves that $\text{Range}(\Lambda)$ is dense in $\mathbf{L}^2(\mathbb{R}_+)$. Hence, item (3) is proven, which completes the proof of Proposition 4.3.

Corollary 4.7. For any $f : \mathbb{Z}_+ \rightarrow \mathbb{R}$ with $f \geq 0$, we have

$$Q_t^\phi \Lambda f(x) = \Lambda Q_t^\phi f(x)$$

for all $x \geq 0$.

Proof. For any nonnegative function f , we can find $\{f_n\} \subset \mathbf{C}_c(\mathbb{Z}_+)$ such that $f_n \uparrow f$ pointwise. Then, Proposition 4.3(1) yields, for all $x \geq 0$,

$$Q_t^\phi \Lambda f_n(x) = \Lambda Q_t^\phi f_n(x).$$

Since Λ is a Markov kernel, $\Lambda f_n \uparrow \Lambda f$ as well. Writing $Q_t^\phi g(x) = \mathbb{E}_x[g(X_\phi(t))]$ and $Q_t^\phi \mathbf{g}(n) = \mathbb{E}_n[\mathbf{g}(\mathbb{X}_\phi(t))]$ and invoking the monotone convergence theorem, the proof follows. \square

End of the Proof of Theorem 2.2. From the proof of Proposition 4.3(1), we already have that

$$Q_{\alpha t}^\phi \Lambda f = \Lambda Q_{\alpha t}^\phi f$$

for all $f \in \mathbf{C}_0(\mathbb{Z}_+)$. By a density argument, the above identity extends for all functions in $\mathbf{C}_b(\mathbb{Z}_+)$. Now, for $\alpha \in [0, 1]$, multiplying by d_α both sides of the above equation, we obtain that, for all $f \in \mathbf{C}_b(\mathbb{Z}_+)$,

$$\Lambda Q_t^\phi \mathbb{D}_\alpha f = Q_t^\phi \Lambda \mathbb{D}_\alpha f = Q_t^\phi d_\alpha \Lambda f = d_\alpha Q_{\alpha t}^\phi \Lambda f = d_\alpha \Lambda Q_{\alpha t}^\phi f = \Lambda \mathbb{D}_\alpha Q_{\alpha t}^\phi f$$

where we used the intertwining relationship between d_α and \mathbb{D}_α given in Proposition 4.1. By means of the injectivity of Λ on $\mathbf{C}_b(\mathbb{Z}_+)$, we complete the proof. \square

4.2. Proof of Theorem 2.5. Let \mathbb{G} denote the generator of the discrete self-similar Markov chain \mathbb{X} . Then, from the definition of the discrete self-similarity, for any $\alpha \in [0, 1]$, we have

$$\mathbb{G} \mathbb{D}_\alpha = \alpha \mathbb{D}_\alpha \mathbb{G} \quad \text{on } \mathbf{D}(\mathbb{G}).$$

Recalling that, for any $m, n \in \mathbb{Z}_+$, $\mathbb{D}_\alpha(m, n) = \mathbb{D}_\alpha \delta_n(m) = \binom{m}{n} \alpha^n (1 - \alpha)^{m-n} \mathbb{1}_{\{n \leq m\}}$ we have, for all $l, n \in \mathbb{Z}_+$,

$$(4.21) \quad \sum_{k \geq l} \mathbb{G}(n, k) \binom{k}{l} \alpha^l (1 - \alpha)^{k-l} = \sum_{j \leq n} \alpha^{j+1} (1 - \alpha)^{n-j} \mathbb{G}(j, l).$$

Taking $n = 0$ and $l = 1$ in the above equation, we obtain, for all $k \in \mathbb{Z}_+$, that

$$\sum_{k \geq 1} \mathbb{G}(0, k) \alpha (1 - \alpha)^{k-1} = \alpha \mathbb{G}(0, 1).$$

Using the fact that $\mathbb{G}(0, k) \geq 0$ for all $k > 0$, we conclude that $\mathbb{G}(0, k) = 0$ for all $k \geq 2$. We now use an induction argument to prove that $\mathbb{G}(n, k) = 0$ for all $k \geq n + 2$. Let us assume that for all $n < N \in \mathbb{Z}_+$, $\mathbb{G}(n, k) = 0$ for all $k \geq n + 2$. Now plugging $n = N, l = N + 1$ in (4.21) and using the induction hypothesis, we have

$$\sum_{k \geq N+1} \mathbb{G}(N, k) \binom{k}{N+1} \alpha^{N+1} (1 - \alpha)^{k-N-1} = \alpha^{N+1} \mathbb{G}(N, N+1).$$

Again invoking the nonnegativity of $\mathbb{G}(N, k)$ for $k \neq N$, we conclude that $\mathbb{G}(N, k) = 0$ for all $k \geq N + 2$. This completes the induction step and therefore the theorem is proved. \square

4.3. Factorial moments of discrete self-similar Markov chains. Let us recall that the skip-free Markov chain associated to ϕ is denoted by \mathbb{X}_ϕ . In the spirit of the work of Bertoin and Yor [8, Proposition 1(i)] on the integer moments of the continuous analogues, we provide an explicit formula for the factorial moments of $\mathbb{X}_\phi(t)$. For $z \in \mathbb{C}$, we recall that $\mathbf{p}_z : \mathbb{Z}_+ \rightarrow \mathbb{C}$ is the function defined by

$$(4.22) \quad \mathbf{p}_z(n) = \frac{\Gamma(n+1)}{\Gamma(n+1-z)}.$$

It is well-known that, for any $n, k \in \mathbb{Z}_+$,

$$(4.23) \quad n^k = \sum_{j=0}^k \left\{ \begin{matrix} k \\ j \end{matrix} \right\} \mathbf{p}_j(n)$$

where $\left\{ \begin{matrix} k \\ j \end{matrix} \right\}$ are the Stirling numbers of second kind, see [34, p. 81].

Theorem 4.8. *For any $n, k \in \mathbb{Z}_+$ and $t \geq 0$,*

$$(4.24) \quad \mathbb{E}[\mathbf{p}_k(\mathbb{X}_\phi(t, n))] = \mathbf{Q}_t^\phi \mathbf{p}_k(n) = \sum_{l=0}^k \binom{k}{l} \frac{W_\phi(k+1)}{W_\phi(l+1)} \mathbf{p}_l(n) t^{k-l}$$

where, for all $n \in \mathbb{Z}_+$, $W_\phi(n+1) = \prod_{k=1}^n \phi(k)$ and $W_\phi(1) = 1$.

Proof. Defining $p_k(x) = x^k$, from [8, Proposition 1(i)], we have, for all $k \in \mathbb{Z}_+$ and $t \geq 0$,

$$(4.25) \quad \begin{aligned} Q_t^\phi p_k(x) &= \mathbb{E}[(X_\phi(t, x))^k] = x^k + \sum_{l=1}^k \binom{k}{l} \phi(k) \phi(k-1) \cdots \phi(k-l+1) x^{k-l} t^l \\ &= \sum_{l=0}^k \binom{k}{l} \frac{W_\phi(k+1)}{W_\phi(l+1)} x^l t^{k-l}. \end{aligned}$$

On the other hand, it is easy to see that, for all $x > 0$, $\Lambda \mathbf{p}_k(x) = p_k(x)$. Applying Corollary 4.7 with $f = \mathbf{p}_k$, yields, for all $t, x \geq 0$,

$$Q_t^\phi p_k(x) = Q_t^\phi \Lambda \mathbf{p}_k(x) = \Lambda Q_t^\phi \mathbf{p}_k(x).$$

Recalling that $\nabla = \Lambda^{-1}$, see [26, Lemma 4], we get

$$\mathbb{E}[\mathbf{p}_k(\mathbb{X}_\phi(t, n))] = \mathbf{Q}_t^\phi \mathbf{p}_k(n) = \nabla Q_t^\phi p_k(n) = \frac{d^n}{dx^n} \left(e^x Q_t^\phi p_k(x) \right) (0).$$

Finally using the expression in (4.25) together with the Leibniz rule, the result follows. \square

Remark 4.9. Using (4.23) and the above theorem, $\mathbb{E}\left[\mathbb{X}_\phi^k(t, n)\right]$ can be also computed explicitly for all $n, k \in \mathbb{Z}_+$.

4.4. Proof of Theorem 2.6(2). For showing the weak convergence, we need to check the following two facts. First, the tightness property of the sequence $((Y_n(t) = \frac{1}{n}\mathbb{X}_\phi(nt, \lfloor nx \rfloor))_{t \geq 0})$ and the finite-dimensional convergence of (Y_n) to X_ϕ . For the tightness property, applying [20, Theorem 16.1] together with the strong Markov property of (Y_n) , it is enough to show that $Y_n(h_n) \xrightarrow{P} x$ (in probability) whenever $h_n \rightarrow 0$. We will in fact show that $\mathbb{E}[(Y_n(h_n) - x)^2] \rightarrow 0$ as $n \rightarrow \infty$. From Theorem 4.8, we obtain for all $t \geq 0$,

$$\mathbb{E}[Y_n(t)] = \frac{1}{n}\mathbb{E}[\mathbb{X}_\phi(nt, \lfloor nx \rfloor)] = \frac{1}{n}\mathbb{E}[\mathfrak{p}_1(\mathbb{X}_\phi(nt, \lfloor nx \rfloor))] = \frac{1}{n}(\lfloor nx \rfloor + \phi(1)nt)$$

and

$$\begin{aligned} \mathbb{E}[Y_n^2(t)] &= \frac{1}{n^2}\mathbb{E}[\mathfrak{p}_2(\mathbb{X}_\phi(nt, \lfloor nx \rfloor)) + \mathfrak{p}_1(\mathbb{X}_\phi(nt, \lfloor nx \rfloor))] \\ &= \frac{1}{n^2}(\mathfrak{p}_2(\lfloor nx \rfloor) + 2\phi(2)nt\lfloor nx \rfloor + \phi(1)\phi(2)n^2t^2 + \lfloor nx \rfloor + \phi(1)nt). \end{aligned}$$

Since $h_n \rightarrow 0$, the last two equations imply $\mathbb{E}[Y_n(h_n)] \rightarrow x$ and $\mathbb{E}[Y_n^2(h_n)] \rightarrow x^2$ as $n \rightarrow \infty$. Therefore, $\mathbb{E}[(Y_n(h_n) - x)^2] \rightarrow 0$, which proves the tightness of (Y_n) . Next, to get the finite-dimensional convergence, it is enough to prove that, for all $0 \leq t_1 < t_2 < \dots < t_k$ and $(\alpha_1, \alpha_2, \dots, \alpha_k) \in \mathbb{Z}_+^k$,

(4.26)

$$\lim_{n \rightarrow \infty} \mathbb{E}[Y_n^{\alpha_1}(t_1)Y_n^{\alpha_2}(t_2) \cdots Y_n^{\alpha_k}(t_k)] = \mathbb{E}\left[X_\phi^{\alpha_1}(t_1, x)X_\phi^{\alpha_2}(t_2, x) \cdots X_\phi^{\alpha_k}(t_k, x)\right],$$

as the finite-dimensional distributions of X_ϕ are moment determinate. To prove the above assertion, we need the following lemma.

Lemma 4.10. *For any $t \geq 0$ and $k \in \mathbb{Z}_+$,*

$$(4.27) \quad \lim_{n \rightarrow \infty} \mathbb{E}\left[(Y_n(t) - X_\phi(t, x))^k\right] = 0.$$

Proof. From Theorem 4.8, it is clear that for any $t \geq 0$ and $k \in \mathbb{Z}_+$, the sequence $(Y_n^k(t))_{n \geq 0}$ is uniformly integrable and, for each $k \in \mathbb{Z}_+$,

$$\begin{aligned} \mathbb{E}[Y_n^k(t)] &= \frac{1}{n^k}\mathbb{E}\left[\mathbb{X}_\phi^k(nt, \lfloor nx \rfloor)\right] \\ &= \frac{1}{n^k}\left(\mathbb{E}[\mathfrak{p}_k(\mathbb{X}_\phi(nt, \lfloor nx \rfloor))] + o(n^k)\right). \end{aligned}$$

Therefore,

$$\begin{aligned}
\lim_{n \rightarrow \infty} \mathbb{E}[Y_n^k(t)] &= \lim_{n \rightarrow \infty} \frac{1}{n^k} \mathbb{E}[\mathfrak{p}_k(\mathbb{X}_\phi(nt, \lfloor nx \rfloor))] \\
&= \lim_{n \rightarrow \infty} \frac{1}{n^k} \left(\mathfrak{p}_k(\lfloor nx \rfloor) + \sum_{l=1}^k \binom{k}{l} \frac{W_\phi(k+1)}{W_\phi(k-l+1)} \mathfrak{p}_{k-l}(\lfloor nx \rfloor) n^l t^l \right) \\
&= x^k + \sum_{l=1}^k \binom{k}{l} \frac{W_\phi(k+1)}{W_\phi(k-l+1)} x^{k-l} t^l = \mathbb{E}[X_\phi^k(t, x)].
\end{aligned}$$

Since the random variable $X_\phi(t, x)$ is moment determinate, the above identity indicates that for each $t \geq 0$, as $n \rightarrow \infty$,

$$Y_n(t) \xrightarrow{d} X_\phi(t, x).$$

This together with the uniform integrability mentioned above proves the lemma. \square

Now, coming back to the proof of the main theorem, we prove (4.26) by induction. Indeed, (4.26) is satisfied for $k = 1$, thanks to Lemma 4.10. Moreover, if for some $k \in \mathbb{Z}_+$ and $(\alpha_1, \alpha_2, \dots, \alpha_k) \in \mathbb{Z}_+^k$ (4.26) holds, then, by an uniform integrability argument as in the proof of the lemma, one can show that

$$(4.28) \quad \lim_{n \rightarrow \infty} \mathbb{E} \left[\left(Y_n^{\alpha_1}(t_1) \cdots Y_n^{\alpha_k}(t_k) - X_\phi^{\alpha_1}(t_1, x) \cdots X_\phi^{\alpha_k}(t_k, x) \right)^2 \right] = 0.$$

Now, writing

$$M_{n,k} = Y_n^{\alpha_1}(t_1) \cdots Y_n^{\alpha_k}(t_k), \quad M_k = X_\phi^{\alpha_1}(t_1, x) \cdots X_\phi^{\alpha_k}(t_k, x),$$

for any $(\alpha_1, \alpha_2, \dots, \alpha_{k+1}) \in \mathbb{Z}_+^{k+1}$, we have

$$\begin{aligned}
& \left| \mathbb{E} \left[\prod_{i=1}^{k+1} Y_n^{\alpha_i}(t_i) - \prod_{i=1}^{k+1} X_\phi^{\alpha_i}(t_i, x) \right] \right| \\
&= \left| \mathbb{E} \left[M_{n,k} Y^{\alpha_{k+1}}(t_{k+1}) - M_k X_\phi^{\alpha_{k+1}}(t_{k+1}, x) \right] \right| \\
&= \left| \mathbb{E} \left[M_{n,k} Y^{\alpha_{k+1}} - M_k Y^{\alpha_{k+1}}(t_{k+1}) + M_k Y^{\alpha_{k+1}}(t_{k+1}) - M_k X_\phi^{\alpha_{k+1}}(t_{k+1}, x) \right] \right| \\
(4.29) \quad & \leq \sqrt{\mathbb{E}[(M_{n,k} - M_k)^2] \mathbb{E}[Y^{2\alpha_{k+1}}(t_{k+1})]} \\
& \quad + \sqrt{\mathbb{E}[M_k^2] \mathbb{E}[(Y^{\alpha_{k+1}}(t_{k+1}) - X_\phi^{\alpha_{k+1}}(t_{k+1}, x))^2]}.
\end{aligned}$$

In view of Lemma 4.10, (4.26) and (4.28), the expression on the right-hand side of (4.29) tends to 0 as $n \rightarrow \infty$. This completes the induction step of our

hypothesis, therefore proving (4.26) for $k + 1$. This completes the proof of the finite-dimensional convergence of the process $(Y_n)_{n \geq 0}$, which concludes the proof of the theorem. \square

4.5. Intertwining of the skip-free Laguerre and generalized Laguerre semigroups. In this section we establish the connection between the generalized Laguerre semigroups as defined in [29] and the skip-free Laguerre semigroups introduced therein. From Theorem 1.6(2) in the aforementioned reference, it is known that, for any $\phi \in \mathbf{B}$, the generalized Laguerre semigroup K^ϕ on \mathbb{R}_+ has a unique invariant distribution ν_ϕ that is absolutely continuous and moment determinate. In the next result we show that the intertwining relationship in (4.7) is retained for the Laguerre semigroups as well. In the next proposition, we use the fact that the semigroup \mathbb{K}^ϕ has a unique invariant distribution denoted by \mathbf{n}_ϕ , which is proved in Proposition 4.12 below.

Proposition 4.11. (1) *Let $\phi \in \mathbf{B}$, then we have, for all $t \geq 0$ and $f \in \mathbf{C}_0(\mathbb{Z}_+)$,*

$$(4.30) \quad K_t^\phi \Lambda f = \Lambda \mathbb{K}_t^\phi f$$

where we recall that $\Lambda f(x) = \mathbb{E}[f(\text{Pois}(x))]$.

(2) $\mathbf{n}_\phi = \nu_\phi \Lambda$ is an invariant distribution of \mathbb{K}^ϕ , and, for all $n \in \mathbb{Z}_+$,

$$\mathbf{n}_\phi(n) = \frac{1}{n!} \int_0^{\phi(\infty)} e^{-x} x^n \nu_\phi(x) dx.$$

(3) *The Feller semigroup \mathbb{K}^ϕ extends uniquely to a strongly continuous Markov semigroup on $\ell^2(\mathbf{n}_\phi)$, which is again denoted by \mathbb{K}^ϕ . Furthermore, the operator*

$$\Lambda : \mathbf{C}_0(\mathbb{Z}_+) \rightarrow \mathbf{C}_0(\mathbb{R}_+)$$

has a unique extension in $\mathcal{B}(\ell^2(\mathbf{n}_\phi), \mathbf{L}^2(\nu_\phi))$, and, for all $t \geq 0$ and $f \in \ell^2(\mathbf{n}_\phi)$,

$$(4.31) \quad K_t^\phi \Lambda f = \Lambda \mathbb{K}_t^\phi f.$$

Moreover, taking the adjoint in the above identity, one gets, for all $t \geq 0$ and $f \in \mathbf{L}^2(\nu_\phi)$,

$$(4.32) \quad \widehat{\mathbb{K}}_t^\phi \widehat{\Lambda}_\phi f = \widehat{\Lambda}_\phi \widehat{K}_t^\phi f$$

where $\widehat{\Lambda}_\phi : \mathbf{L}^2(\nu_\phi) \rightarrow \ell^2(\mathbf{n}_\phi)$ is the adjoint of Λ , and, for all $f \in \mathbf{L}^2(\nu_\phi)$,

$$(4.33) \quad \widehat{\Lambda}_\phi f(n) = \frac{1}{n! \mathbf{n}_\phi(n)} \int_0^\infty e^{-x} x^n \nu_\phi(x) f(x) dx.$$

(4) \mathbb{K}^ϕ is self-adjoint in $\ell^2(\mathbf{n}_\phi)$ if and only if $\phi(u) = \sigma^2 u + m$ for some $\sigma^2, m \geq 0$.

Proof. Since $\mathbb{L}^\phi = \mathbb{G}_\phi + n\partial_-$ and $L^\phi = G_\phi - x\frac{d}{dx}$, it suffices to show that, for all $f \in \mathbf{C}_c(\mathbb{Z}_+)$,

$$(4.34) \quad -x \frac{d}{dx} \Lambda f(x) = \Lambda(n\partial_-) f(x).$$

From [26, Lemma 23], (4.34) readily follows by considering the reverse intertwining relationship (i.e., taking the inverse of Δ in (45) of the aforementioned reference). Thus, we conclude that, for all $f \in \mathbf{C}_c(\mathbb{Z}_+)$,

$$L^\phi \Lambda f = \Lambda \mathbb{L}^\phi f.$$

The rest of the proof follows similarly as in the proof of Theorem 4.3(1).

Next, from the intertwining relation in (1), we deduce that, for all $f \in \mathbf{C}_0(\mathbb{Z}_+)$,

$$\nu_\phi \Lambda \mathbb{K}_t^\phi f = \nu_\phi K_t^\phi \Lambda f = \nu_\phi \Lambda f$$

implying that $\mathbf{n}_\phi = \nu_\phi \Lambda$ is an invariant finite measure for \mathbb{K}^ϕ . Now, for any $n \in \mathbb{Z}_+$,

$$(4.35) \quad \mathbf{n}_\phi(n) = \nu_\phi \Lambda \delta_n = \frac{1}{n!} \int_0^\infty e^{-x} x^n \nu_\phi(x) dx > 0$$

and

$$\sum_{n=0}^\infty \mathbf{n}_\phi(n) = \int_0^\infty \nu_\phi(x) dx = 1.$$

Hence, \mathbf{n}_ϕ is the invariant distribution of \mathbb{K}^ϕ . The uniqueness of the invariant distribution will be proved in Proposition 4.12(1). To prove (3), we note that since $\mathbf{C}_0(\mathbb{Z}_+)$ is dense in $\ell^2(\mathbf{n}_\phi)$ and $\Lambda : \mathbf{C}_0(\mathbb{Z}_+) \rightarrow \mathbf{C}_0(\mathbb{Z}_+)$ is a Markov kernel, Λ can be uniquely extended to a bounded operator in $\mathcal{B}(\ell^2(\mathbf{n}_\phi), \mathbf{L}^2(\nu_\phi))$. Also, using the density of $\mathbf{C}_0(\mathbb{Z}_+)$ in $\ell^2(\mathbf{n}_\phi)$ and item (1), the identity (4.31) follows. Now, to compute the adjoint $\widehat{\Lambda}_\phi$ of Λ , let us first show that the right-hand side of (4.33) as a function of n belongs to $\ell^2(\mathbf{n}_\phi)$.

Using Young's inequality and item (2) one has

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{1}{\mathbf{n}_\phi(n)^2} \left(\int_0^\infty e^{-x} \frac{x^n}{n!} \nu_\phi(x) f(x) dx \right)^2 \mathbf{n}_\phi(n) &\leq \sum_{n=0}^{\infty} \int_0^\infty e^{-x} \frac{x^n}{n!} f^2(x) \nu_\phi(x) dx \\ &= \|f\|_{\mathbf{L}^2(\nu_\phi)}^2. \end{aligned}$$

Now, writing $f(n) = \sum_{n=0}^{\infty} \frac{1}{n \mathbf{n}_\phi(n)} \int_0^\infty e^{-x} x^n f(x) \nu_\phi(x) dx$, we have, for all $\mathbf{g} \in \ell^2(\mathbf{n}_\phi)$,

$$\begin{aligned} \langle \mathbf{g}, f \rangle_{\mathbf{n}_\phi} &= \sum_{n=0}^{\infty} f(n) \mathbf{g}(n) \mathbf{n}_\phi(n) = \sum_{n=0}^{\infty} \frac{1}{n!} \mathbf{g}(n) \int_0^\infty e^{-x} x^n f(x) \nu_\phi(x) dx \\ &= \int_0^\infty \Lambda \mathbf{g}(x) f(x) \nu_\phi(x) dx \end{aligned}$$

where the third equality is justified by Fubini theorem. This shows that $\widehat{\Lambda}_\phi f = f$ which proves (4.33). Finally, to justify (4), we note that, for a $\phi \in \mathbf{B}$, \mathbb{K}^ϕ is self-adjoint in $\ell^2(\mathbf{n}_\phi)$ if and only if, for all $l, n \in \mathbb{Z}_+$,

$$(4.36) \quad \mathbb{L}^\phi(n, l) \mathbf{n}_\phi(n) = \mathbb{L}^\phi(l, n) \mathbf{n}_\phi(l).$$

Since $\mathbb{L}^\phi(n, l) = 0$ whenever $l \geq n + 2$ and $\mathbf{n}_\phi(n) > 0$ for all $n \in \mathbb{Z}_+$ (see the proof of item (2)), the above identity holds only if $\mathbb{L}^\phi(n, l) = 0$ for all $l \neq n, n - 1, n + 1$. This happens only if $\phi(u) = \sigma^2 u + m$ for some $\sigma^2, m \geq 0$. \square

4.6. Proof of Theorem 2.8(2). First, we recall that, for all $t \geq 0$ and $f \in \mathbf{C}_0(\mathbb{R}_+)$, $K_t^\phi f = Q_{e^t-1}^\phi d_{e^{-t}} f$, see e.g. [29]. Then, from (4.30), we have for all $t \geq 0$ and $f \in \mathbf{C}_0(\mathbb{Z}_+)$,

$$\Lambda \mathbb{K}_t^\phi f = K_t^\phi \Lambda f = Q_{e^t-1}^\phi d_{e^{-t}} \Lambda f = Q_{e^t-1}^\phi \Lambda \mathbb{D}_{e^{-t}} f = \Lambda Q_{e^t-1}^\phi \mathbb{D}_{e^{-t}} f$$

where we used, from the third identity onwards, successively (4.7) and (4.1). We conclude the proof by invoking the Feller property of the semigroups as well as the injectivity of Λ on $\mathbf{C}_0(\mathbb{Z}_+)$, see [26, Lemma 4(4)].

4.7. The invariant distribution of the skip-free Laguerre semigroup.

We now show that the invariant distribution \mathbf{n}_ϕ in Proposition 4.11(3) is unique and provide several useful representations. We recall that, for any $\phi \in \mathbf{B}$, W_ϕ is the so-called Bernstein-gamma function which is defined as a solution to the following functional equation

$$W_\phi(z+1) = \phi(z) W_\phi(z) \quad \forall z \in \mathbb{C}_+, \quad W_\phi(1) = 1.$$

The above functional equation has a unique solution in the class of Mellin transforms of probability measures on \mathbb{R}_+ . For a detailed account of these functions, we refer to [30].

Proposition 4.12. (1) For all $\phi \in \mathbf{B}$, the invariant distribution \mathbf{n}_ϕ of \mathbb{K}^ϕ is unique and is determined by its factorial moments

$$(4.37) \quad \mathbf{n}_\phi \mathbf{p}_k = W_\phi(k+1)$$

where \mathbf{p}_k is defined in (4.22).

(2) For any $n \in \mathbb{Z}_+$ and $0 < c < n + 1 + \mathbf{d}_\phi$,

$$(4.38) \quad \mathbf{n}_\phi(n) = \frac{1}{n!} \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \Gamma(z) W_\phi(n-z+1) dz$$

where $\mathbf{d}_\phi = \min\{u \geq 0; \phi(-u) = -\infty, \phi(-u) = 0\} \in [0, \infty]$.

(3) If $0 \leq \sigma^2 < 1$, then, for any $n \in \mathbb{Z}_+$,

$$(4.39) \quad \mathbf{n}_\phi(n) = \frac{1}{n!} \sum_{r=0}^{\infty} (-1)^r \frac{W_\phi(n+r+1)}{r!}.$$

Let us first derive the factorial moments of the skip-free Laguerre chains.

Lemma 4.13. For any $t \geq 0$ and $k \in \mathbb{Z}_+$,

$$(4.40) \quad \mathbb{K}_t^\phi \mathbf{p}_k(n) = \sum_{l=0}^k \binom{k}{l} \frac{W_\phi(k+1)}{W_\phi(l+1)} \mathbf{p}_l(n) e^{-tl} (1 - e^{-t})^{k-l}.$$

Proof. Let us recall that Q^ϕ denote the spectrally negative self-similar semi-group associated to ϕ , and, for any $t \geq 0$, $x > 0$ and $f \geq 0$,

$$K_t^\phi f(x) = Q_{1-e^{-t}}^\phi d_{e^{-t}} f(x),$$

and, writing $p_k(x) = x^k$, $x > 0$, we have, from [8], that, for all $k \in \mathbb{Z}_+$,

$$Q_t^\phi p_k(x) = \sum_{l=0}^k \binom{k}{l} \frac{W_\phi(k+1)}{W_\phi(l+1)} x^l t^{k-l}.$$

Therefore,

$$K_t^\phi p_k(x) = \sum_{l=0}^k \binom{k}{l} \frac{W_\phi(k+1)}{W_\phi(l+1)} e^{-tl} (1 - e^{-t})^{k-l} p_l(x).$$

Recalling that $\nabla = \Lambda^{-1}$ and $\nabla p_l = \mathbf{p}_l$ for all $l \in \mathbb{Z}_+$, it follows that

$$\begin{aligned} \mathbb{K}_t^\phi \mathbf{p}_k(n) &= \mathbb{K}_t^\phi \nabla p_k(n) = \nabla K_t^\phi p_k(n) \\ &= \sum_{l=0}^k \binom{k}{l} \frac{W_\phi(k+1)}{W_\phi(l+1)} e^{-tl} (1 - e^{-t})^{k-l} \nabla p_l(n) \\ &= \sum_{l=0}^k \binom{k}{l} \frac{W_\phi(k+1)}{W_\phi(l+1)} e^{-tl} (1 - e^{-t})^{k-l} \mathbf{p}_l(n) \end{aligned}$$

which completes the proof. \square

4.8. Proof of Proposition 4.12. From Lemma 4.13, we observe that, for all $k, n \in \mathbb{Z}_+$,

$$\lim_{t \rightarrow \infty} \mathbb{K}_t^\phi \mathbf{p}_k(n) = W_\phi(k+1).$$

On the other hand, recalling that $\Lambda \mathbf{p}_k = p_k$ where $p_k(x) = x^k$ and $\nu_\phi p_k = W_\phi(k+1)$, see [29, Proposition 2.6(1)], we get

$$\mathbf{n}_\phi \mathbf{p}_k = \nu_\phi \Lambda \mathbf{p}_k = \nu_\phi p_k = W_\phi(k+1).$$

Now it remains to show that \mathbf{n}_ϕ is determined by its moments. Let us write $\mathbf{e}_a(n) = e^{an}$. Then, applying Tonelli theorem we get

$$\mathbf{e}_a \mathbf{n}_\phi = \sum_{n=0}^{\infty} e^{an} \mathbf{n}_\phi(n) = \int_0^{\infty} e^{-x} \sum_{n=0}^{\infty} \frac{(e^a x)^n}{n!} \nu_\phi(x) dx = \int_0^{\infty} e^{(e^a - 1)x} \nu_\phi(x) dx.$$

Next, we have

$$\int_0^{\infty} e^{(e^a - 1)x} \nu_\phi(x) dx = \sum_{r=0}^{\infty} W_\phi(r+1) \frac{(e^a - 1)^r}{r!}$$

where we used the fact that $\int_0^{\phi(\infty)} x^n \nu_\phi(x) dx = W_\phi(n+1)$, see [29, Proposition 2.6(1)], and thus $\mathbf{e}_a \mathbf{n}_\phi < \infty$ as soon as $(e^a - 1) < \sigma^{-2}$, that is for at least any $0 < a < \log(1 + \sigma^{-2})$. This provides the moment determinacy of \mathbf{n}_ϕ . Next, to prove (2) and (3), we observe, from Proposition 4.14(2), that for all $n \in \mathbb{Z}_+$,

$$\mathbf{n}_\phi(n) = \int_0^{\phi(\infty)} e^{-x} x^n \nu_\phi(x) dx.$$

Expanding the exponential function in the identity above and using a classical Fubini argument, see e.g. [35, Section 1.77], combined with the expression (4.37) of the moment of ν_ϕ , we get

$$(4.41) \quad \mathbf{n}_\phi(n) = \frac{1}{n!} \int_0^{\infty} e^{-x} x^n \nu_\phi(x) dx = \frac{1}{n!} \sum_{r=0}^{\infty} W_\phi(n+r+1) \frac{(-1)^r}{r!}$$

where the series is absolutely convergent as soon as

$$\lim_{k \rightarrow \infty} \frac{\phi(k+n)}{k} = \lim_{k \rightarrow \infty} \frac{\phi(k)}{k} = \sigma^2 < 1.$$

To justify the contour integral representation in (4.38), we consider two cases. Assume first that $\sigma^2 > 0$ and we recall that for large $|\operatorname{Im}(z)|$,

$$(4.42) \quad |\Gamma(z)| \sim C_{\operatorname{Re}(z)} |\operatorname{Im}(z)|^{\operatorname{Re}(z)-\frac{1}{2}} e^{-\frac{\pi}{2}|\operatorname{Im}(z)|}, \quad \operatorname{Re}(z) > 0,$$

$$|W_\phi(n-z+1)| \leq C_{n-\operatorname{Re}(z)} e^{-\frac{\pi}{2}|\operatorname{Im}(z)|}, \quad \operatorname{Re}(z) < n+1+d_\phi,$$

where here and below $C_{\operatorname{Re}(z)} > 0$ is a constant depending only on $\operatorname{Re}(z) > 0$. Note that the first estimate is the classical Stirling formula, see e.g. [28, (2.1.8)], whereas the second bound follows from [29, Theorem 6.2(2b)]. Therefore, the mappings $z \mapsto \Gamma(z)$ and $z \mapsto W_\phi(n-z+1)$ are both in $\mathbf{L}^2(\mathbb{R})$ and holomorphic in the strip $0 < \operatorname{Re}(z) < n+1+d_\phi$, see [29, Theorem 6.1(2)]. Moreover $z \mapsto W_\phi(n+z)$ and $z \mapsto \Gamma(z)$ are the Mellin transform of $x \mapsto x^n \nu_\phi(x)$, see [29], and $x \mapsto e^{-x}$, respectively. Consequently, both of these functions are in $\mathbf{L}^2(\mathbb{R}_+)$. An application of Parseval identity for the Mellin transform yields

$$\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \Gamma(z) W_\phi(n-z+1) dz = \int_0^\infty e^{-x} x^n \nu_\phi(x) dx,$$

from where we easily derive the expression (4.38) for $\sigma^2 > 0$. Next, we consider the other case, that is $\sigma^2 = 0$, which ensures that the series representation (4.41) of $\mathbf{n}_\phi(n)$ is valid for all $n \in \mathbb{Z}_+$. Then, using the facts that the mappings $z \mapsto \Gamma(z)$ and $z \mapsto W_\phi(n-z+1)$ are both holomorphic in the strip $0 < \operatorname{Re}(z) < n+1+d_\phi$ and within this strip, (4.42) still holds and

$$|W_\phi(n-z+1)| \leq C_{(n-\operatorname{Re}(z))}.$$

This implies that for all $n \in \mathbb{N}$, the integral (4.38) is absolutely convergent and an application of Cauchy theorem, see [28] for the detailed computation, yields that the contour integral can be expanded as follows

$$\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \Gamma(z) W_\phi(n-z+1) dz = \sum_{r=0}^{\infty} W_\phi(n+r+1) \frac{(-1)^r}{r!},$$

which completes the proof of (4.39). \square

4.9. Intertwining between skip-free Laguerre semigroups. It has been shown in [29, Theorem 2.1] that for any $\phi \in \mathbf{B}$, the generalized (non self-adjoint) Laguerre semigroup K^ϕ is intertwined with the diffusive (self-adjoint) Laguerre semigroup K (when $\phi(u) = u$) and the intertwining operator is a multiplicative Markov kernel corresponding to the exponential functional of the subordinator associated with ϕ . An analogous result holds

for the skip-free Laguerre semigroups as well (see Theorem 4.14 below), although, we prove it under the assumption that σ^2 in (2.5) is positive. The following proposition describes the intertwining operator \mathbb{I}_ϕ that links the semigroups corresponding to skip-free (non-reversible) and the reversible Laguerre chains respectively. Let $\mathcal{P} = \text{Span} \{\mathbf{p}_k, k \in \mathbb{Z}_+\}$ and for any $\phi \in \mathbf{B}$ associated with the triplet (m, σ^2, Π) , let $\mathbb{I}_\phi : \mathcal{P} \mapsto \mathcal{P}$ be defined by

$$(4.43) \quad \mathbb{I}_\phi \mathbf{f}(n) = \mathbb{E} [\mathbf{f}(\mathbf{B}(I_{\sigma_1}, n))] = \sum_{r=0}^n \mathbf{f}(r) \binom{n}{r} \mathbb{E} [I_{\sigma_1}^r (1 - I_{\sigma_1})^{n-r}]$$

with $\mathbb{E} [I_{\sigma_1}^k] = \frac{\sigma_1^k k!}{W_\phi(k+1)}$ for all $k \in \mathbb{Z}_+$, where σ_1 is defined in (2.19) and we recall that $W_\phi(k+1) = \prod_{r=1}^k \phi(r)$, $W_\phi(1) = 1$.

Theorem 4.14. (1) For any $\phi \in \mathbf{B}$, we have the intertwining relation on \mathcal{P}

$$(4.44) \quad \mathbb{K}_t^\phi \mathbb{I}_\phi = \mathbb{I}_\phi \mathbb{K}_t^{\sigma_1}$$

where $\mathbb{K}^{\sigma_1} = \mathbb{K}^\phi$ with $\phi(u) = \sigma_1 u$.

(2) Moreover, if $\sigma^2 > 0$, then $\mathbb{I}_\phi : \ell^2(\mathbf{n}_{\sigma^2}) \mapsto \ell^2(\mathbf{n}_\phi)$ is a linear operator that is bounded, injective with a dense range and for all $\mathbf{f} \in \ell^2(\mathbf{n}_{\sigma^2})$,

$$(4.45) \quad \|\mathbb{I}_\phi \mathbf{f}\|_{\ell^2(\mathbf{n}_\phi)} \leq \|\mathbf{f}\|_{\ell^2(\mathbf{n}_{\sigma^2})}$$

where \mathbf{n}_{σ^2} is the unique invariant distribution of \mathbb{K}^{σ^2} , and, for all $t \geq 0$, $\mathbf{f} \in \ell^2(\mathbf{n}_{\sigma^2})$,

$$(4.46) \quad \mathbb{K}_t^\phi \mathbb{I}_\phi \mathbf{f} = \mathbb{I}_\phi \mathbb{K}_t^{\sigma^2} \mathbf{f}.$$

As a consequence of the above theorem, we obtain the intertwining relationship among the class of discrete self-similar Markov semigroups.

Corollary 4.15. For $\phi \in \mathbf{B}$ with $\sigma^2 > 0$, we have

$$\mathbb{Q}_t^\phi \mathbb{I}_\phi = \mathbb{I}_\phi \mathbb{Q}_{\sigma^2 t}$$

both on $\mathbf{C}_0(\mathbb{Z}_+)$ and $\ell^2(\mathbb{Z}_+)$, where we recall that \mathbb{Q}^ϕ (resp. \mathbb{Q}) is the discrete self-similar semigroup corresponding to the Bernstein function ϕ (resp. $\phi(u) = u$).

We need the following lemma to prove the above theorem.

Lemma 4.16. Recall the definition of \mathbf{p}_z in (4.22). Then, for all $k, n \in \mathbb{Z}_+$, we have

$$\mathbb{I}_\phi \mathbf{p}_k(n) = \frac{\sigma_1^k k!}{W_\phi(k+1)} \mathbf{p}_k(n)$$

where σ_1 is defined in (2.19).

Proof. Recalling the definition of \mathbb{I}_ϕ in (4.43), we have, for all $k, n \in \mathbb{Z}_+$,

$$(4.47) \quad \mathbb{I}_\phi \mathbf{p}_k(r) = \sum_{r=0}^n \mathbf{p}_k(r) \binom{n}{r} \mathbb{E} [I_{\sigma_1}^r (1 - I_{\sigma_1})^{n-r}] = \mathbb{E} [\mathbf{p}_k(\mathbf{B}(n, I_{\sigma_1}))]$$

where \mathbf{B} denotes the Binomial random variable with the parameters written in the parentheses and the moments of I_{σ_1} are given in (4.43). Also, invoking the definition of the discrete dilation operator \mathbb{D} , we can write the above quantity as

$$\mathbb{E} [\mathbf{p}_k(\mathbf{B}(n, I_{\sigma_1}))] = \mathbb{E} [\mathbb{D}_{I_{\sigma_1}} \mathbf{p}_k(n)] = \mathbf{p}_k(n) \mathbb{E} [I_{\sigma_1}^k] = \frac{\sigma_1^k k!}{W_\phi(k+1)} \mathbf{p}_k(n)$$

where the last equality follows from (4.43). This proves the lemma. \square

4.10. Proof of Theorem 4.14. From Lemma 4.13 and Lemma 4.16, we have, for all $t \geq 0$ and $k \in \mathbb{N}$,

$$\begin{aligned} \mathbb{K}_t^\phi \mathbb{I}_\phi \mathbf{p}_k &= \frac{\sigma_1^k k!}{W_\phi(k+1)} \mathbb{K}_t^\phi \mathbf{p}_k \\ &= \frac{\sigma_1^k k!}{W_\phi(k+1)} \sum_{l=0}^k e^{-tl} (1 - e^{-t})^{k-l} \binom{k}{l} \frac{W_\phi(k+1)}{W_\phi(l+1)} \mathbf{p}_l \\ &= \sigma_1^k k! \sum_{l=0}^k e^{-tl} (1 - e^{-t})^{k-l} \binom{k}{l} \frac{1}{W_\phi(l+1)} \mathbf{p}_l(n). \end{aligned}$$

On the other hand, recalling that $\mathbb{K}^{\sigma_1} = \mathbb{K}^\phi$ with $\phi(u) = \sigma_1 u$, we have

$$\begin{aligned} \mathbb{I}_\phi \mathbb{K}_t^{\sigma_1} \mathbf{p}_k &= \sum_{l=0}^k e^{-tl} (1 - e^{-t})^{k-l} \sigma_1^{k-l} \binom{k}{l} \frac{k!}{l!} \mathbb{I}_\phi \mathbf{p}_l \\ &= k! \sum_{l=0}^k e^{-tl} (1 - e^{-t})^{k-l} \sigma_1^{k-l} \binom{k}{l} \frac{\sigma_1^l}{W_\phi(l+1)}, \end{aligned}$$

which shows that for all $k \in \mathbb{N}$,

$$\mathbb{K}_t^\phi \mathbb{I}_\phi \mathbf{p}_k = \mathbb{I}_\phi \mathbb{K}_t^{\sigma_1} \mathbf{p}_k$$

and therefore, on \mathcal{P} ,

$$\mathbb{K}_t^\phi \mathbb{I}_\phi = \mathbb{I}_\phi \mathbb{K}_t^{\sigma_1}.$$

To prove now (2), it is plain, from Lemma 4.16, that $\mathbb{I}_\phi(\mathcal{P}) = \mathcal{P}$. Then, under the condition $\sigma^2 > 0$, we have that $\mathbb{P}(I_{\sigma^2} \in [0, 1]) = 1$, see [29, Proposition 6.7] (note that $I_{\sigma^2} = \sigma^2 I_\phi$, where I_ϕ is the exponential functional

defined in the aforementioned paper) and thus \mathbb{I}_ϕ is a Markov operator. By means of Hölder's inequality, one obtains, for any $f \in \ell^2(\mathbf{n}_{\sigma^2})$,

$$(4.48) \quad \|\mathbb{I}_\phi f\|_{\ell^2(\mathbf{n}_\phi)} \leq \|\mathbb{I}_\phi f^2\|_{\ell^1(\mathbf{n}_\phi)} = \mathbf{n}_\phi \mathbb{I}_\phi f^2.$$

Now, for all $k \in \mathbb{Z}_+$, using Lemma 4.16 and Proposition 4.12(1), we obtain

$$\mathbf{n}_\phi \mathbb{I}_\phi \mathbf{p}_k = \sum_{n \in \mathbb{Z}_+} \mathbf{n}_\phi(n) \mathbb{I}_\phi \mathbf{p}_k(n) = \sum_{n \in \mathbb{Z}_+} \mathbf{n}_\phi(n) \mathbf{p}_k(n) \frac{\sigma^{2k} k!}{W_\phi(k+1)} = \sigma^{2k} k! = \mathbf{p}_k \mathbf{n}_{\sigma^2}$$

which shows that $\mathbf{n}_\phi \mathbb{I}_\phi = \mathbf{n}_{\sigma^2}$ as $\mathbf{n}_\phi, \mathbf{n}_{\sigma^2}$ are moment determinate. Therefore, (4.48) entails that \mathbb{I}_ϕ is a bounded operator from $\ell^2(\mathbf{n}_{\sigma^2})$ to $\ell^2(\mathbf{n}_\phi)$ when $\sigma^2 > 0$. Hence, by the density of \mathcal{P} in $\ell^2(\mathbf{n}_{\sigma^2})$, the intertwining relation given by (4.44) extends to $\ell^2(\mathbf{n}_{\sigma^2})$. This completes the proof of the proposition. \square

4.11. Hilbert sequences and spectral expansion. In this section, we introduce a few notions from non classical harmonic analysis which have been shown recently to be central in the understanding of the spectral expansions of non self-adjoint operators in Hilbert spaces, see e.g. [29]. Two sequences $(\mathbf{P}_k)_{k \geq 0}$ and $(\mathbf{V}_k)_{k \geq 0}$ are said to be biorthogonal in the Hilbert space $\ell^2(\mathbf{m})$ if for any $k, l \in \mathbb{Z}_+$,

$$(4.49) \quad \langle \mathbf{P}_k, \mathbf{V}_l \rangle_{\mathbf{m}} = \mathbb{1}_{\{k=l\}}.$$

Moreover, a sequence that admits a biorthogonal sequence will be called **minimal** and a sequence that is both minimal and complete, in the sense that its linear span is dense in $\ell^2(\mathbf{m})$, will be called **exact**. It is easy to show that a sequence $(\mathbf{P}_k)_{k \geq 0}$ is minimal if and only if none of its elements can be approximated by linear combinations of the others. If this is the case, then a biorthogonal sequence will be uniquely determined if and only if $(\mathbf{P}_k)_{k \geq 0}$ is complete. We proceed with some basic notions related to the concept of frames in Hilbert spaces. A recent and thorough account on these Hilbert space sequences can be found in the book of Christensen [14]. A sequence $(\mathbf{P}_k)_{k \geq 0}$ in $\ell^2(\mathbf{m})$ is a frame if there exist $A, B > 0$ such that the frame inequalities

$$(4.50) \quad A \|f\|_{\ell^2(\mathbf{m})}^2 \leq \sum_{k=0}^{\infty} |\langle f, \mathbf{P}_k \rangle_{\mathbf{m}}|^2 \leq B \|f\|_{\ell^2(\mathbf{m})}^2$$

hold, for all $f \in \ell^2(\mathbf{m})$. If only the upper bound exists, $(\mathbf{P}_k)_{k \geq 0}$ is called a Bessel sequence. A frame sequence is always complete in the Hilbert space and when it is minimal, it is called a Riesz sequence. The latter are very useful objects as they share substantial properties with orthonormal sequences. Indeed, a Riesz sequence always admits a unique biorthogonal sequence $(\mathbf{V}_k)_{k \geq 0}$ which is also a Riesz sequence and both together form the

so-called Riesz basis. Moreover, the expansion in terms of the Riesz basis of any element of the Hilbert space is unique and convergent in the topology of the norm. When $(\mathbf{P}_k)_{k \geq 0}$ is merely a Bessel sequence, that is only the upper frame condition in (4.50) is satisfied, then the so-called synthesis operator, that is the linear operator $\mathcal{S} : \ell^2(\mathbb{Z}_+) \rightarrow \ell^2(\mathbf{m})$ defined by

$$(4.51) \quad \mathcal{S} : \underline{c} = (c_k)_{k \geq 0} \mapsto \mathcal{S}(\underline{c}) = \sum_{k=0}^{\infty} c_k \mathbf{P}_k$$

is a bounded operator with (operator) norm $\|\mathcal{S}\|_{\mathbf{m}} \leq \sqrt{B}$, that is, the series is norm convergent for any sequence in $\ell^2(\mathbb{Z}_+)$. However, \mathcal{S} is not in principle onto as the $(\mathbf{P}_k)_{k \geq 0}$ does not form in general a basis of the Hilbert space.

Proposition 4.17. *Let $\phi \in \mathbf{B}$.*

(1) *For any $k \in \mathbb{Z}_+$, $\mathbf{P}_k^\phi \in \ell^2(\mathbf{n}_\phi)$, and, for any $t > 0$,*

$$\mathbb{K}_t^\phi \mathbf{P}_k^\phi = e^{-kt} \mathbf{P}_k^\phi.$$

Moreover, $\text{Span}\{\mathbf{P}_k^\phi, k > 0\} = \mathcal{P}$ which is dense in $\ell^2(\mathbf{n}_\phi)$.

(2) *Assume that $\sigma^2 > 0$. Then, $(\mathbf{P}_k^\phi)_{k \geq 0}$ is an exact Bessel sequence in $\ell^2(\mathbf{n}_\phi)$ with bound 1 and for any $k \in \mathbb{Z}_+$,*

$$(4.52) \quad \left\| \mathbf{P}_k^\phi \right\|_{\ell^2(\mathbf{n}_\phi)} \leq 1.$$

(3) *If $\sigma^2 > 0$ and $d_\phi > 0$. Then, $(\sqrt{\mathbf{c}_k(d_\phi)} \mathbf{P}_k^\phi)_{k \geq 0}$ is a Bessel sequence with, for all $k \in \mathbb{Z}_+$,*

$$(4.53) \quad \left\| \mathbf{P}_k^\phi \right\|_{\ell^2(\mathbf{n}_\phi)} \leq \frac{1}{\sqrt{\mathbf{c}_k(d_\phi)}}$$

$$\text{where } \mathbf{c}_k(d_\phi) = \frac{\Gamma(k+d_\phi+1)}{\Gamma(k+1)\Gamma(d_\phi+1)}.$$

Proof. Let $k \in \mathbb{Z}_+$, then it is plain, from Proposition 1, that, as a polynomial, $\mathbf{P}_k^\phi \in \ell^2(\mathbf{n}_\phi)$. Then, we recall, from [29, Theorem 7.3] (after multiplying both sides of the next identity by $(1 + \sigma_1^{-1})^{-\frac{k}{2}}$) that, for any $t > 0$ and $k \in \mathbb{Z}_+$,

$$K_t^\phi \mathcal{P}_k^\phi = e^{-kt} \mathcal{P}_k^\phi$$

where we have set

$$\mathcal{P}_k^\phi(x) = (1 + \sigma_1^{-1})^{-\frac{k}{2}} \sum_{r=0}^k (-1)^r \binom{k}{r} \frac{x^r}{W_\phi(r+1)} \in \mathbf{L}^2(\nu_\phi).$$

Thus, since Λ is injective on $\mathcal{P} \subset \ell^2(\mathbf{n}_\phi)$, the algebra of polynomials, we have $\Lambda^{-1}p_k(n) = \mathbf{p}_k(n)$ and thus, by linearity

$$(4.54) \quad \Lambda^{-1}\mathcal{P}_k^\phi(n) = (1 + \sigma_1^{-1})^{-\frac{k}{2}} \sum_{r=0}^k (-1)^r \frac{\binom{k}{r}}{W_\phi(r+1)} \Lambda^{-1}p_r(n) = \mathbf{P}_k^\phi(n).$$

Finally, we observe from the gateway relationship (4.31) and the linearity of the operators, that

$$\Lambda \mathbb{K}_t^\phi \mathbf{P}_k^\phi = K_t^\phi \Lambda \Lambda^{-1} \mathcal{P}_k^\phi = K_t^\phi \mathcal{P}_k^\phi = e^{-kt} \mathcal{P}_k^\phi = \Lambda e^{-kt} \mathbf{P}_k^\phi.$$

The injectivity of Λ on \mathcal{P} yields the eigenfunction property. To complete the proof of (1), we recall the moment determinacy of \mathbf{n}_ϕ , stated in Proposition 4.12(1), which entails, from classical results on the moment problem, the density property of the algebra of polynomials in the weighted Hilbert space, see [2]. Next, when $\sigma^2 > 0$ and $\phi(u) = \sigma^2 u$, we recall, from Example 3.1, that $(\mathbf{P}_k^{\sigma^2})_{k \geq 0}$ is an orthonormal sequence of eigenfunctions of $\mathbb{K}_t^{\sigma^2}$ associated to the eigenvalues $\{e^{-kt}\}_{k \geq 0}$. Now, from Lemma 4.16, it is easily seen, from the definition of \mathbf{P}_k^ϕ , that, for all $k \geq 0$,

$$\mathbb{I}_\phi \mathbf{P}_k^{\sigma^2} = \mathbf{P}_k^\phi.$$

Since $\mathbb{I}_\phi \in \mathcal{B}(\ell^2(\mathbf{n}_{\sigma^2}), \ell^2(\mathbf{n}_\phi))$ whenever $\sigma^2 > 0$, see (4.45), it follows that, for all $k \geq 0$, one has

$$(4.55) \quad \|\mathbf{P}_k^\phi\|_{\ell^2(\mathbf{n}_\phi)} \leq \|\mathbf{P}_k^{\sigma^2}\|_{\ell^2(\mathbf{n}_{\sigma^2})} \leq 1.$$

After recalling that $(\mathbf{P}_k^{\sigma^2})_{k \geq 0}$ is a complete orthonormal sequence in $\ell^2(\mathbf{n}_{\sigma^2})$, we observe that, for any $\mathbf{f} \in \ell^2(\mathbf{n}_\phi)$,

$$\sum_{k=0}^{\infty} \langle \mathbf{f}, \mathbf{P}_k^\phi \rangle_{\mathbf{n}_\phi} = \sum_{k=0}^{\infty} \langle \widehat{\mathbb{I}}_\phi \mathbf{f}, \mathbf{P}_k^{\sigma^2} \rangle_{\mathbf{n}_{\sigma^2}} = \|\widehat{\mathbb{I}}_\phi \mathbf{f}\|_{\ell^2(\mathbf{n}_{\sigma^2})}^2 \leq \|\mathbf{f}\|_{\ell^2(\mathbf{n}_\phi)}^2.$$

This shows that $(\mathbf{P}_k^\phi)_{k \geq 0}$ is a Bessel sequence in $\ell^2(\mathbf{n}_\phi)$. Combining item (1) with the existence of a biorthogonal sequence, see (4.60) below, we get that $(\mathbf{P}_k^\phi)_{k \geq 0}$ is exact, which proves (2). Finally, to prove (3), let $d_\epsilon = d_\phi - \epsilon$ for some $0 < \epsilon < d_\phi$ and define $\phi_{d_\epsilon}(u) = \frac{u\phi(u)}{u+d_\epsilon}$. From [29, Lemma 10.3], it follows that $\phi_{d_\epsilon} \in \mathbf{B}$ and

$$\lim_{u \rightarrow \infty} \frac{\phi_{d_\epsilon}(u)}{u} = \sigma^2.$$

Now, we need the following whose proof can be carried out by following a line of reasoning similar to the one of Theorem 4.14.

Lemma 4.18. For all $t \geq 0$,

$$(4.56) \quad \mathbb{K}_t^\phi \mathbb{I}_{\phi_{d_\epsilon}} = \mathbb{I}_{\phi_{d_\epsilon}} \mathbb{K}_t^{(d_\epsilon, \sigma^2)} \text{ on } \ell^2(\mathbf{n}_{d_\epsilon, \sigma^2})$$

where $\mathbb{K}^{(d_\epsilon, \sigma^2)}$ is the discrete Laguerre semigroup associated to $\phi(u) = \sigma^2(u + d_\epsilon)$ and $\mathbf{n}_{d_\epsilon, \sigma^2}$ denotes its invariant distribution.

Then, the proof of item (1) ensures that

$$\mathbf{P}_k^{(d_\epsilon, \sigma^2)}(n) = (1 + \sigma^{-2})^{-\frac{k}{2}} \Gamma(d_\epsilon + 1) \sum_{r=0}^k (-1)^r \binom{k}{r} \frac{\mathbf{p}_r(n)}{\Gamma(r + d_\epsilon + 1)}$$

is an eigenfunction of $\mathbb{K}_t^{(d_\epsilon, \sigma^2)}$ corresponding to the eigenvalue e^{-kt} . Therefore, using Lemma 4.18, we have that $\mathbb{I}_{\phi_{d_\epsilon}} \mathbf{P}_k^{(d_\epsilon, \sigma^2)}$ is an eigenfunction of \mathbb{K}_t^ϕ corresponding to the eigenvalue e^{-kt} , and, in fact,

$$\begin{aligned} \mathbb{I}_{\phi_{d_\epsilon}} \mathbf{P}_k^{(d_\epsilon, \sigma^2)}(n) &= (1 + \sigma^{-2})^{-\frac{k}{2}} \Gamma(d_\epsilon + 1) \sum_{r=0}^k \binom{k}{r} \frac{(-1)^r \mathbb{I}_{\phi_{d_\epsilon}} \mathbf{p}_r(n)}{\Gamma(r + d_\epsilon + 1)} \\ &= (1 + \sigma^{-2})^{-\frac{k}{2}} \sum_{r=0}^k \binom{k}{r} \frac{(-1)^r \mathbf{p}_r(n)}{W_\phi(r + 1)} = \mathbf{P}_k^\phi(n). \end{aligned}$$

Since the sequence $\left(\sqrt{\mathbf{c}_k(d_\epsilon)} \mathbf{P}_k^{(d_\epsilon, \sigma^2)} \right)_{k \geq 0}$ is an orthonormal sequence in $\ell^2(\mathbf{n}_{d_\epsilon, \sigma^2})$, see [21, equation (7)] or Example 3.1, and $\mathbb{I}_{\phi_{d_\epsilon}}$ is bounded, we deduce that $\left(\sqrt{\mathbf{c}_k(d_\epsilon)} \mathbf{P}_k^\phi \right)_{k \geq 0}$ is a Bessel sequence in $\ell^2(\mathbf{n}_\phi)$ and $\|\mathbf{P}_k^\phi\|_{\mathbf{n}_\phi} \leq \frac{1}{\sqrt{\mathbf{c}_k(d_\epsilon)}}$. Letting $\epsilon \downarrow 0$, the proof of (3) follows. \square

Proposition 4.19. Let $\phi \in \mathbf{B}$, and, for $k \in \mathbb{Z}_+$, \mathbf{V}_k^ϕ be defined as in (2.21). Then, the following holds.

(1) For all $k \in \mathbb{Z}_+$, $\mathbf{V}_k^\phi \in \ell^2(\mathbf{n}_\phi)$ and, for all $t \geq 0$,

$$\widehat{\mathbb{K}}_t^\phi \mathbf{V}_k^\phi = e^{-kt} \mathbf{V}_k^\phi.$$

(2) For all $k, l \in \mathbb{Z}_+$,

$$\left\langle \mathbf{P}_k^\phi, \mathbf{V}_l^\phi \right\rangle_{\mathbf{n}_\phi} = \mathbb{1}_{\{k=l\}}.$$

(3) If $\sigma^2 > 0$ and $\overline{\overline{\Pi}}(0) < \infty$, then $\left((1 + \sigma^{-2})^{-\frac{k}{2}} \frac{\mathbf{V}_k^\phi}{\sqrt{\mathbf{c}_k(\mathbf{m}_\phi)}} \right)_{k \geq 0}$ is a Bessel sequence in $\ell^2(\mathbf{n}_\phi)$ where we recall that $\mathbf{m}_\phi = \frac{m + \overline{\overline{\Pi}}(0)}{\sigma^2}$ and $\mathbf{c}_k(\mathbf{m}_\phi) = \frac{\Gamma(k + \mathbf{m}_\phi + 1)}{\Gamma(\mathbf{m}_\phi + 1)\Gamma(k + 1)}$.

Proof. Let us recall that for $\phi \in \mathbf{B}$ defined by

$$\phi(u) = m + \sigma^2 u + \int_0^\infty (1 - e^{-uy}) \bar{\Pi}(y) dy,$$

one has

$$\phi(\infty) = \lim_{u \rightarrow \infty} \phi(u) = \infty \mathbb{1}_{\{\sigma^2 > 0\}} + \left(\bar{\Pi}(0) + m \right) \mathbb{1}_{\{\sigma^2 = 0\}}$$

where $0 \leq \bar{\Pi}(0) = \int_0^\infty \bar{\Pi}(y) dy$. Finally, let $\mathbf{k}_\phi = \infty \mathbb{1}_{\{\sigma^2 > 0\}} + \frac{\bar{\Pi}(0)}{2\phi(\infty)}$, and define the set

$$(4.57) \quad \mathbb{Z}_\phi = \begin{cases} \mathbb{Z}_+ & \text{if } \mathbf{k}_\phi = \infty \\ \{k \in \mathbb{Z}_+; k < \mathbf{k}_\phi\} & \text{otherwise.} \end{cases}$$

When both $\bar{\Pi}(0) = \infty$ and $\phi(\infty) = \infty$, we have set $\frac{\bar{\Pi}(0)}{\phi(\infty)} = \infty$. Also, the condition (2.4) on Π implies that

$$\int_0^\infty (1 \wedge y) \bar{\Pi}(y) dy < \infty$$

and as a consequence, $\bar{\Pi}(0) < \infty$ whenever $\bar{\Pi}(0) < \infty$. Thus, $\mathbf{k}_\phi < \infty$ only when $\sigma^2 = 0$ and $\bar{\Pi}(0) < \infty$. It is shown in [29, Theorem 5.2], that $\nu_\phi \in \mathbf{C}_0^{|\mathbf{k}_\phi| - 1}(\mathbb{R}_+)$ and in [29, Theorem 1.11] that, for any $k \in \mathbb{Z}_\phi$, $\mathbf{V}_k^\phi \in \mathbf{L}^2(\nu_\phi)$, where

$$\mathbf{V}_k^\phi(x) = \frac{(1 + \sigma_1^{-1})^{\frac{k}{2}}}{k!} \frac{d^k}{dx^k} (x^k \nu_\phi(x)).$$

We now assume that $k \in \mathbb{Z}_\phi$ and recall from [29, Theorem 8.1], that, for all $t > 0$,

$$\widehat{K}_t^\phi \mathbf{V}_k^\phi = e^{-kt} \mathbf{V}_k^\phi.$$

Now, the intertwining relationship (4.32) entails that, for any $t > 0$,

$$\widehat{\mathbb{K}}_t^\phi \widehat{\Lambda}_\phi \mathbf{V}_k^\phi = \widehat{\Lambda}_\phi \widehat{K}_t^\phi \mathbf{V}_k^\phi = e^{-kt} \widehat{\Lambda}_\phi \mathbf{V}_k^\phi.$$

Let us now characterize the quantity $\widehat{\Lambda}_\phi \mathbf{V}_k^\phi$ when $k \in \mathbb{Z}_\phi$. From (4.33) it can be easily checked that $\widehat{\Lambda}_\phi \mathbf{V}_0^\phi(n) = \widehat{\Lambda}_\phi \mathbf{1}(n) = \mathbf{1}$. Writing $\varrho_1 = \frac{1}{2} \log(1 + \sigma_1^{-1})$,

for any $n, k \in \mathbb{N}$ we have

$$\begin{aligned}
e^{-k\varrho_1} \widehat{\Lambda}_\phi \mathbf{V}_k^\phi(n) &= \frac{1}{n! \mathbf{n}_\phi(n)} \int_0^\infty e^{-x} x^n \frac{d^k}{dx^k} (x^k \nu_\phi(x))}{k!} dx \\
(4.58) \quad &= \frac{(-1)^k}{n! \mathbf{n}_\phi(n) k!} \int_0^\infty \frac{d^k}{dx^k} (e^{-x} x^n) x^k \nu_\phi(x) dx \\
&= \frac{(-1)^k}{n! \mathbf{n}_\phi(n) k!} \sum_{j=0}^{k \wedge n} (-1)^{k-j} \binom{k}{j} \frac{n!}{(n-j)!} \int_0^\infty e^{-x} x^{k+n-j} \nu_\phi(x) dx \\
&= \frac{1}{\mathbf{n}_\phi(n)} \sum_{j=0}^{k \wedge n} (-1)^j \frac{(k+n-j)!}{(k-j)! (n-j)! j!} \mathbf{n}_\phi(k+n-j) \\
(4.59) \quad &= e^{-k\varrho_1} \mathbf{V}_k^\phi(n)
\end{aligned}$$

where we used, for the second identity, the fact that, for all $j = 1, \dots, k$,

$$\lim_{x \rightarrow 0, \phi(\infty)} \frac{d^{k-j}}{dx^{k-j}} (x^k \nu_\phi(x)) \frac{d^j}{dx^j} (e^{-x} x^n) = 0.$$

Indeed, these asymptotic behaviors are deduced easily from [29, Lemma 5.22], which states that for any $x > 0$, $0 \leq j \leq k$ and $a < d_\phi$, $\frac{d^{k-j}}{dx^{k-j}} (x^k \nu_\phi(x)) \leq Cx^{j+a}$ for some constant $C > 0$. Since $\widehat{\Lambda}_\phi : \mathbf{L}^2(\nu_\phi) \mapsto \ell^2(\mathbf{n}_\phi)$ is a bounded linear operator, see Proposition 4.11(3), we have that $\mathbf{V}_k^\phi = \widehat{\Lambda}_\phi \mathbf{V}_k^\phi \in \ell^2(\mathbf{n}_\phi)$ and this concludes the proof of (1) when $k \in \mathbb{Z}_\phi$. Now, let $\sigma^2 > 0$. Then, for any $k, l \in \mathbb{Z}_+$, we have, from Propositions 4.17 and the previous computation that both $\mathbf{P}_k^\phi, \mathbf{V}_l^\phi \in \ell^2(\mathbf{n}_\phi)$ and using (4.54) and (4.59), we obtain

$$(4.60) \quad \left\langle \mathbf{P}_k^\phi, \mathbf{V}_l^\phi \right\rangle_{\mathbf{n}_\phi} = \left\langle \mathbf{P}_k^\phi, \widehat{\Lambda}_\phi \mathbf{V}_l^\phi \right\rangle_{\mathbf{n}_\phi} = \left\langle \mathcal{P}_k^\phi, \mathbf{V}_l^\phi \right\rangle_{\nu_\phi} = \mathbb{1}_{\{k=l\}}$$

where we used that $(\mathcal{P}_k^\phi, \mathbf{V}_k^\phi)_{k \geq 0}$ is a biorthogonal sequence in $\mathbf{L}^2(\nu_\phi)$, see [29, Theorem 1.22(2)], recall that with the notation of this paper, $\mathcal{P}_k^\phi = (1 + \sigma^{-2})^{-\frac{k}{2}} \mathcal{P}_k$ and $\mathbf{V}_k^\phi = (1 + \sigma^{-2})^{\frac{k}{2}} \mathcal{V}_k$. This proves (2) when $\sigma^2 > 0$.

Next, assume that $k \notin \mathbb{Z}_\phi$ which implies that $\mathbf{k}_\phi < \infty$ and thus $\phi(\infty) < \infty$. This entails that the following two-sided bounds hold for any $n \in \mathbb{Z}_+$

$$(4.61) \quad e^{-\phi(\infty)} W_\phi(n+1) \leq \int_0^{\phi(\infty)} e^{-x} x^n \nu_\phi(x) dx \leq W_\phi(n+1) \leq \phi(\infty)^n$$

where the last inequality follows since ϕ is non-decreasing. Thus, we have

$$(4.62) \quad e^{-\phi(\infty)} \frac{W_\phi(n+1)}{n!} \leq \mathbf{n}_\phi(n) \leq \frac{W_\phi(n+1)}{n!} \leq \frac{\phi(\infty)^n}{n!}.$$

Hence, for any $k \in \mathbb{Z}_+$ fixed, with

$$S(k) = \sum_{n=0}^k \frac{1}{\mathbf{n}_\phi(n)} \left(\sum_{j=0}^n (-1)^j \frac{(k+n-j)!}{(k-j)!(n-j)!j!} \mathbf{n}_\phi(k+n-j) \right)^2 < \infty,$$

we have

$$\begin{aligned} e^{-2k\varrho_1} \|\mathbf{V}_k^\phi\|_{\ell^2(\mathbf{n}_\phi)}^2 &= S(k) + \sum_{n=k}^{\infty} \frac{1}{\mathbf{n}_\phi(n)} \left(\sum_{j=0}^k (-1)^j \frac{(k+n-j)!}{(k-j)!(n-j)!j!} \mathbf{n}_\phi(k+n-j) \right)^2 \\ &\leq S(k) + e^{\phi(\infty)} \sum_{n=k}^{\infty} \frac{n!}{W_\phi(n+1)} \left(\sum_{j=0}^k \frac{(k+n-j)!}{(k-j)!(n-j)!j!} \frac{\phi(\infty)^{k+n-j}}{(k+n-j)!} \right)^2 \\ &\leq S(k) + e^{\phi(\infty)} \sum_{n=k}^{\infty} \frac{n! \phi(\infty)^{2n}}{W_\phi(n+1) ((n-k)!)^2} \left(\sum_{j=0}^k \frac{\phi(\infty)^{k-j}}{(k-j)!j!} \right)^2 \\ (4.63) \quad &\leq S(k) + e^{\phi(\infty)} \left(\sum_{j=0}^k \frac{\phi(\infty)^{k-j}}{(k-j)!j!} \right)^2 \sum_{n=k}^{\infty} \frac{n! \phi(\infty)^{2n}}{W_\phi(n+1) ((n-k)!)^2} < \infty \end{aligned}$$

where the last inequality follows after observing that,

$$\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = \lim_{n \rightarrow \infty} \frac{(n+1)\phi(\infty)^2}{\phi(n+1)(n+1-k)^2} = 0$$

where the a_n 's are the coefficient of the last series. When $\phi \in \mathbf{B}$ is such that $\phi(u) = m + \int_0^\infty (1 - e^{-uy}) \bar{\Pi}(y) dy$, let us define $\phi_\epsilon \in \mathbf{B}$ as $\phi_\epsilon(u) = \epsilon u + \phi(u)$ with $\epsilon > 0$. Then, from Proposition 4.12, it follows that for small values of ϵ and for all $n \in \mathbb{Z}_+$,

$$\begin{aligned} \mathbf{n}_{\phi_\epsilon}(n) &= \frac{1}{n!} \sum_{r=0}^{\infty} (-1)^r \frac{W_{\phi_\epsilon}(n+r+1)}{r!}, \\ \mathbf{n}_\phi(n) &= \frac{1}{n!} \sum_{r=0}^{\infty} (-1)^r \frac{W_\phi(n+r+1)}{r!}. \end{aligned}$$

As $\phi_\epsilon(u) \geq \phi(u)$ for all ϵ and $u \geq 0$, it follows that $W_{\phi_\epsilon}(n) \geq W_\phi(n)$ for all $n \in \mathbb{N}$. Also, $W_{\phi_\epsilon} \downarrow W_\phi$ pointwise as $\epsilon \rightarrow 0$. Since, for small values of ϵ (e.g. $0 \leq \epsilon < 1$),

$$\sum_{r=0}^{\infty} \frac{W_{\phi_\epsilon}(r+n+1)}{r!} < \infty,$$

the dominated convergence theorem yields the following pointwise convergence as $\epsilon \rightarrow 0$,

$$(4.64) \quad \mathbf{n}_{\phi_\epsilon} \rightarrow \mathbf{n}_\phi.$$

Hence, for any $j, k \in \mathbb{Z}_+$,

$$(4.65)$$

$$\begin{aligned} \lim_{\epsilon \rightarrow 0} \langle V_k^{\phi_\epsilon}, \mathbf{p}_j \rangle_{\mathbf{n}_{\phi_\epsilon}} &= \lim_{\epsilon \rightarrow 0} \sum_{n=0}^{\infty} \mathbf{p}_j(n) V_k^{\phi_\epsilon}(n) \mathbf{n}_{\phi_\epsilon}(n) \\ &= \lim_{\epsilon \rightarrow 0} \sum_{n=0}^{\infty} \mathbf{p}_j(n) \sum_{j=0}^{k \wedge n} (-1)^{k-j} \frac{(k+n-j)!}{(k-j)!(n-j)!j!} \mathbf{n}_{\phi_\epsilon}(k+n-j) \\ (4.66) \quad &= \lim_{\epsilon \rightarrow 0} \sum_{n=0}^k \mathbf{p}_j(n) \sum_{j=0}^n (-1)^{k-j} \frac{(k+n-j)!}{(k-j)!(n-j)!j!} \mathbf{n}_{\phi_\epsilon}(k+n-j) \\ &\quad + \lim_{\epsilon \rightarrow 0} \sum_{n=k}^{\infty} \mathbf{p}_j(n) \sum_{j=0}^k (-1)^{k-j} \frac{(k+n-j)!}{(k-j)!(n-j)!j!} \mathbf{n}_{\phi_\epsilon}(k+n-j). \end{aligned}$$

In (4.66), the first term is a finite sum and therefore

$$\begin{aligned} &\lim_{\epsilon \rightarrow 0} \sum_{n=0}^k \mathbf{p}_j(n) \sum_{j=0}^n (-1)^{k-j} \frac{(k+n-j)!}{(k-j)!(n-j)!j!} \mathbf{n}_{\phi_\epsilon}(k+n-j) \\ &= \sum_{n=0}^k \mathbf{p}_j(n) \sum_{j=0}^n (-1)^{k-j} \frac{(k+n-j)!}{(k-j)!(n-j)!j!} \mathbf{n}_\phi(k+n-j). \end{aligned}$$

For the second term in (4.66), we have

$$\begin{aligned} &\sum_{n=k}^{\infty} \mathbf{p}_j(n) \sum_{j=0}^k (-1)^{k-j} \frac{(k+n-j)!}{(k-j)!(n-j)!j!} \mathbf{n}_{\phi_\epsilon}(k+n-j) \\ &= \sum_{j=0}^k (-1)^{k-j} \frac{1}{j!(k-j)!} \sum_{n=k}^{\infty} \mathbf{p}_j(n) \frac{(k+n-j)!}{(n-j)!} \mathbf{n}_{\phi_\epsilon}(k+n-j). \end{aligned}$$

Since $\mathbf{n}_{\phi_\epsilon} \rightarrow \mathbf{n}_\phi$ pointwise as $\epsilon \rightarrow 0$, the distribution $\mathbf{n}_{\phi_\epsilon}$ converges to \mathbf{n}_ϕ weakly. Also, for any $k \in \mathbb{Z}_+$, as $\epsilon \rightarrow 0$,

$$\sum_{n=0}^{\infty} \mathbf{p}_k(n) \mathbf{n}_{\phi_\epsilon}(n) = W_{\phi_\epsilon}(k+1) \rightarrow W_\phi(k+1) = \sum_{n=0}^{\infty} \mathbf{p}_k(n) \mathbf{n}_\phi(n).$$

Applying (4.23) on the previous identity we obtain that, for all $k \in \mathbb{Z}_+$, as $\epsilon \rightarrow 0$,

$$(4.67) \quad \sum_{n=0}^{\infty} n^k \mathbf{n}_{\phi_\epsilon}(n) \rightarrow \sum_{n=0}^{\infty} n^k \mathbf{n}_\phi(n).$$

Since, for any $k \in \mathbb{Z}_+$ and $j \leq k$,

$$\frac{(k+n-j)!}{(n-j)!} = O(n^k)$$

uniformly with respect to j , using a dominated convergence theorem one can show that for each $j \leq k$,

$$\lim_{\epsilon \rightarrow 0} \sum_{n=k}^{\infty} \mathbf{p}_j(n) \frac{(k+n-j)!}{(n-j)!} \mathbf{n}_{\phi_\epsilon}(k+n-j) = \sum_{n=k}^{\infty} \mathbf{p}_j(n) \frac{(k+n-j)!}{(n-j)!} \mathbf{n}_\phi(k+n-j).$$

Thus, (4.66) yields

$$(4.68) \quad \lim_{\epsilon \rightarrow 0} \left\langle \mathbf{V}_k^{\phi_\epsilon}, \mathbf{p}_j \right\rangle_{\mathbf{n}_{\phi_\epsilon}} = \left\langle \mathbf{V}_k^\phi, \mathbf{p}_j \right\rangle_{\mathbf{n}_\phi}.$$

Now, if $\sigma^2 = 0$, since, for any $k \in \mathbb{Z}_+$, $\mathbf{P}_k^\phi \in \mathcal{P} = \text{Span}\{\mathbf{p}_j, j \in \mathbb{Z}_+\}$ and the coefficient of \mathbf{p}_j in $\mathbf{P}_k^{\phi_\epsilon}$ converges to the same in \mathbf{P}_k^ϕ for all $j \in \mathbb{Z}_+$, as $\epsilon \rightarrow 0$, applying (4.68) it follows that, for all $k, l \in \mathbb{Z}_+$,

$$\mathbb{1}_{\{k=l\}} = \lim_{\epsilon \rightarrow 0} \left\langle \mathbf{P}_k^{\phi_\epsilon}, \mathbf{V}_l^{\phi_\epsilon} \right\rangle_{\mathbf{n}_{\phi_\epsilon}} = \left\langle \mathbf{P}_k^\phi, \mathbf{V}_l^\phi \right\rangle_{\mathbf{n}_\phi}$$

where $\phi_\epsilon(z) = \epsilon u + \phi(u)$. This proves (2) for all $\sigma^2 \geq 0$ hence for all $\phi \in \mathbf{B}$.

To show that \mathbf{V}_k^ϕ is a co-eigenfunction of \mathbb{K}^ϕ when $\sigma^2 = 0$, we proceed as follows. Proposition 4.17(1) and (4.60) yield that, for $l, k \in \mathbb{Z}_+$, and $t > 0$,

$$\left\langle \widehat{\mathbb{K}}_t^\phi \mathbf{V}_k^\phi, \mathbf{P}_l^\phi \right\rangle_{\mathbf{n}_\phi} = \left\langle \mathbf{V}_k^\phi, \mathbb{K}_t^\phi \mathbf{P}_l^\phi \right\rangle_{\mathbf{n}_\phi} = e^{-tk} \left\langle \mathbf{V}_k^\phi, \mathbf{P}_l^\phi \right\rangle_{\mathbf{n}_\phi} = e^{-tk} \mathbb{1}_{\{k=l\}}.$$

Therefore, for all $\phi \in \mathbf{B}$, $t > 0$ and $k, l \in \mathbb{Z}_+$, we get

$$\left\langle \widehat{\mathbb{K}}_t^\phi \mathbf{V}_k^\phi - e^{-tk} \mathbf{V}_k^\phi, \mathbf{P}_l^\phi \right\rangle_{\mathbf{n}_\phi} = 0.$$

Since $(\mathbf{P}_k^\phi)_{k \geq 0}$ is dense in $\ell^2(\mathbf{n}_\phi)$, we deduce that, for all $t \geq 0$ and $k \in \mathbb{Z}_+$,

$$(4.69) \quad e^{tk} \widehat{\mathbb{K}}_t^\phi \mathbf{V}_k^\phi = \mathbf{V}_k^\phi,$$

which proves (1) for all $\phi \in \mathbf{B}$.

To prove item (3), it is known from [29, Theorem 10.1(1)] (after multiplying by the factor $(1 + \sigma^{-2})^{-\frac{k}{2}}$) that, when $\sigma^2 > 0$ and $\overline{\Pi}(0) < \infty$,

$\left((1 + \sigma^{-2})^{-\frac{k}{2}} \frac{V_k^\phi}{\sqrt{c_k(\mathbf{m}_\phi)}} \right)_{k \geq 0}$ is a Bessel sequence in $\mathbf{L}^2(\nu_\phi)$. Recalling that, for any $k \geq 0$, $\widehat{\Lambda}_\phi V_k^\phi = V_k^\phi$, see (4.59), and $\widehat{\Lambda}_\phi : \mathbf{L}^2(\nu_\phi) \rightarrow \ell^2(\mathbf{n}_\phi)$ is a contraction, we conclude that $\left((1 + \sigma^{-2})^{-\frac{k}{2}} \frac{V_k^\phi}{\sqrt{c_k(\mathbf{m}_\phi)}} \right)_{k \geq 0}$ is a Bessel sequence in $\ell^2(\mathbf{n}_\phi)$ and for all $k \in \mathbb{Z}_+$,

$$\|V_k^\phi\|_{\ell^2(\mathbf{n}_\phi)} \leq (1 + \sigma^{-2})^{\frac{k}{2}} \sqrt{c_k(\mathbf{m}_\phi)}.$$

□

4.12. Proof of Theorem 2.10. The proof of the item (1) and (2) follows directly from Proposition 4.17(1) and Proposition 4.19(1),(2).

Finally for the proof of (3), we now assume that $\sigma^2 > 0$. We recall from (2.19) that $\sigma_1 = \sigma^2$ in this case. Then, for all $\mathbf{f} \in \ell^2(\mathbf{n}_{\sigma^2})$ and $t > 0$, the intertwining relation (4.46) yields that

$$\begin{aligned} \mathbb{K}_t^\phi \mathbb{I}_\phi \mathbf{f} &= \mathbb{I}_\phi \mathbb{K}_t^{\sigma^2} \mathbf{f} \\ &= \mathbb{I}_\phi \sum_{k=0}^{\infty} e^{-kt} \langle \mathbf{f}, \mathbf{P}_k^{\sigma^2} \rangle_{\mathbf{n}_{\sigma^2}} \mathbf{P}_k^{\sigma^2} \\ (4.70) \quad &= \sum_{k=0}^{\infty} e^{-kt} \langle \mathbf{f}, \mathbf{P}_k^{\sigma^2} \rangle_{\mathbf{n}_{\sigma^2}} \mathbf{P}_k^\phi \end{aligned}$$

where the second identity relies on the spectral decomposition of the reversible birth-death chain, see Example 3.1 with $\phi(u) = \sigma^2 u$, whereas the last one is justified as follows. First, since $\sigma^2 > 0$, $\mathbb{I}_\phi : \ell^2(\mathbf{n}_{\sigma^2}) \mapsto \ell^2(\mathbf{n}_\phi)$ is a bounded linear operator, and, with the help of Lemma 4.16 and the definition of \mathbf{P}_k^ϕ in (2.20), it follows that $\mathbb{I}_\phi \mathbf{P}_k^{\sigma^2} = \mathbf{P}_k^\phi$. Moreover, from Proposition 4.17, we have that the sequence $(\mathbf{P}_k^\phi)_{k \geq 0}$ is a Bessel sequence and thus its associated synthesis operator $\mathcal{S} : \ell^2(\mathbb{Z}_+) \mapsto \ell^2(\mathbf{n}_\phi)$, see (4.51) for definition, is bounded. Since $(\mathbf{P}_k^{\sigma^2})_{k \geq 0}$ is an orthonormal sequence in $\ell^2(\mathbf{n}_{\sigma^2})$, it implies that for all $t \geq 0$,

$$\left(e^{-kt} \langle \mathbf{f}, \mathbf{P}_k^{\sigma^2} \rangle_{\mathbf{n}_{\sigma^2}} \right)_{k \geq 0} \in \ell^2(\mathbb{Z}_+)$$

and hence the series on the right-hand side of (4.70) is in $\ell^2(\mathbf{n}_\phi)$. Next, as noted before, we have that $\mathbb{I}_\phi \mathbf{P}_k^{\sigma^2} = \mathbf{P}_k^\phi$ for all $k \in \mathbb{Z}_+$. Now, recalling that $(\mathbf{P}_k^\phi, \mathbf{V}_k^\phi)_{k \geq 0}$ is biorthogonal in $\ell^2(\mathbf{n}_\phi)$, see Proposition 4.19(2), we have for

any $l, k \in \mathbb{Z}_+$,

$$\left\langle \mathbf{P}_l^{\sigma^2}, \widehat{\mathbb{I}}_\phi \mathbf{V}_k^\phi \right\rangle_{\mathbf{n}_{\sigma^2}} = \left\langle \mathbb{I}_\phi \mathbf{P}_l^{\sigma^2}, \mathbf{V}_k^\phi \right\rangle_{\mathbf{n}_\phi} = \left\langle \mathbf{P}_l^\phi, \mathbf{V}_k^\phi \right\rangle_{\mathbf{n}_\phi} = \mathbb{1}_{\{k=l\}}$$

As $(\mathbf{P}_k^{\sigma^2})_{k \geq 0}$ is orthonormal in $\ell^2(\mathbf{n}_{\sigma^2})$ (hence biorthogonal to itself), by uniqueness of biorthogonal sequence we conclude that $\widehat{\mathbb{I}}_\phi \mathbf{V}_k^\phi = \mathbf{P}_k^{\sigma^2}$ for all $k \in \mathbb{Z}_+$. Therefore, writing $\mathbf{g} = \mathbb{I}_\phi \mathbf{f} \in \ell^2(\mathbf{n}_\phi)$, we have, for all $k \in \mathbb{Z}_+$,

$$\left\langle \mathbf{g}, \mathbf{V}_k^\phi \right\rangle_{\mathbf{n}_\phi} = \left\langle \mathbf{f}, \widehat{\mathbb{I}}_\phi \mathbf{V}_k^\phi \right\rangle_{\mathbf{n}_{\sigma^2}} = \left\langle \mathbf{f}, \mathbf{P}_k^{\sigma^2} \right\rangle_{\mathbf{n}_{\sigma^2}}$$

Thus, from (4.70), for $\mathbf{g} \in \text{Ran}(\mathbb{I}_\phi)$, the range of \mathbb{I}_ϕ , one gets

$$\mathbb{K}_t^\phi \mathbf{g} = \sum_{k=0}^{\infty} e^{-kt} \left\langle \mathbf{g}, \mathbf{V}_k^\phi \right\rangle_{\mathbf{n}_\phi} \mathbf{P}_k^\phi = \mathbb{S}_t \mathbf{g}$$

where the last identity serves as defining the spectral operator. Note that since $\langle \mathbb{K}_t^\phi \mathbf{g}, \mathbf{V}_k^\phi \rangle_{\mathbf{n}_\phi} = \langle \mathbf{g}, \widehat{\mathbb{K}}_t^\phi \mathbf{V}_k^\phi \rangle_{\mathbf{n}_{\sigma^2}} = e^{-kt} \langle \mathbf{g}, \mathbf{V}_k^\phi \rangle_{\mathbf{n}_\phi}$, we deduce that

$$\mathbb{S}_t \mathbf{g} = \sum_{k=0}^{\infty} \left\langle \mathbb{K}_t^\phi \mathbf{g}, \mathbf{V}_k^\phi \right\rangle_{\mathbf{n}_\phi} \mathbf{P}_k^\phi.$$

Moreover, as the closure (in $\ell^2(\mathbf{n}_\phi)$) of $\text{Ran}(\mathbb{I}_\phi)$ is $\ell^2(\mathbf{n}_\phi)$, by the bounded linear transformation theorem, \mathbb{K}_t^ϕ is the unique continuous extension of the continuous operator $\mathbb{S}_t : \text{Ran}(\mathbb{I}_\phi) \mapsto \ell^2(\mathbf{n}_\phi)$. We now extend the domain of \mathbb{S}_t to $\ell^2(\mathbf{n}_\phi)$. First, by means of Cauchy-Schwartz inequality, we have, for any $\mathbf{g} \in \ell^2(\mathbf{n}_\phi)$ and $k \in \mathbb{N}$,

$$\begin{aligned} \left| \left\langle \mathbf{g}, \mathbf{V}_k^\phi \right\rangle_{\mathbf{n}_\phi} \right| &\leq \|\mathbf{g}\|_{\ell^2(\mathbf{n}_\phi)} \left\| \mathbf{V}_k^\phi \right\|_{\ell^2(\mathbf{n}_\phi)} = \|\mathbf{g}\|_{\ell^2(\mathbf{n}_\phi)} \|\widehat{\Lambda}_\phi \mathbf{V}_k^\phi\|_{\ell^2(\mathbf{n}_\phi)} \\ &\leq \|\mathbf{g}\|_{\ell^2(\mathbf{n}_\phi)} \left\| \mathbf{V}_k^\phi \right\|_{\mathbf{L}^2(\nu_\phi)} \end{aligned}$$

where we used Proposition 4.19 and the fact that $\widehat{\Lambda}_\phi$ is a bounded operator. Next, since from [29, Theorem 10.1], we have for k large enough and all $\epsilon > 0$, $\|\mathbf{V}_k^\phi\|_{\mathbf{L}^2(\nu_\phi)} \leq C_\epsilon (1 + \sigma^{-2})^{\frac{k}{2}} e^{\epsilon k}$, with $C_\epsilon > 0$, this implies that for all $\mathbf{g} \in \ell^2(\mathbf{n}_\phi)$ and $t > \frac{1}{2} \log(1 + \sigma^{-2})$,

$$\left(e^{-kt} \left\langle \mathbf{g}, \mathbf{V}_k^\phi \right\rangle_{\mathbf{n}_\phi} \right)_{k \geq 0} \in \ell^2(\mathbb{Z}_+).$$

Finally, the Bessel property of the sequence $(\mathbf{P}_k^\phi)_{k \geq 0}$ entails that $\mathbb{S}_t \mathbf{g} \in \ell^2(\mathbf{n}_\phi)$, which completes the proof.

For the item (4), we recall from [29, Theorem 10.1] and the proof of Theorem 2.10(3) that, for all $\epsilon > 0$ and $k \in \mathbb{Z}_+$, there exists $C_\epsilon > 0$ such

that

$$(4.71) \quad \left\| \mathbf{V}_k^\phi \right\|_{\ell^2(\mathbf{n}_\phi)} \leq \left\| \mathbf{V}_k^\phi \right\|_{\mathbf{L}^2(\nu_\phi)} \leq C_\epsilon (1 + \sigma^{-2})^{\frac{k}{2}} e^{\epsilon k}$$

whenever $\sigma^2 > 0$. Moreover, for all $k \in \mathbb{Z}_+$, $\|\mathbf{P}_k^\phi\|_{\ell^2(\mathbf{n}_\phi)} \leq 1$. Therefore, (2.22) entails that for all $t > \frac{1}{2} \log(1 + \sigma^{-2})$, the operator \mathbb{K}_t^ϕ can be approximated by the sequence of finite dimensional operators

$$\mathbf{f} \mapsto \sum_{k=0}^N e^{-kt} \left\langle \mathbf{f}, \mathbf{V}_k^\phi \right\rangle_{\mathbf{n}_\phi} \mathbf{P}_k^\phi, \quad N \geq 1,$$

which proves the compactness of the semigroup. Finally, for $l \in \mathbb{Z}_+$, let us choose $\mathbf{f} = \delta_l$ in (2.22). Then, for all $\sigma^2 > 0, t > 0$ and $n \in \mathbb{Z}_+$, we have

$$(4.72) \quad \mathbb{K}_t^\phi(n, l) = \mathbb{K}_t^\phi \delta_l(n) = \sum_{k=0}^{\infty} e^{-kt} \mathbf{P}_k^\phi(n) \mathbf{V}_k^\phi(l) \mathbf{n}_\phi(l)$$

where the last identity holds in $\ell^2(\mathbf{n}_\phi)$. Now, from Proposition 4.17(2), we have that, for all $k, n \in \mathbb{Z}_+$,

$$\mathbf{P}_k^\phi(n)^2 \mathbf{n}_\phi(n) \leq \|\mathbf{P}_k^\phi\|_{\ell^2(\mathbf{n}_\phi)}^2 \leq 1$$

while, from Jensen's inequality and (4.71), we get, for all $k, l \in \mathbb{Z}_+$, that there exists a uniform constant $C_\epsilon > 0$ such that for all $\epsilon > 0$,

$$(4.73) \quad |\mathbf{V}_k^\phi(l) \mathbf{n}_\phi(l)| \leq \|\mathbf{V}_k^\phi\|_{\ell^1(\mathbf{n}_\phi)} \leq \|\mathbf{V}_k^\phi\|_{\ell^2(\mathbf{n}_\phi)} \leq C_\epsilon (1 + \sigma^{-2})^{\frac{k}{2}} e^{\epsilon k}.$$

Since $\mathbf{n}_\phi(n) > 0$ for all $n \in \mathbb{Z}_+$, see (4.35), for all $\frac{1}{2} \log(1 + \sigma^{-2}) + \epsilon < t$, we have

$$(4.74) \quad \sum_{k=0}^{\infty} e^{-kt} |\mathbf{P}_k^\phi(n)| |\mathbf{V}_k^\phi(l)| \mathbf{n}_\phi(l) \leq C_\epsilon \sum_{k=0}^{\infty} e^{-kt} \frac{e^{\epsilon k}}{\sqrt{\mathbf{n}_\phi(n)}} < \infty.$$

As $\epsilon > 0$ is arbitrary, the proof of the item (5) is completed.

4.13. Proof of Theorem 2.12(1). From [29, Lemma 10.4], we get that

$$m_\phi = \lim_{n \rightarrow \infty} \frac{\phi(n) - \sigma^2 n}{\sigma^2} = \frac{m + \int_0^\infty \Pi(y, \infty) dy}{\sigma^2} > d_\epsilon = (d_\phi - \epsilon) \mathbb{1}_{\{d_\phi - \epsilon > 0\}}.$$

Let us write $\varrho = \frac{1}{2} \log(1 + \sigma^{-2})$. Then, using (2.22) along with the fact that $\mathbf{P}_0^\phi \equiv 1$, we obtain, for all $\mathbf{f} \in \ell_0^2(\mathbf{n}_\phi) = \{\mathbf{g} \in \ell^2(\mathbf{n}_\phi); \mathbf{n}_\phi \mathbf{g} = 0\}$,

$$(4.75) \quad \begin{aligned} \mathbb{K}_t^\phi \mathbf{f} &= \sum_{k=1}^{\infty} e^{-kt} \left\langle \mathbf{f}, \mathbf{V}_k^\phi \right\rangle_{\mathbf{n}_\phi} \mathbf{P}_k^\phi \\ &= \sum_{k=1}^{\infty} e^{-kt} \sqrt{\frac{\mathbf{c}_k(\mathbf{m}_\phi)}{\mathbf{c}_k(\mathbf{d}_\phi)}} e^{k\varrho} \left\langle \mathbf{f}, e^{-k\varrho} \frac{\mathbf{V}_k^\phi}{\sqrt{\mathbf{c}_k(\mathbf{m}_\phi)}} \right\rangle_{\mathbf{n}_\phi} \sqrt{\mathbf{c}_k(\mathbf{d}_\phi)} \mathbf{P}_k^\phi. \end{aligned}$$

Since $(\sqrt{\mathbf{c}_k(\mathbf{d}_\phi)}\mathbf{P}_k^\phi)_{k \geq 1}$ is a Bessel sequence with bound 1, we obtain from the boundedness of the synthesis operator, see (4.51), and (4.75) that, writing $\overline{\mathbf{V}}_k^\phi = \frac{\mathbf{V}_k^\phi}{\sqrt{\mathbf{c}_k(\mathbf{m}_\phi)}}$, for all $\mathbf{f} \in \ell_0^2(\mathbf{n}_\phi)$,

$$(4.76) \quad \begin{aligned} \|\mathbb{K}_t^\phi \mathbf{f}\|_{\ell^2(\mathbf{n}_\phi)}^2 &\leq \sum_{k=1}^{\infty} e^{-2kt} \frac{\mathbf{c}_k(\mathbf{m}_\phi)}{\mathbf{c}_k(\mathbf{d}_\phi)} e^{2k\varrho} \left| \left\langle \mathbf{f}, e^{-k\varrho} \overline{\mathbf{V}}_k^\phi \right\rangle_{\mathbf{n}_\phi} \right|^2 \\ &= e^{-2t} \frac{\mathbf{c}_1(\mathbf{m}_\phi)}{\mathbf{c}_1(\mathbf{d}_\phi)} e^{2\varrho} \sum_{k=1}^{\infty} e^{-2(k-1)(t-\varrho)} \frac{\mathbf{c}_k(\mathbf{m}_\phi)\mathbf{c}_1(\mathbf{d}_\phi)}{\mathbf{c}_k(\mathbf{d}_\phi)\mathbf{c}_1(\mathbf{m}_\phi)} \left| \left\langle \mathbf{f}, e^{-k\varrho} \overline{\mathbf{V}}_k^\phi \right\rangle_{\mathbf{n}_\phi} \right|^2. \end{aligned}$$

Now, from the proof of [29, Theorem 1.18(3)], we know that

$$\sup_{k \geq 1} e^{-2(k-1)(t-\varrho)} \frac{\mathbf{c}_k(\mathbf{m}_\phi)\mathbf{c}_1(\mathbf{d}_\phi)}{\mathbf{c}_k(\mathbf{d}_\phi)\mathbf{c}_1(\mathbf{m}_\phi)} \leq 1 \iff t > T = \frac{1}{2} \log \left(\frac{\mathbf{m}_\phi + 2}{\mathbf{d}_\phi + 2} \right) + \varrho.$$

Thus, using this bound, the fact that $(e^{-k\varrho} \overline{\mathbf{V}}_k^\phi)_{k \geq 1}$ is also a Bessel sequence (with bound 1) in $\ell^2(\mathbf{n}_\phi)$ and the second inequality in (4.50), we deduce from (4.76) that, for all $\mathbf{f} \in \ell_0^2(\mathbf{n}_\phi)$ and $t > T$,

$$(4.77) \quad \|\mathbb{K}_t^\phi \mathbf{f}\|_{\ell^2(\mathbf{n}_\phi)}^2 \leq e^{2t\sigma_1^2} \frac{\mathbf{c}_1(\mathbf{m}_\phi)}{\mathbf{c}_1(\mathbf{d}_\phi)} e^{-2t} \|\mathbf{f}\|_{\ell^2(\mathbf{n}_\phi)}^2 = \frac{1 + \sigma_1^2 \mathbf{m}_\phi + 1}{\sigma_1^2 \mathbf{d}_\phi + 1} e^{-2t} \|\mathbf{f}\|_{\ell^2(\mathbf{n}_\phi)}^2.$$

When $t \leq T$, $\frac{1 + \sigma_1^2 \mathbf{m}_\phi + 1}{\sigma_1^2 \mathbf{d}_\phi + 1} e^{-2t} \geq \frac{\mathbf{m}_\phi + 1}{\mathbf{d}_\phi + 1} \frac{\mathbf{d}_\phi + 2}{\mathbf{m}_\phi + 2} \geq 1$ as $\mathbf{m}_\phi \geq \mathbf{d}_\phi$. Therefore, (4.77) holds for all $t > 0$ as \mathbb{K}^ϕ is a contraction semigroup. Finally, noting that, for any $\mathbf{f} \in \ell^2(\mathbf{n}_\phi)$, $\mathbf{f} - \mathbf{n}_\phi \mathbf{f} \in \ell_0^2(\mathbf{n}_\phi)$, the proof of the theorem follows.

4.14. Interweaving between skip-free and continuous Laguerre semigroups. Following [27], for two Markov semigroups P, P' defined on two Banach spaces \mathbf{B}, \mathbf{B}' respectively, we say that P has an **interweaving relation** with P' if there exist two Markov kernels $\Lambda : \mathbf{B}' \rightarrow \mathbf{B}$ and $\Lambda' : \mathbf{B} \rightarrow \mathbf{B}'$, and a non-negative random variable τ such that

$$\begin{aligned} P\Lambda &= \Lambda P' \text{ on } \mathbf{B}' \\ P'\Lambda' &= \Lambda' P \text{ on } \mathbf{B} \text{ and} \\ \Lambda\Lambda' &= P_\tau = \int_0^\infty P_t \mathbb{P}(\tau \in dt). \end{aligned}$$

We call τ the **warm-up time** or the **delay** and we write $P \leftarrow P'$ or $P \overset{\tau}{\leftarrow} P'$ to emphasize the dependence on τ . Note that when $\tau = \delta_{t_0}$ is the degenerate random variable at $t_0 > 0$, we may simply write, when there is no confusion, $P \overset{t_0}{\leftarrow} P'$.

When τ is in addition infinitely divisible we say that P admits an **interweaving relation with an infinitely divisible** warm-up time with P' and we write $P \overset{\tau}{\leftarrow} P'$. Finally, when we also have

$$(4.78) \quad \Lambda\Lambda' = P'_\tau$$

we say that there is a **symmetric interweaving relation** between P and P' and we write $P \overset{\tau}{\leftrightarrow} P'$. We refer to [27] for a thorough study and several applications of this concept that refines the one of intertwining relations.

Let now \mathbb{K}^ϕ be the skip-free Laguerre semigroup corresponding to the Bernstein function ϕ associated with the triplet (m, σ^2, Π) , see (2.5), and K^{σ^2} be the diffusive Laguerre semigroup with generator

$$(4.79) \quad L^{\sigma^2} = \sigma^2 x \frac{d^2}{dx^2} + (\sigma^2 - x) \frac{d}{dx}.$$

Theorem 4.20. *If $\sigma^2 > 0$ and $\bar{\Pi}(0) = \int_0^\infty \Pi(y, \infty) dy < \infty$, then for all $\beta > m_\phi = \frac{m + \bar{\Pi}(0)}{\sigma^2}$,*

$$\mathbb{K}^\phi \overset{\tau_\beta}{\leftrightarrow} K^{\sigma^2}$$

where τ_β is an infinite divisible random variable characterized, for any $u > 0$, by

$$(4.80) \quad \begin{aligned} \int_0^\infty e^{-ut} \mathbb{P}(\tau_\beta \in dt) &= \left(\frac{\sigma^2}{1 + \sigma^2} \right)^u \frac{\Gamma(1 + \beta) \Gamma(u + 1)}{\Gamma(u + \beta + 1)} \\ &= \left(\frac{\sigma^2}{1 + \sigma^2} \right)^u e^{-\phi_\beta(u)}. \end{aligned}$$

Before proving the theorem, let us show the following lemma.

Lemma 4.21. *Let K^{σ^2} be the semigroup defined as above and K be the semigroup with generator*

$$L = x \frac{d^2}{dx^2} + (1 - x) \frac{d}{dx}.$$

Then, for all $t \geq 0$,

$$K_t^{\sigma^2} d_{\frac{1}{\sigma^2}} = d_{\frac{1}{\sigma^2}} K_t \text{ on } \mathbf{L}^2(\nu)$$

where for $\alpha > 0$, $d_\alpha f(x) = f(\alpha x)$ is the dilation operator on \mathbb{R}_+ and $\nu(x) dx = e^{-x} dx, x > 0$, is the unique invariant distribution of the semigroup K .

Proof. It can be easily checked that if f is a polynomial, then

$$(4.81) \quad L^{\sigma^2} d_{\frac{1}{\sigma^2}} f = d_{\frac{1}{\sigma^2}} Lf.$$

Next, we recall from [29, Theorem 1.6(3)] that the set of all polynomials form a core for L in $\mathbf{L}^2(\nu)$ and $d_{\frac{1}{\sigma^2}} \nu_{\sigma^2} = \nu$ where ν_{σ^2} is the invariant distribution of K^{σ^2} . Since

$$d_{\frac{1}{\sigma^2}} : \mathbf{L}^2(\nu) \rightarrow \mathbf{L}^2(\nu_{\sigma^2})$$

is an invertible operator, (4.81) extends at the level of the corresponding semigroups, which proves the lemma. \square

4.15. Proof of Theorem 4.20. Let $K^{(\beta)}$ be the self-adjoint Laguerre semigroup with the generator

$$L^{(\beta)} = x \frac{d^2}{dx^2} + (1 + \beta - x) \frac{d}{dx}.$$

From [27, Proposition 26], it is known that, for all $\beta > m_\phi$,

$$K^\phi \overset{\tau^{(\beta)}}{\rightsquigarrow} K^{(\beta)}$$

where $\tau^{(\beta)}$ is an infinitely divisible random variable with Laplace transform given by

$$\int_0^\infty e^{-us} \mathbb{P}(\tau^{(\beta)} \in ds) = \frac{\Gamma(1 + \beta) \Gamma(u + 1)}{\Gamma(u + \beta + 1)}, \quad u > 0.$$

More precisely, for all $t \geq 0$,

$$\begin{aligned} K_t^\phi \widehat{\mathbf{I}}_\phi \widehat{\mathbf{B}}_\beta &= \mathbf{I}_\phi \widehat{\mathbf{B}}_\beta K_t^{(\beta)} && \text{on } \mathbf{L}^2(\nu_\beta) \\ K_t^{(\beta)} \mathbf{V}_\beta &= \mathbf{V}_\beta K_t^\phi && \text{on } \mathbf{L}^2(\nu_\phi) \end{aligned}$$

with

$$(4.82) \quad \mathbf{I}_\phi f(x) = \mathbb{E}[f(xI_\phi)]$$

and, for all $k \in \mathbb{Z}_+$, $\mathbb{E}[I_\phi^k] = \frac{k!}{W_\phi(k+1)}$. V_β is another multiplicative Markov kernel associated with the random variable Y_β whose law is determined by its moment sequence given, for all $k \in \mathbb{Z}_+$, by

$$\mathbb{E}[Y_\beta^k] = \Gamma(1 + \beta) \frac{W_\phi(k + 1)}{\Gamma(k + 1 + \beta)}.$$

Finally, we have $\widehat{\mathbf{B}}_\beta f(x) = \frac{x^\beta}{\Gamma(\beta)} \int_0^\infty f((1 + y)x) y^{\beta-1} e^{-yx} dy$, $x > 0$, and $\mathbf{I}_\phi \widehat{\mathbf{B}}_\beta \mathbf{V}_\beta = K_{\tau^{(\beta)}}^\phi$. Now, from Proposition 4.14, we know that

$$(4.83) \quad \mathbf{K}_t^\phi \mathbf{I}_\phi = \mathbf{I}_\phi \mathbf{K}_t^{\sigma^2} \quad \text{on } \ell^2(\mathbf{n}_{\sigma^2})$$

where $\mathbb{K}^{\sigma^2} = \mathbb{K}^\phi$ with $\phi(u) = \sigma^2 u$. On the other hand, from Proposition 4.11, it is known that

$$(4.84) \quad K_t^\phi \Lambda = \Lambda \mathbb{K}_t^\phi \text{ on } \ell^2(\mathbf{n}_\phi)$$

and from [26, Proposition 25] along with [27, Proposition 30] and Lemma 4.21, we have

$$\begin{aligned} \mathbb{K}_t^{\sigma^2} \widehat{\Lambda}_{\sigma^2} &= \widehat{\Lambda}_{\sigma^2} K_t^{\sigma^2} && \text{on } \mathbf{L}^2(\nu_{\sigma^2}) \\ K_t^{\sigma^2} d_{\frac{1}{\sigma^2}} \widehat{\mathbb{B}}_\beta &= d_{\frac{1}{\sigma^2}} \widehat{\mathbb{B}}_\beta K_t^{(\beta)} && \text{on } \mathbf{L}^2(\nu_\beta) \\ K_t^{(\beta)} \mathbb{V}_\beta &= \mathbb{V}_\beta K_t^\phi && \text{on } \mathbf{L}^2(\nu_\phi) \end{aligned}$$

where ν_{σ^2} (resp. ν_β) equals ν_ϕ (the invariant distribution of the semigroup K^ϕ , see Proposition 4.11(2)) with $\phi(u) = \sigma^2 u$ (resp. $\phi(u) = u + \beta$), $d_\alpha f(x) = f(\alpha x)$ is the dilation operator and $\widehat{\Lambda}_{\sigma^2} : \mathbf{L}^2(\nu_{\sigma^2}) \rightarrow \ell^2(\mathbf{n}_{\sigma^2})$ is a Markov operator defined by

$$\widehat{\Lambda}_{\sigma^2}(n, dx) = \frac{\sigma^{2(n-1)}}{(1 + \sigma^2)^n} \frac{x^n}{n+1} e^{-x(1+\sigma^{-2})} dx.$$

By transitivity of the intertwining relation, it follows that

$$(4.85) \quad \mathbb{K}_t^\phi \mathbb{I}_\phi \widehat{\Lambda}_{\sigma^2} = \mathbb{I}_\phi \widehat{\Lambda}_{\sigma^2} K_t^{\sigma^2} \text{ on } \mathbf{L}^2(\nu_{\sigma^2})$$

$$(4.86) \quad K_t^{\sigma^2} \Upsilon = \Upsilon \mathbb{K}_t^\phi \text{ on } \ell^2(\mathbf{n}_\phi)$$

where $\Upsilon = d_{\frac{1}{\sigma^2}} \widehat{\mathbb{B}}_\beta \mathbb{V}_\beta \Lambda$. Now, from (4.85) and (4.86), it remains to show that $\mathbb{I}_\phi \widehat{\Lambda}_{\sigma^2} \Upsilon = \mathbb{K}_{\tau_\beta}^\phi$, where τ_β is defined as in the proposition.

Lemma 4.22. *The operator \mathbb{I}_ϕ in (4.82) commutes with the dilation operator d . Moreover, if $\sigma^2 > 0$ then $d_{\sigma^2} \mathbb{I}_\phi \Lambda = \mathbb{I}_\phi \Lambda$ on $\ell^2(\mathbf{n}_\phi)$.*

Proof. Since \mathbb{I}_ϕ is a multiplicative Markov kernel, commutation with the dilation operator follows readily. Now, for the intertwining relationship, by density of $\mathcal{P} = \text{Span}\{\mathbf{p}_k; k \in \mathbb{Z}_+\}$ in $\ell^2(\mathbf{n}_\phi)$, it suffices to show that, for all $k \in \mathbb{Z}_+$,

$$d_{\sigma^2} \mathbb{I}_\phi \Lambda \mathbf{p}_k = \Lambda \mathbb{I}_\phi \mathbf{p}_k.$$

However, $\mathbb{I}_\phi \mathbf{p}_k = \frac{\sigma^{2k} k!}{W_\phi(k+1)}$ and $d_{\sigma^2} \mathbb{I}_\phi \Lambda \mathbf{p}_k = d_{\sigma^2} \mathbb{I}_\phi p_k = \frac{\sigma^{2k} k!}{W_\phi(k+1)}$, which proves the lemma. \square

Coming back to the main proof, by an application of Lemma 4.21 and Lemma 4.22, we obtain

$$(4.87) \quad \Lambda \mathbb{I}_\phi \widehat{\Lambda}_{\sigma^2} \Upsilon = d_{\sigma^2} \mathbb{I}_\phi \Lambda \widehat{\Lambda}_{\sigma^2} \Upsilon = d_{\sigma^2} \mathbb{I}_\phi \Lambda \widehat{\Lambda}_{\sigma^2} d_{\frac{1}{\sigma^2}} \widehat{\mathbb{B}}_\beta \mathbb{V}_\beta \Lambda.$$

Again invoking [27, Proposition 25], we have $\Lambda\widehat{\Lambda}_{\sigma^2} = K_{\log(1+\sigma^{-2})}^{\sigma^2}$. Writing $\varrho = \frac{1}{2}\log(1+\sigma^{-2})$ as before, (4.87) yields

$$\begin{aligned}\Lambda\mathbb{I}_\phi\Upsilon &= d_{\sigma^2}\mathbb{I}_\phi K_\gamma^{\sigma^2} d_{\frac{1}{\sigma^2}}\widehat{\mathbb{B}}_\beta V_\beta\Lambda \\ &= d_{\sigma^2}\mathbb{I}_\phi d_{\frac{1}{\sigma^2}}\widehat{\mathbb{B}}_\beta K_\gamma^{(\beta)} V_\beta\Lambda \\ &= \mathbb{I}_\phi\widehat{\mathbb{B}}_\beta V_\beta K_{2\varrho}^\phi\Lambda \\ &= K_{\tau^{(\beta)}}^\phi K_{2\varrho}^\phi\Lambda \\ &= \Lambda\mathbb{K}_{\tau_\beta}^\phi\end{aligned}$$

where in the last line of the above equation, we used the fact that $\tau_\beta = \tau^{(\beta)} + 2\varrho$. By injectivity of Λ , it follows that $\mathbb{I}_\phi\Upsilon = \mathbb{K}_{\tau_\beta}^\phi$. This proves the proposition. \square

4.16. Proof of Theorem 2.12(2). We recall that L^{σ^2} is the generator of the self-adjoint Laguerre diffusion defined in (4.79) whose invariant distribution is $\nu_{\sigma^2}(x)dx = \frac{1}{\sigma^2}\nu(x/\sigma^2) = \frac{1}{\sigma^2}e^{-x/\sigma^2}dx$, $x > 0$. Let us first prove the Φ -entropy decay for K^{σ^2} , the semigroup generated by L^{σ^2} , that is, for all admissible function Φ and $f \in \mathbf{L}^1(\nu_{\sigma^2})$ with $\Phi(f) \in \mathbf{L}^1(\nu_{\sigma^2})$ one has

$$(4.88) \quad \text{Ent}_{\nu_{\sigma^2}}^\Phi(K_t^{\sigma^2}f) \leq e^{-t}\text{Ent}_{\nu_{\sigma^2}}^\Phi(f).$$

In Lemma 4.21, we have shown that the semigroups K^{σ^2} and K are equivalent via the similarity transform induced by the dilation operator d_{σ^2} . We first claim that it is enough to prove the exponential entropy decay in (4.88) replacing K^{σ^2} by K . To see why, we note that for any $\sigma^2 > 0$, $f \in \mathbf{L}^1(\nu_{\sigma^2})$ with $\Phi(f) \in \mathbf{L}^1(\nu_{\sigma^2})$, one has by the change of variable along with Lemma 4.21 that,

$$\begin{aligned}\int_0^\infty \Phi(K_t^{\sigma^2}f(x))\nu_{\sigma^2}(x)dx &= \int_0^\infty \frac{1}{\sigma^2}\Phi\left(K_t d_{\sigma^2}f\left(\frac{x}{\sigma^2}\right)\right)\nu\left(\frac{x}{\sigma^2}\right)dx \\ &= \int_0^\infty \Phi(K_t d_{\sigma^2}f(x))\nu(x)dx.\end{aligned}$$

We also observe by the change of variable that for any $f \in \mathbf{L}^1(\nu_{\sigma^2})$, one has $\Phi(\nu_{\sigma^2}f) = \Phi(\nu d_{\sigma^2}f)$. As a result, we have

$$\text{Ent}_{\nu_{\sigma^2}}^\Phi(K_t^{\sigma^2}f) = \text{Ent}_\nu^\Phi(K_t d_{\sigma^2}f)$$

which proves our claim. Next, we state the following result regarding the exponential entropy decay of the semigroup K generated by L .

Lemma 4.23. *For any Φ as above and $f \in \mathbf{L}^1(\nu)$ with $\Phi(f) \in \mathbf{L}^1(\nu)$, one has*

$$\text{Ent}_\nu^\Phi(K_t f) \leq e^{-t} \text{Ent}_\nu^\Phi(f).$$

Proof. Since L is a diffusion operator, from [11, Equations (6) and (7)], it suffices to show that for an admissible function Φ and $f \in \mathbf{L}^1(\nu)$ with $\Phi(f) \in \mathbf{L}^1(\nu)$, one has the following Φ -entropy inequality

$$(4.89) \quad \text{Ent}_\mu^\Phi(f) \leq \mu(\Phi''(f)\Gamma(f))$$

where Γ is the carré-du-champ operator, see [5, Section 1.4.2] associated to L , that is, for smooth functions

$$\Gamma(f) = L(f^2) - 2fLf.$$

From [13, Theorem 2.1(2)] it follows that (4.89) is equivalent to the fact that the operator L satisfies the curvature dimension condition $CD(\frac{1}{2}, \infty)$, which is indeed true from [5, Section 2.7.3]. Hence the lemma is proved. \square

Now coming back to the proof of Theorem 2.12(2), due to the interweaving relation in Theorem 4.20 and the estimate in (4.88), the proof of this theorem follows directly from [27, Theorem 8]. \square

4.17. Proof of Theorem 2.14. For ergodic self-adjoint diffusion semigroups, we know from [5, Theorem 5.2.3] that the hypercontractivity can be interpreted in terms of the log-Sobolev constants corresponding to the semigroups. Let us consider the self-adjoint Laguerre semigroup K^{σ^2} defined in Proposition 4.20. For this semigroup, the invariant distribution is $\nu_{\sigma^2}(x)dx = \frac{1}{\sigma^2}e^{-x/\sigma^2}dx$, $x > 0$, and the log-Sobolev constant is

$$(4.90) \quad c_{LS} = \inf_{f \in \mathbf{C}_b^1(\mathbb{R}_+): \|f\|_{\mathbf{L}^2(\nu_{\sigma^2})} = 1} \frac{4 \int_{\mathbb{R}_+} x f'(x)^2 \nu_{\sigma^2}(dx)}{\int_{\mathbb{R}_+} f(x)^2 \log(f(x)^2) \nu_{\sigma^2}(dx)}.$$

The numerator in the above expression is four times the Dirichlet energy associated to L^{σ^2} defined by

$$\mathcal{E}(f, f) = -\langle L^{\sigma^2} f, f \rangle_\nu = \int_{\mathbb{R}_+} x f'(x)^2 \nu_{\sigma_1^2}(dx).$$

It was shown by Bakry [4] that $c_{LS} = 1$. Hence, by applying [5, Theorem 5.2.3], we infer that for all $t \geq 0$,

$$\|K_t^{\sigma^2}\|_{\mathbf{L}^2(\nu_{\sigma^2}) \rightarrow \mathbf{L}^{p(t)}(\nu_{\sigma^2})} \leq 1$$

where $p(t) = 1 + e^t$ and $\nu(dx) = e^{-x}dx$, $x > 0$. Having the above hypercontractivity estimate, the rest of the proof follows from [27, Theorem 9] and Theorem 4.20. \square

4.18. **Proof of Theorem 2.15.** First, we note that the semigroup $\mathbb{K}^{\phi, \tau_\beta}$ has the same invariant distribution \mathbf{n}_ϕ as \mathbb{K}^ϕ . Let us recall that for $\sigma^2 > 0$, $\varrho = \frac{1}{2} \log(1 + \sigma^{-2})$. If $t > \frac{1}{2}$, Theorem 2.10(3) entails that, for all $s > 0$ and $\mathbf{f} \in \ell^2(\mathbf{n}_\phi)$, we have

$$(4.91) \quad \mathbb{K}_{s+2\varrho t}^\phi \mathbf{f} = \sum_{k=0}^{\infty} e^{-k\varrho t} e^{-ks} \langle \mathbf{f}, \mathbf{V}_k^\phi \rangle_{\mathbf{n}_\phi} \mathbf{P}_k^\phi \quad \text{in } \ell^2(\mathbf{n}_\phi).$$

For $t \geq 0$, let us define the random variable $\tilde{\tau}_\beta(t)$ such that $\tau_\beta(t) = 2\varrho t + \tilde{\tau}_\beta(t)$. Indeed, from (2.29) it follows that for all $t \geq 0$,

$$\log \mathbb{E} \left[e^{-u\tilde{\tau}_\beta(t)} \right] = -\log \left(\frac{\Gamma(u + \beta + 1)}{\Gamma(1 + \beta)\Gamma(u + 1)} \right).$$

Then, integrating both sides of (4.91) with respect to $\mathbb{P}(\tilde{\tau}_\beta(t) \in ds)$ with $t > \frac{1}{2}$ we obtain

$$\begin{aligned} \mathbb{K}_t^{\phi, \tau_\beta} \mathbf{f} &= \int_0^\infty \mathbb{K}_s^\phi \mathbf{f} \mathbb{P}(\tau_\beta(t) \in ds) = \int_0^\infty \mathbb{K}_{s+2\varrho t}^\phi \mathbf{f} \mathbb{P}(\tilde{\tau}_\beta(t) \in ds) \\ &= \int_0^\infty \left(\sum_{k=0}^{\infty} e^{-2k\varrho t} e^{-ks} \langle \mathbf{f}, \mathbf{V}_k^\phi \rangle_{\mathbf{n}_\phi} \mathbf{P}_k^\phi \right) \mathbb{P}(\tilde{\tau}_\beta(t) \in ds) \\ &= \sum_{k=0}^{\infty} e^{-2k\varrho t} \langle \mathbf{f}, \mathbf{V}_k^\phi \rangle_{\mathbf{n}_\phi} \mathbf{P}_k^\phi \int_0^\infty e^{-ks} \mathbb{P}(\tilde{\tau}_\beta(t) \in ds) \end{aligned}$$

where the last equality follows due to Fubini theorem with the help of the estimates $\|\mathbf{P}_k^\phi\|_{\ell^2(\mathbf{n}_\phi)} \leq 1$, $\|\mathbf{V}_k^\phi\|_{\ell^2(\mathbf{n}_\phi)} \leq C_\epsilon e^{k(\varrho + \epsilon)}$ for arbitrary $\epsilon > 0$ and $k \in \mathbb{Z}_+$, see Proposition 4.17 and the proof of Theorem 2.10(3). The proof of this item is concluded by recalling that, for all $k \in \mathbb{Z}_+$,

$$e^{-2\varrho t} \int_0^\infty e^{-ks} \mathbb{P}(\tilde{\tau}_\beta(t) \in ds) = e^{-t\phi_\beta(k)}.$$

For the next item, applying Jensen's inequality we observe that, for all $\beta, t > 0$,

$$\text{Ent}_{\mathbf{n}_\phi}^\Phi \left(\mathbb{K}_{t+1}^{\phi, \tau_\beta} \mathbf{f} \right) \leq \int_0^\infty \text{Ent}_{\mathbf{n}_\phi}^\Phi \left(\mathbb{K}_{s+\tau_\beta}^\phi \mathbf{f} \right) \mathbb{P}(\tau_\beta(t) \in ds).$$

Using Theorem 2, when $\sigma^2 > 0$ and $\beta > m_\phi$, the right-hand side of the above inequality is bounded above by

$$\int_0^\infty e^{-s} \text{Ent}_{\mathbf{n}_\phi}^\Phi(\mathbf{f}) \mathbb{P}(\tau_\beta(t) \in ds) = e^{-t\phi_\beta(1)} \text{Ent}_{\mathbf{n}_\phi}^\Phi(\mathbf{f}).$$

This proves (2). Finally, for any $\mathbf{f} \in \ell^2(\mathbf{n}_\phi)$, we have by the triangle inequality

$$\begin{aligned} \left\| \mathbb{K}_{t+1}^{\phi, \tau_\beta} \mathbf{f} \right\|_{\ell^2(\mathbf{n}_\phi)} &= \left\| \int_0^\infty \mathbb{K}_{s+\tau_\beta}^\phi \mathbf{f} \mathbb{P}(\tau_\beta(t) \in ds) \right\|_{\ell^2(\mathbf{n}_\phi)} \\ &\leq \int_0^\infty \left\| \mathbb{K}_{s+\tau_\beta}^\phi \mathbf{f} \right\|_{\ell^2(\mathbf{n}_\phi)} \mathbb{P}(\tau_\beta(t) \in ds). \end{aligned}$$

Invoking Theorem 2.14, the right-hand side of the above inequality is bounded above by

$$\|\mathbf{f}\|_{\ell^2(\mathbf{n}_\phi)} \int_0^\infty \mathbb{P}(\tau_\beta(t) \in ds) = \|\mathbf{f}\|_{\ell^2(\mathbf{n}_\phi)}$$

which completes the proof.

REFERENCES

- [1] F. Achleitner, A. Arnold, and E. A. Carlen. On multi-dimensional hypocoercive BGK models. *Kinet. Relat. Models*, 11(4):953–1009, 2018.
- [2] N.I. Akhiezer. *The Classical Moment Problem*. Oliver and Boyd, 1965.
- [3] T. Assiotis. On a gateway between the Laguerre process and dynamics on partitions. *Latin American Journal of Probability and Mathematical Statistics*, 2019.
- [4] D. Bakry. Remarques sur les semigroupes de Jacobi Astérisque, (236):23-39, 1996. Hommage à P.A. Meyer et J. Neveu.
- [5] D. Bakry, I. Gentil, and M. Ledoux. *Analysis and geometry of Markov diffusion operators*, volume 348 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Cham, 2014.
- [6] F. Baudoin. Bakry–Émery meet Villani. *J. Funct. Anal.*, 273(7):2275–2291, 2017.
- [7] J. Bertoin. The asymptotic behavior of fragmentation processes *J. Europ. Math. Soc.*, 5:395-416, 2003.
- [8] J. Bertoin and M. Yor, On the entire moments of self-similar Markov processes and exponential functionals of Lévy processes *Ann. Fac. Sci. Toulouse Math. (6)*, 11:33–45, 2002.
- [9] J. Bertoin, I. Kortchemski, Self-similar scaling limits of Markov chains on the positive integers *Ann. Appl. Probab.*, 26:2556-2595, 2016.
- [10] A. Borodin and G. Olshanski. Markov dynamics on the Thoma cone: a model of time-dependent determinantal processes with infinitely many particles. *Electron. J. Probab.*, 18:no. 75, 43, 2013.
- [11] F. Bolley, I. Gentil. Phi-entropy inequalities and Fokker-Planck equations, *Progress in analysis and its applications* 463-469, 2010.
- [12] G. Carinci, C. Franceschini, C. Giardinà, F. Redig, W. Groenevelt, Orthogonal dualities of Markov processes and unitary symmetries. *SIGMA Symmetry Integrability Geom. Methods Appl.* 15, no. 53, 2019.
- [13] D. Chafai. Entropies, convexity, and functional inequalities, *J. Math. Kyoto. Univ.*, 44-2(2004), 325-363
- [14] O. Christensen. *An Introduction to Frames and Riesz Bases*. Birkhäuser, 2003.
- [15] J. Dolbeault, C. Mouhot, and C. Schmeiser. Hypocoercivity for linear kinetic equations conserving mass. *Trans. Amer. Math. Soc.*, 367(6):3807–3828, 2015.

- [16] E. B. Dynkin. *Markov processes. Vols. I, II*, volume 122 of *Translated with the authorization and assistance of the author by J. Fabius, V. Greenberg, A. Maitra, G. Majone. Die Grundlehren der Mathematischen Wissenschaften, Bände 121*. Academic Press Inc., Publishers, New York, 1965.
- [17] Ethier, Stewart N. and Kurtz, Thomas G. *Markov processes: Characterization and convergence*.
- [18] W. Groenevelt, Orthogonal stochastic duality functions from Lie algebra representations, *J. Stat. Phys.* 174, 97–119, 2019.
- [19] S. Jansen and N. Kurt, On the notion(s) of duality for Markov processes, *Probability Surveys*, 11, 59 – 120, 2014.
- [20] O. Kallenberg, *Foundations of modern probability*, second edition, Springer-Verlag, New York, 2002.
- [21] S. Karlin, J. McGregor Linear growth birth and death processes. *J. Math. Mech.*, 7:643-662, 1958
- [22] R. Koekoek, P.A. Lesky, R.F. Swarttouw *Hypergeometric orthogonal polynomials and their q-analogues* Springer-Verlag, Berlin, 2010
- [23] J. Lamperti. Semi-stable stochastic processes. *Trans. Amer. Math. Soc.*, 104:62–78, 1962.
- [24] J. Lamperti. Semi-stable Markov processes. I. *Z. Wahrsch. Verw. Geb.*, 22:205–225, 1972.
- [25] J.F. Le-Gall, G. Miermont Scaling limits of random planar maps with large faces *Ann. Probab.* 39, no. 1, 1–69, 2011.
- [26] L. Miclo, P. Patie, On a gateway between continuous and discrete Bessel and Laguerre processes. *Annales Henri Lebesgue*, 2:59–98, 2019.
- [27] L. Miclo and P. Patie, On interweaving relations, *J. Funct. Anal.*, 280, no. 3, 53pp., 2021.
- [28] R. B. Paris and D. Kaminski. *Asymptotics and Mellin-Barnes integrals*, volume 85 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2001.
- [29] P. Patie and M. Savov. Spectral expansion of non-self-adjoint generalized Laguerre semigroups, *Mem. Amer. Math. Soc.*, 272, no. 1336, vii+182 pp, 2021.
- [30] P. Patie and M. Savov. Bernstein-gamma functions and exponential functionals of Lévy processes, *Electron. J. Probab.*, vol. 23, p. 101 pp., 2018.
- [31] P. Patie and A. Vaidyanathan, A spectral theoretical approach for hypocoercivity applied to some degenerate hypoelliptic, and non-local operators, *Kinetic and Related Models*, 13(3): 479-506, 2020.
- [32] J. Pitman and L.G. Rogers. Markov functions. *Ann. Probab.*, 9:573–582, 1981.
- [33] F. Redig and F. Sau. Factorized duality, stationary product measures and generating functions. *J. Stat. Phys.* 172(4): 980-1008, 2018.
- [34] R.P. Stanley. *Enumerative Combinatorics: Vol 1, Second edition*
- [35] E.C. Titchmarsh. *The theory of functions*. Oxford University Press, Oxford, 1939.
- [36] C. Villani. Hypocoercivity. *Mem. Amer. Math. Soc.*, 202(950):iv+141, 2009.
- [37] Robert M. Young. *An introduction to nonharmonic Fourier series*. Academic Press, Inc., San Diego, CA, first edition, 2001.

INSTITUT DE MATHÉMATIQUES DE TOULOUSE AND TOULOUSE SCHOOL OF ECONOMICS,
UNIVERSITÉ DE TOULOUSE AND CNRS, FRANCE

Email address: `miclo@math.cnrs.fr`

SCHOOL OF OPERATIONS RESEARCH AND INFORMATION ENGINEERING, CORNELL UNI-
VERSITY, ITHACA, NY 14853.

Email address: `pp396@cornell.edu`

SCHOOL OF OPERATIONS RESEARCH AND INFORMATION ENGINEERING, CORNELL UNI-
VERSITY, ITHACA, NY 14853.

Email address: `rs2466@cornell.edu`