



Groups of order 8 and 16

J Lapuyade-Lahorgue

► To cite this version:

| J Lapuyade-Lahorgue. Groups of order 8 and 16. Master. France. 2018. cel-01841041

HAL Id: cel-01841041

<https://hal.science/cel-01841041>

Submitted on 17 Jul 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Groups of order 8 and 16

J. Lapuyade-Lahorgue

1 Introduction

This document is inspired of the work of David Clausen (University of Puget Sound, USA) with simplification in the proofs. The reader has to have basic knowledge on group theory including:

- Abelian and cyclic groups.
- Lagrange theorem.
- Normal subgroup and quotient group.
- Direct and semi-direct products.
- Cauchy theorem.
- Operation of groups on a set.

2 Preliminaries

Definition 1 (Center of a group). *Let G be a group, the center of G is defined as:*

$$Z(G) = \{h : \forall g, hg = gh\}$$

Definition 2 (Commutator subgroup). *Let G be a group, the commutator of G is the smallest subgroup containing the commutators $[g_1, g_2] = g_1g_2g_1^{-1}g_2^{-1}$*

Proposition 1. *If the quotient group $G/Z(G)$ is cyclic then G is Abelian and consequently $G/Z(G) \cong \{e\}$.*

Proof. Suppose $G/Z(G) = \{Z(G), gZ(G), g^2Z(G), \dots, g^{n-1}Z(G)\}$. Let $g^i x_i$ and $g^j x_j$, with $x_i, x_j \in Z(G)$ be two elements of G . Then:

$$\begin{aligned} g^i x_i g^j x_j &= g^i g^j x_i x_j \\ &= x_j g^{i+j} x_i \\ &= x_j g^j g^i x_i \\ &= g^j x_j g^i x_i \end{aligned}$$

□

Proposition 2. *If $|G| = p^n$, where p is a prime number, then $|Z(G)| = p^k$ for $k \geq 1$.*

Proof. G operates on itself by conjugaison $g.h = ghg^{-1}$. For any h , the stabilizer of h is the centralizer $S_h = \{g : gh = hg\}$ and its orbital is the set $\mathcal{O}(h) = \{ghg^{-1} : g \in G\}$. We recall that $\mathcal{O}(h)$ is in bijection with the quotient set $(G/S_h)_l = \{gS_h : g \in G\}$ and consequently has the same cardinal. It is easy to show that the cardinal $|\mathcal{O}(h)| = 1$ if and only if $h \in Z(G)$, consequently:

$$|G| = |Z(G)| + \sum_{h \notin Z(G)} \frac{|G|}{|S_h|},$$

where the sum is indexed for the h which are not in the same orbital.

The term $\sum_{h \notin Z(G)} \frac{|G|}{|S_h|}$ is a multiple of p , consequently $|Z(G)|$ also. Using the Lagrange's theorem, we deduce the result. □

Theorem 1 (Burnside's theorem). *If the order of a group G is equal to p^2 , with p a prime number; then G is abelian.*

Proof. Easy: use the two previous propositions. □

Proposition 3. *The commutator subgroup, denoted $D(G)$, is the smallest normal subgroup such that $G/D(G)$ is Abelian.*

Proof. First step: $D(G)$ is a normal subgroup:

$$\begin{aligned} h[g_1, g_2]h^{-1} &= hg_1g_2g_1^{-1}g_2^{-1}h^{-1} \\ &= hg_1g_2g_1^{-1}(hg_2)^{-1} \\ &= hg_1h^{-1}(hg_2)g_1^{-1}(hg_2)^{-1} \\ &= hg_1h^{-1}g_1^{-1}g_1(hg_2)g_1^{-1}(hg_2)^{-1} \\ &= [h, g_1][g_1, hg_2] \in D(G) \end{aligned}$$

Second step: $G/D(G)$ is Abelian:

It is trivial by construction of $D(G)$. Indeed, $g_1^{-1}g_2^{-1}g_1g_2 \in D(G)$, consequently, $g_1g_2D(G) = g_2g_1D(G)$. **Third step:** If H is a normal subgroup such that G/H is Abelian, then H contains the commutator. The conclusion is trivial. \square

Theorem 2 (Correspondance theorem). *Let H be a normal subgroup of a group G , then there is a bijection between the set of the subgroups S of G containing H and the set of the subgroups S/H of G/H .*

Proposition 4. *Let H and K be two subgroups of a group G . We define the set $HK = \{hk : h \in H, k \in K\}$, which is not necessarily a group. Then:*

$$|HK| = \frac{|H| \times |K|}{|H \cap K|}$$

3 Groups of order 8

We now classify all groups of order 8. The neutral element will be denoted e .

3.1 Abelian groups of order 8

Name	Character presentation
\mathbb{Z}_8	$a^8 = e$
$\mathbb{Z}_4 \times \mathbb{Z}_2$	$a^4 = b^2 = e$ and $D(G) = \{e\}$
$(\mathbb{Z}_2)^3$	$a^2 = b^2 = c^2 = e$ and $D(G) = \{e\}$.

3.2 No Abelian groups of order 8

Proposition 5. *We have necessarily:*

$$Z(G) \cong \mathbb{Z}_2$$

and:

$$G/Z(G) \cong \mathbb{Z}_2 \times \mathbb{Z}_2$$

Proof. As $G/Z(G)$ cannot be cyclic and $|Z(G)|$ is a non trivial power of 2 different of 8, we deduce the result. \square

From the previous proposition, as G is not Abelian and $G/Z(G)$ is Abelian; we deduce that $D(G) = Z(G)$.

Moreover, we remark that:

$$\mathbb{Z}_2 \times \mathbb{Z}_2 = \{(0, 0), (1, 0), (0, 1), (1, 1)\},$$

has three subgroups of order 2. Consequently, by the correspondance theorem, G has three subgroups of order 4, denoted G_1, G_2, G_3 , containing the center.

Proposition 6. *If $g_1 \in G_1 \setminus Z(G)$ and $g_2 \in G_2 \setminus Z(G)$, then $g_1 g_2 \in G_3 \setminus Z(G)$.*

Proof. Suppose, by the correspondance theorem, that G_1 (resp. G_2) corresponds to the subgroup $\{(0, 0), (1, 0)\}$ (resp. $\{(0, 0), (0, 1)\}$). Then, g_1 corresponds to $(1, 0)$ and g_2 to $(0, 1)$. As $(1, 0) + (0, 1) = (1, 1)$ which corresponds to $g_1 g_2$; we deduce the conclusion. \square

Proposition 7. *If $g_1 \in G_1 \setminus Z(G)$ and $g_2 \in G_2 \setminus Z(G)$, then $g_1 g_2 \neq g_2 g_1$.*

Proof. Suppose the converse. Then g_1 commutes with the elements of $Z(G)$, of $g_1 Z(G)$ and of $g_2 Z(G)$. Consequently, the centralizer of g_1 has at least 6 elements. And, as the centralizer is a subgroup of G then it has exactly 8 elements. We deduce that the centralizer of g_1 is G , so $g_1 \in Z(G)$; which is contradictory. \square

Proposition 8. *Let G_i, G_j be two distinct subgroups amongst G_1, G_2, G_3 , then:*

$$\begin{aligned} G &= G_i G_j \\ G_i \cap G_j &= Z(G) \end{aligned}$$

Proof. Use the formula:

$$|G_i G_j| = \frac{|G_i| \times |G_j|}{|G_i \cap G_j|} = \frac{16}{|G_i \cap G_j|},$$

and the fact that $G_i G_j$ cannot have more elements than G . \square

As the sub-groups G_i has order equal to 4, they are either isomorphic to \mathbb{Z}_4 or $\mathbb{Z}_2 \times \mathbb{Z}_2$.

In the following, let $Z(G) = \{e, \alpha\}$ with $\alpha^2 = e$.

3.2.1 $G_1, G_2, G_3 \cong \mathbb{Z}_2 \times \mathbb{Z}_2$

In this case, there are three elements a, b, c in $G \setminus Z(G)$ such that:

- $G_1 = \{e, a, \alpha, a\alpha\}$.
- $G_2 = \{e, b, \alpha, b\alpha\}$.
- $G_3 = \{e, c, \alpha, c\alpha\}$.

From the Proposition 6., $ba \in G_3$ and consequently has order 2; so $baba = e$. But, as a and b have order 2, then $baab = e$. Consequently, $ba = ab$, which contradicts Proposition 7.

3.2.2 $G_1 \cong \mathbb{Z}_4, G_2, G_3 \cong \mathbb{Z}_2 \times \mathbb{Z}_2$

There exists three elements a, b, c in $G \setminus Z(G)$ such that:

- $G_1 = \{e, a, a^2 = \alpha, a^3\}$.
- $G_2 = \{e, b, \alpha, b\alpha\}$.
- $G_3 = \{e, c, \alpha, c\alpha\}$.

ba has order 2, so $baba = e$. a has order 4 and b has order 2, so $baa^3b = e$. Consequently, $ba = a^3b$. As $G = G_1G_2$, G is generated by a and b . Moreover, $|\langle a \rangle| \times |\langle b \rangle| = 4 \times 2 = 8 = |G|$ and $\langle a \rangle \cap \langle b \rangle = \{e\}$; G is the semi-direct product $\langle a \rangle \rtimes \langle b \rangle \cong \mathbb{Z}_4 \rtimes \mathbb{Z}_2$. It is the diedral group D_4 . Its character representation is $a^4 = b^2 = e$ and $ba = a^3b$.

3.2.3 $G_1, G_2 \cong \mathbb{Z}_4, G_3 \cong \mathbb{Z}_2 \times \mathbb{Z}_2$

There exists three elements a, b, c in $G \setminus Z(G)$ such that:

- $G_1 = \{e, a, a^2 = \alpha, a^3\}$.
- $G_2 = \{e, b, b^2 = \alpha, b^3\}$.
- $G_3 = \{e, c, \alpha, c\alpha\}$.

We have also $baba = e$ and $baa^3b^3 = e$. Consequently $ba = a^3b^3 = a^2abb^2 = \alpha aba = \alpha^2 ab = ab$ because $\alpha \in Z(G)$ and has order 2. $ba = ab$ leads to a contradiction.

3.2.4 $G_1, G_2, G_3 \cong \mathbb{Z}_4$

There exists three elements a, b, c in $G \setminus Z(G)$ such that:

- $G_1 = \{e, a, a^2 = \alpha, a^3\}$.
- $G_2 = \{e, b, b^2 = \alpha, b^3\}$.
- $G_3 = \{e, c, c^2 = \alpha, c^3\}$.

We have $ba(ba)^3 = e$ and $baa^3b^3 = e$. Consequently $(ba)^3 = a^3b^3 = ab$; so $(ba)^2ba = ab$; so $\alpha ba = ab$; so $ba = \alpha ab = a^3b$. It is the group of quaternions denoted \mathbb{H} . It is not a semi-direct product and its character representation is $a^4 = b^4 = e$, $a^2 = b^2$ and $ba = a^3b$. It is frequent that a and b are denoted i and j , ab is denoted k and a^2 is denoted -1 . With these notation $\mathbb{H} = \{1, -1, i, -i, j, -j, k, -k\}$ with $i^2 = j^2 = -1$, $(-1)^2 = 1$, $ji = -ij$, $ij = k$.

We have classified all groups of order 8. The following table gives the non-Abelian groups of order 8.

Name	Character presentation
$D_4 \cong \mathbb{Z}_4 \rtimes \mathbb{Z}_2$	$a^4 = b^2 = e, ba = a^3b$
\mathbb{H}	$a^4 = b^4 = e, a^2 = b^2, ba = a^3b$

4 Groups of order 16

4.1 Abelian groups of order 16

Name	Character presentation
\mathbb{Z}_{16}	$a^{16} = e$
$\mathbb{Z}_8 \times \mathbb{Z}_2$	$a^8 = b^2 = e, D(G) = \{e\}$
$\mathbb{Z}_4 \times \mathbb{Z}_4$	$a^4 = b^4 = e, D(G) = \{e\}$
$\mathbb{Z}_4 \times (\mathbb{Z}_2)^2$	$a^4 = b^2 = c^2 = e, D(G) = \{e\}$
$(\mathbb{Z}_2)^4$	$a^2 = b^2 = c^2 = d^2 = e, D(G) = \{e\}$

4.2 Non-Abelian groups of order 16

As $G/Z(G)$ cannot be cyclic, we show easily that $|Z(G)| \in \{2, 4\}$.

4.2.1 First case: $|Z(G)| = 4$

In this case, as $G/Z(G)$ is not cyclic, $G/Z(G) \cong \mathbb{Z}_2 \times \mathbb{Z}_2$. As $G/Z(G)$ is Abelian, then $D(G) \triangleleft Z(G)$.

As it has been done for the groups of order 8, we show that G has three sub-groups G_1, G_2, G_3 of order 8 which contain the center.

Proposition 9. *The subgroups G_1, G_2, G_3 are Abelian.*

Proof. Indeed, as $Z(G) < G_i$, we deduce easily that $Z(G) < Z(G_i)$. Consequently, $Z(G_i)$ has at least 4 elements. And as the center of a group of order 8 has either 2 elements or 8 elements, then $|Z(G_i)| = 8$ and we deduce that G_i is Abelian. \square

The following propositions can be proved in a similar way as for the groups of order 8.

Proposition 10. *If $g_1 \in G_1 \setminus Z(G)$ and $g_2 \in G_2 \setminus Z(G)$, then $g_1 g_2 \in G_3 \setminus Z(G)$.*

Proposition 11. *If $g_1 \in G_1 \setminus Z(G)$ and $g_2 \in G_2 \setminus Z(G)$, then $g_1 g_2 \neq g_2 g_1$.*

Proposition 12. *Let G_i, G_j be two distinct subgroups amongst G_1, G_2, G_3 , then:*

$$\begin{aligned} G &= G_i G_j \\ G_i \cap G_j &= Z(G) \end{aligned}$$

A) First sub-case: $Z(G) \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ Let $Z(G) = \{e, \alpha, \beta, \gamma\}$ with $\alpha^2 = \beta^2 = \gamma^2 = e$.

The Abelian groups of order 8 which have $\mathbb{Z}_2 \times \mathbb{Z}_2$ for subgroup are:

- $\mathbb{Z}_4 \times \mathbb{Z}_2 = \{(0, 0), (1, 0), (2, 0), (3, 0), (0, 1), (1, 1), (2, 1), (3, 1)\}$ with:
 - $\{(0, 0), (2, 0), (0, 1), (2, 1)\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.
- $(\mathbb{Z}_2)^3 = \{(0, 0, 0), (0, 0, 1), (0, 1, 0), (0, 1, 1), (1, 0, 0), (1, 0, 1), (1, 1, 0), (1, 1, 1)\}$ with:
 - $\{(0, 0, 0), (0, 0, 1), (0, 1, 0), (0, 1, 1)\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.
 - $\{(0, 0, 0), (0, 0, 1), (1, 0, 0), (1, 0, 1)\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.
 - $\{(0, 0, 0), (0, 0, 1), (1, 1, 0), (1, 1, 1)\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.
 - $\{(0, 0, 0), (0, 1, 0), (1, 0, 0), (1, 1, 0)\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.
 - $\{(0, 0, 0), (0, 1, 0), (1, 0, 1), (1, 1, 1)\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.
 - $\{(0, 0, 0), (0, 1, 1), (1, 0, 0), (1, 1, 1)\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.
 - $\{(0, 0, 0), (0, 1, 1), (1, 0, 1), (1, 1, 0)\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.

A1) $G_1, G_2, G_3 \cong (\mathbb{Z}_2)^3$: There exists three elements a, b, c which are not in the center $Z(G)$ such that:

- $G_1 = \{e, \alpha, \beta, \gamma, a, a\alpha, a\beta, a\gamma\}$.
- $G_2 = \{e, \alpha, \beta, \gamma, b, b\alpha, b\beta, b\gamma\}$.
- $G_3 = \{e, \alpha, \beta, \gamma, c, c\alpha, c\beta, c\gamma\}$.

ba has order 2, so $baba = e$. But, a and b have order 2, so $baab = e$. Consequently $ba = ab$, which leads to a contradiction.

A2) $G_1 \cong \mathbb{Z}_4 \times \mathbb{Z}_2, G_2, G_3 \cong (\mathbb{Z}_2)^3$: There exists three elements a, b, c which are not in the center $Z(G)$ such that:

- $G_1 = \{e, a, a^2 = \alpha, a^3, \beta, a\beta, a^2\beta = \gamma, a^3\beta\}$.
- $G_2 = \{e, \alpha, \beta, \gamma, b, b\alpha, b\beta, b\gamma\}$.
- $G_3 = \{e, \alpha, \beta, \gamma, c, c\alpha, c\beta, c\gamma\}$.

We show easily that $ba = a^3b$ (ba has order 2, a has order 4 and b has order 2). As $G = G_1G_2$, G is generated by a, b, β . We have $|\langle a \rangle| \times |\langle b \rangle| = 8$ and $\langle a \rangle \cap \langle b \rangle = \{e\}$. Consequently $\langle a, b \rangle \cong D_4$. Moreover, $\beta a = a\beta$ and $\beta b = b\beta$, as $\beta \in Z(G)$. $|\langle a, b \rangle| \times |\langle \beta \rangle| = 16 = |G|$ and $\langle a, b \rangle \cap \langle \beta \rangle = \{e\}$. Consequently, the group G is:

$$G \cong D_4 \times \mathbb{Z}_2$$

Its character presentation is $x^4 = y^2 = z^2, yx = x^3y, zx = xz, zy = yz$. (In order to avoid confusion, we will prefer character presentation with letter x, y, z, w, \dots rather than a, b, c, \dots)

A3) $G_1, G_2 \cong \mathbb{Z}_4 \times \mathbb{Z}_2, G_3 \cong (\mathbb{Z}_2)^3$: We consider two cases:

First case:

There exists three elements a, b, c which are not in the center $Z(G)$ such that:

- $G_1 = \{e, a, a^2 = \alpha, a^3, \beta, a\beta, a^2\beta = \gamma, a^3\beta\}$.
- $G_2 = \{e, b, b^2 = \alpha, b^3, \beta, b\beta, b^2\beta = \gamma, b^3\beta\}$.
- $G_3 = \{e, \alpha, \beta, \gamma, c, c\alpha, c\beta, c\gamma\}$.

Then we show by considering the order of elements that $ba = a^3b^3$. Consequently, $ba = a^3b^3 = a^2abb^2 = \alpha ab\alpha = \alpha^2ab = ab$, which leads to a contradiction.

Second case:

There exists three elements a, b, c which are not in the center $Z(G)$ such that:

- $G_1 = \{e, a, a^2 = \alpha, a^3, \beta, a\beta, a^2\beta = \gamma, a^3\beta\}$.
- $G_2 = \{e, b, b^2 = \beta, b^3, \alpha, b\alpha, b^2\alpha = \gamma, b^3\alpha\} = \{e, b, b^2 = \beta, b^3, \gamma, b\gamma, b^2\gamma = \alpha, b^3\gamma = b\alpha\}$.
- $G_3 = \{e, \alpha, \beta, \gamma, c, c\alpha, c\beta, c\gamma\}$.

We show that $ba = a^3b^3 = \alpha ab\beta = \gamma ab$. Without loss of generality, one can suppose that $ab = c$, so $ba = \gamma c$. We have $b\gamma = \gamma b$ and $bc = bab = \gamma cb$, $\langle \gamma, c \rangle \cong \mathbb{Z}_2 \times \mathbb{Z}_2$, $\langle b \rangle \cong \mathbb{Z}_4$, $|\langle \gamma, c \rangle| \times |\langle b \rangle| = 4 \times 4 = 16 = |G|$ and $\langle \gamma, c \rangle \cap \langle b \rangle = \{e\}$. Consequently, G is the semi-direct product:

$$G \cong (\mathbb{Z}_2 \times \mathbb{Z}_2) \rtimes_{\varphi} \mathbb{Z}_4,$$

with:

$$\begin{aligned} \varphi : \mathbb{Z}_4 &\rightarrow \text{Aut}(\mathbb{Z}_2 \times \mathbb{Z}_2) \\ 1 &\rightarrow \begin{pmatrix} (1, 0) \rightarrow (1, 0) \\ (0, 1) \rightarrow (1, 1) \end{pmatrix} \end{aligned}$$

Its minimal character presentation is $x^4 = y^4 = e$, $yx = x^3y^3$, $x^2, y^2 \in Z(G)$. (another possible presentation can be $x^2 = y^2 = z^4 = e$, $yx = xy$, $zx = xyz$, $zy = yz$ but needs more generators).

A4) $G_1, G_2, G_3 \cong \mathbb{Z}_4 \times \mathbb{Z}_2$: We consider two cases:

First case:

There exists three elements a, b, c which are not in the center $Z(G)$ such that:

- $G_1 = \{e, a, a^2 = \alpha, a^3, \beta, a\beta, a^2\beta = \gamma, a^3\beta\}$.
- $G_2 = \{e, b, b^2 = \beta, b^3, \alpha, b\alpha, b^2\alpha = \gamma, b^3\alpha\}$.
- $G_3 = \{e, c, c^2, c^3, \dots\}$ (the structure of G_3 will be precised latter).

As ba is not in the center but in G_3 then ba has order 4. In all case, $(ba)^2 = c^2$. Indeed, in $\mathbb{Z}_4 \times \mathbb{Z}_2$, the square of all elements of order 4 is $(2, 0)$. We show that $(ba)^2ba = a^3b^3 = a^2abb^2 = \alpha ab\beta = \gamma ab$. Consequently, $(ba)^2 \neq \gamma$. Without loss of generality, one can suppose that $c^2 = (ba)^2 = \beta$; which gives the entire structure of

G_3 . Consequently, $ba = \beta\gamma ab = \alpha ab = a^2 ab = a^3 b$. We have $\langle a \rangle \cong \mathbb{Z}_4$ (idem for $\langle b \rangle$), $|\langle a \rangle| \times |\langle b \rangle| = 16 = |G|$ and $\langle a \rangle \cap \langle b \rangle = \{e\}$. Consequently, G is the semi-direct product:

$$G \cong \mathbb{Z}_4 \rtimes \mathbb{Z}_4$$

Its character presentation is $x^4 = y^4 = e, yx = x^3y$.

Second case:

From the first case, one can see that amongst the square a^2, b^2, c^2 , at least two are equal. The second case is when $a^2 = b^2 = c^2$. Without loss of generality:

- $G_1 = \{e, a, a^2 = \alpha, a^3, \beta, a\beta, a^2\beta = \gamma, a^3\beta\}$.
- $G_2 = \{e, b, b^2 = \alpha, b^3, \beta, b\beta, b^2\beta = \gamma, b^3\beta\}$.
- $G_3 = \{e, c, c^2 = \alpha, c^3, \beta, c\beta, c^2\beta = \gamma, c^3\beta\}$

We show that $(ba)^3 = a^3b^3 = a^2abb^2 = ab$, so $(ba)^2ba = ab$, so $ba = a^3b$. As the character presentation of $\langle a, b \rangle$ is $a^4 = b^4 = e, a^2 = b^2, ba = a^3b$, we deduce that $\langle a, b \rangle \cong \mathbb{H}$. Moreover, $G = G_1G_2 = \langle a, b, \beta \rangle$. As $\beta \in Z(G)$, $|\langle \beta \rangle| \times |\langle a, b \rangle| = 2 \times 8 = 16$ and $\langle \beta \rangle \cap \langle a, b \rangle = \{e\}$, we deduce that G is the direct product:

$$\mathbb{H} \times \mathbb{Z}_2$$

Its character presentation is $x^4 = y^4 = z^2 = e, x^2 = y^2, yx = x^3y, zx = xz, zy = yz$.

We have classified all groups of order 16 such that $Z(G) \cong \mathbb{Z}_2 \times \mathbb{Z}_2$:

Name	Character presentation
$D_4 \times \mathbb{Z}_2$	$x^4 = y^2 = z^2 = e, yx = x^3y, zx = xz, zy = yz$
$(\mathbb{Z}_2 \times \mathbb{Z}_2) \rtimes \mathbb{Z}_4$	$x^4 = y^4 = e, yx = x^3y^3, x^2, y^2 \in Z(G)$
$\mathbb{Z}_4 \rtimes \mathbb{Z}_4$	$x^4 = y^4 = e, yx = x^3y$
$\mathbb{H} \times \mathbb{Z}_2$	$x^4 = y^4 = z^2 = e, x^2 = y^2, yx = x^3y, zx = xz, zy = yz$

Groups of order 16 with $Z(G) \cong \mathbb{Z}_2 \times \mathbb{Z}_2$

B) Second sub-case: $Z(G) \cong \mathbb{Z}_4$.

Let $Z(G) = \{e, \alpha, \alpha^2, \alpha^3\}$.

The Abelian groups of order 8 which have \mathbb{Z}_4 for subgroups are:

- $\mathbb{Z}_8 = \{0, 1, 2, 3, 4, 5, 6, 7\}$ with:
 - $\{0, 2, 4, 6\} \cong \mathbb{Z}_4$
- $\mathbb{Z}_4 \times \mathbb{Z}_2 = \{(0, 0), (1, 0), (2, 0), (3, 0), (0, 1), (1, 1), (2, 1), (3, 1)\}$ with:

- $\{(0, 0), (1, 0), (2, 0), (3, 0)\} \cong \mathbb{Z}_4$
- $\{(0, 0), (1, 1), (2, 0), (3, 1)\} \cong \mathbb{Z}_4$

We remark that the square of any element of \mathbb{Z}_8 is in the subgroup of order 4 and the square of any element of $\mathbb{Z}_4 \times \mathbb{Z}_2$ is equal to $(0, 0)$ or $(2, 0)$ and is in its two subgroups of order 4. We deduce that the square of any element of G_i is in the center. From this fact, we deduce the following proposition.

Proposition 13. *For $i \neq j$, the commutator $[g_i, g_j] = g_i g_j g_i^{-1} g_j^{-1}$ between two elements respectively from $G_i \setminus Z(G)$ and $G_j \setminus Z(G)$ has an order equal to 2.*

Proof. As $g_i^2 \in Z(G)$ and using $D(G) < Z(G)$, we deduce $g_j g_i^2 = g_i^2 g_j = g_i g_i g_j = g_i [g_i, g_j] g_j g_i = [g_i, g_j] g_i g_j g_i = [g_i, g_j]^2 g_j g_i^2$ and we deduce the conclusion. \square

B1) $G_1, G_2, G_3 \cong \mathbb{Z}_4 \times \mathbb{Z}_2$: By considering how $\mathbb{Z}_4 \times \mathbb{Z}_2$ is built from one of its subgroup of order 4 and an element of order 2 which is not in the considered subgroup, we deduce that there exists a, b, c not in the center and of order 2 such that:

- $G_1 = \{e, \alpha, \alpha^2, \alpha^3, a, a\alpha, a\alpha^2, a\alpha^3\}$ with $|a\alpha| = 4$, $|a\alpha^2| = 2$ and $|a\alpha^3| = 4$.
- $G_2 = \{e, \alpha, \alpha^2, \alpha^3, b, b\alpha, b\alpha^2, b\alpha^3\}$ with $|b\alpha| = 4$, $|b\alpha^2| = 2$ and $|b\alpha^3| = 4$.
- $G_3 = \{e, \alpha, \alpha^2, \alpha^3, c, c\alpha, c\alpha^2, c\alpha^3\}$ with $|c\alpha| = 4$, $|c\alpha^2| = 2$ and $|c\alpha^3| = 4$.

The order of ba is either 2 or 4.

Suppose that $|ba| = 2$. Then $baba = e$ and as a and b have order 2, then $baab = e$.

We deduce $ba = ab$, which leads to a contradiction.

Suppose that $|ba| = 4$. Then $ba(ba)^3 = e$ and $baab = e$. Consequently, $(ba)^3 = ab$. From the structure of $\mathbb{Z}_4 \times \mathbb{Z}_2$, we show easily that $(ba)^2 = \alpha^2$, so $ba = \alpha^2 ab$. As $\langle \alpha, a \rangle = G_1 \cong \mathbb{Z}_4 \times \mathbb{Z}_2$, $\langle b \rangle \cong \mathbb{Z}_2$, $|\langle \alpha, a \rangle| \times |\langle b \rangle| = 8 \times 2 = 16$ and $\langle \alpha, a \rangle \cap \langle b \rangle = \{e\}$; G is the semi-direct product:

$$G \cong (\mathbb{Z}_4 \times \mathbb{Z}_2) \rtimes_{\varphi} \mathbb{Z}_2$$

with:

$$\begin{aligned} \varphi : \mathbb{Z}_2 &\rightarrow \text{Aut}(\mathbb{Z}_4 \times \mathbb{Z}_2) \\ 1 &\rightarrow \begin{pmatrix} (1, 0) \rightarrow (1, 0) \\ (0, 1) \rightarrow (2, 1) \end{pmatrix} \end{aligned}$$

Its character presentation is $x^4 = y^2 = z^2 = e$, $xy = yx$, $zx = xz$, $zy = x^2 yz$.

B2) $G_1 \cong \mathbb{Z}_8, G_2, G_3 \cong \mathbb{Z}_4 \times \mathbb{Z}_2$: There exists an element a not in the center and of order 8 and two elements b, c not in the center of order 2 such that:

- $G_1 = \{e, a, a^2 = \alpha, a^3, a^4 = \alpha^2, \dots\}$.
- $G_2 = \{e, \alpha, \alpha^2, \alpha^3, b, b\alpha, b\alpha^2, b\alpha^3\}$ with $|b\alpha| = 4, |b\alpha^2| = 2$ and $|b\alpha^3| = 4$.
- $G_3 = \{e, \alpha, \alpha^2, \alpha^3, c, c\alpha, c\alpha^2, c\alpha^3\}$ with $|c\alpha| = 4, |c\alpha^2| = 2$ and $|c\alpha^3| = 4$.

Let us consider that $|ba| = 2$, then $baba = e$ and $baa^7b = e$. Consequently, $ba = a^7b = a^6ab = \alpha^3ab$. But α^3 , which is the commutator $[b, a]$ is of order 4; which is contradictory.

Consider $|ba| = 4$, then we show that $(ba)^2ba = a^7b = \alpha^3ab$. As $(ba)^2 = \alpha^2$; then $ba = \alpha ab$; which is also contradictory.

B3) $G_1, G_2 \cong \mathbb{Z}_8, G_3 \cong \mathbb{Z}_4 \times \mathbb{Z}_2$: There exists two elements a, b which are not in the center and of order 8 and an element c not in the center of order 2 such that:

- $G_1 = \{e, a, a^2 = \alpha, a^3, a^4 = \alpha^2, \dots\}$.
- $G_2 = \{e, b, b^2 = \alpha, b^3, b^4 = \alpha^2, \dots\}$.
- $G_3 = \{e, \alpha, \alpha^2, \alpha^3, c, c\alpha, c\alpha^2, c\alpha^3\}$ with $|c\alpha| = 4, |c\alpha^2| = 2$ and $|c\alpha^3| = 4$.

If $|ba| = 2$, then $ba = a^7b^7 = a^6abb^6 = \alpha^3aba^3 = \alpha^2ab$. We have also $|ab| = 2$. Without loss of generality, one can suppose $ab = c$. We deduce $ca = aba = a\alpha^2ab = a\alpha^2c = a^5c$. $\langle a \rangle \cong \mathbb{Z}_8, \langle c \rangle \cong \mathbb{Z}_2, |\langle a \rangle| \times |\langle c \rangle| = 8 \times 2 = 16$ and $\langle a \rangle \cap \langle c \rangle = \{e\}$. Consequently, G is the semi-direct product:

$$G \cong \mathbb{Z}_8 \rtimes_{\varphi_1} \mathbb{Z}_2,$$

where:

$$\begin{aligned} \varphi_1 : \mathbb{Z}_2 &\rightarrow \text{Aut}(\mathbb{Z}_8) \\ 1 &\rightarrow (1 \rightarrow 5) \end{aligned}$$

Its character presentation is $x^8 = y^2 = e, yx = x^5y$.

If $|ba| = 4$, then $(ba)^3 = \alpha^2ab$. As $(ba)^2 = \alpha^2$, then $ba = ab$, which is contradictory.

B4) $G_1, G_2, G_3 \cong \mathbb{Z}_8$: Then, there exists a, b not in the center of order 8 and such that ba is also of order 8. We deduce $(ba)^7 = a^7b^7 = \alpha^3ab\alpha^3 = \alpha^2ab$. As $(ba)^6 = \alpha^2$ then $ba = ab$; which leads to a contradiction.

We have classified all groups of order 16 such that $Z(G) \cong \mathbb{Z}_4$:

Name	Character presentation
$(\mathbb{Z}_4 \times \mathbb{Z}_2) \rtimes \mathbb{Z}_2$	$x^4 = y^2 = z^2 = e, xy = yx, zx = xz, zy = x^2yz$
$\mathbb{Z}_8 \rtimes_{\varphi_1} \mathbb{Z}_2$	$x^8 = y^2 = e, yx = x^5y$

Groups of order 16 with $Z(G) \cong \mathbb{Z}_4$

4.2.2 Second case: $|Z(G)| = 2$

This case is more difficult as $G/Z(G)$ is not necessarily Abelian.

In the following, we will denote $Z(G) = \{e, z\}$.

We have to study the different cases for $G/Z(G)$. $G/Z(G)$ is not cyclic. The different cases are presented below.

A) $G/Z(G) \cong \mathbb{Z}_4 \times \mathbb{Z}_2$:

The maximal subgroups of $\mathbb{Z}_4 \times \mathbb{Z}_2 = \{(0, 0), (1, 0), (2, 0), (3, 0), (0, 1), (1, 1), (2, 1), (3, 1)\}$ are:

- $\{(0, 0), (1, 0), (2, 0), (3, 0)\} \cong \mathbb{Z}_4$ and has only one subgroup of order 2.
- $\{(0, 0), (1, 1), (2, 0), (3, 1)\} \cong \mathbb{Z}_4$ and has only one subgroup of order 2.
- $\{(0, 0), (0, 1), (2, 0), (2, 1)\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ and has three subgroups of order 2.

Consequently, G has 3 subgroups G_1, G_2, G_3 of order 8 containing the center such that G_1, G_2 has only one subgroup of order 4 containing the center and G_3 has three subgroups of order 4 containing the center. We show that G_1 (resp. G_2) is Abelian. Indeed, $Z(G) < Z(G_1)$. If G_1 were not Abelian, then it has three subgroups of order 4 containing $Z(G_1)$ and consequently containing $Z(G)$; which is contradictory.

The following proposition conducts to a contradiction:

Proposition 14. *If G has at least two Abelian subgroups of order 8 then $|Z(G)| \geq 4$.*

Proof. Using:

$$|G_1G_2| = \frac{|G_1||G_2|}{|G_1 \cap G_2|} = \frac{64}{|G_1 \cap G_2|},$$

we deduce that $G = G_1G_2$ and $|G_1 \cap G_2| = 4$.

If G_1 and G_2 are Abelian, then taking $g \in G_1 \cap G_2$ and $g_1g_2 \in G$. $gg_1g_2 = g_1gg_2 = g_1g_2g$. Consequently, $G_1 \cap G_2 \subset Z(G)$. We deduce the result. \square

B) $G/Z(G) \cong (\mathbb{Z}_2)^3$: In this case, for any $g \in G$, $g^2 \in Z(G)$. As G is not Abelian, then there exists an element g such that $g^2 = z$, so $|g| = 4$. The group G has no element of order 8 nor 16. Let $\langle g \rangle = \{e, g, g^2 = z, g^3\} \cong \mathbb{Z}_4$. It is a subgroup of G containing $Z(G)$. Consequently, by the correspondance theorem, $\langle g \rangle / Z(G)$ is a subgroup of order 2 of $G/Z(G)$. Without loss of generality, one can suppose that this subgroup is identified to the subgroup $\{(0, 0, 0), (0, 0, 1)\}$ of $(\mathbb{Z}_2)^3$. This subgroup is contained to the following three subgroups of order 4:

- $\{(0, 0, 0), (0, 0, 1), (0, 1, 0), (0, 1, 1)\}$.
- $\{(0, 0, 0), (0, 0, 1), (1, 0, 0), (1, 0, 1)\}$.
- $\{(0, 0, 0), (0, 0, 1), (1, 1, 0), (1, 1, 1)\}$.

Consequently, G has three subgroups G_1, G_2, G_3 of order 8 containing $\langle g \rangle$. One can easily show, using the correspondance theorem, that for $i \neq j$ and for $g_i \in G_i \setminus \langle g \rangle$, $g_j \in G_j \setminus \langle g \rangle$ then $g_i g_j \in G_k \setminus \langle g \rangle$ for $k \notin \{i, j\}$. From the proposition 14, the maximum number of Abelian subgroups of order 8 is 1. Consequently, one can suppose that G_2 and G_3 are non-Abelian. Consequently, one can take $g_2 \in G_2 \setminus \langle g \rangle$ (resp. $g_3 \in G_3 \setminus \langle g \rangle$) which does not commute with g . As $G/Z(G)$ is Abelian, then $D(G) = Z(G)$. Consequently, $gg_2 = zg_2g$ and $gg_3 = zg_3g$. We deduce $gg_2g_3 = zg_2gg_3 = zg_2zg_3g = z^2g_2g_3g = g_2g_3g$. Consequently, g commutes with all elements of $G_1 \setminus \langle g \rangle$; and as a consequence, $g \in Z(G_1)$. $Z(G_1)$ contains $\{e, z, g\}$. We deduce that G_1 is Abelian. As G does not contains element of order 8 and as $g \in G_1$ is of order 4 then $G_1 \cong \mathbb{Z}_4 \times \mathbb{Z}_2$. Let us denote:

$G_1 = \{e, g, g^2 = z, g^3, g_1, gg_1, g^2g_1, g^3g_1\}$. Let $g_2 \in G_2 \setminus \langle g \rangle$ (corresponds to a symmetry in D_4 or one of the four other elements of order 4 in \mathbb{H}). If one of the 4 elements of $G_1 \setminus \langle g \rangle$ commutes with one of the elements of $G_2 \setminus \langle g \rangle$, then it commutes with all elements of G_2 and the center of G_2 has at least 3 elements, which is contradictory. Consequently; none element of $G_1 \setminus \langle g \rangle$ commutes with one element of $G_2 \setminus \langle g \rangle$.

Let $g_1 \in G_1 \setminus \langle g \rangle$ and $g_2 \in G_2 \setminus \langle g \rangle$, then $g_1g_2 = zg_2g_1$. We show that $gg_1g_2 = gzg_2g_1 = zgg_2g_1 = z^2g_2gg_1 = g_2gg_1$. Consequently, g_2 commutes with gg_1 ; it leads to a contradiction.

C) $G/Z(G) \cong \mathbb{H}$:

The maximal subgroups of $\mathbb{H} = \{1, -1, i, -i, j, -j, k, -k\}$ are:

- $\{1, i, -1, -i\} \cong \mathbb{Z}_4$, which has only one subgroup of order 2.
- $\{1, j, -1, -j\} \cong \mathbb{Z}_4$, which has only one subgroup of order 2.

- $\{1, k, -1, -k\} \cong \mathbb{Z}_4$, which has only one subgroup of order 2.

Consequently, G has three Abelian subgroups of order 8. Using the proposition 14, it leads to a contradiction.

D) $G/Z(G) \cong D_4$:

The maximal subgroups of $D_4 = \{Id, r, r^2, r^3, s, rs, r^2s, r^3s\}$ are:

- $\{Id, r, r^2, r^3\} \cong \mathbb{Z}_4$ which has only one subgroup of order 2.
- $\{Id, s, r^2, r^3s\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ which has three subgroups of order 2.
- $\{Id, rs, r^2, r^3s\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ which has three subgroups of order 2.

Consequently, G has three subgroups of order 8, G_1, G_2, G_3 which contain the center such that G_1 has only one subgroup of order 4 which contains the center and G_2, G_3 have three subgroups of order 4 which contain the center. G_1 is Abelian and G_2, G_3 are necessarily non-Abelian. We show easily $Z(G) = Z(G_2) = Z(G_3)$ as the center of a non-Abelian group of order 8 has two elements. If $G_2 \cong D_4$, as $r^2 = rsr^3s$, then $r^2 \in D(G)$. If $G_2 \cong \mathbb{H}$, as $-1 = ij(-i)(-j)$, then $-1 \in D(G)$. In all cases, $Z(G) < D(G)$ and the inclusion is strict as $G/Z(G)$ is non-Abelian. As, in D_4 , s and rs do not commute; then there exists $g_2 \in G_2$ and $g_3 \in G_3$ such that $g_2Z(G)$ (identified to s) and $g_3Z(G)$ (identified to rs) do not commute. Consequently, there exists $g' \in G$ (such $g'Z(G)$ is identified to r^2) such $g_2g_3Z(G) = g_3g_2g'Z(G)$. We deduce that there exists $z_0 \in Z(G)$ such $g_2g_3 = g_3g_2g'z_0$. Consequently, $g'z_0 \in D(G)$ and as $z_0 \in D(G)$, we deduce that $g' \in D(G)$.

We have $\{e, g', z, g'z\} \subset D(G)$ and forms a group. Let us show that this group is a normal subgroup of G . We have $hg'zh^{-1}Z(G) = hg'h^{-1}Z(G) = g'Z(G)$ as $g'Z(G) \in Z(G/Z(G))$. Consequently $hg'h^{-1}, hg'zh^{-1} \in \{g', g'z\}$; we deduce that $\{e, g', z, g'z\}$ is a normal subgroup of G . As $G/\{e, g', z, g'z\}$ is of order 4, it is necessarily Abelian.

Consequently, $D(G) = \{e, g', z, g'z\}$. Moreover, $D(G)/Z(G) = Z(G/Z(G))$ and corresponds to $\{Id, r^2\}$. We deduce that $D(G) < G_i$ for any $i = 1, 2, 3$. Using the formula $|G_iG_j| = \frac{|G_i||G_j|}{|G_i \cap G_j|}$, we deduce $G = G_iG_j$ and $G_i \cap G_j = D(G)$.

D1) $D(G) \cong \mathbb{Z}_2 \times \mathbb{Z}_2$:

G_1 cannot be isomorphic to $(\mathbb{Z}_2)^3$, as any subgroup of order 2 is included in three subgroups of order 4. The only possibility is $G_1 \cong \mathbb{Z}_4 \times \mathbb{Z}_2$. The unique subgroup of order 4 containing the center is then identified to $\{(0, 0), (2, 0), (0, 1), (2, 1)\}$. As

the square of any element of $\mathbb{Z}_4 \times \mathbb{Z}_2$ is either $(0, 0)$ or $(2, 0)$ and $(2, 0)$ is in three subgroups of order 4; we deduce that there exists an element a of order 4 in $G \setminus D(G)$ such that:

- $G_1 = \{e, a, a^2 = g', a^3, z, az, a^2z = g'z, a^3z\} \cong \mathbb{Z}_4 \times \mathbb{Z}_2$.

As \mathbb{H} has no subgroup isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2$; we deduce $G_2, G_3 \cong D_4$. As $Z(G) = Z(G_2) = Z(G_3)$, then there exists b, c in $G \setminus D(G)$ of order 4 such that:

- $G_2 = \{e, b, b^2 = z, b^3, g', bg' = g'b^3, b^2g' = zg', b^3g' = g'b\}$.

- $G_3 = \{e, c, c^2 = z, c^3, g', cg' = g'c^3, c^2g' = zg', c^3g' = g'c\}$.

By correspondance theorem, we show that $baZ(G) = abg'Z(G) = g'abZ(G)$.

Suppose that $ba = g'ab$. If $|ba| = 2$, then $ba = a^3b^3 = a^2abb^2 = g'abz = g'zab$. Consequently, $g' = g'z$, which is contradictory. We have $ab = g'ba$. If we suppose that $|ba| = 4$, as g' is a symmetry, then $|ab| = 2$ and it conducts to the same contradiction.

Suppose that $ba = g'zab$. If $|ba| = 4$, then $(ba)^2ba = a^3b^3 = g'abz = g'zab$. As $(ba)^2 = z$ then $ba = g'ab$ which is contradictory. We have $ab = g'zba$. If $|ba| = 2$, as ba is not in $D(G)$, we deduce $|ab| = 4$ which leads to the same contradiction.

Consequently, it is impossible that $D(G) \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.

D2) $D(G) \cong \mathbb{Z}_4$:

In this case, $D(G) = \{e, g', g'^2 = z, g'^3\}$. We have seen that G_1 cannot be isomorphic to $(\mathbb{Z}_2)^3$. However, as $\mathbb{Z}_4 \times \mathbb{Z}_2$ has two groups isomorphic to \mathbb{Z}_4 which contains a common subgroup of order 2; G_1 cannot be isomorphic to $\mathbb{Z}_4 \times \mathbb{Z}_2$ (as the center will be included in two subgroups of order 4). Consequently, $G_1 \cong \mathbb{Z}_8$ and there exists $a \in G \setminus D(G)$ of order 8 such that:

- $G_1 = \{e, a, a^2 = g', a^3, a^4 = z, \dots\}$.

D2-1) $G_2, G_3 \cong D_4$:

There exists b, c not in $D(G)$ and of order 2 such that:

- $G_2 = \{e, g', g'^2 = z, g'^3, b, g'b = bg'^3, g'^2b, g'^3b = bg'\}$.

- $G_3 = \{e, g', g'^2 = z, g'^3, c, g'c = cg'^3, g'^2c, g'^3c = cg'\}$.

By the correspondance theorem and the structure of the maximal subgroups of D_4 , we deduce that $ba \in G^3 \setminus D(G)$. Consequently, $|ba| = 2$ and $ba = a^7b$. $\langle a \rangle = G_1 \cong \mathbb{Z}_8$, $\langle b \rangle \cong \mathbb{Z}_2$, $|\langle a \rangle \times \langle b \rangle| = 8 \times 2 = 16$ and $\langle a \rangle \cap \langle b \rangle = \{e\}$. Consequently, G is the semi-direct product, also called diedral group D_8 :

$$G \cong \mathbb{Z}_8 \rtimes_{\varphi_2} \mathbb{Z}_2,$$

with:

$$\begin{aligned} \varphi_2 : \mathbb{Z}_2 &\rightarrow \text{Aut}(\mathbb{Z}_8) \\ 1 &\rightarrow (1 \rightarrow 7) \end{aligned}$$

Its character presentation is $x^8 = y^2 = e$, $yx = x^7y$.

D2-2) $G_2 \cong D_4$, $G_3 \cong \mathbb{H}$:

There exists b, c not in $D(G)$ respectively of order 2 and 4 such that:

- $G_2 = \{e, g', g'^2 = z, g'^3, b, g'b = bg'^3, g'^2b, g'^3b = bg'\}$.
- $G_3 = \{e, g', g'^2 = z, g'^3, c, g'c, g'^2c, g'^3c\}$.

$ba \in G_3 \setminus D(G)$, then $|ba| = 4$. Consequently, $(ba)^2ba = a^7b$. As $(ba)^2 = g'^2 = z = a^4$, then $ba = a^3b$. With the same method as previously, we deduce that G is the semi-direct product (also called semi-diedral group SD_8):

$$G \cong \mathbb{Z}_8 \rtimes_{\varphi_3} \mathbb{Z}_2,$$

with:

$$\begin{aligned} \varphi_3 : \mathbb{Z}_2 &\rightarrow \text{Aut}(\mathbb{Z}_8) \\ 1 &\rightarrow (1 \rightarrow 3) \end{aligned}$$

Its character presentation is $x^8 = y^2 = e$, $yx = x^3y$.

D2-3) $G_2, G_3 \cong \mathbb{H}$:

There exists b and c not in $D(G)$ of order 4 such that:

- $G_2 = \{e, g', g'^2 = z, g'^3, b, g'b, g'^2b, g'^3b\}$.
- $G_3 = \{e, g', g'^2 = z, g'^3, c, g'c, g'^2c, g'^3c\}$.

$|ba| = 4$ then $(ba)^2ba = a^7b^3$. As $(ba)^2 = z = a^4$ then $ba = a^3b^3$. It is not a semi-direct product. Its character presentation is $x^8 = y^4 = e, x^4 = y^2, yx = x^3y^3$.

The following Table gives the groups of order 16 such that $Z(G) \cong \mathbb{Z}_2$:

Name	Character presentation
$\mathbb{Z}_8 \rtimes_{\varphi_2} \mathbb{Z}_2$	$x^8 = y^2 = e, yx = x^7y$
$\mathbb{Z}_8 \rtimes_{\varphi_3} \mathbb{Z}_2$	$x^8 = y^2 = e, yx = x^3y$
Generalized Quaternions	$x^8 = y^4 = e, x^4 = y^2, yx = x^3y^3$

5 Conclusion

The groups of order 16 are given in the following table:

Name	Character presentation	Center
\mathbb{Z}_{16}	$a^{16} = e$	$Z(G) = G$
$\mathbb{Z}_8 \times \mathbb{Z}_2$	$a^8 = b^2 = e, ba = ab$	$Z(G) = G$
$\mathbb{Z}_4 \times \mathbb{Z}_4$	$a^4 = b^4 = e, ba = ab$	$Z(G) = G$
$\mathbb{Z}_4 \times (\mathbb{Z}_2)^2$	$a^4 = b^2 = c^2 = e, D(G) = \{e\}$	$Z(G) = G$
$(\mathbb{Z}_2)^4$	$a^2 = b^2 = c^2 = d^2 = e, D(G) = \{e\}$	$Z(G) = G$
$D_4 \times \mathbb{Z}_2$	$x^4 = y^2 = z^2 = e, yx = x^3y, zx = xz, zy = yz$	$\{e, x^2, z, zx^2\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$
$(\mathbb{Z}_2 \times \mathbb{Z}_2) \rtimes \mathbb{Z}_4$	$x^4 = y^4 = e, yx = x^3y^3$	$\{e, x^2, y^2, x^2y^2\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$
$\mathbb{Z}_4 \rtimes \mathbb{Z}_4$	$x^4 = y^4 = e, yx = x^3y$	$\{e, x^2, y^2, x^2y^2\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$
$\mathbb{H} \times \mathbb{Z}_2$	$x^4 = y^4 = z^2 = e, x^2 = y^2, yx = x^3y, zx = xz, zy = yz$	$\{e, x^2, z, x^2z\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$
$(\mathbb{Z}_4 \times \mathbb{Z}_2) \rtimes \mathbb{Z}_2$	$x^4 = y^2 = z^2 = e, xy = yx, zx = xz, zy = x^2yz$	$\{e, x, x^2, x^3\} \cong \mathbb{Z}_4$
$\mathbb{Z}_8 \rtimes_{\varphi_1} \mathbb{Z}_2$	$x^8 = y^2 = e, yx = x^5y$	$\{e, x^2, x^4, x^6\} \cong \mathbb{Z}_4$
$\mathbb{Z}_8 \rtimes_{\varphi_2} \mathbb{Z}_2$	$x^8 = y^2 = e, yx = x^7y$	$\{e, x^4\} \cong \mathbb{Z}_2$
$\mathbb{Z}_8 \rtimes_{\varphi_3} \mathbb{Z}_2$	$x^8 = y^2 = e, yx = x^3y$	$\{e, x^4\} \cong \mathbb{Z}_2$
Gen. Quat.	$x^8 = y^4 = e, x^4 = y^2, yx = x^3y^3$	$\{e, x^4\} \cong \mathbb{Z}_2$