

# A short proof of the Strong Normalization of Classical Natural Deduction with Disjunctio

René David, Karim Nour

► **To cite this version:**

René David, Karim Nour. A short proof of the Strong Normalization of Classical Natural Deduction with Disjunctio. *The Journal of Symbolic Logic*, 2003, 68 (4), pp.1277-1288. hal-00381601

**HAL Id: hal-00381601**

**<https://hal.archives-ouvertes.fr/hal-00381601>**

Submitted on 6 May 2009

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A short proof of the Strong Normalization of Classical Natural Deduction with Disjunction

René David and Karim NOUR  
LAMA - Equipe de Logique  
Université de Chambéry  
73376 Le Bourget du Lac  
e-mail: {david,nour}@univ-savoie.fr

## Abstract

We give a direct, purely arithmetical and elementary proof of the strong normalization of the cut-elimination procedure for full (i.e. in presence of all the usual connectives) classical natural deduction.

## 1 Introduction

This paper gives a direct proof of the strong normalization of the cut-elimination procedure for full propositional classical logic. By full we mean that all the connectives ( $\rightarrow$ ,  $\wedge$  and  $\vee$ ) and  $\perp$  (for the absurdity) are considered as primitive and they have their intuitionistic meaning. As usual, the negation is defined by  $\neg A = A \rightarrow \perp$ .

It is well known that, when the underlying logic is the classical one (i.e. the absurdity rule is allowed) these connectives are redundant since, for example,  $\vee$  and  $\wedge$  can be coded by using  $\rightarrow$  and  $\perp$ . From a logical point of view, considering the full logic is thus somehow useless. However, from the computer science point of view, considering the full logic is interesting because, by the so-called Curry-Howard isomorphism, formulas can be seen as types for functional programming languages and correct programs can be extracted from proofs. For that reason various systems have been studied in the last decades (see, for example, [2, 4, 5, 12, 14, 15, 17, 22]) both for intuitionistic and classical logic. The connectives  $\wedge$  and  $\vee$  have a functional counter-part ( $\wedge$  corresponds to a product and  $\vee$  to a co-product, i.e. a *case of*) and it is thus useful to have them as primitive.

Until very recently (see the introduction of [7] for a brief history), no proof of the strong normalization of the cut-elimination procedure was known for full logic. In [7], de Groote gives such a proof by using a CPS-style transformation from full classical logic to intuitionistic logic with  $\rightarrow$  as the only connective, i.e. the simply typed  $\lambda$ -calculus. A very elegant and direct proof of the strong normalization of the full logic is given in [11] but only the intuitionistic case is given.

We give here another proof of de Groote's result. This proof is based on a proof of the strong normalization of the simply typed  $\lambda$ -calculus due to the first author (see [8]) which, itself, is a simplification of the one given by Matthes in [11]. After this paper had been written we were told by Curien and some others that this kind of technique was already present in van Daalen (see [27]) and Levy (see [13]). The same idea is used in [10] to give a short proof of the strong normalization of the simply typed  $\lambda\mu$ -calculus of [17]. Apart the fact that this proof is direct (i.e. uses no translation into an other system whose strong normalization is known) and corresponds to the intuition (the main argument of the proof is an induction on the complexity of the cut-formula) we believe that our technique is quite general and may be used in other circumstances. A crucial lemma of our proof is used in [16] to give a semantical proof of the strong normalization. Finally [9] uses

the same technique to give an elementary proof of the strong normalization of a typed  $\lambda$ -calculus with explicit substitutions which, from the logical point of view, correspond to explicit cuts and weakenings.

## 2 The typed system

We code proofs by using a set of terms (denoted  $\mathcal{T}$ ) which extends the  $\lambda\mu$ -terms of Parigot [17] and is given by the following grammar where  $x, y, \dots$  are (intuitionistic) variables and  $a, b, \dots$  are (classical) variables:

$$\begin{aligned} \mathcal{T} ::= & x \mid \lambda x \mathcal{T} \mid (\mathcal{T} \ \mathcal{E}) \mid \langle \mathcal{T}, \mathcal{T} \rangle \mid \omega_1 \mathcal{T} \mid \omega_2 \mathcal{T} \mid \mu a \mathcal{T} \mid (a \ \mathcal{T}) \\ \mathcal{E} ::= & \mathcal{T} \mid \pi_1 \mid \pi_2 \mid [x.\mathcal{T}, y.\mathcal{T}] \end{aligned}$$

The meaning of the new constructors is given by the typing rules of figure 1 below where  $\Gamma$  is a context, i.e. a set of declarations of the form  $x : A$  and  $a : \neg A$  where  $x$  is an intuitionistic variable,  $a$  is a classical variable and  $A$  is a formula.

Note that, since we only are concerned with the logical point of view, we should only consider typed terms, i.e. use a  $\lambda$ -calculus à la Church. However, for the simplicity of notation, the set of terms has been given in an untyped formalism i.e. we use a  $\lambda$ -calculus à la Curry.

$$\begin{array}{c} \frac{}{\Gamma, x : A \vdash x : A} \text{ax} \quad \frac{\Gamma_1 \vdash M : A \quad \Gamma_2 \vdash N : B}{\Gamma_1, \Gamma_2 \vdash \langle M, N \rangle : A \wedge B} \wedge_i \quad \frac{\Gamma \vdash M : A_1 \wedge A_2}{\Gamma \vdash (M \ \pi_i) : A_i} \wedge_e \\ \frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash \lambda x M : A \rightarrow B} \rightarrow_i \quad \frac{\Gamma_1 \vdash M : A \rightarrow B \quad \Gamma_2 \vdash N : A}{\Gamma_1, \Gamma_2 \vdash (M \ N) : B} \rightarrow_e \\ \frac{\Gamma \vdash M : A_i}{\Gamma \vdash \omega_i M : A_1 \vee A_2} \vee_i \\ \frac{\Gamma \vdash M : A_1 \vee A_2 \quad \Gamma_1, x_1 : A_1 \vdash N_1 : C \quad \Gamma_2, x_2 : A_2 \vdash N_2 : C}{\Gamma, \Gamma_1, \Gamma_2 \vdash (M [x_1.N_1, x_2.N_2]) : C} \vee_e \\ \frac{\Gamma, a : \neg A \vdash M : \perp}{\Gamma \vdash \mu a M : A} \text{abs}_i \quad \frac{\Gamma, a : \neg A \vdash M : A}{\Gamma \vdash (a \ M) : \perp} \text{abs}_e \end{array}$$

Figure 1.

This coding is essentially the same as in [1] and [7]. We have adopted the notations of [1] which are also used by [11]: what is written  $\pi_i M$  in [7] is written  $(M \ \pi_i)$  here and  $\delta(M, x_1.N_1, x_2.N_2)$  in [7] is written  $(M [x_1.N_1, x_2.N_2])$  here. These notations have the advantage of making the permutative and classical reduction rules more uniform and thus simplifies the proofs.

The cut-elimination procedure corresponds to the reduction rules given below. There are three kinds of cuts.

*Logical cuts:* they appear when the introduction of a connective ( $\rightarrow$ ,  $\wedge$  and  $\vee$ ) is immediately followed by its elimination. The corresponding rules are:

- $(\lambda x M \ N) \triangleright M[x := N]$
- $(\langle M_1, M_2 \rangle \ \pi_i) \triangleright M_i$
- $(\omega_i M [x_1.N_1, x_2.N_2]) \triangleright N_i[x_i := M]$

*Permutative cuts :* they appear when the elimination rule of the disjunction is followed by the elimination rule of a connective. They are considered as cuts because a logical cut may be hidden by the  $\vee_e$  rule. Considering these cuts is necessary to get the sub-formula property. The corresponding rule is:

- $(M [x_1.N_1, x_2.N_2] \varepsilon) \triangleright (M [x_1.(N_1 \varepsilon), x_2.(N_2 \varepsilon)])$

*Classical cuts* : they appear when the classical rule is followed by the elimination rule of a connective. The corresponding rule is:

- $(\mu a M \varepsilon) \triangleright \mu a M[a :=^* \varepsilon]$  where  $M[a :=^* \varepsilon]$  is obtained by replacing each sub-term of  $M$  of the form  $(a N)$  by  $(a (N \varepsilon))$ .

**Notation 2.1** *Let  $M$  be in  $\mathcal{E}$ .*

1.  $M \triangleright M'$  means that  $M$  reduces to  $M'$  by using one step of the reduction rules given above. As usual,  $\triangleright^+$  (resp.  $\triangleright^*$ ) is the transitive (resp. reflexive and transitive) closure of  $\triangleright$ .
2.  $M$  is strongly normalizable (this is denoted by  $M \in SN$ ) if there is no infinite sequence of  $\triangleright$  reductions.

*Remark* If  $M = [y_1.N_1, y_2.N_2]$ ,  $M \triangleright M'$  means that  $M'$  is either  $[y_1.N'_1, y_2.N_2]$  or  $[y_1.N_1, y_2.N'_2]$  where  $N_1 \triangleright N'_1$  or  $N_2 \triangleright N'_2$ . It is thus clear that  $M \in SN$  iff  $N_1, N_2 \in SN$ .

The following result is straightforward.

**Lemma 2.1 (Subject reduction)** *If  $\Gamma \vdash M : A$  and  $M \triangleright^* N$  then  $\Gamma \vdash N : A$ .*

The goal of this paper is the proof of theorem 2.1 below.

**Theorem 2.1** *Every typed term is strongly normalizable.*

The proof is an immediate corollary of theorem 4.1: if  $M, N \in SN$ , then  $M[x := N] \in SN$ .

The proof of theorem 4.1 uses a characterization of strongly normalizable (theorem 3.1): a term is in  $SN$  iff its arguments and head reduct (see definition 3.3) are in  $SN$ . This theorem needs another result (theorem 3.2) which is, intuitively, very clear but whose formal proof needs some work.

The main difficulties are the following:

- The first one is minor: in the  $\lambda$ -calculus, each term has a unique *head*, either a head variable or a head redex. Due to the connective  $\wedge$ , this is no longer true here and a term may have both a head variable and a head redex. This is treated by showing that it is enough to consider only the simple terms (see definition 3.1).

- The second one is crucial and due to the presence of critical pairs such as  $(\mu a M [y_1.N_1, y_2.N_2] \varepsilon)$ . We can choose as head redex either the classical one or the permutative one. If we choose the classical one, the proof of theorem 3.1 will be easy but the proof of theorem 4.1 does not work because, in the rule  $\vee_e$ , the type of the main hypothesis has nothing to do with the type of its conclusion. We thus have to choose, as head redex, the permutative one but then, theorem 3.1 needs the difficult theorem 3.2. For the same reason (the rule  $\vee_e$ ), the proof of theorem 4.1 needs a rather complex induction: we use a 5-tuple of integers. Note that E. Tahhan Bittar [3] has given a proof of the strong normalization of the *sequent calculus* by using essentially the same 5-tuple of integers.

*Remark* It is also for simplicity of proofs that, in the totality of this section, we only consider typed terms and thus, for example, that terms such as  $(\lambda x M [x_1.N_1, x_2.N_2])$  are not allowed because they, obviously, cannot be typed since the type of  $\lambda x M$  must be an implication. Actually, theorems 3.1 and 3.2 would also be true for untyped terms i.e. even if terms as  $(\lambda x M [x_1.N_1, x_2.N_2])$  were allowed and its proof will be essentially the same since such a term is strongly normalizable iff  $M, N_1, N_2$  also are strongly normalizable.

### 3 Characterization of strongly normalizable terms

**Definition 3.1** 1. A term  $M$  is simple if  $M$  either is a variable or an application.

2. The set of contexts is given by the following grammar:

$$C := *_i \mid \lambda x C \mid \omega_i C \mid \langle C_1, C_2 \rangle \mid \mu a C$$

3. If  $C$  is a context with holes  $*_1, \dots, *_n$  and  $M_1, \dots, M_n$  are terms,  $C[M_1, \dots, M_n]$  is the term obtained by replacing each  $*_i$  by  $M_i$ .

**Lemma 3.1** Each term  $M$  can be uniquely written as  $C[M_1, \dots, M_n]$  where  $C$  is a context and  $M_1, \dots, M_n$  are simple terms.

**Proof** By induction on  $M$ .  $\square$

**Lemma 3.2** Let  $C$  be a context and  $M_1, \dots, M_n$  be terms. Then  $C[M_1, \dots, M_n] \in SN$  iff  $M_1, \dots, M_n \in SN$ .

**Proof** By induction on  $C$ .  $\square$

**Definition 3.2 (and notation)** A (possibly empty) sequence  $\vec{N} = N_1, \dots, N_n$  of elements of  $\mathcal{E}$  is nice if each  $N_i \in \mathcal{T} \cup \{\pi_1, \pi_2\}$  except possibly for  $i = n$ . If  $M$  is a term,  $(M N_1 \dots N_n)$  will be denoted as  $(M \vec{N})$ .

**Lemma 3.3 (and definition)** Let  $M$  be a simple term.

1. Then  $M$  can be uniquely written as one of the cases of the figure below where  $\vec{T} = T_1, \dots, T_n$  is a nice sequence and, in case (4) and (5),  $\varepsilon \vec{T}$  is also nice, i.e. if  $\varepsilon = [y_1.N_1, y_2.N_2]$  then  $\vec{T}$  is empty.
2. The set of arguments of  $M$  (denoted as  $\text{arg}(M)$ ) and the head of  $M$  (denoted as  $\text{hd}(M)$ ), either a redex or a variable, are defined by the figure below.
3. The head reduct of  $M$  (denoted as  $\text{hred}(M)$ ) is the term obtained by reducing, if any, the head redex of  $M$ .

	$M$	$\text{hd}(M)$	$\text{arg}(M)$
0	$(x \vec{T})$ or $(a T)$	$x$ or $a$	$\{T_1, \dots, T_n\}$ or $T$
1	$(\lambda x N O \vec{T})$	$(\lambda x N O)$	$\{O\}$
2	$(\langle N_1, N_2 \rangle \pi_i \vec{T})$	$(\langle N_1, N_2 \rangle \pi_i)$	$\{N_1, N_2\}$
3	$(\omega_i N [x_1.O_1, x_2.O_2])$	$M$	$\{N, O_1, O_2\}$
4	$(\mu a N \varepsilon \vec{T})$	$(\mu a N \varepsilon)$	$\{\varepsilon\}$
5	$(N [x_1.O_1, x_2.O_2] \varepsilon \vec{T})$	$(N [x_1.O_1, x_2.O_2] \varepsilon)$	$\emptyset$

**Proof** Since  $M$  is simple, and for trivial typing reasons, it looks like either (a)  $(x \vec{S})$  or (b)  $(a S)$  or (c)  $(\lambda x N O \vec{S})$  or (d)  $(\langle N_1, N_2 \rangle \pi_i \vec{S})$  or (e)  $(\omega_i N [x_1.O_1, x_2.O_2] \vec{S})$  or (f)  $(\mu a N \varepsilon \vec{S})$ . If  $\vec{S}$  is empty the result is clear.

Otherwise, assume first  $\vec{S}$  is nice. The cases (a), (b) and (c) are clear. Case (d) gives (5). Case (e) gives (5) if  $\varepsilon = [y_1.N_1, y_2.N_2]$  or (4) otherwise.

Assume finally  $\vec{S}$  is not nice. Then  $\vec{S}$  can be written as  $\vec{S}_1[y_1.N_1, y_2.N_2]\vec{S}_2$  where  $\vec{S}_2$  is nice and non empty. It is then easy to see that, in all cases, this gives (5) where  $\vec{S}_2 = \varepsilon \vec{T}$ .

For uniqueness, check easily (by looking whether  $\vec{T}$  has an  $[y_1.N_1, y_2.N_2]$  or not) that if  $M$  is in case 0 to 4 it cannot also be in case 5.  $\square$

**Theorem 3.1** *Let  $M$  be a simple term. If  $M$  has an head redex, then  $M \in SN$  iff  $arg(M) \subset SN$  and  $hred(M) \in SN$ . Otherwise,  $M \in SN$  iff  $arg(M) \subset SN$ .*

**Proof** The case of an head variable is trivial. Case 1 of the figure of lemma 3.3 is done as follows. Since  $hred(M) \in SN$ ,  $N$  and  $\vec{T}$  are in  $SN$ . Thus and since  $\vec{T}$  is nice, an infinite reduction of  $M$  must look like:  $M \triangleright^* (\lambda x N_1 O_1 \vec{T}_1) \triangleright (N_1[x := O_1] \vec{T}_1) \triangleright \dots$ . The contradiction comes from the fact (see lemma 3.4 below) that  $hred(M) \triangleright^* (N_1[x := O_1] \vec{T}_1)$ . Cases 2, 3, 4 are similar.

Case 5 is theorem 3.2 below.  $\square$

**Lemma 3.4** *Let  $M, N \in \mathcal{T}$ . Assume  $M \triangleright M'$  and  $N \triangleright N'$ . Let  $\sigma$  (resp.  $\sigma'$ ) be either  $[x := N]$  or  $[a :=^* N]$  (resp.  $[x := N']$  or  $[a :=^* N']$ ). Then  $M[\sigma] \triangleright M'[\sigma]$  and  $M[\sigma] \triangleright^* M'[\sigma']$ .*

**Proof** Straightforward.  $\square$

**Theorem 3.2** *Assume the sequence  $\varepsilon \vec{V}$  is nice and  $S_2 = (N [x_1. (N_1 \varepsilon), x_2. (N_2 \varepsilon)] \vec{V}) \in SN$ . Then  $S_1 = (M [x_1. N_1, x_2. N_2] \varepsilon \vec{V}) \in SN$ .*

**Proof** See section 5.  $\square$

## 4 Proof of theorem 2.1

By induction on  $M$ . The cases  $x$ ,  $\lambda x N$ ,  $\langle N, O \rangle$ ,  $\omega_i N$ ,  $(a N)$  and  $\mu a N$  are immediate. The last case is  $M = (N \varepsilon) = (x \varepsilon)[x := N]$  where  $x$  is a fresh variable and the result follows from the induction hypothesis and theorem 4.1 below.  $\square$

**Definition 4.1** *Let  $M$  be a term. Then,  $cxy(M)$  is the number of symbols occurring in  $M$  and, if  $M \in SN$ ,  $\eta(M)$  is the length of the longest reduction of  $M$ .*

In lemma 4.1 and theorem 4.1 below,  $\sigma$  denotes a substitution of the form  $[x_i := N_i / i = 1 \dots n]$ , i.e. we substitute only intuitionistic variables.

**Lemma 4.1** *Let  $M$  be a simple term with an head redex and  $\sigma$  be a substitution. Then,  $hd(M[\sigma]) = hd(M)[\sigma]$ ,  $arg(M[\sigma]) = arg(M)[\sigma]$  and  $hred(M[\sigma]) = hred(M)[\sigma]$ .*

**Proof** Immediate.  $\square$

**Theorem 4.1** *Let  $M \in SN$  be a term and  $\sigma$  be a substitution. Assume that the substituted variables all have the same type and, for all  $x$ ,  $\sigma(x) \in SN$ . Then  $M[\sigma] \in SN$ .*

**Proof**

The proof is by induction on  $(lgt(\sigma), \eta(M), cxy(M), \eta(\sigma), cxy(\sigma))$  where  $lgt(\sigma)$  is the number of connectives in the type of the substituted variables and  $\eta(\sigma)$  (resp.  $cxy(\sigma)$ ) is the sum of the  $\eta(N)$  (resp.  $cxy(N)$ ) for the  $N$  that are actually substituted, i.e. for example if  $\sigma = [x := N]$  and  $x$  occurs  $n$  times in  $M$ , then  $\eta(\sigma) = n \cdot \eta(N)$  and  $cxy(\sigma) = n \cdot cxy(N)$ . The induction hypothesis will be abbreviated as *IH*.

By the *IH* and lemmas 3.2 and 3.1 we may assume that  $M$  is simple. Consider then the various cases of lemma 3.3.

- If  $M$  has an head redex: by lemma 4.1 and the *IH*,  $arg(M[\sigma]) \subseteq SN$  since for each  $N \in arg(M)$ ,  $cxy(N) < cxy(M)$ . By lemma 4.1,  $hred(M[\sigma]) = hred(M)[\sigma]$  and thus, since  $\eta(hred(M)) < \eta(M)$ ,  $hred(M[\sigma]) \in SN$  follows from the *IH*.

- Otherwise, if the head variable is a classical variable or an intuitionistic variable not in the domain of  $\sigma$ , the result is trivial.
- Otherwise, i.e  $M = (x \overrightarrow{T})$ 
  - If  $hd(M[\sigma])$  is a variable, the result is trivial.
  - If  $hd(M[\sigma]) = hd(\sigma(x))$ : let  $M' = z \overrightarrow{T}$  where  $z$  is a fresh variable and  $\sigma'$  be the substitution defined as follows  $\sigma'(z) = hred(\sigma(x))$  and, for the variables  $y$  occurring in  $\overrightarrow{T}$ ,  $\sigma'(y) = \sigma(y)$ . Then,  $hred(M[\sigma]) = M'[\sigma']$  and thus, by the *IH*,  $hred(M[\sigma]) \in SN$  since  $\eta(\sigma') < \eta(\sigma)$ .
  - Otherwise, the head redex has been created by the substitution. The various cases are:
    1.  $M = (x O \overrightarrow{S})$  and  $\sigma(x) = \lambda y N$ . By the *IH*,  $arg(M[\sigma]) \subseteq SN$  and thus, by theorem 3.1, we have to show that  $P = (N[y := O[\sigma]] \overrightarrow{S[\sigma]}) \in SN$ . By the *IH*,  $(z \overrightarrow{S[\sigma]}) \in SN$  and since  $lgt(O[\sigma]) < lgt(\lambda y N)$ ,  $N[y := O[\sigma]] \in SN$ . Thus  $P = (z \overrightarrow{S[\sigma]})[z := N[y := O[\sigma]]] \in SN$  since  $lgt(N[y := O[\sigma]]) < lgt(\lambda y N)$ .
    2.  $M = (x \pi_i \overrightarrow{S})$  and  $\sigma(x) = \langle N_1, N_2 \rangle$  or  $M = (x [x_1.M_1, x_2.M_2])$  and  $\sigma(x) = \omega_i N$ . The proof is similar.
    3.  $M = (x [x_1.M_1, x_2.M_2])$  and  $\sigma(x) = \mu a N$ . By the *IH*,  $arg(M[\sigma]) \subseteq SN$  and thus (by theorem 3.1) we have to show  $P = \mu a N[a :=^* [x_1.P_1, x_2.P_2]] \in SN$  where, for  $i = 1, 2$ ,  $P_i = M_i[\sigma]$ . Since  $cxy(M_i) < cxy(M)$ , the fact that  $P_i \in SN$  follows from the *IH*. The result is thus a particular case of the claim below.

*Claim* Let  $P_1, P_2, T \in SN$  and  $a_1, \dots, a_n$  be variables of type  $\neg(A \vee B)$ . Let  $T[\tau]$  denotes  $T[a_i :=^* [P] / i = 1..n]$  where  $[P]$  is an abbreviation for  $[x_1.P_1, x_2.P_2]$ . Then  $T[\tau] \in SN$ .

*Proof* By induction on  $(\eta(T), cxy(T))$ . We may assume that  $T$  is simple. Consider the various cases of lemma 3.3.

- If  $T$  has an head redex, the result follows immediately from *IH* and lemma 4.1.
- Otherwise and if the head variable of  $T$  is not in  $\tau$ , the result is trivial.
- Otherwise and because of the type of the  $a_i$ ,  $T = (a V)$  where  $V \in \mathcal{T}$ . It is thus enough to prove that  $(V[\tau] [P]) \in SN$  and, for that, it is enough to show that its head reduct  $Q \in SN$ . The various cases are:
  - $V = \omega_i W$  and  $Q = P_i[x_i := W[\tau]]$ . By the *IH*,  $W[\tau] \in SN$  since  $cxy(W) < cxy(T)$  and thus, since  $lgt(W) < lgt(N)$ ,  $Q \in SN$  follows from the main *IH* (recall we are “inside” the proof of theorem 4.1,  $type(W) = A$  or  $type(W) = B$  and  $type(N) = A \vee B$ ).
  - $V = \mu b W$  and  $Q = \mu b W[\tau][b :=^* [P]] = \mu b W[\tau']$  where  $\tau' = \tau \cup [b :=^* [P]]$ . Since  $cxy(W) < cxy(T)$ , the result follows from the *IH*.
  - $V = (W \varepsilon)$  and  $\varepsilon$  is not in the form  $[x_1.W_1, x_2.W_2]$ . Then, the head redex of  $(V[\tau] [P])$  must come from  $V$  and  $Q = (V'[\tau] [P])$  for some  $V'$  such that  $V \triangleright V'$ . Let  $T' = (a V')$ . Since  $\eta(T') < \eta(T)$ ,  $T'[\tau] \in SN$ . But  $T'[\tau] \triangleright Q$  and thus  $Q \in SN$ .
  - $V = (W [x_1.W_1, x_2.W_2])$  and  $Q = (W[\tau] [x_1.(W_1[\tau] [P]), x_2.(W_2[\tau] [P])])$ . Let  $T_j = (a W_j)$ . Since  $cxy(T_j) < cxy(T)$ , by the *IH*,  $T_j[\tau] \in SN$

and thus  $(W_j[\tau] [P]) \in SN$  since  $T_j[\tau] \triangleright (W_j[\tau] [P])$ . By the *IH*, since  $cxy(W) < cxy(T)$ ,  $W[\tau] \in SN$ . By theorem 3.1, it is thus enough to show that  $Q' = hred(Q) \in SN$ .

If  $hd(Q)$  comes from  $W$ , the result follows from the *IH*. Otherwise, the various cases are:

- \*  $W = \omega_i W'$  and  $Q' = (W_i[\tau] [P])[x_i := W'[\tau]]$ . Let  $T' = (a W_i[x_i := W'])$ . Then  $T = (a (\omega_i W' [x_1.W_1, x_2.W_2])) \triangleright T'$ . By the *IH*,  $T'[\tau] \in SN$  and the result follows from the fact that  $T'[\tau] \rightarrow (W_i [P])[x_i := W'[\tau]] = Q'$ .
- \* If  $W = \mu b W'$  or  $W = (W' [x_1.W'_1, x_2.W'_2])$ : the proof is similar.

4.  $M = (x \varepsilon \vec{T})$ ,  $\varepsilon \neq [x_1.M_1, x_2.M_2]$  and  $\sigma(x) = \mu a N$ . We prove exactly as in case 3 that  $(\mu a N \varepsilon[\sigma]) \in SN$ . To prove that  $M[\sigma] \in SN$ , it is enough to use the same trick as in case 1:  $M[\sigma] = (z \vec{T}[\sigma])[z := (\mu a N \varepsilon[\sigma])]$  where  $z$  is a fresh variable and the *IH* gives the result since  $lgt(z) < lgt(x)$ .
5.  $M = (x [x_1.M_1, x_2.M_2])$  and  $\sigma(x) = (N_3 [y_1.N_1, y_2.N_2])$ . By theorem 3.1, it is enough to show  $P = (N_3 [y_1.(N_1 [P]), y_2.(N_2 [P])]) \in SN$  where, for  $i = 1, 2$ ,  $P_i = M_i[\sigma]$  and  $(N_i [P])$  is a notation for  $(N_i [x_1.P_1, x_2.P_2])$ . Let  $M' = (z [x_1.M_1, x_2.M_2])$  where  $z$  be a fresh variable. For  $i = 1, 2$ , let  $\sigma_i = \sigma \cup [z := N_i]$ . By the *IH*,  $M'[\sigma_i] \in SN$  since  $\eta(\sigma') \leq \eta(\sigma)$  and  $cxy(\sigma') < cxy(\sigma)$ . Then  $(N_i [P]) \in SN$  since  $M'[\sigma_i] \triangleright (N_i [P])$ . By theorem 3.1, it is thus enough to show that  $Q = hred(P) \in SN$ .

If  $hd(P)$  comes from  $W$ , the result follows from the *IH*. Otherwise, the various cases are:

- $N_3 = \omega_i N'_3$  and  $Q = (N_i[x_i := N'_3] [P])$ . Let  $M' = (z [x_1.M_1, x_2.M_2])$  where  $z$  is a fresh variable and  $\sigma' = \sigma \cup \{[z := N_i[x_i := N'_3]]\}$ . Then  $Q = M'[\sigma'] \in SN$  since  $\eta(\sigma') \leq \eta(\sigma)$  and  $cxy(\sigma') < cxy(\sigma)$ .
  - $N_3 = \mu a N'_3$  or  $N_3 = (Q_3 [y_1.Q_1, y_2.Q_2])$ . The proof is similar.
6. If  $M = (x \varepsilon \vec{T})$ ,  $\varepsilon \neq [x_1.M_1, x_2.M_2]$  and  $\sigma(x) = (N_3 [x_1.N_1, x_2.N_2])$ . We prove exactly as in case 5 that  $(N_3 [x_1.N_1, x_2.N_2] \varepsilon[\sigma]) \in SN$ . To prove that  $M[\sigma] \in SN$ , it is enough to use the same trick as in case 1:  $M[\sigma] = (z \vec{T}[\sigma])[z := (N_3 [x_1.N_1, x_2.N_2] \varepsilon[\sigma])]$  where  $z$  is a fresh variable and the *IH* gives the result since  $lgt(z) < lgt(x)$ . □

## 5 Proof of theorem 3.2

The idea of the proof is the following: we show that an infinite reduction of  $S_1$  can be translated into an infinite reduction of  $S_2$ . These reductions are the same except that, in  $S_1$ ,  $\varepsilon$  can be far away from the  $N_i$ . We mark  $\varepsilon$  and the  $N_i$  to keep their trace. This gives the set of marked terms  $\mathcal{T}'$  of definition 5.1. The correct terms of definition 5.3 intuitively are the marked terms for which each marked  $N_i$  *knows* who is the corresponding marked  $\varepsilon$ . Concretely, being correct is a sufficient condition to ensure that a reduction in the marked  $S_1$  can be translated to the corresponding  $S_2$ .

The main difficulty of the proof consists in writing precise definitions. The proofs of the lemmas consist in easy but tedious verifications.

*Important remark.* The proof is uniform in the sequence  $\varepsilon \vec{V}$ . In definition 5.1 below, we implicitly assume the following: if we are proving theorem 3.2 for  $\varepsilon \in \mathcal{T}$

(resp.  $\varepsilon = \pi_i$ ,  $\varepsilon = [y_1. M_1, y_2. M_2]$ ) then, in the sub-terms of the form  $\circ_{\varepsilon'}$ , we necessarily have  $\varepsilon' \in \mathcal{T}$  (resp.  $\varepsilon' = \pi_i$ ,  $\varepsilon' = [y_1. Q_1, y_2. Q_2]$ ). Note that we could also assume that  $\varepsilon'$  is a reduct of  $\varepsilon$  but this does not really matter for the proof. However, in the case  $\varepsilon = [y_1. P_1, y_2. P_2]$ , since the sequence  $\varepsilon \vec{V}$  is nice  $\vec{V}$  is empty and this must appear in the proof. We will do the proof only for  $\varepsilon \in \mathcal{T}$  or  $\varepsilon = \pi_i$ . The proof for the case  $\varepsilon = [y_1. P_1, y_2. P_2]$  is essentially the same: we just have to add a third condition in definition 5.3 and check in the lemmas that this condition is preserved. This new condition is given in the final remark of this section.

**Definition 5.1** 1. Let  $\mathcal{T}'$  be the set of terms obtained from  $\mathcal{T}$  by adding new constructors:  $\diamond_N$  and  $\circ_\varepsilon$  where  $N \in \mathcal{T}$  and  $\varepsilon \in \mathcal{E}$  are closed.

2. The reduction rules for  $\mathcal{T}'$  are the ones of  $\mathcal{T}$  plus the following:

- If  $N \triangleright N'$  then  $\diamond_N \triangleright \diamond_{N'}$  and  $\circ_N \triangleright \circ_{N'}$ .
- $(\diamond_N \circ_\varepsilon) \triangleright (N \varepsilon)$ .

3. Let  $\blacktriangleright$  be the congruence defined by the following reduction rules:

- $(M [x_1.N_1, x_2.N_2] \circ_\varepsilon) \blacktriangleright (M [x_1.(N_1 \circ_\varepsilon), x_2.(N_2 \circ_\varepsilon)])$
- $(\mu a M \circ_\varepsilon) \blacktriangleright \mu a M[a :=^* \circ_\varepsilon]$

*Comments*

An element of  $\mathcal{T}'$  is a term in  $\mathcal{T}$  where some sub-terms have been replaced by terms as  $\diamond_N$  or  $\circ_\varepsilon$  where  $N \in \mathcal{T}$  and  $\varepsilon \in \mathcal{E}$  and, in particular, have no sub-terms as  $\diamond_{N'}$  or  $\circ_{\varepsilon'}$ . It is assumed, in the definition, that the  $N$  and  $\varepsilon$  occurring in  $\diamond_N$  or  $\circ_\varepsilon$  are closed. In fact, they are allowed to have free variables (both intuitionistic and classical) but it is assumed that these variables will never be captured and thus act as constants.

**Definition 5.2** Let  $M \in \mathcal{T}'$ .

1.  $M$  is acceptable iff  $M = \diamond_N$  or  $M = \mu a M_1$  and, for each sub-term of  $M$  of the form  $(a N)$ ,  $N$  is acceptable or  $M = (N [x_1.N_1, x_2.N_2])$  and  $N_1, N_2$  are acceptable.
2. If  $M$  is acceptable, the set  $st(M)$  of terms is defined by:  $st(\diamond_N) = \{\diamond_N\}$ ,  $st(\mu a M_1) = \cup \{st(S) / (a S) \text{ sub-term of } M_1\}$  and  $st((N [x_1.N_1, x_2.N_2])) = st(N_1) \cup st(N_2)$ .

**Lemma 5.1** Let  $M \in \mathcal{T}'$  be an acceptable term.

1. If  $\sigma$  is a substitution either of the form  $[x := N]$  or  $[a :=^* N]$ , then  $M[\sigma]$  is acceptable and  $st(M[\sigma]) = st(M)$ .
2. If  $M \triangleright M'$ , then  $M'$  is acceptable and  $st(M') \subseteq st(M)$ .

**Proof** By induction on  $M$ . (1) trivial. For (2) use (1). □

**Definition 5.3** A term  $M \in \mathcal{T}'$  is correct if the following conditions hold.

1. Each occurrence of a term of the form  $\circ_\varepsilon$  appears as  $(U \circ_\varepsilon)$  for some acceptable term  $U$ .
2. For each sub-term of  $M$  of the form  $\diamond_N$  there is a sub-term (necessarily unique) of the form  $(U \circ_\varepsilon)$  such that  $\diamond_N$  belongs to  $st(U)$ . The corresponding  $\varepsilon$  is denoted as  $eps(N)$

## Examples

- Assume  $M, N, O, P, \varepsilon$  are closed terms. Then  $A = (M [x_1.\diamond_N, x_2.\diamond_O] \circ_\varepsilon P)$  is correct.
- Assume  $M, N, O, P, Q, R, S, \varepsilon_1, \varepsilon_2$  are closed terms. Then  $B = (M [x_1.(N [y_1.\diamond_O, y_2.\mu aP] \circ_{\varepsilon_1}), x_2.(\mu b(b \mu c(c (Q [z_1.\mu dR, z_2.\diamond_S]))) \circ_{\varepsilon_2})])$  is correct.

**Lemma 5.2** *If  $M$  is correct and  $M \triangleright M'$ , then  $M'$  is correct.*

**Proof** Let  $(U \circ_\varepsilon)$  be a sub-term of  $M$ . A reduction can be, either in  $\varepsilon$  or in  $U$  or between  $U$  and  $\varepsilon$  or, finally, above  $(U \circ_\varepsilon)$ . Since  $U$  is acceptable and by using lemma 5.1 it is easy to check that, in each case the conditions of correctness are still satisfied.  $\square$

**Lemma 5.3** *Let  $M$  be a correct term.*

1.  $M$  has no sub-term of the form  $(O \diamond_N)$ .
2. If  $(\diamond_N O)$  is a sub-term of  $M$ , then  $O = \circ_\varepsilon$  for some  $\varepsilon$ .

**Proof** Otherwise, let  $(U \circ_\varepsilon)$  be the sub-term such that  $\diamond_N \in st(U)$ . The result follows easily from the fact that  $U$  is acceptable.  $\square$

**Definition 5.4** *Let  $M \in \mathcal{T}'$ .*

1.  $T_1(M)$  is the term obtained by replacing  $\diamond_N$  by  $N$  and  $\circ_\varepsilon$  by  $\varepsilon$ .
2. If  $M$  is a sub-term of a correct term,  $T_2(M)$  is the term obtained by replacing each occurrence of  $(U \circ_\varepsilon)$  by  $U'$  where  $U'$  is obtained from  $U$  by replacing each occurrence of  $\diamond_N$  such that  $\varepsilon = eps(N)$  is a sub-term of  $M$  by  $(N \varepsilon)$ .

## Comments and examples

1. If  $M$  itself is correct,  $T_2(M) \in \mathcal{T}$ . Otherwise, some  $\diamond_N$  that are related to a  $\circ_\varepsilon$  outside  $M$  are not replaced. We need this more general definition for the proof of lemma 5.4.
2. If  $M$  is correct,  $T_1(M) \triangleright^* T_2(M)$ . More precisely  $T_2(M) = T_1(M')$  where  $M'$  is the normal form of  $M$  for the rules  $\blacktriangleright$ . Since we will not use this result, we do not prove it.

3. Let  $A, B$  be the terms of the previous example. Then

$$T_1(A) = (M [x_1.N, x_2.O] \varepsilon P) \text{ and } T_2(A) = (M [x_1.(N \varepsilon), x_2.(O \varepsilon)] P).$$

$$T_1(B) = (M [x_1.(N [y_1.O, y_2.\mu aP] \varepsilon_1), x_2.(\mu b(b \mu c(c (Q [z_1.\mu dR, z_2.S]))) \varepsilon_2)])$$

$$\text{and } T_2(B) = (M [x_1.(N [y_1.(O \varepsilon_1), y_2.\mu aP]), x_2.\mu b(b \mu c(c (Q [z_1.\mu dR, z_2.(S \varepsilon_2)])))]).$$

**Lemma 5.4** *Let  $M \in \mathcal{T}'$  be correct. If  $T_1(M) \triangleright N$ , there is a correct term  $M'$  such that  $M \triangleright^+ M'$  and  $T_1(M') = N$ .*

**Proof** Let  $R$  be the redex that has been reduced. By lemma 5.3, the only cases to consider are:

- There is a redex  $S$  in  $M$  such that  $R = T_1(S)$ . The result follows then from lemma 5.3.
- There is a sub-term of  $M$  of the form  $\diamond_N$  or  $\circ_\varepsilon$ , such that  $R$  is a sub-term of  $N$  or  $\varepsilon$ . The result is then trivial.
- Finally,  $R = (T_1(U) \varepsilon)$  where  $(U \circ_\varepsilon)$  is a sub-term of  $M$ , the result follows from the fact that  $U$  is acceptable.  $\square$

**Lemma 5.5** *Let  $P = (M O)$  be a sub-term of a correct term. Assume  $O \neq \circ_\varepsilon$  and  $P \triangleright P'$  by reducing the redex  $P$ . Then,  $T_2(P) \triangleright^* T_2(P')$ .*

**Proof** It is, for example, straightforward to check that, if  $M = \lambda x M_1$ , then  $T_2(P) = (\lambda x T_2(M_1) T_2(O))$  and  $T_2(P') = T_2(M_1)[x := T_2(O)]$ . The other cases are similar.  $\square$

**Lemma 5.6** *Assume  $M = (N \circ_\varepsilon)$  is a sub-term of a correct term. Then,*

- $T_2(M) = T_2((T_2(N) \circ_\varepsilon))$ .
- If  $N$  has no sub-terms of the form  $\circ_{\varepsilon'}$  and  $N \triangleright N'$  then,  $T_2(M) \triangleright^* T_2((N' \circ_\varepsilon))$ .

**Proof** Straightforward.  $\square$

**Lemma 5.7** *Let  $M$  is a sub-term of a correct term. If  $M \triangleright M'$ , then  $T_2(M) \triangleright^* T_2(M')$ . Moreover, if  $T_2(M) = T_2(M')$ , then  $M \blacktriangleright M'$ .*

**Proof** By induction on  $(nb(M), cxy(M))$  where  $nb(M)$  is the number of sub-terms of the form  $\circ_\varepsilon$  in  $M$ . The only non trivial case is  $M = (N O)$ .

- Assume  $O \neq \circ_\varepsilon$ . If  $M' = (N' O)$  where  $N \triangleright N'$  or  $M' = (N O')$  where  $O \triangleright O'$ , the result is trivial. Otherwise,  $M$  itself is the reduced redex and the result comes from lemma 5.5.
- Assume  $O = \circ_\varepsilon$ . If  $M$  itself is the reduced redex then  $T_2(M) = T_2(M')$  and  $M \blacktriangleright M'$ . If  $M' = (N \circ_{\varepsilon'})$  where  $\varepsilon \triangleright \varepsilon'$ , the result is trivial. Otherwise, i.e.  $M' = (N' \circ_\varepsilon)$  where  $N \triangleright N'$ . If  $nb(N) = 0$ , the result follows from lemma 5.6(2). Otherwise, by the induction hypothesis,  $T_2(N) \triangleright^* T_2(N')$  and the result comes from lemma 5.6.  $\square$

**Lemma 5.8** *Let  $M \in \mathcal{T}'$  be a correct term. Then  $M$  is strongly normalizable for the  $\blacktriangleright$  reduction.*

**Proof** If  $M$  is correct, let  $lg(M)$  be the sum of the length of the path (i.e. the number of nodes in the tree representing  $M$ ) relating the  $\diamond_N$  to the corresponding  $\circ_\varepsilon$ . It is easy to see that, if  $M \blacktriangleright M'$ , then  $lg(M') < lg(M)$ .  $\square$

### Proof of theorem 3.2

Assume  $S_2 \in SN$  and  $S_1 \notin SN$ . Let  $(U_i)$  be a sequence of terms such that  $U_0 = S_1$  and, for each  $i$ ,  $U_i \triangleright U_{i+1}$ . Let  $M = (N [x_1. \diamond_{N_1}, x_2. \diamond_{N_2}] \circ_\varepsilon \vec{V})$ . By using lemma 5.4, we get a sequence of correct terms  $M_i$  such that, for each  $i$ ,  $M_i \triangleright^+ M_{i+1}$  and  $T_1(M_i) = U_i$ . By lemma 5.7,  $T_2(M_i) \triangleright^* T_2(M_{i+1})$ . Since  $S_2 = T_2(M) \in SN$ , there is an  $i_0$  such that, for  $i \geq i_0$ ,  $T_2(M_i) = T_2(M_{i+1})$  and thus, by lemma 5.7,  $M_i \triangleright^+ M_{i+1}$ . This contradicts lemma 5.8.  $\square$

### Remark.

Assume that, in theorem 3.2,  $\varepsilon = [y_1.Q_1, y_2.Q_2]$ . If  $\vec{V}$  were not empty, the proof of lemma 5.4 would not work because a redex could be created by the transformation  $T_1$ . Here is an example: let  $M = (P \circ_\varepsilon V)$  be correct and assume  $T_1(M) = (P [y_1.Q_1, y_2.Q_2] V) \triangleright N = (P [y_1.(Q_1 V), y_2.(Q_2 V)])$ . There is no way to find  $M'$  such that  $M \triangleright M'$  and  $T_1(M') = N$  because  $(P \circ_\varepsilon V)$  is not a redex.

We do not know wether theorem 3.2 remains true if the sequence  $\varepsilon \vec{V}$  is not nice: to prove it,  $(P \circ_\varepsilon V)$  should then be considered as a redex but  $\mathcal{T}'$  becomes much more complicated. Since  $\varepsilon \vec{V}$  is nice, it is simpler to add a new condition in the definition 5.3 of correctness to ensure that this situation (of the creation of a redex by the transformation  $T_1$ ) does not appear. This condition is the following:

3.  $M$  is good wrt the set of all its sub-terms of the form  $(U \circ_\varepsilon)$ .

where, if  $E$  is a subset of  $\mathcal{T}'$ ,  $M$  is good wrt to  $E$  is defined by:  $M \in E$  or  $M = \mu a N$  and for each occurrence of  $(a N)$  in  $M$ ,  $N$  is good wrt to  $E$  or  $M = (N [x_1.N_1, x_2.N_2])$  and  $N_1, N_2$  are good wrt to  $E$ .

This condition implies in particular that, in a correct term, there is no sub-term of the form  $(N \circ_\varepsilon U)$  and thus that lemma 5.4 remains valid. It is not difficult to check that the other lemmas remain also valid.

## References

- [1] Y. Andou. *Church-Russer property of simple reduction for full first-order classical natural deduction*. Annals of Pure and Applied logic 119 (2003) 225-237.
- [2] F. Barbanera and S. Berardi *A symmetric lambda-calculus for classical program extraction*. In M. Hagiya and J.C. Mitchell, editors, Proceedings of theoretical aspects of computer software, TACS'94. LNCS (789), pp. 495-515. Springer Verlag, 1994.
- [3] E. T. Bittar. *Strong Normalisation Proofs for Cut-Elimination in Gentzen's Sequent Calculi*. Logic, Algebra and Computer Science, 46, Banach Center Publications, pp. 179-225, 1999.
- [4] R. Constable and C. Murthy *Finding computational content in classical proofs*. In G. Huet and G. Plotkin, editors, Logical Frameworks, pp. 341-362, Cambridge University Press, 1991.
- [5] J.-Y. Girard *A new constructive logic: classical logic*. MSCS (1), pp. 255-296, 1991.
- [6] P. de Groote. *On the strong normalization of natural deduction with permutation-conversions*. In 10th International Conference on Typed Lambda Calculi and Applications, TLCA'95. LNCS (902) pp. 201-215. Springer Verlag, 1995.
- [7] P. de Groote. *Strong normalization of classical natural deduction with disjunction*. In 5th International Conference on typed lambda calculi and applications, TLCA'01. LNCS (2044), pp. 182-196. Springer Verlag, 2001.
- [8] R. David. *Normalization without reducibility*. Annals of Pure and Applied Logic (107), p. 121-130, 2001.
- [9] R. David and B. Guillaume. *Strong normalization of the typed  $\lambda_{ws}$ -calculus*. Accepted in the CSL'03 colloquium.
- [10] R. David and K. Nour. *A short proof of the strong normalization of the simply typed  $\lambda\mu$ -calculus*. To appear in Schedae Informaticae vol.12, 2003.
- [11] F. Joachimski and R. Matthes. *Short proofs of normalization for the simply-typed lambda-calculus, permutative conversions and Gödel's T*. Archive for Mathematical Logic 42, pp 59-87 (2003).
- [12] J.-L. Krivine. *Classical logic, storage operators and 2nd order lambda-calculus*. Annals of Pure and Applied Logic (68), pp. 53-78, 1994.
- [13] J.J. Levy. *Rductions correctes et optimales dans le lambda-calcul*. PhD thesis Paris 7, 1978.

- [14] C.R. Murthy. *An evaluation semantics for classical proofs*. In Proceedings of the sixth annual IEEE symposium on logic in computer science, pp. 96-107, 1991.
- [15] C.R. Murthy. *A computational analysis of Girard's translation and LC*. In Proceedings of the seventh annual IEEE symposium on logic in computer science, pp. 90-101, 1992.
- [16] K. Nour and K. Saber. *A semantical proof of the strong normalization theorem of full propositionnal classical natural deduction*. Manuscript 2003.
- [17] M. Parigot  *$\lambda\mu$ -calculus: An algorithm interpretation of classical natural deduction*. Lecture Notes in Artificial Intelligence (624), pp. 190-201. Springer Verlag 1992.
- [18] M. Parigot. *Proofs of strong normalization for second order classical natural deduction*. Journal of Symbolic Logic, 62 (4), pp. 1461-1479, 1997.
- [19] D. Prawitz. *Natural deduction, a proof-theoretical study*. Almqvist & Wiksell, Stockholm, 1965.
- [20] D. Prawitz. *Ideas and results in proof-theory*. In J.E. Fenstad, editor, Proceedings of the second scandinavian logic symposium, pp. 237-309. North-Holland, 1971.
- [21] D. Pym and E. Ritter. *On the semantics of classical disjunction*. Journal of Pure and Applied Algebra, To appear.
- [22] N.J. Rehof and M.H. Sorensen. *The  $\lambda_{\Delta}$ -calculus*. In M. Hagiya and J.C. Mitchell, editors, Proceedings of the international symposium on theoretical aspects of computer software, TACS'94, LNCS (789), pp. 516-542. Springer Verlag, 1994.
- [23] E. Ritter, D. Pym and L. Wallen. *On the intuitionistic force of classical search*. Theoretical Computer Science (232), pp. 299-333, 2000.
- [24] E. Ritter, D. Pym and L. Wallen. *Proof-terms for classical and intuitionistic resolution*. Journal of logic and computation, 10 (2), pp. 173-207, 2000.
- [25] J. Seldin. *On the proof theory of the intermediate logic MH*. Journal of symbolic logic, 51 (3), pp. 626-647, 1986.
- [26] G. Stalmarck. *Normalization theorems for full first-order classical natural deduction*. Journal of symbolic logic, 56 (1), pp. 129-149, 1991.
- [27] D. van Daalen *The language theory of Automath*. PhD Thesis. Eindhoven 1977.