



HAL
open science

Material symmetry group of the non-linear polar-elastic continuum

Victor A. Eremeyev, Wojciech Pietraszkiewicz

► **To cite this version:**

Victor A. Eremeyev, Wojciech Pietraszkiewicz. Material symmetry group of the non-linear polar-elastic continuum. *International Journal of Solids and Structures*, 2012, 49 (14), pp.1993-2005. hal-00835636

HAL Id: hal-00835636

<https://hal.science/hal-00835636>

Submitted on 19 Jun 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Material symmetry group of the non-linear polar-elastic continuum

Victor A. Eremeyev^{a,b}, Wojciech Pietraszkiewicz^{c,*}

^a Otto-von-Guericke-University Magdeburg, Universitätplatz 2, 39106 Magdeburg, Germany

^b South Scientific Center, RASci & South Federal University, Rostov on Don, Russia

^c Institute of Fluid-Flow Machinery, PASci, ul. Gen. J. Fiszerza 14, 80-952 Gdańsk, Poland

A B S T R A C T

We extend the material symmetry group of the non linear polar elastic continuum by taking into account microstructure curvature tensors as well as different transformation properties of polar and axial tensors. The group consists of an ordered triple of tensors which makes the strain energy density of polar elastic continuum invariant under change of reference placement. An analog of the Noll rule is established. Four simple specific cases of the group with corresponding reduced forms of the strain energy density are discussed. Definitions of polar elastic fluids, solids, liquid crystals and subfluids are given in terms of members of the symmetry group. Within polar elastic solids we discuss in more detail isotropic, hemitropic, cubic symmetric, transversely isotropic, and orthotropic materials and give explicitly corresponding reduced representations of the strain energy density. For physically linear polar elastic solids, when the density becomes a quadratic function of strain measures, reduced representations of the density are established for monoclinic, orthotropic, cubic symmetric, hemitropic and isotropic materials in terms of appropriate joint scalar invariants of stretch, wryness and undeformed structure curvature tensors.

1. Introduction

Material symmetry group of the non linear micropolar continuum was first characterized by Eringen and Kafadar (1976). They discussed all density preserving deformations and all microrotations of reference placement of the continuum that cannot be experimentally detected. Simple micropolar solids and simple micropolar fluids in the spatial description were defined in terms of members of the group. After 35 years we are still not aware of any detailed discussion of the material symmetry group of non linear micropolar continuum and of appropriate reduction of constitutive equations in terms of members of the group.

In the present report we restrict ourselves to the non linear polar elastic continuum, which material behavior is described by the strain energy density W per unit reference volume. As compared with Eringen and Kafadar (1976) we introduce three modifications:

1. At each material point our density W , satisfying the principle of material frame indifference, depends explicitly not only on natural Lagrangian stretch \mathbf{E} and wryness $\mathbf{\Gamma}$ tensors, but additionally upon the reference microstructure curvature tensor \mathbf{B} as a

parametric tensor. Eringen and Kafadar (1976) used similar strain measures in W , but introduced referential mass density ρ_x and microinertia tensor \mathbf{J}_x as parametric quantities.

2. In discussing invariance properties of W we take into account that, while \mathbf{E} is a polar tensor, $\mathbf{\Gamma}$ and \mathbf{B} are axial tensors which change signs under inversion transformation (mirror reflection) of 3D space. Eringen and Kafadar (1976) did not take into account that their $\mathbf{\Gamma}$ was the axial tensor.
3. Our material symmetry group \mathcal{G}_x consists of ordered triple of tensors: unimodular \mathbf{P} , orthogonal \mathbf{R} , and second order \mathbf{L} . These tensors appear from transformation of \mathbf{E} , $\mathbf{\Gamma}$ and \mathbf{B} under an arbitrary change of reference placement of the micropolar body. The transformation properties of \mathbf{B} are quite different from those of \mathbf{J}_x .

As a result of these modifications, our material symmetry group \mathcal{G}_x does not coincide with the group introduced by Eringen and Kafadar (1976).

The tensor \mathbf{B} appears naturally during description of the reference placement, see Pietraszkiewicz and Eremeyev (2009). The case $\mathbf{B} \neq \mathbf{0}$ indicates non uniform distribution of directors in the reference placement. From the mathematical point of view the case $\mathbf{B} \neq \mathbf{0}$ relates to non Euclidean geometry of polar material because the directors can be considered as a nonholonomic basis. From the physical point of view necessity of taking into account \mathbf{B} in constitutive equations corresponds to proper description of microstructure of materials. \mathbf{B} can be considered as an analog of

* Corresponding author. Tel.: +48 58 69 95 263; fax: +48 58 341 61 44.

E-mail addresses: eremeyev.victor@gmail.com (V.A. Eremeyev), pietrasz@imp.gda.pl, pietrasz@karol.imp.gda.pl (W. Pietraszkiewicz).

the curvature tensor in the theory of shells or of material parameters describing helical substructures of rods, see for example Eremeyev and Pietraszkiewicz (2006), Lauderdale and O'Reilly (2007).

Within Cosserat type theories of shells and rods various definitions of material symmetry groups were proposed for example by Murdoch and Cohen (1979), Eremeyev and Pietraszkiewicz (2006), Luo and O'Reilly (2000), Lauderdale and O'Reilly (2007). The material symmetry group of the second grade elastic materials was considered for example by Huang and Smith (1967), Elżanowski and Epstein (1992). Huang and Smith (1967) took into account difference between axial and polar tensors in construction of the material symmetry group. Let us note that the invariance of the strain energy density under transformations of the reference placement is similar but not the same as the uniformity and homogeneity properties considered by Epstein and de Leon (1996, 1998).

The paper is organized as follows. In Section 2 we recall after Pietraszkiewicz and Eremeyev (2009) some basic relations of non linear micropolar continuum. In particular, we remind definitions of stretch \mathbf{E} and wryness $\mathbf{\Gamma}$ tensors, of undeformed structure curvature tensor \mathbf{B} , of strain energy density W satisfying the principle of material frame indifference, of referential \mathbf{S} , \mathbf{K} and Eulerian \mathbf{T} , \mathbf{M} stress and couple stress tensors, as well as appropriate forms of equilibrium conditions.

Transformation properties of various fields under change of reference placement are discussed in Section 3. Then in Section 4 invariance requirements for W under change of reference placement are analyzed. This allows one to derive the material symmetry group \mathcal{G}_x of the polar elastic continuum. The group \mathcal{G}_x consists of an ordered triple of tensors which make W invariant under change of reference placement. For \mathcal{G}_x we establish an analog of Noll's rule, see Noll (1958).

Four specific cases of \mathcal{G}_x with corresponding considerably simplified forms of W are discussed in Section 5. After introducing in Section 6 property of isotropy of the polar elastic material, polar elastic fluids are defined by requiring \mathcal{G}_x to be the maximal group. It is found that polar elastic fluids are always isotropic and their strain energy density W is the hemitropic scalar valued even function of the structure curvature tensor \mathbf{C} of deformed placement. By the representation theorem of Smith and Smith (1971) the density ρ is reduced to the scalar valued function of the deformed density ρ and of six scalar invariants of \mathbf{C} . The constitutive equations for \mathbf{T} and \mathbf{M} are derived, which generalize the ones proposed earlier for polar elastic fluids for example by Aero et al. (1965), Eringen (1966), Eringen (1997), Allen et al. (1967).

In Section 8 we define the group \mathcal{G}_x of polar elastic solids using an additional hypothesis that elements of \mathcal{G}_x are described by orthogonal tensors which are the same for the position vector and the directors. Polar elastic liquid crystals and subfluids are then defined in Section 9 by \mathcal{G}_x which contains some elements not belonging to the orthogonal group. Examples of Kelvin's medium defined by Grekova and Zhilin (2001) and of Ericksen's liquid crystal proposed by Eringen (1997, 2001) are briefly discussed.

Simplified representations of W for some forms of anisotropy of polar elastic solids are discussed in Section 10.

Applying representation theorems of the scalar valued tensor functions derived by Spencer (1965, 1971), we analyze possible reduction of W for isotropic, hemitropic, cubic symmetric, orthotropic, and transversely isotropic non linear polar elastic solids. As in Section 6 we apply the theory of representation of scalar valued functions of several tensorial arguments summarized in Spencer (1971), Boehler (1987), Smith (1994), Zheng (1994) where other references can be found. For polar elastic solids the material symmetry group is described by orthogonal transformations alone. This gives possibility to represent W in terms of joint invariants of the strain measures \mathbf{E} and $\mathbf{\Gamma}$ as well as of the structure curvature tensor \mathbf{B} .

Finally, in Sections 11 and 12 physically linear polar elastic solids are analyzed under mirror reflection as well as under monoclinic, orthotropic, cubic, hemitropic, and isotropic symmetry conditions. For each case of symmetries reduced forms of W in terms of appropriate joint scalar invariants of tensors \mathbf{E} , $\mathbf{\Gamma}$, and \mathbf{B} are given. In Appendix A we present table with invariants for the isotropic symmetry.

2. Some relations of the non-linear micropolar elastic continuum

Let the body \mathcal{B} consisting of material particles X, Y, \dots deform in three dimensional (3D) Euclidean physical space \mathcal{E} which translation vector space is E . According to Cosserat and Cosserat (1909), Truesdell and Toupin (1960), Eringen and Kafadar (1976) for example, each material particle $X \in \mathcal{B}$ of the polar elastic continuum has six degrees of freedom of rigid body.

In the reference (undeformed) placement $\chi(\mathcal{B}) : B_x \subset \mathcal{E}$ position $x \in \mathcal{E}$ of $X \in \mathcal{B}$ is given by the vector $\mathbf{x} \in E$ relative to origin $o \in \mathcal{E}$ of an inertial frame (o, \mathbf{i}_a) , where $\mathbf{i}_a \in E$, $a = 1, 2, 3$, is a right handed triple of orthonormal vectors. Orientation of $X \in \mathcal{B}$ in E is fixed by the right handed triple of orthonormal directors $\mathbf{h}_a \in E$, so that $\mathbf{h}_a = \mathbf{H}\mathbf{i}_a$, where $\mathbf{H} = \mathbf{h}_a \otimes \mathbf{i}_a \in Orth^+$ is the proper orthogonal structure tensor of B_x , $\mathbf{H}^{-1} = \mathbf{H}^T$, $\det \mathbf{H} = +1$, and \otimes denotes the tensor product.

In the actual (deformed) placement $\gamma(\mathcal{B}) : B_y \subset \mathcal{E}$, $\chi(B_x) \in \mathcal{E}$, $\gamma \circ \chi^{-1}$, position $y \in B_y$ of $X \in \mathcal{B}$ becomes defined by the vector $\mathbf{y} \in E$ taken here relative to the same origin $o \in \mathcal{E}$. Orientation of X becomes fixed by the right handed triple of orthonormal directors $\mathbf{d}_a \in E$ or by the structure tensor $\mathbf{D} = \mathbf{d}_a \otimes \mathbf{i}_a \in Orth^+$ of B_y . As a result, the finite displacement of polar elastic continuum can be described by two smooth mappings (see Fig. 1):

$$\mathbf{y} = \chi(\mathbf{x}) = \mathbf{x} + \mathbf{u}(\mathbf{x}), \quad \mathbf{d}_a = \mathbf{Q}(\mathbf{x})\mathbf{h}_a, \quad (1)$$

where $\mathbf{u} \in E$ is the translation vector and $\mathbf{Q} = \mathbf{D}\mathbf{H}^T = \mathbf{d}_a \otimes \mathbf{h}_a \in Orth^+$ is the proper orthogonal microrotation tensor, $\mathbf{Q}^{-1} = \mathbf{Q}^T$, $\det \mathbf{Q} = +1$. Two independent fields $\mathbf{u}(\mathbf{x})$ and $\mathbf{Q}(\mathbf{x})$ describe translational and rotational degrees of freedom of polar elastic continuum.

Natural Lagrangian relative stretch and wryness (or change of microstructure orientation) tensors \mathbf{E} and $\mathbf{\Gamma}$ are defined according to Pietraszkiewicz and Eremeyev (2009) as

$$\mathbf{E} = \mathbf{Q}^T \mathbf{F} - \mathbf{I}, \quad \mathbf{\Gamma} = \frac{1}{2} \mathbf{E} : (\mathbf{Q}^T \text{Grad } \mathbf{Q}). \quad (2)$$

Here $\mathbf{F} = \text{Grad } \mathbf{y}$, $\det \mathbf{F} > 0$, is the classical deformation gradient tensor taken relative to B_x , \mathbf{I} is the identity (metric) tensor of 3D space




Fig. 1. Micropolar body deformation.

E , $\mathbf{E} = \mathbf{I} \times \mathbf{I}$ is the 3rd order skew permutation tensor with \times the vector product, while the double dot product : of two 3rd order tensors \mathbf{A} , \mathbf{B} represented in \mathbf{h}_a is defined as $\mathbf{A}:\mathbf{B} = A_{amnb}B_{mnb}\mathbf{h}_a \otimes \mathbf{h}_b$. The strain measures defined in (2) are not symmetric, in general, $\mathbf{E}^T \neq \mathbf{E}$, $\mathbf{\Gamma}^T \neq \mathbf{\Gamma}$.

Gradient of vector field $\mathbf{v}(\mathbf{x}) \in E$ is a 2nd order tensor field $\text{Grad} \mathbf{v} \in E \otimes E$ and gradient of 2nd order tensor field $\mathbf{A} \in E \otimes E$ is a 3rd order tensor field $\text{Grad} \mathbf{A} \in E \otimes E \otimes E$ defined by (see for example Ogden (1984))

$$\begin{aligned} [\text{Grad} \mathbf{v}(\mathbf{x})] \mathbf{a} &= \frac{d}{dt} \mathbf{v}(\mathbf{x} + t\mathbf{a})|_{t=0}, \\ [\text{Grad} \mathbf{A}(\mathbf{x})] \mathbf{a} &= \frac{d}{dt} \mathbf{A}(\mathbf{x} + t\mathbf{a})|_{t=0}, \quad \text{for any } t \in R, \mathbf{a} \in E. \end{aligned} \quad (3)$$

In particular, gradient of product of 2nd order tensor $\mathbf{A}(\mathbf{x})$ and vector $\mathbf{v}(\mathbf{x})$ fields is calculated according to

$$\text{Grad} (\mathbf{A}\mathbf{v}) = \mathbf{v} \text{Grad} \mathbf{A}^T + \mathbf{A} \text{Grad} \mathbf{v}. \quad (4)$$

The wryness tensor $\mathbf{\Gamma}$ can also be expressed in several equivalent forms, see Pietraszkiewicz and Eremeyev (2009), for example

$$\mathbf{\Gamma} = \frac{1}{2} \mathbf{h}_a \times (\mathbf{h}_a \mathbf{Q}^T \text{Grad} \mathbf{Q}) - \mathbf{Q}^T \mathbf{C} \mathbf{F} - \mathbf{B}, \quad (5)$$

where \mathbf{B} and \mathbf{C} are the respective microstructure curvature tensors of polar continuum in the reference and actual placements defined by

$$\mathbf{B} = \frac{1}{2} \mathbf{h}_a \times \text{Grad} \mathbf{h}_a, \quad \mathbf{C} = \frac{1}{2} \mathbf{d}_a \times \text{grad} \mathbf{d}_a, \quad (6)$$

with the operator grad being taken in the deformed placement B_γ .

In what follows \mathbf{B} and \mathbf{C} play an important role because they characterize non uniform distributions of directors \mathbf{h}_a and \mathbf{d}_a in the reference and actual placements, respectively. In particular, if \mathbf{h}_a are constant in space then $\mathbf{B} = \mathbf{0}$.

Material behavior of the micropolar (hyper)elastic continuum is described by the strain energy density W_κ per unit volume of undeformed placement B_κ . The density W_κ satisfying the principle of material frame indifference takes the reduced form

$$W_\kappa = \widehat{W}_\kappa(\mathbf{E}, \mathbf{\Gamma}; \mathbf{x}, \mathbf{B}). \quad (7)$$

We call the polar elastic continuum *homogeneous* if there exists a reference placement B_κ such that W_κ does not depend on \mathbf{x} and *materially uniform* if W_κ does not depend on \mathbf{B} or $\mathbf{B} = \mathbf{0}$.

Constitutive equations for referential stress \mathbf{S} and couple stress \mathbf{K} tensors introduced in Pietraszkiewicz and Eremeyev (2009) are now defined as

$$\mathbf{S} = \frac{\partial W_\kappa}{\partial \mathbf{E}}, \quad \mathbf{K} = \frac{\partial W_\kappa}{\partial \mathbf{\Gamma}}. \quad (8)$$

It is apparent that \mathbf{S} and \mathbf{K} are not symmetric, in general, $\mathbf{S}^T \neq \mathbf{S}$, $\mathbf{K}^T \neq \mathbf{K}$.

The local Lagrangian equilibrium equations as well as kinematic and dynamic boundary conditions of the non linear theory of micropolar continuum are, see Pietraszkiewicz and Eremeyev (2009), Appendix A,

$$\text{Div} \mathbf{T}_N + \rho_\kappa \mathbf{f} = \mathbf{0}, \quad (9)$$

$$\text{Div} \mathbf{M}_N + \text{ax}(\mathbf{T}_N^T \mathbf{F}^T - \mathbf{F} \mathbf{T}_N) + \rho_\kappa \mathbf{c} = \mathbf{0} \quad \text{in } B_\kappa, \quad (10)$$

$$\mathbf{x} = \mathbf{x}^0, \quad \mathbf{Q} = \mathbf{Q}^0 = \mathbf{0} \quad \text{along } \partial B_{\kappa d}. \quad (11)$$

$$\mathbf{n} \mathbf{T}_N = \mathbf{t}^0 = \mathbf{0}, \quad \mathbf{n} \mathbf{M}_N = \mathbf{m}^0 = \mathbf{0} \quad \text{along } \partial B_{\kappa f}. \quad (12)$$

In Eqs. (9) and (10), divergence operator Div in B_κ is defined as, for example, in Billington (1986) by

$$[\text{Div} \mathbf{A}(\mathbf{x})] \mathbf{a} = \text{Div}[\mathbf{A}(\mathbf{x}) \mathbf{a}], \quad \forall \mathbf{A} \in E \otimes E, \quad \forall \mathbf{a} \in E. \quad (13)$$

In Eqs. (9) (12), $\mathbf{T}_N, \mathbf{M}_N \in E \otimes E$ are nominal stress and couple stress tensors following from the generalized Cauchy theorem $\mathbf{t}_{(n)} = \mathbf{n} \mathbf{T}_N$, $\mathbf{m}_{(n)} = \mathbf{n} \mathbf{M}_N$, in which $\mathbf{t}_{(n)}$ and $\mathbf{m}_{(n)}$ are surface traction and moment vectors applied at any point of boundary ∂P_γ of $P_\gamma \subset B_\gamma$, but measured per unit area of ∂P_κ , respectively, with \mathbf{n} being the unit vector externally normal to ∂P_κ . The nominal tensors $\mathbf{T}_N, \mathbf{M}_N$ are related to the referential tensors \mathbf{S} and \mathbf{K} given in (8) by

$$\mathbf{T}_N = \mathbf{S}^T \mathbf{Q}^T, \quad \mathbf{M}_N = \mathbf{K}^T \mathbf{Q}^T. \quad (14)$$

Additionally in (9) (12), $\mathbf{f}, \mathbf{c} \in E$ are the external force and couple vectors applied at any point of B_γ but measured per unit mass of B_κ , ρ_κ is the mass density in B_κ , while $\mathbf{t}^0, \mathbf{m}^0 \in E$ are the external boundary force and couple vectors applied on part of the deformed boundary $\partial B_{\gamma f}$ but measured per unit area of $\partial B_{\kappa f}$, respectively. The vector \mathbf{x}^0 and the tensor \mathbf{Q}^0 given on $\partial B_{\kappa d}$ describe translation and rotation of particles prescribed on $\partial B_{\kappa d}$, respectively, where $\partial B_\kappa = \partial B_{\kappa d} \cup \partial B_{\kappa f}$, $\partial B_{\kappa d} \cap \partial B_{\kappa f} = \emptyset$.

Formally different than (9) (12) but equivalent coordinate free forms of local equilibrium conditions follow by applying alternative definitions of gradient and divergence operators as well as of the Cauchy theorem, see for example Maugin (1974, 1998), Scarpetta (1989), Lurie (1990), Zubov (1997), Yeremeyev and Zubov (1999), Dai (2003).

The local Eulerian equilibrium equations and dynamic boundary conditions in the actual placement B_γ corresponding to (9) (12) are

$$\text{div} \mathbf{T} + \rho \mathbf{f} = \mathbf{0}, \quad \text{div} \mathbf{M} + \text{ax}(\mathbf{T}^T - \mathbf{T}) + \rho \mathbf{c} = \mathbf{0} \quad \text{in } B_\gamma, \quad (15)$$

$$\mathbf{n} \mathbf{T} = \mathbf{t}^0 = \mathbf{0}, \quad \mathbf{n} \mathbf{M} = \mathbf{m}^0 = \mathbf{0} \quad \text{along } \partial B_{\gamma f}. \quad (16)$$

In (15) and (16), \mathbf{T} and \mathbf{M} are Cauchy type stress and couple stress tensors which are related to the referential stress measures by

$$\mathbf{T} = (\det \mathbf{F})^{-1} \mathbf{F} \mathbf{T}_N, \quad (\det \mathbf{F})^{-1} \mathbf{F} \mathbf{S}^T \mathbf{Q}^T, \quad (17)$$

$$\mathbf{M} = (\det \mathbf{F})^{-1} \mathbf{F} \mathbf{M}_N, \quad (\det \mathbf{F})^{-1} \mathbf{F} \mathbf{K}^T \mathbf{Q}^T, \quad (18)$$

ρ is the mass density in B_γ , div is the divergence operator in B_γ , defined as in (13), $\mathbf{t}^0, \mathbf{m}^0$ are measured per unit area of $\partial B_{\gamma f}$, with \mathbf{n} being the unit vector externally normal to ∂B_γ .

3. Change of reference placement

Let us introduce another reference placement $\kappa_*(\mathcal{B}) \subset B_\kappa \in \mathcal{E}$ of \mathcal{B} , in which position $\kappa_* \in B_*$ of $X \in \mathcal{B}$ is given by the vector \mathbf{x}_* relative to the same origin $o \in \mathcal{E}$ and its orientation is fixed by three right handed orthonormal directors \mathbf{h}_{*a} (Fig. 2). Let \mathbf{P} , $\det \mathbf{P} > 0$, be the deformation gradient transforming $d\mathbf{x}$ into $d\mathbf{x}_*$, and $\mathbf{R} \in \text{Orth}^+$ be the rotation tensor transforming \mathbf{h}_a into \mathbf{h}_{*a} , so that

$$d\mathbf{x}_* = \mathbf{P} d\mathbf{x}, \quad \mathbf{h}_{*a} = \mathbf{R} \mathbf{h}_a. \quad (19)$$

In what follows all fields associated with deformation relative to B_* will be marked by lower index $*$.

Let us analyze how transform the strain measures \mathbf{E} and $\mathbf{\Gamma}$ under change of reference placement $B_\kappa \rightarrow B_*$.

Since $d\mathbf{y} = \mathbf{F} d\mathbf{x} = \mathbf{F}_* d\mathbf{x}_*$, where $\mathbf{F}_* = \text{Grad}_{*} \mathbf{y}(\mathbf{x}_*)$, $\det \mathbf{F}_* > 0$, is the deformation gradient relative to B_* and rotation \mathbf{h}_{*a} into \mathbf{d}_a is given by $\mathbf{Q}_* = \mathbf{d}_a \otimes \mathbf{h}_{*a} \in \text{Orth}^+$, from (1) and (19) it follows that

$$\mathbf{F} = \mathbf{F}_* \mathbf{P}, \quad \mathbf{Q} = \mathbf{Q}_* \mathbf{R}. \quad (20)$$

Thus, from (2)₁ and (20) we immediately obtain

$$\mathbf{E}_* = \mathbf{Q}_*^T \mathbf{F}_* - \mathbf{I} = \mathbf{R} \mathbf{E} \mathbf{P}^{-1} + \mathbf{R} \mathbf{P}^{-1} - \mathbf{I} = \mathbf{R}(\mathbf{E} + \mathbf{I}) \mathbf{P}^{-1} - \mathbf{I}. \quad (21)$$

To find $\mathbf{\Gamma}_*$ one needs first express \mathbf{B}_* through \mathbf{B} . Using (4), (6)₁, (19) and applying the chain rule $\text{Grad}_{*} \mathbf{h}_{*a} = (\text{Grad} \mathbf{h}_{*a}) \mathbf{P}^{-1}$, we obtain




Fig. 2. Change of reference placement.

$$\begin{aligned} \mathbf{B}_* &= \frac{1}{2} \mathbf{h}_{*a} \times \text{Grad}_* \mathbf{h}_{*a} - \frac{1}{2} [(\mathbf{R} \mathbf{h}_a) \text{Grad} (\mathbf{R} \mathbf{h}_a)] \mathbf{P}^{-1} \\ &= \mathbf{R} \left[\frac{1}{2} \mathbf{h}_a \times \mathbf{R}^T (\mathbf{h}_a \text{Grad} \mathbf{R}^T + \mathbf{R} \text{Grad} \mathbf{h}_a) \right] \mathbf{P}^{-1} \\ &= \mathbf{R} \left[\frac{1}{2} \mathbf{h}_a \times (\mathbf{h}_a \mathbf{R}^T \text{Grad} \mathbf{R}^T + \text{Grad} \mathbf{h}_a) \right] \mathbf{P}^{-1} = \mathbf{R} \mathbf{B} \mathbf{P}^{-1} = \mathbf{L}, \quad (22) \end{aligned}$$

$$\begin{aligned} \mathbf{L} &= \mathbf{R} \left[\frac{1}{2} \mathbf{h}_a \times (\mathbf{h}_a \mathbf{R} \text{Grad} \mathbf{R}^T) \right] \mathbf{P}^{-1} \\ &= \mathbf{R} \left[\frac{1}{2} \mathbf{E} : (\mathbf{R} \text{Grad} \mathbf{R}^T) \right] \mathbf{P}^{-1} = \mathbf{R} \mathbf{Z} \mathbf{P}^{-1}, \\ \mathbf{Z} &= \frac{1}{2} \mathbf{E} : (\mathbf{R} \text{Grad} \mathbf{R}^T). \quad (23) \end{aligned}$$

As a result,

$$\mathbf{F}_* = \mathbf{Q}_*^T \mathbf{C}_* \mathbf{B}_* = \mathbf{R} \mathbf{Q}^T \mathbf{C} \mathbf{P}^{-1} = \mathbf{R} \mathbf{B} \mathbf{P}^{-1} + \mathbf{L} = \mathbf{R} \mathbf{F} \mathbf{P}^{-1} + \mathbf{L}. \quad (24)$$

Let us note that the 2nd order tensors \mathbf{B} , \mathbf{C} , $\mathbf{\Gamma}$, \mathbf{M}_N and \mathbf{K} are axial tensors (pseudotensors), not usual (polar) ones. Axial tensors differ from polar ones in that they change signs under inversion transformation \mathbf{I} of 3D space E . Pseudoscalars, pseudovectors and pseudotensors are widely used in modern physics, see for example Nye (1957), Korn and Korn (1968), Feynman et al. (1977), Arfken and Weber (2000). An example of pseudovector is the vector product $\mathbf{a} \times \mathbf{b}$ of two polar vectors \mathbf{a} , $\mathbf{b} \in E$. If $\mathbf{O} \in \text{Orth}$ is an orthogonal tensor then $(\mathbf{O}\mathbf{a}) \times (\mathbf{O}\mathbf{b}) = (\det \mathbf{O}) \mathbf{O}(\mathbf{a} \times \mathbf{b})$ and the vector product changes sign if $\mathbf{O} = -\mathbf{I}$. An example of axial 2nd order tensor is the skew tensor $\mathbf{\Omega} = \boldsymbol{\omega} \times \mathbf{I}$ with the polar vector $\boldsymbol{\omega} \in E$. An example of axial 3rd order tensor is the permutation tensor \mathbf{E} .

Within classical (or Cauchy type) continuum discussed for example in Truesdell and Noll (1965), Truesdell (1966, 1991) it is enough to consider polar 2nd order tensors when defining the material symmetry group of Green elastic body. However, for the micropolar (or Cosserat type) continuum one has to take into account that $\mathbf{\Gamma}$ and \mathbf{B} appearing in the density W_x defined in (7) are axial 2nd order tensors, not the polar ones. Thus one has to extend B_* by allowing \mathbf{R} in (7) to be an orthogonal tensor $\mathbf{R} \in \text{Orth}$ and

by requiring \mathbf{P} to be non singular tensor, $\det \mathbf{P} \neq 0$. Additional inversion of space orientation does not correspond to any real deformation of the reference placement. It simply reflects necessary invariance of constitutive equations under mirror reflection of the reference placement or, equivalently, under change of orientation of the base vectors \mathbf{h}_a from right hand to left hand one. Then the refined transformations (22) and (24) become

$$\mathbf{B}_* = (\det \mathbf{R}) \mathbf{R} \mathbf{B} \mathbf{P}^{-1} = \mathbf{L}, \quad \mathbf{\Gamma}_* = (\det \mathbf{R}) \mathbf{R} \mathbf{\Gamma} \mathbf{P}^{-1} + \mathbf{L}. \quad (25)$$

4. Definition of material symmetry group

The form of elastic strain energy density W_x of the micropolar body at any particle $X \in \mathcal{B}$ depends upon the choice of reference placement, in general. Particularly important are sets of reference placements which leave unchanged the form of energy density. Transformations of reference placement under which the energy density remains unchanged are called here invariant transformations. Knowledge of all invariant transformations allows one to precisely define fluid, solid, liquid crystal or subfluid as well as to introduce notions of isotropic, hemitropic or orthotropic polar elastic continua, for example. Similar approach is used in classical continuum mechanics and in non linear elasticity by Truesdell (1964), Truesdell and Noll (1965), Wang and Truesdell (1973), Truesdell (1991), Rivlin (1980), as well as in non linear theories of shells by Wang (1973), Murdoch and Cohen (1979), Eremeyev and Pietraszkiewicz (2006).

The elastic strain energy density W_* relative to B_* depends in each point $x_* \in B_*$ on the stretch tensor \mathbf{E}_* , the wryness tensor $\mathbf{\Gamma}_*$, and also upon the structure curvature tensor \mathbf{B}_* . This dependence may, in general, be different than that of $W_x(\mathbf{E}, \mathbf{\Gamma}; \mathbf{x}, \mathbf{B})$. However, the strain energy of any part of continuum should be conserved, so that

$$\iiint_{P_x} W_x dv_x = \iiint_{P_*} W_* dv_* \quad (26)$$

for any part $P_x \subset B_x$ corresponding to $P_* \subset B_*$, because the functions W_x and W_* describe strain energy densities of the same deformed state of $P_\gamma \subset B_\gamma = \chi(P_x) = \chi_*(P_*)$, where χ_* is deformation function from B_* to B_γ .

Changing variables $\mathbf{x}_* \rightarrow \mathbf{x}$ in the right hand integral of (26), we obtain

$$\iiint_{P_*} W_*[\mathbf{E}_*(\mathbf{x}_*), \mathbf{\Gamma}_*(\mathbf{x}_*); \mathbf{x}_*, \mathbf{B}_*(\mathbf{x}_*)] dv_* \\ \iiint_{P_x} |\det \mathbf{P}| W_*[\mathbf{E}_*(\mathbf{x}), \mathbf{\Gamma}_*(\mathbf{x}); \mathbf{x}, \mathbf{B}_*(\mathbf{x})] dv_x.$$

Thus, from (26) it follows that W_* and W_x are related by

$$|\det \mathbf{P}| W_*(\mathbf{E}_*, \mathbf{\Gamma}_*; \mathbf{x}, \mathbf{B}_*) = W_x(\mathbf{E}, \mathbf{\Gamma}; \mathbf{x}, \mathbf{B}).$$

Here \mathbf{E}_* , $\mathbf{\Gamma}_*$, and \mathbf{B}_* are expressed as in (21) and (25).

From physical reasons invariant transformations of the reference placement should preserve elementary volume of B_x . Hence, the transformation tensor \mathbf{P} should belong to the unimodular group for which $|\det \mathbf{P}| = 1$.

The assumption that the constitutive relation be insensitive to the change of reference placement means that explicit forms of W_x and W_* should coincide, that is

$$W_x(\mathbf{E}, \mathbf{\Gamma}; \mathbf{x}, \mathbf{B}) = W_*(\mathbf{E}_*, \mathbf{\Gamma}_*; \mathbf{x}, \mathbf{B}_*).$$

In other words, one may use the same function for the strain energy density independently upon the choice of B_x or B_* , but with different expressions for stretch and wryness tensors as well as for the microstructure curvature tensor. In what follows we not always explicitly indicate that all functions depend also on the position vector \mathbf{x} and W is taken relative to the undeformed placement B_x .

Using (21) and (25) we obtain the following invariance requirement for W under change of reference placement:

$$W(\mathbf{E}, \mathbf{\Gamma}; \mathbf{B}) = W[\mathbf{R}\mathbf{E}\mathbf{P}^1 + \mathbf{R}\mathbf{P}^1 \quad \mathbf{I}, (\det \mathbf{R})\mathbf{R}\mathbf{\Gamma}\mathbf{P}^1 + \mathbf{L}; (\det \mathbf{R})\mathbf{R}\mathbf{B}\mathbf{P}^1 \quad \mathbf{L}]. \quad (27)$$

The relation (27) holds locally, i.e. it should be satisfied at any \mathbf{x} and \mathbf{B} , and tensors \mathbf{P} , \mathbf{R} , \mathbf{L} are treated as independent here. As a result, local invariance of W under change of reference placement is described by triple of tensors $(\mathbf{P}, \mathbf{R}, \mathbf{L})$.

In what follows we shall use the following nomenclature:

$Orth = \{\mathbf{O}: \mathbf{O}^{-1} = \mathbf{O}^T, \det \mathbf{O} = \pm 1\}$ the group of orthogonal tensors;
 $Orth^+ = \{\mathbf{O}: \mathbf{O} \in Orth, \det \mathbf{O} = 1\}$ the group of rotation tensors;
 $Unim = \{\mathbf{P}: \mathbf{P} \in E \otimes E, \det \mathbf{P} = \pm 1\}$ the unimodular group;
 $Lin = \{\mathbf{L} \in E \otimes E\}$ the linear group.

Here $Orth$ and $Unim$ are groups with regard to multiplication, and Lin is the group with regard to addition.

Now we are able to introduce the following definition:

Definition 1. By the *material symmetry group* \mathcal{G}_x at x and \mathbf{B} of the polar elastic continuum we call all sets of ordered triples of tensors

$$\mathbb{X} = (\mathbf{P} \in Unim, \mathbf{R} \in Orth, \mathbf{L} \in Lin), \quad (28)$$

satisfying the relation

$$W(\mathbf{E}, \mathbf{\Gamma}; \mathbf{B}) = W[\mathbf{R}\mathbf{E}\mathbf{P}^1 + \mathbf{R}\mathbf{P}^1 \quad \mathbf{I}, (\det \mathbf{R})\mathbf{R}\mathbf{\Gamma}\mathbf{P}^1 + \mathbf{L}; (\det \mathbf{R})\mathbf{R}\mathbf{B}\mathbf{P}^1 \quad \mathbf{L}] \quad (29)$$

for any tensors \mathbf{E} , $\mathbf{\Gamma}$, \mathbf{B} in domain of definition of function W .

The set \mathcal{G}_x is group relative to the group operation \circ defined by

$$(\mathbf{P}_1, \mathbf{R}_1, \mathbf{L}_1) \circ (\mathbf{P}_2, \mathbf{R}_2, \mathbf{L}_2) = [\mathbf{P}_1\mathbf{P}_2, \mathbf{R}_1\mathbf{R}_2, \mathbf{L}_1 + (\det \mathbf{R}_1)\mathbf{R}_1\mathbf{L}_2\mathbf{P}_1^1].$$

Let us check that if $\mathbb{X}_1 \equiv (\mathbf{P}_1, \mathbf{R}_1, \mathbf{L}_1) \in \mathcal{G}_x$ and $\mathbb{X}_2 \equiv (\mathbf{P}_2, \mathbf{R}_2, \mathbf{L}_2) \in \mathcal{G}_x$, then also $\mathbb{X}_1 \circ \mathbb{X}_2 \in \mathcal{G}_x$. Indeed, if $\mathbb{X}_1 \in \mathcal{G}_x$ and $\mathbb{X}_2 \in \mathcal{G}_x$ then

$$W(\mathbf{E}, \mathbf{\Gamma}; \mathbf{B}) = W[\mathbf{R}_1\mathbf{E}\mathbf{P}_1^1 + \mathbf{R}_1\mathbf{P}_1^1 \quad \mathbf{I}, (\det \mathbf{R}_1)\mathbf{R}_1\mathbf{\Gamma}\mathbf{P}_1^1 + \mathbf{L}_1; (\det \mathbf{R}_1)\mathbf{R}_1\mathbf{B}\mathbf{P}_1^1 \quad \mathbf{L}_1] \\ = W[\mathbf{R}_2\mathbf{E}\mathbf{P}_2^1 + \mathbf{R}_2\mathbf{P}_2^1 \quad \mathbf{I}, (\det \mathbf{R}_2)\mathbf{R}_2\mathbf{\Gamma}\mathbf{P}_2^1 + \mathbf{L}_2; (\det \mathbf{R}_2)\mathbf{R}_2\mathbf{B}\mathbf{P}_2^1 \quad \mathbf{L}_2].$$

Taking these relations into account we have

$$W[\mathbf{R}_1\mathbf{R}_2\mathbf{E}\mathbf{P}_2^1\mathbf{P}_1^1 + \mathbf{R}_1\mathbf{R}_2\mathbf{P}_2^1\mathbf{P}_1^1 \quad \mathbf{I}, \\ (\det \mathbf{R}_1)(\det \mathbf{R}_2)\mathbf{R}_1\mathbf{R}_2\mathbf{\Gamma}\mathbf{P}_2^1\mathbf{P}_1^1 + \mathbf{L}_1 + (\det \mathbf{R}_1)\mathbf{R}_1\mathbf{L}_2\mathbf{P}_1^1; \\ (\det \mathbf{R}_1)(\det \mathbf{R}_2)\mathbf{R}_1\mathbf{R}_2\mathbf{B}\mathbf{P}_2^1\mathbf{P}_1^1 \quad \mathbf{L}_1 \quad (\det \mathbf{R}_1)\mathbf{R}_1\mathbf{L}_2\mathbf{P}_1^1] \\ W\{\mathbf{R}_1(\mathbf{R}_2\mathbf{E}\mathbf{P}_2^1 + \mathbf{R}_2\mathbf{P}_2^1 \quad \mathbf{I})\mathbf{P}_1^1 + \mathbf{R}_1\mathbf{P}_1^1 \quad \mathbf{I}, \\ (\det \mathbf{R}_1)\mathbf{R}_1[(\det \mathbf{R}_2)\mathbf{R}_2\mathbf{\Gamma}\mathbf{P}_2^1 + \mathbf{L}_2]\mathbf{P}_1^1 + \mathbf{L}_1; \\ (\det \mathbf{R}_1)\mathbf{R}_1[(\det \mathbf{R}_2)\mathbf{R}_2\mathbf{B}\mathbf{P}_2^1 \quad \mathbf{L}_2]\mathbf{P}_1^1 \quad \mathbf{L}_1\} \\ W[\mathbf{R}_2\mathbf{E}\mathbf{P}_2^1 + \mathbf{R}_2\mathbf{P}_2^1 \quad \mathbf{I}, (\det \mathbf{R}_2)\mathbf{R}_2\mathbf{\Gamma}\mathbf{P}_2^1 + \mathbf{L}_2; (\det \mathbf{R}_2)\mathbf{R}_2\mathbf{B}\mathbf{P}_2^1 \quad \mathbf{L}_2] \\ W(\mathbf{E}, \mathbf{\Gamma}; \mathbf{B}),$$

which proves that $\mathbb{X}_1 \circ \mathbb{X}_2$ belongs to the group \mathcal{G}_x indeed.

The unit element of \mathcal{G}_x is $\mathbb{1} = (\mathbf{I}, \mathbf{I}, \mathbf{0})$, and the inverse element to $\mathbb{X} \in \mathcal{G}_x$ is given by

$$\mathbb{X}^{-1} \equiv (\mathbf{P}, \mathbf{R}, \mathbf{L})^{-1} = [\mathbf{P}^{-1}, \mathbf{R}^T, (\det \mathbf{R})\mathbf{R}^T\mathbf{L}\mathbf{P}].$$

Indeed,

$$\mathbb{X} \circ \mathbb{X}^{-1} \equiv (\mathbf{P}, \mathbf{R}, \mathbf{L}) \circ (\mathbf{P}, \mathbf{R}, \mathbf{L})^{-1} \\ = [\mathbf{P}\mathbf{P}^{-1}, \mathbf{R}\mathbf{R}^T, \mathbf{L} + (\det \mathbf{R})^2\mathbf{R}\mathbf{R}^T\mathbf{L}\mathbf{P}\mathbf{P}^{-1}] = (\mathbf{I}, \mathbf{I}, \mathbf{0}).$$

Our symmetry group \mathcal{G}_x differs from the one of micropolar elastic materials proposed by Eringen and Kafadar (1976). We take into account explicit dependence of W upon \mathbf{B} as parametric tensor, while in Eringen and Kafadar (1976) dependence of \mathbf{T} , \mathbf{M} and other quantities on the microinertia tensor \mathbf{J}_x is assumed. In our notation, definition of \mathcal{G}_x by Eringen and Kafadar (1976) is based on the formula

$$W(\mathbf{E}, \mathbf{\Gamma}; \mathbf{J}_x) = W[\mathbf{R}\mathbf{E}\mathbf{P}^1 + \mathbf{R}\mathbf{P}^1 \quad \mathbf{I}, \mathbf{R}\mathbf{\Gamma}\mathbf{P}^1 + \mathbf{L}; \mathbf{R}\mathbf{J}_x\mathbf{R}^T]. \quad (30)$$

The microinertia tensor \mathbf{J}_x appearing in (30) has another nature and transforms by another rule than \mathbf{B} under change of reference placement. Moreover, in what follows we take into account that $\mathbf{\Gamma}$ and \mathbf{B} as arguments of W are axial tensors. Eringen and Kafadar (1976) did not take into account that $\mathbf{\Gamma}$ was the axial tensor. The mentioned differences in definitions of \mathcal{G}_x lead to different restrictions of constitutive relations of polar elastic continua.

Let us show that Definition 1 allows one to establish an analog of Noll's rule given for classical simple material continuum by Noll (1958). Since the material symmetry group depends not only on particle $X \in \mathcal{B}$ but also upon choice of reference placement, let us analyze how symmetry groups corresponding to different reference placements are related. Let B_1 and B_2 be two different reference placements, and \mathcal{G}_1 and \mathcal{G}_2 be material symmetry groups relative to these reference placements, respectively. In what follows quantities described in the placements B_1 and B_2 are marked by the respective lower indices 1 and 2.

Let now \mathbf{P} be a non singular deformation gradient, $\det \mathbf{P} \neq 0$, \mathbf{R} be an orthogonal tensor associated with the transformation $B_1 \rightarrow B_2$, as well as \mathbf{P}^{-1} and \mathbf{R}^T be an inverse deformation gradient and an inverse orthogonal tensor associated with an inverse transformation $B_2 \rightarrow B_1$, respectively. Cases $\det \mathbf{P} < 1$ and $\det \mathbf{R} = -1$ correspond to change from right handed frame in B_1 to left handed one in B_2 . Then, by analogy to (21) and (25) we can relate the strain

measures \mathbf{E}_1 and \mathbf{E}_2 , $\mathbf{\Gamma}_1$ and $\mathbf{\Gamma}_2$, as well as the structure curvature tensors \mathbf{B}_1 and \mathbf{B}_2 defined relative to two different reference placements by

$$\mathbf{E}_2 = \mathbf{R}\mathbf{E}_1\mathbf{P}^{-1} + \mathbf{R}\mathbf{P}^{-1} \mathbf{I}, \quad \mathbf{\Gamma}_2 = (\det \mathbf{R})\mathbf{R}\mathbf{\Gamma}_1\mathbf{P}^{-1} + \mathbf{L}. \quad (31)$$

$$\mathbf{B}_2 = (\det \mathbf{R})\mathbf{R}\mathbf{B}_1\mathbf{P}^{-1} \mathbf{L}, \quad \mathbf{L} = (\det \mathbf{R})\mathbf{R}\mathbf{Z}\mathbf{P}^{-1}. \quad (32)$$

Let W_1 and W_2 be strain energy densities defined relative to the two reference placements. From (26) it follows that W_2 and W_1 are related by

$$|\det \mathbf{P}| W_2(\mathbf{E}_2, \mathbf{\Gamma}_2; \mathbf{B}_2) = W_1(\mathbf{E}_1, \mathbf{\Gamma}_1; \mathbf{B}_1).$$

Taking into account (31) and (32) we have

$$|\det \mathbf{P}| W_2 \left[\mathbf{R}\mathbf{E}_1\mathbf{P}^{-1} + \mathbf{R}\mathbf{P}^{-1} \mathbf{I}, (\det \mathbf{R})\mathbf{R}\mathbf{\Gamma}_1\mathbf{P}^{-1} + \mathbf{L}; (\det \mathbf{R})\mathbf{R}\mathbf{B}_1\mathbf{P}^{-1} \mathbf{L} \right] = W_1(\mathbf{E}_1, \mathbf{\Gamma}_1; \mathbf{B}_1). \quad (33)$$

Let the element $\mathbb{X}_1 \equiv (\mathbf{P}_1, \mathbf{R}_1, \mathbf{L}_1) \in \mathcal{G}_1$. Then using (33) we obtain

$$\begin{aligned} |\det \mathbf{P}| W_2(\mathbf{E}_2, \mathbf{\Gamma}_2; \mathbf{B}_2) &= W_1(\mathbf{E}_1, \mathbf{\Gamma}_1; \mathbf{B}_1) \\ &= W_1[\mathbf{R}_1\mathbf{E}_1\mathbf{P}_1^{-1} + \mathbf{R}_1\mathbf{I}\mathbf{P}_1^{-1} \mathbf{I}, \\ & \quad (\det \mathbf{R}_1)\mathbf{R}_1\mathbf{\Gamma}_1\mathbf{P}_1^{-1} + \mathbf{L}_1; \\ & \quad (\det \mathbf{R}_1)\mathbf{R}_1\mathbf{B}_1\mathbf{P}_1^{-1} \mathbf{L}_1] \\ &= |\det \mathbf{P}| W_2[\mathbf{R}\mathbf{R}_1\mathbf{E}_1\mathbf{P}_1^{-1}\mathbf{P}^{-1} + \mathbf{R}\mathbf{R}_1\mathbf{I}\mathbf{P}_1^{-1}\mathbf{P}^{-1} \mathbf{I}, \\ & \quad (\det \mathbf{R})(\det \mathbf{R}_1)\mathbf{R}\mathbf{R}_1\mathbf{\Gamma}_1\mathbf{P}_1^{-1}\mathbf{P}^{-1} \\ & \quad + (\det \mathbf{R})\mathbf{R}\mathbf{L}_1\mathbf{P}^{-1} + \mathbf{L}; \\ & \quad (\det \mathbf{R})(\det \mathbf{R}_1)\mathbf{R}\mathbf{R}_1\mathbf{B}_1\mathbf{P}_1^{-1}\mathbf{P}^{-1} \\ & \quad (\det \mathbf{R})\mathbf{R}\mathbf{L}_1\mathbf{P}^{-1} \mathbf{L}] \\ &= |\det \mathbf{P}| W_2[\mathbf{R}\mathbf{R}_1\mathbf{R}^T\mathbf{E}_2\mathbf{P}\mathbf{P}_1^{-1}\mathbf{P}^{-1} \\ & \quad + \mathbf{R}\mathbf{R}_1\mathbf{R}^T\mathbf{I}\mathbf{P}\mathbf{P}_1^{-1}\mathbf{P}^{-1} \mathbf{I}, \\ & \quad (\det \mathbf{R}_1)\mathbf{R}\mathbf{R}_1\mathbf{R}^T\mathbf{\Gamma}_2\mathbf{P}\mathbf{P}_1^{-1}\mathbf{P}^{-1} \\ & \quad (\det \mathbf{R}_1)\mathbf{R}\mathbf{R}_1\mathbf{R}^T\mathbf{L}\mathbf{P}\mathbf{P}_1^{-1}\mathbf{P}^{-1} \\ & \quad + (\det \mathbf{R}_1)\mathbf{R}\mathbf{L}_1\mathbf{P}^{-1} + \mathbf{L}; \\ & \quad (\det \mathbf{R}_1)\mathbf{R}\mathbf{R}_1\mathbf{R}^T\mathbf{B}_2\mathbf{P}\mathbf{P}_1^{-1}\mathbf{P}^{-1} \\ & \quad + (\det \mathbf{R}_1)\mathbf{R}\mathbf{R}_1\mathbf{R}^T\mathbf{L}\mathbf{P}\mathbf{P}_1^{-1}\mathbf{P}^{-1} \\ & \quad (\det \mathbf{R}_1)\mathbf{R}\mathbf{L}_1\mathbf{P}^{-1} \mathbf{L}]. \end{aligned} \quad (34)$$

From (34) it follows that the element $\mathbb{X}_2 \equiv (\mathbf{P}_2, \mathbf{R}_2, \mathbf{L}_2) \in \mathcal{G}_2$, where

$$\begin{aligned} \mathbf{P}_2 &= \mathbf{P}\mathbf{P}_1\mathbf{P}^{-1}, \quad \mathbf{R}_2 = \mathbf{R}\mathbf{R}_1\mathbf{R}^T, \\ \mathbf{L}_2 &= \mathbf{L} + (\det \mathbf{R}_1)\mathbf{R}\mathbf{L}_1\mathbf{P}^{-1} - (\det \mathbf{R}_1)\mathbf{R}\mathbf{R}_1\mathbf{R}^T\mathbf{L}\mathbf{P}\mathbf{P}_1^{-1}\mathbf{P}^{-1}. \end{aligned}$$

It is easy to show that $\mathbb{X}_2 = \mathbb{P} \circ \mathbb{X}_1 \circ \mathbb{P}^{-1}$, where $\mathbb{P} \equiv (\mathbf{P}, \mathbf{R}, \mathbf{L})$. Indeed,

$$\mathbb{P} \circ \mathbb{X}_1 \equiv (\mathbf{P}, \mathbf{R}, \mathbf{L}) \circ (\mathbf{P}_1, \mathbf{R}_1, \mathbf{L}_1) = \left[\mathbf{P}\mathbf{P}_1, \mathbf{R}\mathbf{R}_1, \mathbf{L} + (\det \mathbf{R})\mathbf{R}^T\mathbf{L}\mathbf{P} \right].$$

Taking into account that $\mathbb{P}^{-1} = [\mathbf{P}^{-1}, \mathbf{R}^T, (\det \mathbf{R})\mathbf{R}^T\mathbf{L}\mathbf{P}]$, we obtain

$$\begin{aligned} \mathbb{P} \circ \mathbb{X}_1 \circ \mathbb{P}^{-1} &= \left[\mathbf{P}\mathbf{P}_1\mathbf{P}^{-1}, \mathbf{R}\mathbf{R}_1\mathbf{R}^T, \mathbf{L} + (\det \mathbf{R})^2(\det \mathbf{R}_1)\mathbf{R}\mathbf{L}_1\mathbf{P}^{-1} \right. \\ & \quad \left. (\det \mathbf{R})^2(\det \mathbf{R}_1)\mathbf{R}\mathbf{R}_1\mathbf{R}^T\mathbf{L}\mathbf{P}\mathbf{P}_1^{-1}\mathbf{P}^{-1} \right], \end{aligned}$$

from which follows the sought result.

Thus the material symmetry group under change of reference placement transforms according to

$$\mathcal{G}_2 = \mathbb{P} \circ \mathcal{G}_1 \circ \mathbb{P}^{-1}. \quad (35)$$

The transformation (35) is a counterpart in non linear micropolar elasticity of the well known Noll rule for symmetry groups of simple materials in classical continuum mechanics, see Noll (1958), Truesdell and Noll (1965), Wang and Truesdell (1973).

5. Material symmetry and reduced forms of strain energy density

The structure of \mathcal{G}_κ puts some constraints on the form of W which allow us to considerably simplify this form.

In this Section we consider some simple specific cases of \mathcal{G}_κ .

Case 1. Let us discuss the trivial symmetry group \mathcal{G}_κ consisting of two elements: \mathbb{I} and $\mathbb{I}(\mathbf{I}, \mathbf{I}, \mathbf{0})$. Applying \mathbb{I} , from definition of \mathcal{G}_κ it follows that W should be an even function of $\mathbf{\Gamma}$ and \mathbf{B} , that is

$$W(\mathbf{E}, \mathbf{\Gamma}; \mathbf{B}) = W(\mathbf{E}, -\mathbf{\Gamma}; \mathbf{B}). \quad (36)$$

The condition (36) means, in particular, that explicit expression of $W(\mathbf{E}, \mathbf{\Gamma}; \mathbf{B})$ cannot have terms linear in $\mathbf{\Gamma}$ alone such as $\text{tr}(\mathbf{E}^T\mathbf{\Gamma})$. Only when explicit dependence of W on \mathbf{B} is taken into account, terms linear in $\mathbf{\Gamma}$ of the type $\text{tr}(\mathbf{B}^T\mathbf{\Gamma})$ or $\text{tr}(\mathbf{B}^T\mathbf{\Gamma}\mathbf{E})$ are allowed.

Case 2. If \mathcal{G}_κ consists of tensor triples containing two identities and an arbitrary tensor $\mathbf{L} \in \text{Lin}$, then the number of arguments in W can be reduced.

Indeed, let $\mathbb{X} = (\mathbf{I}, \mathbf{I}, \mathbf{L}) \in \mathcal{G}_\kappa$. Then

$$W(\mathbf{E}, \mathbf{\Gamma}; \mathbf{B}) = W(\mathbf{E}, \mathbf{\Gamma} + \mathbf{L}; \mathbf{B} - \mathbf{L}), \quad \forall \mathbf{L} \in \text{Lin}. \quad (37)$$

Introducing one parameter family of transformations

$$W(\mathbf{E}, \mathbf{\Gamma}; \mathbf{B}) = W(\mathbf{E}, \mathbf{\Gamma} + t\mathbf{L}; \mathbf{B} - t\mathbf{L}), \quad \forall \mathbf{L} \in \text{Lin}, \quad \forall t \in \mathbb{R},$$

and differentiating the latter equation with regard to t at $t = 0$, we find that

$$0 = \frac{\partial W_\kappa}{\partial \mathbf{\Gamma}} : \mathbf{L} - \frac{\partial W_\kappa}{\partial \mathbf{B}} : \mathbf{L}, \quad \forall \mathbf{L} \in \text{Lin}.$$

This equation is satisfied if and only if

$$W = W(\mathbf{E}, \mathbf{\Gamma} + \mathbf{B}) = \widetilde{W}(\mathbf{E}, \mathbf{\Pi}),$$

where $\mathbf{\Pi} = \mathbf{\Gamma} + \mathbf{B}$. Here $:$ denotes the double dot (scalar) product of two 2nd order tensors, $\mathbf{A}:\mathbf{B} = \text{tr}(\mathbf{A}^T\mathbf{B}) = A_{mn}B_{mn}$.

Alternatively, since (37) should be satisfied by any \mathbf{L} let us take \mathbf{L} to be equal \mathbf{B} . Then from (37) it directly follows that

$$W(\mathbf{E}, \mathbf{\Gamma}; \mathbf{B}) = W(\mathbf{E}, \mathbf{\Gamma} + \mathbf{B}; \mathbf{0}) = \widetilde{W}(\mathbf{E}, \mathbf{\Pi}).$$

Case 3. Let us consider the reduced form $W = W(\mathbf{E}, \mathbf{\Gamma})$, i.e. let us neglect its explicit dependence on \mathbf{B} . This form is widely used in the literature, see for example Maugin (1998), Nikitin and Zubov (1998), Eringen (1999), Ramezani and Naghdabadi (2007), Ramezani et al. (2009).

If $\mathcal{G}_\kappa = \{\mathbb{X} : \mathbb{X} = (\mathbf{I}, \mathbf{I}, \mathbf{L})\}$ contains an arbitrary tensor $\mathbf{L} \in \text{Lin}$, we have $W(\mathbf{E}, \mathbf{\Gamma}) = W(\mathbf{E}, \mathbf{\Gamma} + \mathbf{L})$. Then we can again introduce one parameter family of transformations

$$W(\mathbf{E}, \mathbf{\Gamma}) = W(\mathbf{E}, \mathbf{\Gamma} + t\mathbf{L}), \quad \forall \mathbf{L} \in \text{Lin}, \quad \forall t \in \mathbb{R}.$$

Differentiating this equation with regard to t at $t = 0$, we obtain

$$\frac{\partial W}{\partial \mathbf{\Gamma}} : \mathbf{L} = \mathbf{0}, \quad \forall \mathbf{L} \in \text{Lin},$$

from which it follows that $\partial W / \partial \mathbf{\Gamma} = \mathbf{0}$. Thus in this case W takes the reduced form

$$W = W(\mathbf{E}). \quad (38)$$

Such a significant reduction of $W = W(\mathbf{E}, \mathbf{\Gamma})$ follows directly from assumption that the material symmetry group contains an arbitrary tensor \mathbf{L} . To keep the second argument $\mathbf{\Gamma}$ in W it is necessary to avoid using material symmetry groups containing an arbitrary tensor \mathbf{L} .

The form (38) of W corresponds to so called reduced elastic Cosserat continuum in which the couple stress tensor \mathbf{K} vanishes

while the stress tensor \mathbf{S} still remains non symmetric, in general. The couple equilibrium equations (10) and (15)₂ reduce to

$$\text{ax}(\mathbf{T}_N^T \mathbf{F}^T \quad \mathbf{F} \mathbf{T}_N) + \rho_x \mathbf{c} = \mathbf{0} \quad \text{in } B_x, \quad (39)$$

$$\text{ax}(\mathbf{T}^T \quad \mathbf{T}) + \rho \mathbf{c} = \mathbf{0} \quad \text{in } B_\gamma. \quad (40)$$

These equations become non trivial here and can be used to find the field of rotation \mathbf{Q} . However, since (39) and (40) constitute the system of non linear arbitrary equations with respect to \mathbf{Q} , it is not possible to assume arbitrary rotation \mathbf{Q} at the boundary $\partial B_{x,d}$.

The linear version of reduced Cosserat elastic continuum was used to describe wave propagation in soils and rocks, see for example Grekova et al. (2009).

Case 4. Let the material symmetry group of reduced Cosserat continuum (Case 3) be wider: $\mathcal{G}_x = \{\mathbb{X} : \mathbb{X} \in \{\mathbf{I}, \mathbf{R} \in \text{Orth}^+, \mathbf{L} \in \text{Lin}\}\}$. Then it describes the classical non linear Green elastic material (Truesdell and Noll, 1965; Ogden, 1984) for which $W = W(\mathbf{U})$, where $\mathbf{U} = (\mathbf{F}^T \mathbf{F})^{1/2}$.

Within the reduced Cosserat continuum tensors \mathbf{P} , \mathbf{R} and \mathbf{L} belong to \mathcal{G}_x if

$$W(\mathbf{E}) = W(\mathbf{R} \mathbf{E} \mathbf{P}^1 + \mathbf{R} \mathbf{P}^1 \mathbf{I}) = W(\mathbf{R} \mathbf{Q}^T \mathbf{F} \mathbf{P}^1 \mathbf{I}).$$

Let us substitute here $\mathbf{P} = \mathbf{I}$ and $\mathbf{R} = \mathbf{A}^T \mathbf{Q}$, where \mathbf{A} is the proper orthogonal tensor following from the polar decomposition $\mathbf{F} = \mathbf{A} \mathbf{U}$ of deformation gradient \mathbf{F} and $\mathbf{Q} \in \text{Orth}^+$. Then we obtain

$$W(\mathbf{E}) = W(\mathbf{A}^T \mathbf{Q} \mathbf{Q}^T \mathbf{F} \mathbf{I}) = W(\mathbf{A}^T \mathbf{F} \mathbf{I}) = W(\mathbf{U} \mathbf{I}) = W^\times(\mathbf{U}).$$

In this case the corresponding Cauchy stress tensor \mathbf{T} becomes symmetric, $\mathbf{T} = \mathbf{T}^T$. The proof is based on straightforward calculation of \mathbf{S} using $W = W^\times(\mathbf{U})$ which gives us the formula

$$\mathbf{T} = (\det \mathbf{F})^{-1} \mathbf{A} \mathbf{U} \mathbf{S}^\times \mathbf{A}^T,$$

where $\mathbf{S}^\times = dW^\times/d\mathbf{U}$, and $\mathbf{S}^{\times T} = \mathbf{S}^\times$.

Since $\mathbf{K} = \mathbf{0}$, the couple equilibrium equations (10) or (15)₂ are identically satisfied for vanishing couple vector \mathbf{c} . Material symmetry groups and representations of constitutive equations of non linear elastic continuum were discussed for example in Truesdell and Noll (1965), Wang and Truesdell (1973), Ogden (1984).

Four simple specific cases discussed above demonstrate that knowledge of elements of the material symmetry group allows one to substantially simplify the form of W . In next sections we consider some cases of \mathcal{G}_x which lead to definitions of polar elastic solids, fluids, liquid crystals, or subfluids, and to their reduced constitutive equations.

6. Polar-elastic isotropic material

As in case of non polar elastic materials, property of isotropy of the polar elastic material is expressed in terms of orthogonal group.

Definition 2. The micropolar elastic continuum is called *isotropic* at x and \mathbf{B} if there exists a reference placement B_x , called undistorted, such that the material symmetry group relative to B_x contains the group \mathcal{S}_x ,

$$\mathcal{S}_x \subset \mathcal{G}_x, \quad \mathcal{S}_x \equiv \{(\mathbf{P} \quad \mathbf{O}, \mathbf{O}, \mathbf{O}) : \mathbf{O} \in \text{Orth}\}. \quad (41)$$

From physical point of view this definition means that uniform rotations and mirror reflections of the undistorted reference placement B_x cannot be recognized by any experiment.

7. Polar-elastic fluids

The strain energy density of micropolar elastic fluids should be insensitive to any change of reference placement, i.e. the Eq. (27) should be satisfied by any triple of tensors $\mathbf{P} \in \text{Unim}$, $\mathbf{R} \in \text{Orth}$, $\mathbf{L} \in \text{Lin}$.

Definition 3. The micropolar elastic continuum is called the *polar elastic fluid* at x and \mathbf{B} if there exists a reference placement B_x , called undistorted, such that the material symmetry group relative to B_x is given by

$$\mathcal{G}_x = \mathcal{U}_x \equiv \{(\mathbf{P} \in \text{Unim}, \mathbf{R} \in \text{Orth}, \mathbf{L} \in \text{Lin})\}. \quad (42)$$

Hence, the strain energy density of the polar elastic fluid satisfies the relation

$$\begin{aligned} W_x(\mathbf{E}, \mathbf{F}; \mathbf{B}) &= W_x[\mathbf{R} \mathbf{E} \mathbf{P}^1 + \mathbf{R} \mathbf{P}^1 \mathbf{I}, (\det \mathbf{R}) \mathbf{R} \mathbf{F} \mathbf{P}^1 + \mathbf{L}; \\ &(\det \mathbf{R}) \mathbf{R} \mathbf{P}^1 \mathbf{L}], \\ &\forall \mathbf{P} \in \text{Unim}, \quad \forall \mathbf{R} \in \text{Orth}, \quad \forall \mathbf{L} \in \text{Lin}. \end{aligned} \quad (43)$$

From Noll's rule (35) it is easy to verify that any reference placement of the polar elastic fluid becomes undistorted, similarly as it is for the non polar elastic fluid, because the symmetry group becomes here maximal. Obviously, any polar elastic fluid is also isotropic.

Strain energy density of polar elastic fluid satisfying (43) takes the form

$$W = W(\mathbf{E}, \mathbf{C}) = W^\times(\det \mathbf{F}, \mathbf{C}), \quad (44)$$

where \mathbf{C} is the structure curvature tensor of the deformed placement B_γ defined in (6)₂. Since \mathbf{C} does not depend on the choice of reference placements, it is easy to check that (44) satisfies the requirement (43).

The strain energy density (44) describes polar elastic continuum which is insensitive to arbitrary deformations preserving an elementary volume element. However, it is still sensitive to change of orientation of particles.

By the principle of material frame indifference (invariance under superposed rigid body deformation) the function W^\times satisfies the condition

$$W(\det \mathbf{F}, \mathbf{C}) = W^\times[\det(\mathbf{Q}^T \mathbf{F}), \mathbf{O}^T \mathbf{C} \mathbf{O}], \quad \forall \mathbf{O} \in \text{Orth}^+,$$

i.e. W^\times is the hemitropic function with regard to \mathbf{C} .

Using general representations of isotropic and hemitropic scalar valued functions of one non symmetric tensor \mathbf{C} given by Spencer and Rivlin (1962), Smith (1965), Smith (1994), Spencer (1965), Spencer (1971), Smith and Smith (1971) with the help of algebraic theory of the invariants, W^\times can be constructed as a function of six invariants j_n , $n = 1, \dots, 6$,

$$W = W^\times(\det \mathbf{F}, j_1, j_2, \dots, j_6) \quad (45)$$

where

$$\begin{aligned} j_1 &= \text{tr } \mathbf{C}, \quad j_2 = \text{tr } \mathbf{C}^2, \quad j_3 = \text{tr } \mathbf{C}^3, \quad j_4 = \text{tr}(\mathbf{C} \mathbf{C}^T), \\ j_5 &= \text{tr}(\mathbf{C}^2 \mathbf{C}^T), \quad j_6 = \text{tr}(\mathbf{C}^2 \mathbf{C}^{T2}) \end{aligned} \quad (46)$$

as was used for example in Kafadar and Eringen (1971).

Since $\det \mathbf{F} = \rho_x / \rho$, the strain energy density of the polar elastic fluid may be given in another form more convenient in hydrodynamics,

$$W = \bar{W}(\rho, \mathbf{C}) = \bar{W}(\rho, j_1, j_2, \dots, j_6). \quad (47)$$

Because \mathbf{C} is the axial tensor, it changes sign under change of frame orientation. Thus, \bar{W} should be an even function of \mathbf{C} , $\bar{W}(\rho, \mathbf{C}) = \bar{W}(\rho, -\mathbf{C})$, and its invariant expression should have the property

$$\bar{W}(\rho, j_1, j_2, j_3, j_4, j_5, j_6) = \bar{W}(\rho, j_1, j_2, j_3, j_4, j_5, j_6).$$

As a simple example of the polynomial strain energy density we can propose the following quadratic function:

$$\bar{W}(\rho, \mathbf{C}) = \alpha_0(\rho) + \alpha_1 j_1^2 + \alpha_2 j_2 + \alpha_3 j_4, \quad (48)$$

where α_a , $a = 1, 2, 3$, are assumed to be constants.

Constitutive equations for \mathbf{T} and \mathbf{M} corresponding to (47) are

$$\mathbf{T} = p\mathbf{I} - \mathbf{MC}, \quad \mathbf{M} = \frac{\rho}{\rho_x} \frac{\partial \bar{W}}{\partial \mathbf{C}^T}, \quad p = \frac{\rho^2}{\rho_x} \frac{\partial \bar{W}}{\partial \rho}. \quad (49)$$

For example, with the strain energy density (48) the couple stress tensor \mathbf{M} is given by the constitutive equation

$$\mathbf{M} = 2 \frac{\rho}{\rho_x} (\alpha_1 j_1 \mathbf{I} + \alpha_2 \mathbf{C}^T + \alpha_3 \mathbf{C}).$$

Basic equations of viscous micropolar fluids were proposed by Aero et al. (1965), Eringen (1966). Similar theory of fluids with three rigid directors as primary quantities was introduced by Allen et al. (1967). Within the framework of these models the strain energy density had the form $W = W(\rho)$, which is the particular case of (47). Strain energy density (44) of the polar elastic fluids was established by Zubov and Eremeev (1996), Yeremeyev and Zubov (1999) without introduction of material symmetry group.

8. Polar-elastic solids

Material symmetry group of non polar non linear elastic solids is constructed with the help of orthogonal transformations describing rotations and reflections of reference placement, see Truesdell and Noll (1965), Wang and Truesdell (1973), Ogden (1984). But for polar elastic continua material symmetry group consists of ordered triple of independent tensors, $\mathcal{G}_x = (\mathbf{P}, \mathbf{R}, \mathbf{L})$.

In Definition 3 of the polar elastic fluid tensors \mathbf{P} and \mathbf{R} are entirely independent from each other. Considering directors \mathbf{h}_a as unit orthogonal vectors rigidly attached to rigid particles (e.g. molecules) of the fluid, it is apparent that \mathbf{P} need not coincide with \mathbf{R} . This is also true in the mechanical modeling of sand, powder, etc.

In polar elastic solids both \mathbf{P} and \mathbf{R} describing change of the reference placement should be orthogonal. Then corresponding constitutive equations might be sensitive to differences of both orthogonal tensors \mathbf{P} and \mathbf{R} , which would be difficult to accept. Hence, we assume here that $\mathbf{P} = \mathbf{R} \in Orth$.

The case $\mathbf{L} \neq \mathbf{0}$ corresponds to a non homogeneous field of \mathbf{R} , see (23). It can be shown that the material symmetry group with $\mathbf{L} \neq \mathbf{0}$ would lead to constitutive equations describing polar elastic liquid crystals and subfluids. Hence, for the polar elastic solids the following hypothesis seems to be physically justified:

Hypothesis. The material symmetry group of the polar elastic solids consists of all transformations of reference placement performed by the same orthogonal transformations of position vector and directors. Additionally we assume that $\mathbf{L} = \mathbf{0}$.

The hypothesis requires that $\mathbf{P} = \mathbf{R}$ and values of \mathbf{R} should be long to a subgroup of *Orth*.

Accepting the hypothesis we can propose the following definition:

Definition 4. The micropolar elastic continuum is called *the polar elastic solid* at x and \mathbf{B} if there exists a reference placement B_x , called undistorted, such that the material symmetry group relative to B_x is given by

$$\mathcal{G}_x = \mathcal{R}_x \equiv \{(\mathbf{P}, \mathbf{O}, \mathbf{0}, \mathbf{0}) : \mathbf{O} \in \mathcal{O}_x \subset Orth\}. \quad (50)$$

The group \mathcal{R}_x is fully described by a subgroup \mathcal{O}_x of orthogonal group *Orth*. Invariance requirement of W leads here to finding the subgroup \mathcal{O}_x such that

$$W(\mathbf{E}, \Gamma; \mathbf{B}) = W[\mathbf{OEO}^T, (\det \mathbf{O})\mathbf{O}\Gamma\mathbf{O}^T; (\det \mathbf{O})\mathbf{OBO}^T], \quad \forall \mathbf{O} \in \mathcal{O}_x. \quad (51)$$

Both our definitions of fluids (43) and solids (51) differ from definitions proposed by Eringen and Kafadar (1976) by factors $\det \mathbf{R}$ and $\det \mathbf{O}$, which take into account that our Γ and \mathbf{B} are axial tensors. Additionally, the tensor \mathbf{B} appears instead of \mathbf{J}_x in the list of arguments of our W .

9. Polar-elastic liquid crystals and subfluids

The strain energy density of polar elastic continuum may also admit other material symmetry groups, in general. For example, it is possible to construct material symmetry groups of W in analogy to symmetry groups used to model liquid crystals or subfluids in continuum mechanics of simple materials, see Coleman (1965), Wang (1965), Truesdell and Noll (1965), Wang and Truesdell (1973). However, those mathematical models differ from physical models of liquid crystals used for nematics, smectics and other materials, which are usually based on introducing one rigid director, c.f. Truesdell and Noll (1965), de Gennes and Prost (1993), Ericksen (1998).

Definition 5. The micropolar elastic continuum is called *the polar elastic liquid crystal* at x and \mathbf{B} if the material symmetry group \mathcal{G}_x does not coincide with \mathcal{U}_x , but there exist elements $\mathbb{X} \in \mathcal{G}_x$, which are not members of any group constructed using only orthogonal tensors.

Definition 5 means that a polar elastic liquid crystal is neither a polar elastic fluid nor a polar elastic solid.

Within micropolar elastic continuum the number of material symmetry groups corresponding to polar elastic liquid crystals is larger than in case of non polar elastic liquid crystals, because the structure of \mathcal{G}_x of the former is more complex. The Cases 2-4 considered above belong to polar elastic liquid crystals by their definitions. Below we give other examples of polar elastic liquid crystals.

Grekova and Zhilin (2001) defined the Kelvin medium as a special case of polar elastic solids, which strain energy density $W = W(\mathbf{E}, \Gamma)$ is insensitive to rotation by any angle φ about a fixed axis with unit vector \mathbf{e} . It is possible to prove that the Kelvin medium should in fact be considered as a polar elastic liquid crystal, because its material symmetry group contains non orthogonal tensors.

Within non polar elastic materials Wang developed constitutive equations of 14 classes of the simple subfluids being special cases of non polar liquid crystals, see Wang (1965), Truesdell and Noll (1965), Wang and Truesdell (1973), Cohen and Wang (1983). The material symmetry group of simple subfluids contained only elements of the dilatation group, that is the group of all unimodular tensors having three fixed linearly independent vectors as their proper vectors. The dilatation group consists of tensors

$$\mathcal{D}_x = \{\mathbf{P} : \mathbf{P} \in Unim, \mathbf{P} = p_1 \mathbf{e}_1 \otimes \mathbf{e}_1 + p_2 \mathbf{e}_2 \otimes \mathbf{e}_2 + p_3 \mathbf{e}_3 \otimes \mathbf{e}_3, p_1 p_2 p_3 = \pm 1\}.$$

Here $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ are three orthogonal unit vectors called the preferred basis.

In analogy to Wang (1965) we can define polar elastic subfluids as follows:

Definition 6. The micropolar elastic continuum is called *the polar elastic subfluid* at x and \mathbf{B} if its material symmetry group \mathcal{G}_x contains elements of the dilatation group, that is

$$\mathcal{G}_x = \{(\mathbf{P}, \mathbf{R}, \mathbf{L}) : \mathbf{P} \in \mathcal{D}_x, \mathbf{R} \in Orth, \mathbf{L} \in Lin\}.$$

Discussing application of micropolar elastic continuum for modeling liquid crystals, Eringen (1997) developed constitutive equations based on the strain energy density

$$W = W(\rho, \gamma; \mathbf{j}), \quad (52)$$

where $\mathbf{j} = \mathbf{Q}\mathbf{j}_x\mathbf{Q}^T$, $\gamma = \mathbf{Q}\mathbf{\Gamma}\mathbf{F}^{-1}$ in our notation. The relative Eulerian wryness measure γ determines relative changes of orientation of the polar elastic continuum, see [Pietraszkiewicz and Eremeyev \(2009\)](#). From (5) it follows that

$$\gamma = \mathbf{C} - \mathbf{Q}\mathbf{B}\mathbf{F}^{-1}.$$

Hence, $\gamma = \mathbf{C}$ if and only if $\mathbf{B} = \mathbf{0}$. In case of $\mathbf{B} \neq \mathbf{0}$ the strain energy density W in (52) corresponds to a special type of polar elastic continuum, neither a fluid nor a solid, because W depends on the relative wryness measure γ which is sensitive to change of orientation of continuum particles with respect to the reference placement. Under change of the reference placement γ transforms according to

$$\gamma_* = \gamma + \mathbf{Q}\mathbf{Z}\mathbf{F}^{-1},$$

$$\text{where } \gamma_* = \mathbf{C} - \mathbf{Q}_*\mathbf{B}_*\mathbf{F}_*^{-1}.$$

Using \mathbf{j} as the structural tensor, Eringen derived various types of constitutive equations. Since in our model \mathbf{B} is used as the parametric tensor instead of \mathbf{j} , Eringen's model may be incomparable with ours in sense of the material symmetry group.

Brief discussion stretched above and in Section 5 indicates that polar elastic liquid crystals and subfluids deserve detailed presentation in a separate paper.

10. Some material symmetry groups for polar-elastic solids

Let us discuss simplified forms of W for some particular cases of anisotropy.

Definition 7 (*Isotropic material*). The polar elastic solid is called *isotropic* at x and \mathbf{B} if there exists a reference placement B_x , called undistorted, such that the material symmetry group relative to B_x takes the form

$$\mathcal{G}_x = \mathcal{S}_x \equiv \{(\mathbf{P} \quad \mathbf{O}, \mathbf{O}, \mathbf{O}) : \mathbf{O} \in Orth\}. \quad (53)$$

This definition means that the strain energy density of the polar elastic isotropic solid satisfies the relation

$$W(\mathbf{E}, \mathbf{\Gamma}; \mathbf{B}) = W[\mathbf{O}\mathbf{E}\mathbf{O}^T, (\det \mathbf{O})\mathbf{O}\mathbf{\Gamma}\mathbf{O}^T; (\det \mathbf{O})\mathbf{O}\mathbf{B}\mathbf{O}^T], \quad \forall \mathbf{O} \in Orth.$$

Scalar valued isotropic functions of a few 2nd order tensors can be expressed by so called representation theorems in terms of joint invariants of the tensorial arguments. To construct representation for $W(\mathbf{E}, \mathbf{\Gamma}; \mathbf{B})$ as an isotropic function of joint invariants of \mathbf{E} , $\mathbf{\Gamma}$, and \mathbf{B} , we recall first the results of [Spencer \(1965, 1971\)](#) on isotropic scalar valued functions which are invariant with respect to proper orthogonal tensors $\mathbf{O} \in Orth^+$. Then, we apply additional restriction on the representations by requiring also invariance with respect to mirror reflection, i.e. additionally considering $\mathbf{O} = -\mathbf{I}$.

Decomposing non symmetric tensors \mathbf{E} , $\mathbf{\Gamma}$, and \mathbf{B} into symmetric and skew parts,

$$\begin{aligned} \mathbf{E} &= \mathbf{E}_S + \mathbf{E}_A, & \mathbf{E}_S &= \frac{1}{2}(\mathbf{E} + \mathbf{E}^T), & \mathbf{E}_A &= \frac{1}{2}(\mathbf{E} - \mathbf{E}^T), \\ \mathbf{\Gamma} &= \mathbf{\Gamma}_S + \mathbf{\Gamma}_A, & \mathbf{\Gamma}_S &= \frac{1}{2}(\mathbf{\Gamma} + \mathbf{\Gamma}^T), & \mathbf{\Gamma}_A &= \frac{1}{2}(\mathbf{\Gamma} - \mathbf{\Gamma}^T), \\ \mathbf{B} &= \mathbf{B}_S + \mathbf{B}_A, & \mathbf{B}_S &= \frac{1}{2}(\mathbf{B} + \mathbf{B}^T), & \mathbf{B}_A &= \frac{1}{2}(\mathbf{B} - \mathbf{B}^T), \end{aligned}$$

we represent W as scalar valued function of three symmetric and three skew tensors,

$$W = W(\mathbf{E}_S, \mathbf{E}_A, \mathbf{\Gamma}_S, \mathbf{\Gamma}_A; \mathbf{B}_S, \mathbf{B}_A). \quad (54)$$

Integrity basis for the proper orthogonal group is given by [Spencer](#), see Table 1 in [Spencer \(1965\)](#) or Table II in [Spencer \(1971\)](#). These tables are too lengthy to be presented here. Number of

members of integrity basis of $\mathbf{E}_S, \mathbf{E}_A, \mathbf{\Gamma}_S, \mathbf{\Gamma}_A, \mathbf{B}_S, \mathbf{B}_A$ is much larger than number of components of these tensors. However, there are some polynomial dependencies between elements of the integrity basis of three symmetric and three skew tensors. Hence the number of functionally independent invariants can be reduced, but it is still very large, see [Zheng \(1994\)](#).

For proper orthogonal group there is no difference in transformations of the axial and polar tensors. It is not the case if one considers transformations using the full orthogonal group. Since $\mathbf{\Gamma}_S, \mathbf{\Gamma}_A, \mathbf{B}_S, \mathbf{B}_A$ are axial tensors, not all invariants listed in [Spencer \(1965, 1971\)](#) are absolute invariants under orthogonal transformations, because some of them change sign under non proper orthogonal transformations. Following [Spencer \(1971\)](#), we call such invariants relative invariants. Examples of relative invariants are $\text{tr } \mathbf{\Gamma}_S, \text{tr } \mathbf{\Gamma}_S^3, \text{tr } \mathbf{E}_S \mathbf{\Gamma}_S, \text{tr } \mathbf{E}_S \mathbf{B}_S$, etc. This gives us the following property of W :

$$W(\mathbf{E}_S, \mathbf{E}_A, \mathbf{\Gamma}_S, \mathbf{\Gamma}_A; \mathbf{B}_S, \mathbf{B}_A) = W(\mathbf{E}_S, \mathbf{E}_A, \mathbf{\Gamma}_S, \mathbf{\Gamma}_A; \mathbf{B}_S, \mathbf{B}_A). \quad (55)$$

There are 119 invariants of non symmetric tensors $\mathbf{E}, \mathbf{\Gamma}, \mathbf{B}$ presented in Table A.1, see Appendix A. The strain energy density of polar elastic isotropic solids can be any scalar valued function of these invariants which satisfy (55).

If we neglect explicit dependence of W on \mathbf{B} , or assume that $\mathbf{B} = \mathbf{0}$, then $W = W(\mathbf{E}, \mathbf{\Gamma})$. Integrity basis of two non symmetric tensors under the orthogonal group contains 39 members. Following [Zheng \(1994\)](#), [Ramezani et al. \(2009\)](#) listed these invariants for the non linear polar elastic solids and proposed corresponding constitutive equations. Let us note, however, that not all 39 elements of this integrity basis are functionally independent. [Kafadar and Eringen \(1971\)](#) constructed the functional basis for two non symmetric tensors taking into account these functional dependencies. Table A.1 contains the invariants of [Ramezani et al. \(2009\)](#) and of [Kafadar and Eringen \(1971\)](#) as well as additional joint invariants of $\mathbf{E}, \mathbf{\Gamma}$, and \mathbf{B} . According to [Kafadar and Eringen \(1971\)](#), as the isotropic scalar valued function of two non symmetric tensors \mathbf{E} and $\mathbf{\Gamma}$, W is expressible in terms of 15 invariants,

$$W = W(I_1, I_2, \dots, I_{15}), \quad (56)$$

where I_k are given by

$$\begin{aligned} I_1 &= \text{tr } \mathbf{E}, & I_2 &= \text{tr } \mathbf{E}^2, & I_3 &= \text{tr } \mathbf{E}^3, \\ I_4 &= \text{tr } \mathbf{E}\mathbf{E}^T, & I_5 &= \text{tr } \mathbf{E}^2\mathbf{E}^T, & I_6 &= \text{tr } \mathbf{E}^2\mathbf{E}^T{}^2, \\ I_7 &= \text{tr } \mathbf{E}\mathbf{\Gamma}, & I_8 &= \text{tr } \mathbf{E}^2\mathbf{\Gamma}, & I_9 &= \text{tr } \mathbf{E}\mathbf{\Gamma}^2, \\ I_{10} &= \text{tr } \mathbf{\Gamma}, & I_{11} &= \text{tr } \mathbf{\Gamma}^2, & I_{12} &= \text{tr } \mathbf{\Gamma}^3, \\ I_{13} &= \text{tr } \mathbf{\Gamma}\mathbf{\Gamma}^T, & I_{14} &= \text{tr } \mathbf{\Gamma}^2\mathbf{\Gamma}^T, & I_{15} &= \text{tr } \mathbf{\Gamma}^2\mathbf{\Gamma}^T{}^2. \end{aligned}$$

Taking into account that $W = W(\mathbf{E}, \mathbf{\Gamma})$ is an even function with respect to $\mathbf{\Gamma}$, because in our case the group \mathcal{S}_x contains the reflection \mathbb{I} , W becomes also an even function with respect to some invariants,

$$\begin{aligned} W(I_1, I_2, I_3, I_4, I_5, I_6, I_7, I_8, I_9, I_{10}, I_{11}, I_{12}, I_{13}, I_{14}, I_{15}) \\ = W(I_1, I_2, I_3, I_4, I_5, I_6, I_7, I_8, I_9, I_{10}, I_{11}, I_{12}, I_{13}, I_{14}, I_{15}). \end{aligned} \quad (57)$$

Expanding W into the Taylor series relative to \mathbf{E} and $\mathbf{\Gamma}$, and keeping up to quadratic terms, we obtain the approximate polynomial representation of (57),

$$\begin{aligned} W = w_0 + a_1 I_1 + a_2 I_{10} + b_1 I_1^2 + b_2 I_1 I_{10} + b_3 I_{10}^2 + b_4 I_4 + b_5 I_2 + b_6 I_7 \\ + b_7 I_{11} + b_8 I_{13} + O(\max(\|\mathbf{E}\|^3, \|\mathbf{\Gamma}\|^3)), \end{aligned}$$

where $w_0, a_1, a_2, b_1, \dots, b_8$ are material constants. Applying to the latter expression the property (57), we finally obtain the following polynomial representation of W up to quadratic terms of \mathbf{E} and $\mathbf{\Gamma}$:

$$W = w_0 + a_1 I_1 + b_1 I_1^2 + b_3 I_{10}^2 + b_4 I_4 + b_5 I_2 + b_7 I_{11} + b_8 I_{13}. \quad (58)$$

We may also consider the representation of W as a sum of two scalar functions each depending on one strain measure,

$$W = W_1(\mathbf{E}) + W_2(\mathbf{\Gamma}). \quad (59)$$

The form (59) was used for example by Ramezani et al. (2009) in order to generalize classical neo Hookean and Mooney Rivlin models to polar elastic solids. As in Section 7, one can use here the representation theorem for isotropic scalar valued function of one non symmetric tensor given by Smith and Smith (1971). This leads to the following representation of W :

$$W = \widetilde{W}_1(I_1, \dots, I_6) + \widetilde{W}_2(I_{10}, \dots, I_{15}), \quad (60)$$

where \widetilde{W}_2 is such that

$$\widetilde{W}_2(I_{10}, I_{11}, I_{12}, I_{13}, I_{14}, I_{15}) = \widetilde{W}_2(I_{10}, I_{11}, I_{12}, I_{13}, I_{14}, I_{15}). \quad (61)$$

Expanding (60) with (61) into the Taylor series and keeping up to quadratic terms in \mathbf{E} and $\mathbf{\Gamma}$, W takes the form (59) with

$$W_1 = w_0 + a_1 I_1 + b_1 I_1^2 + b_4 I_4 + b_5 I_2, \quad W_2 = b_3 I_{10}^2 + b_7 I_{11} + b_8 I_{13}.$$

Definition 8 (*Hemitropic material*). The polar elastic solid is called *hemitropic* at x and \mathbf{B} if there exists a reference placement B_x , called undistorted, such that the material symmetry group relative to B_x takes the form

$$\mathcal{G}_x = \mathcal{S}_x^+ \equiv \{(\mathbf{P} \quad \mathbf{O}, \mathbf{O}, \mathbf{O}) : \mathbf{O} \in Orth^+\}. \quad (62)$$

The strain energy density of the hemitropic polar elastic solid satisfies the relation

$$W(\mathbf{E}, \mathbf{\Gamma}; \mathbf{B}) = W(\mathbf{OEO}^T, \mathbf{O}\mathbf{\Gamma}\mathbf{O}^T; \mathbf{OBO}^T), \quad \forall \mathbf{O} \in Orth^+. \quad (63)$$

The hemitropic polar elastic solid is insensitive to change of orientation of the space. In case of reduced strain energy density $W = W(\mathbf{E}, \mathbf{\Gamma})$ the representation of W is given by (56), but the property (57) does not hold, in general. Obviously, the polar elastic isotropic solid is also hemitropic.

Definitions (53) and (63) are somewhat similar to the corresponding definition of the isotropic polar elastic solid proposed by Eringen and Kafadar (1976), who required

$$W(\mathbf{E}, \mathbf{\Gamma}; \mathbf{J}_x) = W(\mathbf{OEO}^T, \mathbf{O}\mathbf{\Gamma}\mathbf{O}^T; \mathbf{OJ}_x\mathbf{O}^T), \quad \forall \mathbf{O} \in Orth. \quad (64)$$

However, properties (57) or (61) do not follow from (64) and additionally dependence of W upon \mathbf{J}_x is disregarded here.

Definition 9 (*Orthotropic material*). The polar elastic solid is called *orthotropic* at x and \mathbf{B} if the material symmetry group for some reference placement B_x takes the form

$$\mathcal{G}_x = \{(\mathbf{P} \quad \mathbf{O}, \mathbf{O}, \mathbf{O}) : \mathbf{O} \in \{\mathbf{I}, \mathbf{I}, 2\mathbf{e}_1 \otimes \mathbf{e}_1 - \mathbf{I}, 2\mathbf{e}_2 \otimes \mathbf{e}_2 - \mathbf{I}, 2\mathbf{e}_3 \otimes \mathbf{e}_3 - \mathbf{I}\}\}, \quad (65)$$

where \mathbf{O} are orthogonal tensors performing mirror reflections and rotations of 180° about three orthonormal vectors \mathbf{e}_k .

Obviously, the polar elastic isotropic solid is also orthotropic. Thus, all invariants in Table A.1 can be used for representation of the strain energy density of the polar elastic orthotropic solid. There are 60 additional absolute and relative invariants responsible for orthotropic properties of polar elastic solids, so that the total number of invariants becomes 179.

Definition 10 (*Cubic symmetry*). The polar elastic solid is called *cubic symmetric* at x and \mathbf{B} if the material symmetry group for some reference placement B_x takes the form

$$\mathcal{G}_x = \{(\mathbf{P} \quad \mathbf{O}, \mathbf{O}, \mathbf{O}) : \mathbf{O} \in \{\mathbf{I}, \mathbf{I}, \mathbf{e}_1 \otimes \mathbf{e}_1 \mp \mathbf{e}_1 \times \mathbf{I}, \mathbf{e}_2 \otimes \mathbf{e}_2 \mp \mathbf{e}_2 \times \mathbf{I}, \mathbf{e}_3 \otimes \mathbf{e}_3 \mp \mathbf{e}_3 \times \mathbf{I}\}\}, \quad (66)$$

where \mathbf{O} are orthogonal tensors performing mirror reflections and rotations of 90° about three orthonormal vectors \mathbf{e}_k .

Definition 11 (*Transversely isotropic material*). The polar elastic solid is called *transversely isotropic* at x and \mathbf{B} with respect to a direction described by \mathbf{e} if the material symmetry group for some reference placement B_x takes the form

$$\mathcal{G}_x = \{(\mathbf{P} \quad \mathbf{O}, \mathbf{O}, \mathbf{O}) : \mathbf{O} \in \{\mathbf{I}, \mathbf{I}, \mathbf{O}(\varphi\mathbf{e}), \forall \varphi\}\}, \quad (67)$$

where $\mathbf{O}(\varphi\mathbf{e}) = (\mathbf{I} - \mathbf{e} \otimes \mathbf{e})\cos\varphi + \mathbf{e} \otimes \mathbf{e} + \mathbf{e} \times \text{Isin}\varphi$ is the rotation tensor with the rotation angle φ about the unit vector \mathbf{e} .

There are 167 absolute and relative invariants for polar elastic transversely isotropic solids.

11. Physically linear polar-elastic solids

Let us consider the polar elastic strain energy density as a quadratic function of \mathbf{E} and $\mathbf{\Gamma}$,

$$W = \frac{1}{2} \mathbf{E} : \mathbb{C} : \mathbf{E} + \mathbf{E} : \mathbb{B} : \mathbf{\Gamma} + \frac{1}{2} \mathbf{\Gamma} : \mathbb{D} : \mathbf{\Gamma}, \quad (68)$$

where \mathbb{C} , \mathbb{B} , and \mathbb{D} are 4th order tensors of elastic moduli of the polar elastic solid. Components of tensors \mathbb{C} and \mathbb{D} have symmetry properties

$$C_{ijmn} = C_{mnij}, \quad D_{ijmn} = D_{mnij}.$$

With (68) corresponding referential stress measures \mathbf{S} and \mathbf{K} follow from (8) to be

$$\mathbf{S} = \mathbb{C} : \mathbf{E} + \mathbb{B} : \mathbf{\Gamma}, \quad \mathbf{K} = \mathbb{B} : \mathbf{E} + \mathbb{D} : \mathbf{\Gamma}.$$

The model based on (68) can be called *the physically linear polar elastic solid*. In this case tensors \mathbb{C} , \mathbb{B} , and \mathbb{D} coincide with the tangent stiffness tensors introduced by Ramezani et al. (2009). Since \mathbf{E} and $\mathbf{\Gamma}$ are non symmetric tensors, W contains $2 \cdot 45 + 81 = 171$ independent material constants, in general. In what follows we find restrictions for \mathbb{C} , \mathbb{B} , and \mathbb{D} corresponding to some particular material symmetry groups.

1. Mirror reflection. Let \mathcal{S}_x contain two elements, $\mathcal{S}_x = \{\mathbb{I}, \mathbb{I}\}$. Substituting \mathbb{I} into (51) we obtain

$$\frac{1}{2} \mathbf{E} : \mathbb{C} : \mathbf{E} + \mathbf{E} : \mathbb{B} : \mathbf{\Gamma} + \frac{1}{2} \mathbf{\Gamma} : \mathbb{D} : \mathbf{\Gamma} = \frac{1}{2} \mathbf{E} : \mathbb{C} : \mathbf{E} - \mathbf{E} : \mathbb{B} : \mathbf{\Gamma} + \frac{1}{2} \mathbf{\Gamma} : \mathbb{D} : \mathbf{\Gamma},$$

from which it follows that either $\mathbb{B} = \mathbb{O}$, where \mathbb{O} is the 4th order zeroth tensor, or as has been shown above (Case 1 in Section 5), W is an even function of \mathbf{E} and $\mathbf{\Gamma}$. In this case one has 90 independent scalar elastic moduli. In what follows we always assume that \mathcal{S}_x contains \mathbb{I} with only one exception for the hemitropic solid. Hence, W , \mathbf{S} , and \mathbf{K} take the reduced form

$$W = \frac{1}{2} \mathbf{E} : \mathbb{C} : \mathbf{E} + \frac{1}{2} \mathbf{\Gamma} : \mathbb{D} : \mathbf{\Gamma}, \quad \mathbf{S} = \mathbb{C} : \mathbf{E}, \quad \mathbf{K} = \mathbb{D} : \mathbf{\Gamma}. \quad (69)$$

In the tensor base comprised of directors \mathbf{h}_a the tensors \mathbb{C} and \mathbb{D} can be expressed as symmetric 9×9 matrices. For example, the matrix of \mathbb{C} is

$$[\mathbb{C}] \begin{pmatrix} C_{1111} & C_{1122} & C_{1133} & C_{1123} & C_{1131} & C_{1112} & C_{1132} & C_{1113} & C_{1121} \\ & C_{2222} & C_{2233} & C_{2223} & C_{2231} & C_{2212} & C_{2232} & C_{2213} & C_{2221} \\ & & C_{3333} & C_{3323} & C_{3331} & C_{3312} & C_{3332} & C_{3313} & C_{3321} \\ & & & C_{2323} & C_{2331} & C_{2312} & C_{2332} & C_{2313} & C_{2321} \\ & & & & C_{3131} & C_{3112} & C_{3132} & C_{3113} & C_{3121} \\ & & & & & C_{1212} & C_{1232} & C_{1213} & C_{1221} \\ & & & & & & C_{3232} & C_{3213} & C_{3221} \\ & & & & & & & C_{1313} & C_{1321} \\ & & & & & & & & C_{2121} \end{pmatrix},$$

Since $[\mathbb{D}]$ has the same structure as $[\mathbb{C}]$, in what follows we restrict ourselves to representation of $[\mathbb{C}]$.

2. Monoclinic symmetry. The corresponding material symmetry group \mathcal{S}_x consists of mirror reflection and rotations of 180° about unit vector \mathbf{e} : $\mathbf{O} = \{\mathbf{I}, \mathbf{I}, 2\mathbf{e} \otimes \mathbf{e} - \mathbf{I}\}$. Using \mathbf{e} as coordinate vector of the 3rd axis, we obtain that matrices $[\mathbb{C}]$ and $[\mathbb{D}]$ should have some zero elements, so that $[\mathbb{C}]$ takes the form

$$[\mathbb{C}] \begin{pmatrix} C_{1111} & C_{1122} & C_{1133} & 0 & 0 & C_{1112} & 0 & 0 & C_{1121} \\ & C_{2222} & C_{2233} & 0 & 0 & C_{2212} & 0 & 0 & C_{2221} \\ & & C_{3333} & 0 & 0 & C_{3312} & 0 & 0 & C_{3321} \\ & & & C_{2323} & C_{2331} & 0 & C_{2332} & C_{2313} & 0 \\ & & & & C_{3131} & 0 & C_{3132} & C_{3113} & 0 \\ & & & & & C_{1212} & 0 & 0 & C_{1221} \\ & & & & & & C_{3232} & C_{3213} & 0 \\ & & & & & & & C_{1313} & 0 \\ & & & & & & & & C_{2121} \end{pmatrix}.$$

The polar elastic physically linear monoclinic solid is described by 50 independent elastic moduli.

3. Orthotropic symmetry. In this case we have 30 independent elastic moduli. The corresponding matrix $[\mathbb{C}]$ is given by

$$[\mathbb{C}] \begin{pmatrix} C_{1111} & C_{1122} & C_{1133} & 0 & 0 & 0 & 0 & 0 & 0 \\ & C_{2222} & C_{2233} & 0 & 0 & 0 & 0 & 0 & 0 \\ & & C_{3333} & 0 & 0 & 0 & 0 & 0 & 0 \\ & & & C_{2323} & 0 & 0 & C_{2332} & 0 & 0 \\ & & & & C_{3131} & 0 & 0 & C_{3113} & 0 \\ & & & & & C_{1212} & 0 & 0 & C_{1221} \\ & & & & & & C_{3232} & 0 & 0 \\ & & & & & & & C_{1313} & 0 \\ & & & & & & & & C_{2121} \end{pmatrix}.$$

4. Rotations by 90° . Additionally to the previous case, let us require invariance of W under rotations of 90° about the unit vector \mathbf{e}_3 : $\mathbf{O} = \{\mathbf{e}_3 \otimes \mathbf{e}_3 \mp \mathbf{e}_3 \times \mathbf{I}\}$. As a result, the solid is described by 18 independent elastic moduli and $[\mathbb{C}]$ is given by

$$[\mathbb{C}] \begin{pmatrix} C_{1111} & C_{1122} & C_{1133} & 0 & 0 & 0 & 0 & 0 & 0 \\ & C_{1111} & C_{1133} & 0 & 0 & 0 & 0 & 0 & 0 \\ & & C_{3333} & 0 & 0 & 0 & 0 & 0 & 0 \\ & & & C_{2323} & 0 & 0 & C_{2332} & 0 & 0 \\ & & & & C_{3131} & 0 & 0 & C_{2332} & 0 \\ & & & & & C_{1212} & 0 & 0 & C_{1221} \\ & & & & & & C_{3131} & 0 & 0 \\ & & & & & & & C_{2323} & 0 \\ & & & & & & & & C_{1212} \end{pmatrix}.$$

5. Cubic symmetry. The material symmetry group \mathcal{S}_x related to cubic symmetry additionally contains rotations of 90° about all orthogonal axes with unit vectors \mathbf{e}_k , $k = 1, 2, 3$. Hence, the matrix $[\mathbb{C}]$ becomes

$$[\mathbb{C}] \begin{pmatrix} C_{1111} & C_{1122} & C_{1122} & 0 & 0 & 0 & 0 & 0 & 0 \\ & C_{1111} & C_{1122} & 0 & 0 & 0 & 0 & 0 & 0 \\ & & C_{1111} & 0 & 0 & 0 & 0 & 0 & 0 \\ & & & C_{1212} & 0 & 0 & C_{1221} & 0 & 0 \\ & & & & C_{1212} & 0 & 0 & C_{1221} & 0 \\ & & & & & C_{1212} & 0 & 0 & C_{1221} \\ & & & & & & C_{1212} & 0 & 0 \\ & & & & & & & C_{1212} & 0 \\ & & & & & & & & C_{1212} \end{pmatrix}.$$

For the physically linear polar elastic solid with cubic symmetry we have eight independent elastic moduli.

6. Isotropy. In this case one can use representation (56) with (57). Taking into account only quadratic terms in \mathbf{E} and $\mathbf{\Gamma}$ we find that

$$2W = \alpha_1 I_1^2 + \alpha_2 I_2 + \alpha_3 I_4 + \beta_1 I_{10}^2 + \beta_2 I_{11} + \beta_3 I_{13}, \quad (70)$$

where α_a, β_a , $a = 1, 2, 3$ are independent elastic moduli. The corresponding tensors \mathbb{C} and \mathbb{D} are

$$\mathbb{C} = \alpha_1 \mathbf{I} \otimes \mathbf{I} + \alpha_2 \mathbf{h}_a \otimes \mathbf{I} \otimes \mathbf{h}_a + \alpha_3 \mathbf{h}_a \otimes \mathbf{h}_b \otimes \mathbf{h}_a \otimes \mathbf{h}_b, \quad (71)$$

$$\mathbb{D} = \beta_1 \mathbf{I} \otimes \mathbf{I} + \beta_2 \mathbf{h}_a \otimes \mathbf{I} \otimes \mathbf{h}_a + \beta_3 \mathbf{h}_a \otimes \mathbf{h}_b \otimes \mathbf{h}_a \otimes \mathbf{h}_b, \quad (72)$$

where \mathbf{h}_a , $a = 1, 2, 3$ is now any Cartesian base.

The strain energy density of the physically linear polar elastic isotropic solid contains only six scalar elastic moduli. This constitutive relation corresponds to the linear isotropic elastic Cosserat continuum widely used in the literature, see for example Nowacki (1986), Eringen (1999). For example, in terms of material constants used by Eringen (1999), formulas 5.3.1, our constants α_a, β_a are expressed as follows: $\alpha_1 = \lambda$, $\alpha_2 = \mu$, $\alpha_3 = \mu + \kappa$, $\beta_1 = \alpha$, $\beta_2 = \beta$, $\beta_3 = \gamma$. Within linear micropolar elasticity the explicit structure of tensors \mathbb{C} and \mathbb{D} was presented by Zheng and Spencer (1993) for 14 symmetry groups.

7. Hemitropy. For the polar elastic hemitropic solid the material symmetry group contains only rotation tensors, see (62). Hence $\mathbb{B} \neq \mathbb{O}$, in general. Tensors \mathbb{C} and \mathbb{D} take here the form (71) while \mathbb{B} is given by

$$\mathbb{B} = \gamma_1 \mathbf{I} \otimes \mathbf{I} + \gamma_2 \mathbf{h}_a \otimes \mathbf{I} \otimes \mathbf{h}_a + \gamma_3 \mathbf{h}_a \otimes \mathbf{h}_b \otimes \mathbf{h}_a \otimes \mathbf{h}_b,$$

where γ_a , $a = 1, 2, 3$, are additional independent scalar elastic moduli. As a result, for the polar elastic physically linear hemitropic solid the strain energy density contains nine independent scalar elastic moduli. In linear micropolar elasticity this representation is given for example in Dyszlewicz (2004).

12. Physically linear polar-elastic solids which elasticity tensors depend on the microstructure curvature tensor

Let us now consider the polar elastic strain energy density as a quadratic function (68) of \mathbf{E} and $\mathbf{\Gamma}$,

$$W = \frac{1}{2} \mathbf{E} : \mathbb{C}(\mathbf{B}) : \mathbf{E} + \mathbf{E} : \mathbb{B}(\mathbf{B}) : \mathbf{\Gamma} + \frac{1}{2} \mathbf{\Gamma} : \mathbb{D}(\mathbf{B}) : \mathbf{\Gamma}. \quad (73)$$

But now \mathbb{C}, \mathbb{B} , and \mathbb{D} are 4th order tensors of elastic moduli assumed to depend on \mathbf{B} . The tensor \mathbf{B} plays here the role of structural tensor used in representation of anisotropic tensor functions, see for example Boehler (1987) and Zheng (1994). Dependence of elasticity

tensors on \mathbf{B} changes significantly representation of (73) and the number of independent elastic moduli.

In what follows, for simplicity, we restrict ourselves to polar elastic hemitropic and isotropic solids. Using Table A.1, we obtain the following 58 quadratic invariants of \mathbf{E} and $\mathbf{\Gamma}$:

$$\begin{aligned}
J_1 & \text{tr}^2 \mathbf{E}_S, & J_2 & \text{tr} \mathbf{E}_S^2, & J_3 & \text{tr} \mathbf{E}_A^2, \\
J_4 & \text{tr}^2 \mathbf{\Gamma}_S, & J_5 & \text{tr} \mathbf{\Gamma}_S^2, & J_6 & \text{tr} \mathbf{\Gamma}_A^2, \\
J_7 & \text{tr} \mathbf{E}_S \mathbf{\Gamma}_S, & & & & \\
J_8 & \text{tr}^2 \mathbf{E}_S \mathbf{B}_S, & J_9 & \text{tr} \mathbf{E}_S^2 \mathbf{B}_S, & & \\
J_{10} & \text{tr}^2 \mathbf{E}_S \mathbf{B}_S^2, & J_{11} & \text{tr} \mathbf{E}_S^2 \mathbf{B}_S^2, & & \\
J_{12} & \text{tr}^2 \mathbf{\Gamma}_S \mathbf{B}_S, & J_{13} & \text{tr} \mathbf{\Gamma}_S^2 \mathbf{B}_S, & & \\
J_{14} & \text{tr}^2 \mathbf{\Gamma}_S \mathbf{B}_S^2, & J_{15} & \text{tr} \mathbf{\Gamma}_S^2 \mathbf{B}_S^2, & J_{16} & \text{tr} \mathbf{E}_S \mathbf{\Gamma}_S \mathbf{B}_S, \\
J_{17} & \text{tr} \mathbf{E}_A \mathbf{\Gamma}_A, & J_{18} & \text{tr}^2 \mathbf{E}_A \mathbf{B}_A, & & \\
J_{19} & \text{tr}^2 \mathbf{\Gamma}_A \mathbf{B}_A, & J_{20} & \text{tr} \mathbf{E}_A \mathbf{\Gamma}_A \mathbf{B}_A, & & \\
J_{21} & \text{tr}^2 \mathbf{E}_S \mathbf{B}_A^2, & J_{22} & \text{tr} \mathbf{E}_S^2 \mathbf{B}_A^2, & & \\
J_{23} & \text{tr}^2 \mathbf{\Gamma}_S \mathbf{B}_A^2, & J_{24} & \text{tr} \mathbf{\Gamma}_S^2 \mathbf{B}_A^2, & & \\
J_{25} & \text{tr} \mathbf{B}_S \mathbf{\Gamma}_A^2, & J_{26} & \text{tr} \mathbf{B}_S^2 \mathbf{\Gamma}_A^2, & & \\
J_{27} & \text{tr} \mathbf{B}_S \mathbf{E}_A^2, & J_{28} & \text{tr} \mathbf{B}_S^2 \mathbf{E}_A^2, & & \\
J_{29} & \text{tr} \mathbf{E}_S \mathbf{\Gamma}_S \mathbf{B}_A, & J_{30} & \text{tr} \mathbf{E}_S \mathbf{B}_S \mathbf{E}_A, & & \\
J_{31} & \text{tr} \mathbf{E}_S \mathbf{B}_S \mathbf{\Gamma}_A, & J_{32} & \text{tr} \mathbf{E}_S \mathbf{B}_S^2 \mathbf{\Gamma}_A, & & \\
J_{33} & \text{tr}^2 \mathbf{E}_S \mathbf{B}_S \mathbf{B}_A, & J_{34} & \text{tr} \mathbf{E}_S^2 \mathbf{B}_S \mathbf{B}_A, & & \\
J_{35} & \text{tr}^2 \mathbf{E}_S \mathbf{B}_S^2 \mathbf{B}_A, & J_{36} & \text{tr}^2 \mathbf{E}_S \mathbf{B}_A^2 \mathbf{B}_S \mathbf{B}_A, & & \\
J_{37} & \text{tr} \mathbf{\Gamma}_S \mathbf{B}_S \mathbf{E}_A, & J_{38} & \text{tr} \mathbf{\Gamma}_S \mathbf{B}_S \mathbf{\Gamma}_A, & J_{39} & \text{tr} \mathbf{\Gamma}_S \mathbf{B}_S^2 \mathbf{\Gamma}_A, \\
J_{40} & \text{tr}^2 \mathbf{\Gamma}_S \mathbf{B}_S \mathbf{B}_A, & J_{41} & \text{tr} \mathbf{\Gamma}_S^2 \mathbf{B}_S \mathbf{B}_A, & & \\
J_{42} & \text{tr}^2 \mathbf{\Gamma}_S \mathbf{B}_S^2 \mathbf{B}_A, & J_{43} & \text{tr}^2 \mathbf{\Gamma}_S \mathbf{B}_A^2 \mathbf{B}_S \mathbf{B}_A, & & \\
J_{44} & \text{tr} \mathbf{E}_S \mathbf{E}_A \mathbf{B}_A, & J_{45} & \text{tr} \mathbf{E}_S \mathbf{E}_A \mathbf{B}_A^2, & & \\
J_{46} & \text{tr} \mathbf{E}_S \mathbf{\Gamma}_A \mathbf{B}_A, & J_{47} & \text{tr} \mathbf{E}_S \mathbf{\Gamma}_A \mathbf{B}_A^2, & & \\
J_{48} & \text{tr} \mathbf{\Gamma}_S \mathbf{E}_A \mathbf{B}_A, & J_{49} & \text{tr} \mathbf{\Gamma}_S \mathbf{E}_A \mathbf{B}_A^2, & & \\
J_{50} & \text{tr} \mathbf{\Gamma}_S \mathbf{\Gamma}_A \mathbf{B}_A, & J_{51} & \text{tr} \mathbf{\Gamma}_S \mathbf{\Gamma}_A \mathbf{B}_A^2, & J_{52} & \text{tr} \mathbf{B}_S \mathbf{E}_A \mathbf{\Gamma}_A, \\
J_{53} & \text{tr}^2 \mathbf{B}_S \mathbf{E}_A \mathbf{B}_A, & J_{54} & \text{tr} \mathbf{B}_S \mathbf{E}_A^2 \mathbf{B}_A, & J_{55} & \text{tr}^2 \mathbf{B}_S \mathbf{E}_A \mathbf{B}_A^2, \\
J_{56} & \text{tr}^2 \mathbf{B}_S \mathbf{\Gamma}_A \mathbf{B}_A, & J_{57} & \text{tr} \mathbf{B}_S \mathbf{\Gamma}_A^2 \mathbf{B}_A, & J_{58} & \text{tr}^2 \mathbf{B}_S \mathbf{\Gamma}_A \mathbf{B}_A^2.
\end{aligned} \tag{74}$$

The strain energy density of linear polar elastic hemitropic solid is now the sum of 58 terms,

$$W = \sum_{i=1}^{58} c_i J_i, \tag{75}$$

where c_i are functions of invariants of \mathbf{B} , in general.

Since underlined terms in (74) are relative invariants, which change signs under change of orientation of the space, for the physically linear polar elastic isotropic solid the strain energy density takes the form (75), but now with $c_7 = c_9 = c_{13} = c_{25} = c_{27} = c_{30} = c_{32} = c_{44} = c_{47} = 0$.

One can derive in the same manner quadratic strain energy densities of physically linear polar elastic orthotropic and transversely isotropic solids. Let us however note that when one takes into account explicit dependence of elasticity tensors on \mathbf{B} one obtains a huge number of material parameters even when one assumes that c_i are constant elastic moduli.

13. Conclusions

We have defined anew the material symmetry group \mathcal{G}_\times of the non linear polar continuum. The group generalizes the one proposed by Eringen and Kafadar (1976) by introducing the undeformed microstructure curvature tensor \mathbf{B} instead of the micr

Table A.1
Invariants in W .

Agencies	Invariants				
\mathbf{E}_S	$\text{tr} \mathbf{E}_S$	$\text{tr} \mathbf{E}_S^2$	$\text{tr} \mathbf{E}_S^3$		
\mathbf{E}_A	$\text{tr} \mathbf{E}_A^2$				
$\mathbf{E}_S, \mathbf{E}_A$	$\text{tr} \mathbf{E}_S \mathbf{E}_A^2$	$\text{tr} \mathbf{E}_S^2 \mathbf{E}_A^2$	$\text{tr} \mathbf{E}_S^2 \mathbf{E}_A^2 \mathbf{E}_S \mathbf{E}_A$		
$\mathbf{\Gamma}_S$	$\text{tr} \mathbf{\Gamma}_S$	$\text{tr} \mathbf{\Gamma}_S^2$	$\text{tr} \mathbf{\Gamma}_S^3$		
$\mathbf{\Gamma}_A$	$\text{tr} \mathbf{\Gamma}_A^2$				
$\mathbf{\Gamma}_S, \mathbf{\Gamma}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{\Gamma}_A^2$	$\text{tr} \mathbf{\Gamma}_S^2 \mathbf{\Gamma}_A^2$	$\text{tr} \mathbf{\Gamma}_S^2 \mathbf{\Gamma}_A^2 \mathbf{\Gamma}_S \mathbf{\Gamma}_A$		
\mathbf{B}_S	$\text{tr} \mathbf{B}_S$	$\text{tr} \mathbf{B}_S^2$	$\text{tr} \mathbf{B}_S^3$		
\mathbf{B}_A	$\text{tr} \mathbf{B}_A^2$				
$\mathbf{B}_S, \mathbf{B}_A$	$\text{tr} \mathbf{B}_S \mathbf{B}_A^2$	$\text{tr} \mathbf{B}_S^2 \mathbf{B}_A^2$	$\text{tr} \mathbf{B}_S^2 \mathbf{B}_A^2 \mathbf{B}_S \mathbf{B}_A$		
$\mathbf{E}_S, \mathbf{\Gamma}_S$	$\text{tr} \mathbf{E}_S \mathbf{\Gamma}_S$	$\text{tr} \mathbf{E}_S^2 \mathbf{\Gamma}_S$	$\text{tr} \mathbf{E}_S \mathbf{\Gamma}_S^2$	$\text{tr} \mathbf{E}_S^2 \mathbf{\Gamma}_S^2$	
$\mathbf{E}_S, \mathbf{B}_S$	$\text{tr} \mathbf{E}_S \mathbf{B}_S$	$\text{tr} \mathbf{E}_S^2 \mathbf{B}_S$	$\text{tr} \mathbf{E}_S \mathbf{B}_S^2$	$\text{tr} \mathbf{E}_S^2 \mathbf{B}_S^2$	
$\mathbf{\Gamma}_S, \mathbf{B}_S$	$\text{tr} \mathbf{\Gamma}_S \mathbf{B}_S$	$\text{tr} \mathbf{\Gamma}_S^2 \mathbf{B}_S$	$\text{tr} \mathbf{\Gamma}_S \mathbf{B}_S^2$	$\text{tr} \mathbf{\Gamma}_S^2 \mathbf{B}_S^2$	
$\mathbf{E}_S, \mathbf{\Gamma}_S, \mathbf{B}_S$	$\text{tr} \mathbf{E}_S \mathbf{\Gamma}_S \mathbf{B}_S$				
$\mathbf{E}_A, \mathbf{\Gamma}_A$	$\text{tr} \mathbf{E}_A \mathbf{\Gamma}_A$				
$\mathbf{E}_A, \mathbf{B}_A$	$\text{tr} \mathbf{E}_A \mathbf{B}_A$				
$\mathbf{\Gamma}_A, \mathbf{B}_A$	$\text{tr} \mathbf{\Gamma}_A \mathbf{B}_A$				
$\mathbf{E}_A, \mathbf{\Gamma}_A, \mathbf{B}_A$	$\text{tr} \mathbf{E}_A \mathbf{\Gamma}_A \mathbf{B}_A$				
$\mathbf{E}_S, \mathbf{\Gamma}_A$	$\text{tr} \mathbf{E}_S \mathbf{\Gamma}_A^2$	$\text{tr} \mathbf{E}_S^2 \mathbf{\Gamma}_A^2$	$\text{tr} \mathbf{E}_S^2 \mathbf{\Gamma}_A^2 \mathbf{E}_S \mathbf{\Gamma}_A$		
$\mathbf{E}_S, \mathbf{B}_A$	$\text{tr} \mathbf{E}_S \mathbf{B}_A^2$	$\text{tr} \mathbf{E}_S^2 \mathbf{B}_A^2$	$\text{tr} \mathbf{E}_S^2 \mathbf{B}_A^2 \mathbf{E}_S \mathbf{B}_A$		
$\mathbf{\Gamma}_S, \mathbf{E}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{E}_A^2$	$\text{tr} \mathbf{\Gamma}_S^2 \mathbf{E}_A^2$	$\text{tr} \mathbf{\Gamma}_S^2 \mathbf{E}_A^2 \mathbf{\Gamma}_S \mathbf{E}_A$		
$\mathbf{\Gamma}_S, \mathbf{B}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{B}_A^2$	$\text{tr} \mathbf{\Gamma}_S^2 \mathbf{B}_A^2$	$\text{tr} \mathbf{\Gamma}_S^2 \mathbf{B}_A^2 \mathbf{\Gamma}_S \mathbf{B}_A$		
$\mathbf{B}_S, \mathbf{\Gamma}_A$	$\text{tr} \mathbf{B}_S \mathbf{\Gamma}_A^2$	$\text{tr} \mathbf{B}_S^2 \mathbf{\Gamma}_A^2$	$\text{tr} \mathbf{E}_S^2 \mathbf{\Gamma}_A^2 \mathbf{B}_S \mathbf{\Gamma}_A$		
$\mathbf{B}_S, \mathbf{E}_A$	$\text{tr} \mathbf{B}_S \mathbf{E}_A^2$	$\text{tr} \mathbf{B}_S^2 \mathbf{E}_A^2$	$\text{tr} \mathbf{B}_S^2 \mathbf{E}_A^2 \mathbf{B}_S \mathbf{E}_A$		
$\mathbf{E}_S, \mathbf{\Gamma}_S, \mathbf{E}_A$	$\text{tr} \mathbf{E}_S \mathbf{\Gamma}_S \mathbf{E}_A$	$\text{tr} \mathbf{E}_S^2 \mathbf{\Gamma}_S \mathbf{E}_A$	$\text{tr} \mathbf{E}_S \mathbf{\Gamma}_S^2 \mathbf{E}_A$	$\text{tr} \mathbf{E}_S \mathbf{E}_A^2 \mathbf{\Gamma}_S \mathbf{E}_A$	
$\mathbf{E}_S, \mathbf{\Gamma}_S, \mathbf{\Gamma}_A$	$\text{tr} \mathbf{E}_S \mathbf{\Gamma}_S \mathbf{\Gamma}_A$	$\text{tr} \mathbf{E}_S^2 \mathbf{\Gamma}_S \mathbf{\Gamma}_A$	$\text{tr} \mathbf{E}_S \mathbf{\Gamma}_S^2 \mathbf{\Gamma}_A$	$\text{tr} \mathbf{E}_S \mathbf{\Gamma}_A^2 \mathbf{\Gamma}_S \mathbf{\Gamma}_A$	
$\mathbf{E}_S, \mathbf{\Gamma}_S, \mathbf{B}_A$	$\text{tr} \mathbf{E}_S \mathbf{\Gamma}_S \mathbf{B}_A$	$\text{tr} \mathbf{E}_S^2 \mathbf{\Gamma}_S \mathbf{B}_A$	$\text{tr} \mathbf{E}_S \mathbf{\Gamma}_S^2 \mathbf{B}_A$	$\text{tr} \mathbf{E}_S \mathbf{B}_A^2 \mathbf{\Gamma}_S \mathbf{B}_A$	
$\mathbf{E}_S, \mathbf{B}_S, \mathbf{E}_A$	$\text{tr} \mathbf{E}_S \mathbf{B}_S \mathbf{E}_A$	$\text{tr} \mathbf{E}_S^2 \mathbf{B}_S \mathbf{E}_A$	$\text{tr} \mathbf{E}_S \mathbf{B}_S^2 \mathbf{E}_A$	$\text{tr} \mathbf{E}_S \mathbf{E}_A^2 \mathbf{B}_S \mathbf{E}_A$	
$\mathbf{E}_S, \mathbf{B}_S, \mathbf{\Gamma}_A$	$\text{tr} \mathbf{E}_S \mathbf{B}_S \mathbf{\Gamma}_A$	$\text{tr} \mathbf{E}_S^2 \mathbf{B}_S \mathbf{\Gamma}_A$	$\text{tr} \mathbf{E}_S \mathbf{B}_S^2 \mathbf{\Gamma}_A$	$\text{tr} \mathbf{E}_S \mathbf{\Gamma}_A^2 \mathbf{B}_S \mathbf{\Gamma}_A$	
$\mathbf{E}_S, \mathbf{B}_S, \mathbf{B}_A$	$\text{tr} \mathbf{E}_S \mathbf{B}_S \mathbf{B}_A$	$\text{tr} \mathbf{E}_S^2 \mathbf{B}_S \mathbf{B}_A$	$\text{tr} \mathbf{E}_S \mathbf{B}_S^2 \mathbf{B}_A$	$\text{tr} \mathbf{E}_S \mathbf{B}_A^2 \mathbf{B}_S \mathbf{B}_A$	
$\mathbf{\Gamma}_S, \mathbf{B}_S, \mathbf{E}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{B}_S \mathbf{E}_A$	$\text{tr} \mathbf{\Gamma}_S^2 \mathbf{B}_S \mathbf{E}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{B}_S^2 \mathbf{E}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{E}_A^2 \mathbf{B}_S \mathbf{E}_A$	
$\mathbf{\Gamma}_S, \mathbf{B}_S, \mathbf{\Gamma}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{B}_S \mathbf{\Gamma}_A$	$\text{tr} \mathbf{\Gamma}_S^2 \mathbf{B}_S \mathbf{\Gamma}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{B}_S^2 \mathbf{\Gamma}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{\Gamma}_A^2 \mathbf{B}_S \mathbf{\Gamma}_A$	
$\mathbf{\Gamma}_S, \mathbf{B}_S, \mathbf{B}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{B}_S \mathbf{B}_A$	$\text{tr} \mathbf{\Gamma}_S^2 \mathbf{B}_S \mathbf{B}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{B}_S^2 \mathbf{B}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{B}_A^2 \mathbf{B}_S \mathbf{B}_A$	
$\mathbf{E}_S, \mathbf{E}_A, \mathbf{\Gamma}_A$	$\text{tr} \mathbf{E}_S \mathbf{E}_A \mathbf{\Gamma}_A$	$\text{tr} \mathbf{E}_S \mathbf{E}_A^2 \mathbf{\Gamma}_A$	$\text{tr} \mathbf{E}_S \mathbf{E}_A \mathbf{\Gamma}_A^2$		
$\mathbf{E}_S, \mathbf{E}_A, \mathbf{B}_A$	$\text{tr} \mathbf{E}_S \mathbf{E}_A \mathbf{B}_A$	$\text{tr} \mathbf{E}_S \mathbf{E}_A^2 \mathbf{B}_A$	$\text{tr} \mathbf{E}_S \mathbf{E}_A \mathbf{B}_A^2$		
$\mathbf{E}_S, \mathbf{\Gamma}_A, \mathbf{B}_A$	$\text{tr} \mathbf{E}_S \mathbf{\Gamma}_A \mathbf{B}_A$	$\text{tr} \mathbf{E}_S \mathbf{\Gamma}_A^2 \mathbf{B}_A$	$\text{tr} \mathbf{E}_S \mathbf{\Gamma}_A \mathbf{B}_A^2$		
$\mathbf{\Gamma}_S, \mathbf{E}_A, \mathbf{\Gamma}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{E}_A \mathbf{\Gamma}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{E}_A^2 \mathbf{\Gamma}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{E}_A \mathbf{\Gamma}_A^2$		
$\mathbf{\Gamma}_S, \mathbf{E}_A, \mathbf{B}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{E}_A \mathbf{B}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{E}_A^2 \mathbf{B}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{E}_A \mathbf{B}_A^2$		
$\mathbf{\Gamma}_S, \mathbf{\Gamma}_A, \mathbf{B}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{\Gamma}_A \mathbf{B}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{\Gamma}_A^2 \mathbf{B}_A$	$\text{tr} \mathbf{\Gamma}_S \mathbf{\Gamma}_A \mathbf{B}_A^2$		
$\mathbf{B}_S, \mathbf{E}_A, \mathbf{\Gamma}_A$	$\text{tr} \mathbf{B}_S \mathbf{E}_A \mathbf{\Gamma}_A$	$\text{tr} \mathbf{B}_S \mathbf{E}_A^2 \mathbf{\Gamma}_A$	$\text{tr} \mathbf{B}_S \mathbf{E}_A \mathbf{\Gamma}_A^2$		
$\mathbf{B}_S, \mathbf{E}_A, \mathbf{B}_A$	$\text{tr} \mathbf{B}_S \mathbf{E}_A \mathbf{B}_A$	$\text{tr} \mathbf{B}_S \mathbf{E}_A^2 \mathbf{B}_A$	$\text{tr} \mathbf{B}_S \mathbf{E}_A \mathbf{B}_A^2$		
$\mathbf{B}_S, \mathbf{\Gamma}_A, \mathbf{B}_A$	$\text{tr} \mathbf{B}_S \mathbf{\Gamma}_A \mathbf{B}_A$	$\text{tr} \mathbf{B}_S \mathbf{\Gamma}_A^2 \mathbf{B}_A$	$\text{tr} \mathbf{B}_S \mathbf{\Gamma}_A \mathbf{B}_A^2$		

oinertia tensor \mathbf{J}_\times as well as by taking into account that \mathbf{B} and the wryness tensor $\mathbf{\Gamma}$ are axial tensors which change signs under inversion transformation of 3D space. Our group \mathcal{G}_\times consists of an ordered triple of tensors which make the strain energy density invariant under change of reference placement. In terms of members of \mathcal{G}_\times polar elastic fluids, solids, liquid crystals, and subfluids

have conveniently been defined. Reduced forms of constitutive equations for non-linear and physically linear polar-elastic solids are given for several particular material symmetry groups.

In many cases discussed here even the reduced constitutive equations of polar-elastic solids still involve a large number of material constants, which should be experimentally or theoretically identified. Difficulties in identifications of material constants pose a serious challenge for wider application of polar-elastic solids in science and technology.

Acknowledgements

Victor A. Eremeyev was supported by the RFBR with the Grant No. 12-01-00038 and by the DFG with the Grant No. AL 341/33-1, while Wojciech Pietraszkiewicz by the Polish Ministry of Science and Higher Education with the Grant No. N506 254 237.

Appendix A. Invariants in strain energy density

Using the representations of Zheng (1994), for isotropic polar-elastic solids in Table A.1 we present the list of 119 irreducible invariants of \mathbf{E} , $\mathbf{\Gamma}$, and \mathbf{B} expressed in terms of their symmetric and skew-symmetric parts.

References

- Aero, E.L., Bulygin, A.N., Kuvshinskii, E.V., 1965. Asymmetric hydromechanics. *Journal of Applied Mathematics and Mechanics* 29 (2), 333–346.
- Allen, S.J., DeSilva, C.N., Kline, K.A., 1967. A theory of simple deformable directed fluids. *Physics of Fluids* 10 (12), 2551–2555.
- Arfken, G.B., Weber, H.J., 2000. *Mathematical Methods for Physicists*. Springer, Berlin.
- Billington, E.W., 1986. *Introduction to the Mechanics and Physics of Solids*. Adam Hilger Ltd., Bristol.
- Boehler, J.P. (Ed.), 1987. *Applications of Tensor Functions in Solid Mechanics*. Springer, Wien.
- Cohen, H., Wang, C.-C., 1983. Principal waves in elastic subfluids. *Archive for Rational Mechanics and Analysis* 83 (2), 139–168.
- Coleman, B.D., 1965. Simple liquid crystals. *Archive for Rational Mechanics and Analysis* 20 (1), 41–58.
- Cosserat, E., Cosserat, F., 1909. *Théorie des corps déformables*. Herman et Fils, Paris.
- Dai, T.-M., 2003. Renewal of basic laws and principles for polar continuum theories (I) – micropolar continua. *Applied Mathematics and Mechanics* 24, 1119–1125.
- de Gennes, P.G., Prost, J., 1993. *The Physics of Liquid Crystals*, second ed. Clarendon Press, Oxford.
- Dyzlewicz, J., 2004. *Micropolar Theory of Elasticity*. Springer, Berlin.
- Elzanowski, M., Epstein, M., 1992. The symmetry group of second-grade materials. *International Journal of Non-Linear Mechanics* 27 (4), 635–638.
- Epstein, M., de Leon, M., 1996. Homogeneity conditions for generalized Cosserat media. *Journal of Elasticity* 43 (3), 189–201.
- Epstein, M., de Leon, M., 1998. Geometrical theory of uniform Cosserat media. *Journal of Geometry and Physics* 26 (1–2), 127–170.
- Eremeyev, V.A., Pietraszkiewicz, W., 2006. Local symmetry group in the general theory of elastic shells. *Journal of Elasticity* 85 (2), 125–152.
- Erickson, J.L., 1998. *Introduction to the Thermodynamics of Solids*. Springer, New York.
- Eringen, A.C., 1966. Theory of micropolar fluids. *Journal of Mathematics and Mechanics* 16 (1), 1–18.
- Eringen, A.C., 1997. A unified continuum theory of liquid crystals. *ARI*, 369–374.
- Eringen, A.C., 1999. *Microcontinuum Field Theory. I. Foundations and Solids*. Springer, New York.
- Eringen, A.C., 2001. *Microcontinuum Field Theory II. Fluent Media*. Springer, New York.
- Eringen, A.C., Kafadar, C.B., 1976. Polar field theories. In: Eringen, A.C. (Ed.), *Continuum Physics, Vol. IV*. Academic Press, New York, pp. 1–75.
- Feynman, R.P., Leighton, R.B., Sands, M., 1977. *The Feynman Lectures on Physics*, 6th ed., vol. 1. Addison-Wesley, Reading, Massachusetts.
- Grekova, E., Zhilin, P., 2001. Basic equations of Kelvins medium and analogy with ferromagnets. *Journal of Elasticity* 64, 29–70.
- Grekova, E.F., Kulesh, M.A., Herrman, G.C., 2009. Waves in linear elastic media with microrotations, Part 2: isotropic reduced cosserat model. *Bulletin of the Seismological Society of America* 99 (2B), 1423–1428.
- Huang, C.L., Smith, G.F., 1967. The energy function for isotropic materials with couple stresses. *ZAMP* 18 (6), 905–909.
- Kafadar, C.B., Eringen, A.C., 1971. Micropolar media – I. The classical theory. *International Journal of Engineering Science* 9, 271–305.
- Korn, G.A., Korn, T.M., 1968. *Mathematical Handbook for Scientists and Engineers: Definitions, Theorems and Formulas for Reference and Review*, second ed. McGraw-Hill, New York.
- Lauderdale, T.A., O'Reilly, O.M., 2007. On the restrictions imposed by non-affine material symmetry groups for elastic rods: application to helical substructures. *European Journal of Mechanics A. Solids* 26 (4), 701–711.
- Luo, C., O'Reilly, O.M., 2000. On the material symmetry of elastic rods. *Journal of Elasticity* 60 (1), 35–56.
- Lurie, A.I., 1990. *Non-linear Theory of Elasticity*. North-Holland, Amsterdam.
- Maugin, G.A., 1974. Acceleration waves in simple and linear viscoelastic micropolar materials. *International Journal Engineering Science* 12, 143–157.
- Maugin, G.A., 1998. On the structure of the theory of polar elasticity. *Philosophical Transactions of Royal Society London A* 356, 1367–1395.
- Murdoch, A.L., Cohen, H., 1979. Symmetry considerations for material surfaces. *Archive for Rational Mechanics and Analysis* 72 (1), 61–98.
- Nikitin, E., Zubov, L.M., 1998. Conservation laws and conjugate solutions in the elasticity of simple materials and materials with couple stress. *Journal of Elasticity* 51, 1–22.
- Noll, W., 1958. A mathematical theory of the mechanical behavior of continuous media. *Archive for Rational Mechanics and Analysis* 2 (1), 197–226.
- Nowacki, W., 1986. *Theory of Asymmetric Elasticity*. Pergamon-Press, Oxford.
- Nye, J.F., 1957. *Physical Properties of Crystals. Their Representation by Tensors and Matrices*. Clarendon Press, Oxford.
- Ogden, R.W., 1984. *Non-linear Elastic Deformations*. Ellis Horwood, Chichester.
- Pietraszkiewicz, W., Eremeyev, V.A., 2009. On natural strain measures of the non-linear micropolar continuum. *International Journal of Solids and Structures* 46 (3–4), 774–787.
- Ramezani, S., Naghdabadi, R., 2007. Energy pairs in the micropolar continuum. *International Journal of Solids and Structures* 44, 4810–4818.
- Ramezani, S., Naghdabadi, R., Sohrabpour, S., 2009. Constitutive equations for micropolar hyper-elastic materials. *International Journal of Solids and Structures* 46 (14–15), 2765–2773.
- Rivlin, R.S., 1980. Material symmetry and constitutive equations. *Ingenieur-Archiv* 49 (5–6), 325–336.
- Scarpetta, E., 1989. Some results on well-posedness for a nonlinear theory of micropolar elastostatics. *Quarterly Journal of Mechanics and Applied Mathematics* 42, 1–11.
- Smith, G.F., 1994. *Constitutive Equations for Anisotropic and Isotropic Materials*. North-Holland, Elsevier, Amsterdam, New York.
- Smith, M.M., Smith, R.F., 1971. Irreducible expressions for isotropic functions of two tensors. *International Journal of Engineering Science* 19 (6), 811–817.
- Smith, R.F., 1965. Material symmetry and constitutive equations. *Archive for Rational Mechanics and Analysis* 18 (5–6), 325–336.
- Spencer, A.J.M., 1965. Isotropic integrity bases for vectors and second-order tensors. Part II. *Archive for Rational Mechanics and Analysis* 18 (1), 51–82.
- Spencer, A.J.M., 1971. Theory of invariants. In: Eringen, A.C. (Ed.), *Continuum Physics, vol. 1*. Academic Press, New-York, pp. 239–353.
- Spencer, A.J.M., Rivlin, R.S., 1962. Isotropic integrity bases for vector second-order tensors. Part 1. *Archive for Rational Mechanics and Analysis* 9 (1), 45–63.
- Truesdell, C., 1964. A theorem on the isotropy groups of a hyperelastic material. *Proceedings of the National Academy of Sciences of the United States of America* 52 (4), 1081–1083.
- Truesdell, C., 1966. *The Elements of Continuum Mechanics*. Springer, New York.
- Truesdell, C., Noll, W., 1965. The nonlinear field theories of mechanics. In: Flügge, S. (Ed.), *Handbuch der Physik, vol. III/3*. Springer, Berlin, pp. 1–602.
- Truesdell, C., Toupin, R., 1960. The classical field theories. In: Flügge, S. (Ed.), *Handbuch der Physik, vol. III/1*. Springer, Berlin, pp. 226–793.
- Truesdell, C.A., 1991. *A first course in rational continuum mechanics, . General Concepts*, 2nd ed., vol. 1. Academic Press, Boston.
- Wang, C.-C., 1965. A general theory of subfluids. *Archive for Rational Mechanics and Analysis* 20 (1), 1–40.
- Wang, C.-C., 1973. On the response functions of isotropic elastic shells. *International Journal of Solids and Structures* 50 (2), 81–98.
- Wang, C.C., Truesdell, C., 1973. *Introduction to Rational Elasticity*. Noordhoff International Publishing, Leyden.
- Yeremeyev, V.A., Zubov, L.M., 1999. The theory of elastic and viscoelastic micropolar liquids. *Journal of Applied Mathematics and Mechanics* 63, 755–767.
- Zheng, Q.S., 1994. Theory of representations for tensor functions – A unified invariant approach to constitutive equations. *Applied Mechanics Reviews* 47 (11), 545–587.
- Zheng, Q.S., Spencer, A.J.M., 1993. On the canonical representations for Kronecker powers of orthogonal tensors with application to material symmetry problems. *International Journal of Engineering Science* 31 (4), 617–635.
- Zubov, L.M., 1997. *Nonlinear Theory of Dislocations and Disclinations in Elastic Bodies*. Springer, Berlin.
- Zubov, L.M., Eremeev, V.A., 1996. Equations for a viscoelastic micropolar fluid. *Doklady Physics* 41 (12), 598–601.