



HAL
open science

Three-dimensional Anosov flag manifolds

Thierry Barbot

► **To cite this version:**

Thierry Barbot. Three-dimensional Anosov flag manifolds. *Geometry and Topology*, 2010, 14 (1), pp.153–191. hal-00004967v3

HAL Id: hal-00004967

<https://hal.science/hal-00004967v3>

Submitted on 27 May 2005

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THREE-DIMENSIONAL ANOSOV FLAG MANIFOLDS

THIERRY BARBOT

ABSTRACT. Let Γ be a surface group of higher genus. Let $\rho_0 : \Gamma \rightarrow \mathrm{PGL}(V)$ be a discrete faithful representation with image contained in the natural embedding of $\mathrm{SL}(2, \mathbb{R})$ in $\mathrm{PGL}(3, \mathbb{R})$ as a group preserving a point and a disjoint projective line in the projective plane. We prove that such a representation is (G, Y) -Anosov (following the terminology of [16]), where Y is the frame bundle. More generally, we prove that all the deformations $\rho : \Gamma \rightarrow \mathrm{PGL}(3, \mathbb{R})$ studied in [1] are (G, Y) -Anosov. As a corollary, we obtain all the main results of [1], and extend them to any small deformation of ρ_0 , not necessarily preserving a point or a projective line in the projective space: in particular, there is a $\rho(\Gamma)$ -invariant solid torus Ω in the flag variety. The quotient space $\rho(\Gamma) \backslash \Omega$ is a flag manifold, naturally equipped with two 1-dimensional transversely projective foliations arising from the projections of the flag variety on the projective plane and its dual; if ρ is strongly irreducible, these foliations are not minimal. More precisely, if one of these foliations is minimal, then it is topologically conjugate to the strong stable foliation of a double covering of a geodesic flow, and ρ preserves a point or a projective line in the projective plane. All these results hold for any (G, Y) -Anosov representation which is not quasi-Fuchsian, i.e., does not preserve a strictly convex domain in the projective plane.

1. INTRODUCTION

A flag is a pair (p, d) where p is the point of the projective plane, and d a projective line containing p . The group $G = \mathrm{PGL}(3, \mathbb{R})$ of projective transformations of the projective plane acts naturally on the flag variety X , i.e., the space of flags.

Let Γ be the fundamental group of a closed surface Σ of higher genus. In [1], we considered representations $\rho : \Gamma \rightarrow G$ near "horocyclic" representations, i.e., obtained from a faithful discrete representation $\Gamma \rightarrow H = \mathrm{SL}(2, \mathbb{R})$ composed with the natural morphism identifying H with the commutator subgroup of the stabilizer in G of a point p_0 and a projective line d_0 in the projective plane, with $p_0 \notin d_0$. Actually, in [1], we only considered some deformations of horocyclic representations for which p_0 is still a global fixed point: we called these representations *hyperbolic representations*. We proved that for such a representation, there is a closed $\rho(\Gamma)$ -invariant simple closed curve Λ in X , and an open $\rho(\Gamma)$ -invariant domain Ω , both depending on ρ , such that:

- Λ is the image of a 1 to 1 continuous Γ -equivariant map from the projective line $\mathbb{R}P^1$ into X , where the action of Γ on $\mathbb{R}P^1$ is the usual projective action (which is unique up to topological conjugacy).
- The action of $\rho(\Gamma)$ on Ω is free and properly discontinuous. The quotient space of this action, called M , is a flag manifold (cf. § 2.2). The first tautological foliation (see § 6.1) is topologically conjugate to the strong stable foliation of a double covering of the geodesic flow on Σ . On the other hand, the second tautological foliation is not minimal, except when $\rho(\Gamma)$ also preserves a projective line.

Date: May 17, 2005. Work supported by CNRS.

Key words and phrases. Flag manifold, Anosov representation.

In the present paper, we extend all these results omitting the assumption that $\rho(\Gamma)$ admits a global fixed point. The key observation is that horocyclic representations are (G, Y) -Anosov in the terminology of [16], where Y is the frame variety, i.e., the space of non-collinear points in the projective plane (see § 2.3).

Typical (G, Y) -Anosov representations are *hyperconvex*, i.e., those preserving a strictly convex domain of the projective plane. S. Choi and W. Goldman proved that every representation $\rho : \Gamma \rightarrow G$ which is quasi-Fuchsian, i.e., which can be continuously deformed to a representation taking values in $\mathrm{SO}_0(1, 2) \subset \mathrm{PGL}(3, \mathbb{R})$, preserves such a strictly convex curve ([8]). It follows easily that they are (G, Y) -Anosov, hence, hyperconvex. F. Labourie has extended this result to the higher-dimensional case ([16]). In [13], O. Guichard proved that conversely, any hyperconvex representation is quasi-Fuchsian.

Here, we consider general (G, Y) -Anosov representations, not necessarily hyperconvex. Each preserves a limit curve Λ , which is a Hölder continuous simple closed curve (see definition 3.4). In § 3, we establish some general results on this limit curve. In addition, we prove that such a representation always preserves an open domain Ω on which it acts freely and properly discontinuously (Theorem 5.1). We insist here on the low regularity of Λ and of the boundary $\partial\Omega$: when ρ is irreducible and not hyperconvex, then they are not Lipschitz regular (see corollary 7.2). Let's mention here that a similar statement is true in the hyperconvex case: the limit curve is in general C^1 with Hölder derivatives, but if the derivatives are Lipschitz, then $\rho(\Gamma)$ preserves a conic, i.e., is conjugate in $\mathrm{PGL}(3, \mathbb{R})$ to a Fuchsian subgroup.

Our first interest is in non-hyperconvex (G, Y) -Anosov representations: we prove that hyperbolic representations are (G, Y) -Anosov (Theorem 4.2, Lemma 4.5). We actually suspect that non-hyperconvex (G, Y) -Anosov representations form a connected space (see Question 2 in § 8). Observe that they all belong to the same connected component of the space of representations of Γ into G : the component of the trivial representation (corollary 6.6), whereas hyperconvex representations are those in the Hitchin component (see § 2.6, Remark 2.5.2).

For any (G, Y) -Anosov representation ρ , the quotient $M = \rho(\Gamma) \backslash \Omega$ is a natural flag manifold, which we call an *Anosov flag manifold*. It is therefore naturally equipped with two transversely projective 1-dimensional foliations: the tautological foliations (see § 6.1). When ρ is hyperconvex, M admits three connected components, with well-understood geometrical features (see remark 3.16), and tautological foliations with well-identified dynamical properties (see Remark 6.2): they are either foliations by circles or doubly covered by the geodesic flow on Σ for any hyperbolic metric.

The situation when ρ is not hyperconvex is completely different (cf. § 6.4): the tautological foliations in this case are never foliations by circles or finitely covered by Anosov flows. The correct picture is hard to capture: in some cases, they are topologically conjugate to the horocyclic flow of some Anosov flow (see Proposition 6.4), but this is not true in general. For example, when ρ is strongly irreducible, the tautological foliations are *not* minimal, in contrast to the horocyclic flow. The dynamical properties of these foliations are quite interesting. We suspect that these tautological foliations never admit periodic orbits (see Question 5 in § 8). If our suspicion is confirmed, it would provide examples of flows with unusual behavior. For example, recall the Seifert Conjecture, asserting that any flow on the three-dimensional sphere admits a periodic orbit. The first smooth counterexamples to this conjecture were found by K. Kuperberg ([14]). Observe moreover that the tautological foliations considered here can be volume-preserving: for example, this is the case for second tautological foliations associated to hyperbolic radial representations for which the morphism $u : \Gamma \rightarrow \mathbb{R}$ is trivial (see § 4.2). As far

as we know, the only known examples of volume preserving flows on the 3-sphere without periodic orbits have regularity at most C^2 ([15], [11]). As a matter of fact, volume preserving flows on 3-manifolds which are not minimal and without periodic orbits are quite uncommon; hence, it seems to us quite interesting to answer our Question 5.

ACKNOWLEDGEMENTS

I would like to thank F. Labourie, who made me aware of the (G, Y) -Anosov character of horocyclic representations. G. Kuperberg has corrected some imprecisions in a previous version of this paper regarding counterexamples to the Seifert conjecture.

2. DEFINITIONS

2.1. The Flag variety. – Let V be the vector space \mathbb{R}^3 , and (e_1, e_2, e_3) its canonical basis. Let V^* be the dual vector space, with the dual canonical basis (e_1^*, e_2^*, e_3^*) . We denote by $\langle v | v^* \rangle$ the evaluation of an element v^* of V^* on an element v of V .

– Let $N(v)$, $N(v^*)$ denote the norms on V , V^* , respectively, for the Euclidean metrics on V , V^* in which the canonical basis is orthonormal.

– $P(V)$ and $P(V^*)$ are the associated projective spaces. Elements of $P(V)$ are denoted $[v]$.

– The *flag variety* X is the closed subset of $P(V) \times P(V^*)$ formed by pairs $([v], [v^*])$ satisfying $\langle v | v^* \rangle = 0$.

– G is the group $\mathrm{SL}(V) \approx \mathrm{PGL}(V)$. The group G acts naturally on V and admits a dual (left) action on V^* uniquely defined by requiring $\langle u | g.v^* \rangle = \langle g^{-1}u | v^* \rangle$ for any u, v^* . For any g in $G = \mathrm{SL}(V)$, we denote by g^* the corresponding element of $\mathrm{SL}(V^*)$. If $\mathrm{SL}(V)$ and $\mathrm{SL}(V^*)$ are identified with $\mathrm{SL}(3, \mathbb{R})$ via the canonical basis, g^* is the inverse of the transpose of g .

– The diagonal action restricts as a natural action of G on X .

Remark 2.1. Every element of $P(V^*)$ defines a projective line in $P(V)$ (the projection to $P(V)$ of its kernel). Dually, every element of $P(V)$ corresponds to a projective line in $P(V^*)$. Hence, we can consider $P(V)$ to be the space of projective lines of $P(V^*)$ and $P(V^*)$ to be the space of projective lines of $P(V)$.

The image we have in mind is to view an element of X as a flag, i.e., a point in the projective plane and a projective line containing this point.

In order to formalize this point of view, we introduce the following notation: if K is a subspace of V (resp. of V^*), we denote by K^\perp its orthogonal, i.e., the subset of V^* (resp. of V) vanishing on K . Hence, for any $[v]$ in $P(V)$, $[v^\perp]$ is the corresponding projective line in $P(V^*)$.

If $[u], [v]$ are distinct elements of $P(V)$, we denote by $[(uv)^*]$ the element $[K^\perp]$ of $P(V^*)$, where K is the 2-dimensional space spanned by u and v . We employ similar notation when $[u], [v]$ belong to $P(V^*)$.

Remark 2.2. Let P^+ (respectively P^-) be the subgroup of G containing the upper (respectively lower) triangular matrices (under the identification of G with $\mathrm{SL}(3, \mathbb{R})$). Let X^\pm be the quotient spaces G/P^\pm : the map $g \mapsto ([ge_1], [g^*e_3^*])$ induces the identification $X^+ = G/P^+ \approx X$, and the map $g \mapsto ([ge_3], [g^*e_1^*])$ induces the identification $X^- = G/P^- \approx X$. These identifications are G -equivariant.

2.2. Flag manifolds. A flag structure is a (G, X) -structure, for (G, X) as above. We briefly present this notion here. For a more complete description of (G, X) -structures, see [19] or [12].

A flag structure on a manifold M is an atlas on M with charts taking values in X and coordinates changes expressed in the charts by restrictions of elements of

G . A typical example of a flag structure is the quotient of an open domain Ω of X by a discrete subgroup of G acting freely and properly discontinuously on Ω ; in particular, the flag variety X , itself, is an example. Less trivial examples are given in § 2.5.5, § 6.

A flag map between flag manifolds is a map which can be locally expressed in the flag charts by restrictions of elements of G . A flag map is always a local homeomorphism. A flag map which is bijective is called a flag isomorphism. A flag manifold is an isomorphism class of flag structures on the manifold.

Let $p : \widetilde{M} \rightarrow M$ be the universal covering and $\widetilde{\Gamma}$ the fundamental group of M , viewed as the group of covering automorphisms of p . For any flag structure on M , there is a map $\mathcal{D} : \widetilde{M} \rightarrow X$, called the *developping map*, and a representation $\rho : \widetilde{\Gamma} \rightarrow G$, called *holonomy representation*, such that:

- the maps p and \mathcal{D} are flag maps,
- \mathcal{D} is $\widetilde{\Gamma}$ -equivariant:

$$\forall \gamma \in \widetilde{\Gamma} \quad \mathcal{D} \circ \gamma = \rho(\gamma) \circ \mathcal{D}$$

2.3. The frame variety. Let Y denote the *frame variety*, i.e., the space of triples $([u], [v], [w])$ of noncollinear elements of $P(V)$. This 6-dimensional space is homogeneous under the diagonal action of G : it can be identified with the right quotient G/Z , where Z is the group of diagonal matrices.

The frame variety admits two natural projections π_{\pm} on X , defined by (see remark 2.1 for the notation):

$$\pi_+([u], [v], [w]) = ([u], [(uv)^*]) \quad ; \quad \pi_-([u], [v], [w]) = ([w], [(wv)^*])$$

These projections are both G -equivariant. Together, they define a map $\Pi : Y \rightarrow X \times X$. We denote by \mathcal{Y} the image of Π .

Lemma 2.3. *The elements of \mathcal{Y} are the pairs $(([u], [u^*]), ([v], [v^*]))$ satisfying:*

- $\langle u \mid u^* \rangle = 0$,
- $\langle v \mid v^* \rangle = 0$,
- $\langle u \mid v^* \rangle \neq 0$,
- $\langle v \mid u^* \rangle \neq 0$. ■

In other words, $[u]$ (respectively $[v]$) must belong to the projective line $[u^*]$ (resp. $[v^*]$) since we are considering elements of X , but it cannot belong to the projective line $[v^*]$ (respectively $[u^*]$).

The fibers of the projections π_{\pm} are the leaves of G -invariant foliations \mathcal{G}^{\pm} of X . Let E^{\pm} be the tangent bundles to these foliations. We obtain a G -invariant decomposition $TY = E^+ \oplus E^-$ of the tangent bundle of Y .

Remark 2.4. The inclusions $Z \subset P^{\pm}$ defines canonical maps $G/Z \rightarrow X^{\pm}$, which, via the identifications $X^{\pm} \approx X$ presented in remark 2.2 and $G/Z \approx Y$, are the maps π_{\pm} .

2.4. The geodesic flow as an Anosov flow. Let Σ be a closed surface with negative Euler characteristic, and Γ the fundamental group of Σ . Select a 1 to 1 morphism $\iota : \Gamma \rightarrow H$ with discrete image, where H denotes the group $SL(2, \mathbb{R})$. This induces a Fuchsian representation $\bar{\iota} : \Gamma \rightarrow \overline{H}$, where \overline{H} denotes the group $PSL(2, \mathbb{R})$ (observe that conversely, any Fuchsian representation lifts to a representation into H , since the associated Euler class is even).

Consider the flow on H induced by the right action of the 1-parameter group $A = \{a^t\}$, where:

$$a^t = \begin{pmatrix} e^t & 0 \\ 0 & e^{-t} \end{pmatrix}$$

This flow induces another flow on $M = \mathfrak{i}(\Gamma)\backslash H$, denoted by Φ^t . Its projection on $\overline{M} = \mathfrak{i}(\Gamma)\backslash \overline{H} \approx T^1\Sigma$ is a flow $\overline{\Phi}^t$.

Remark 2.5. This flow divided by 2, i.e., the flow $p \mapsto \Phi^{t/2}(p)$, is the familiar geodesic flow associated to the Riemannian surface $\mathfrak{i}(\Gamma)\backslash \mathbb{H}^2 \approx \Sigma$, where \mathbb{H}^2 is the hyperbolic plane.

Consider the following 1-parameter subgroups of H :

$$h_+^s = \begin{pmatrix} 1 & s \\ 0 & 1 \end{pmatrix}; \quad h_-^s = \begin{pmatrix} 1 & 0 \\ s & 1 \end{pmatrix}$$

We have the identities:

$$(1) \quad h_+^s a^t = a^t h_+^{\exp(-2t)s}; \quad h_-^s a^t = a^t h_-^{\exp(2t)s}$$

Denote also by h_\pm^s the flow on M induced by the right actions of these 1-parameter subgroups. The flow h_+^s (resp. h_-^s) is called the *stable horocyclic flow* (resp. *unstable horocyclic flow*). The flow Φ^t permutes the orbits of h_\pm^s . Moreover, the orbits of h_+^s are exponentially contracted by Φ^t , whereas the orbits of h_-^s are exponentially expanded by Φ^t . This feature establishes precisely that Φ^t and $\overline{\Phi}^t$ are *Anosov*.

Definition 2.6. *An Anosov flow on a closed manifold M equipped with a Riemannian metric $\|\cdot\|$ is a non-singular flow Φ^t such that the differential of Φ^t preserves a decomposition $TM = \Delta \oplus E^{ss} \oplus E^{uu}$ of the tangent bundle, satisfying the following properties, for some positive constants a, b :*

- *The line bundle Δ is tangent to the flow,*
- *for any vector v in E^{ss} over a point x of M , and for any positive t :*

$$\|D_x \Phi^t(v)\| \leq b e^{-at} \|v\|$$

- *for any vector v in E^{uu} over a point x of M , and for any negative t :*

$$\|D_x \Phi^t(v)\| \leq b e^{at} \|v\|$$

2.4.1. *Stable and unstable leaf spaces.* Denote by \mathcal{A}_\pm the group generated by a^t and h_\pm^s . It is isomorphic to the group of volume preserving affine transformations of the plane. The orbits on M or \overline{M} of \mathcal{A}_+ are called *stable leaves*; the orbits of \mathcal{A}_- are called *unstable leaves*. We denote by $\mathcal{S}_\pm, \overline{\mathcal{S}}_\pm$ the quotient spaces $H/\mathcal{A}_\pm, \overline{H}/\mathcal{A}_\pm$. The latter, $\overline{\mathcal{S}}_\pm$, are both homeomorphic to the projective line $\mathbb{R}P^1$, and \mathcal{S}_\pm are double coverings of $\overline{\mathcal{S}}_\pm$. Moreover, these identifications are H -equivariant, where the H -action on $\overline{\mathcal{S}}_\pm$ is the action induced by left translation, and the H -action on $\mathbb{R}P^1$ is the usual projective action.

2.4.2. *The bifoliated orbit space.* The map $hA \mapsto (h\mathcal{A}_+, h\mathcal{A}_-)$ embeds the orbit space $\overline{Q} = \overline{H}/a^t$ into the torus $\overline{\mathcal{S}}_+ \times \overline{\mathcal{S}}_-$. More precisely, for fixed H -equivariant identifications $\overline{\mathcal{S}}_\pm \approx \mathbb{R}P^1$, the image of this embedding is the complement in $\mathbb{R}P^1 \times \mathbb{R}P^1$ of the diagonal Δ . In other words, every \mathcal{A}_+ -orbit x intersects every \mathcal{A}_- -orbit, *except one*, which we call $\alpha(x)$. Note that we have defined a continuous Γ -equivariant map $\alpha: \overline{\mathcal{S}}_+ \rightarrow \overline{\mathcal{S}}_-$.

We denote by \overline{Q} the image of \overline{Q} in $\overline{\mathcal{S}}_+ \times \overline{\mathcal{S}}_-$; this is the complement of the graph of $\alpha: \overline{\mathcal{S}}_+ \rightarrow \overline{\mathcal{S}}_-$.

2.5. Anosov representations. Let $\rho : \Gamma \rightarrow G$ be any representation, and let $\pi_\rho : E_\rho \rightarrow M$ be the associated flat (G, Y) -bundle: E_ρ is the quotient of $H \times Y$ by the relation identifying each (h, y) with $(\iota(\gamma)h, \rho(\gamma).y)$, for every γ in Γ . The projection $(h, y) \mapsto h$ induces a map π_ρ from E_ρ onto $M = \iota(\Gamma) \backslash H$, which is a G -bundle, with fiber Y .

The (trivial) foliation of $H \times Y$ having as leaves the fibers of $(h, y) \mapsto y$ induces a foliation on M , which we denote by \mathcal{F}_ρ , and call the *horizontal foliation*. The leaves of \mathcal{F}_ρ are transverse to the fibers of π_ρ .

The flow Φ^t lifts uniquely to a horizontal flow Φ_ρ^t in E_ρ , i.e., tangent to the horizontal foliation: just take the flow induced in the quotient by the flow on $H \times Y$ defined by $(h, y) \mapsto (ha^t, y)$.

We have defined foliations \mathcal{G}^\pm on Y . They provide two 3-dimensional foliations on $H \times Y$ and induce on E_ρ two 3-dimensional foliations \mathcal{F}^\pm which are preserved by Φ_ρ^t and tangent to the fibers of π_ρ .

We will mainly consider the tangent bundles E_ρ^\pm of these foliations, which are canonically induced by the bundles E^\pm over Y .

Definition 2.7 ([16]). *A (G, Y) -Anosov structure over (M, Φ^t) is the data of a representation $\rho : \Gamma \rightarrow G$ and a continuous section s of π_ρ with the following properties:*

- the image S of s is Φ_ρ^t -invariant,
- for any norm $\| \cdot \|$ on E_ρ , there are positive constants a and b such that for any p in S , any positive t , any vector v^+ in E_ρ^+ over p , and any vector v^- in E_ρ^- over p , we have:

$$\begin{aligned} \|D_p \Phi_\rho^t(v^+)\| &\leq be^{-at} \|v^+\| \\ \|D_p \Phi_\rho^{-t}(v^-)\| &\leq be^{-at} \|v^-\| \end{aligned}$$

If these conditions are fulfilled, then the representation ρ is said to be (G, Y) -Anosov, or an Anosov flag representation.

Remark 2.8. Here, Y is the frame variety, and G the group $\text{SL}(V)$, but, of course, the definition 2.7 extends to any other pair (G, Y) , and any Anosov flow can play the role played here by the "geodesic flow" Φ^t .

The section s appearing in this definition is not assumed to be differentiable, even nor Lipschitz. The maximal regularity which can be required in general is Hölder continuity.

The main interest of Anosov representations is their stability: it follows from the structural stability of Anosov flows (more precisely, of hyperbolic closed sets) that the set of (G, Y) -Anosov representations is an open domain in the space of representations of Γ into G equipped with its natural topology (see proposition 2.1 of [16]). Observe that the perturbed (G, Y) -Anosov structure covers the *same* Anosov flow (M, Φ^t) .

Another important feature of Anosov representations is that they provide nice Γ -invariant geometric objects, obtained as follows:

The section s of a (G, Y) -Anosov structure lifts to a continuous map $f : H \rightarrow Y$ such that:

- f is Γ -equivariant: $f \circ \iota(\gamma) = \rho(\gamma) \circ f$,
- f is invariant by the lifted flow: $f(ha^t) = f(h)$.

Therefore, f induces a continuous Γ -equivariant map from the orbit space $Q = H/A$ into Y .

Lemma 2.9. *The restriction of $\pi_+ \circ f$ (resp. $\pi_- \circ f$) to any \mathcal{A}_+ -orbit (resp. \mathcal{A}_- -orbit) is constant.*

Proof Let p, p' be two elements of M belonging to the same stable leaf, i.e., on the same (right) \mathcal{A}_+ -orbit: for positive times t , the iterates $\Phi^t(p)$ and $\Phi^t(p')$ remain at a bounded distance apart inside the stable leaf. Then the exponential dilatation along the leaves of \mathcal{F}^- implies that $s(p)$ and $s(p')$ must belong to the same leaf of \mathcal{F}^+ . The lemma follows. \blacksquare

It follows from the lemma 2.9 that $\pi_+ \circ f$ and $\pi_- \circ f$ induce maps $f_+ : \mathcal{S}_+ \rightarrow X$ and $f_- : \mathcal{S}_- \rightarrow X$. Of course, f_{\pm} are both Γ -equivariant: their images (which, as we will see, are the same) are Γ -invariant (*a priori* only immersed) topological circles in the flag variety.

2.5.1. *Splitting the definition.* Consider the flat X -bundle $\pi_{\rho}^X : E_{\rho}(X) \rightarrow M$ associated to ρ . The total space $E_{\rho}(X)$ is the quotient of $H \times X$ by the Γ -action defined by $(h, x) \rightarrow (\gamma h, \rho(\gamma)x)$. The map Π defines a fibered embedding $\Pi_{\rho} : E_{\rho} \rightarrow E_{\rho}(X) \times E_{\rho}(X)$. Let \mathcal{Y}_{ρ} be the image of Π_{ρ} . The flow Φ^t also lifts in a unique way to a horizontal flow Φ_X^t on $E_{\rho}(X)$: $\Phi_X^t(h, x) = (h\alpha^t, x)$. Obviously, Definition 2.7 is equivalent to:

Definition 2.10. A (G, Y) -Anosov structure over (M, Φ^t) is the data of a representation $\rho : \Gamma \rightarrow G$ and two continuous sections s^{\pm} of π_{ρ}^X satisfying the following properties:

- the sections s^{\pm} are preserved by the flows: $\Phi_X^t(s^{\pm}(p)) = s^{\pm}(\Phi^t p)$,
- the image of s^+ (resp. s^-) is a (exponentially) repeller (resp. attractor) for Φ_X^t ,
- for every p in M , the pair $(s_+(p), s_-(p))$ belongs to \mathcal{Y}_{ρ} .

2.5.2. *Quasi-Fuchsian representations.* A very nice family of (G, Y) -Anosov representations is the family of *quasi-Fuchsian* representations (in the terminology of [16]), i.e., the representations $\rho : \Gamma \rightarrow G \approx \mathrm{SL}(3, \mathbb{R})$ which are in the *Hitchin component*, i.e., which can be deformed to *Fuchsian representations*, in other words, to a representation $\rho_0 : \Gamma \rightarrow \mathrm{SO}_0(1, 2)$. Indeed, F. Labourie has proven that quasi-Fuchsian representations are (G, Y) -Anosov.

Let's be a bit more precise: S. Choi and B. Goldman proved in [8] that any representation in the Hitchin component induces an action on $P(V)$ preserving a strictly convex domain C . The set of flags $([u], [v^*])$ where $[u]$ is a point in ∂C and $[v^*]$ a support projective line of C is a $\rho(\Gamma)$ -invariant curve in X ; in fact, this topological circle is equal to the images $f_+(\mathcal{S}_+)$ and $f_-(\mathcal{S}_-)$. In order to provide a simple hint, we just claim that the fact that this kind of representation is flag Anosov can be inferred from the fact that the geodesic flow associated to the Finsler Hilbert metric on $\rho(\Gamma) \backslash C$ is Anosov.

Following F. Labourie, this kind of curve is said to be *hyperconvex*, and a (G, Y) -Anosov representation preserving a hyperconvex curve is said hyperconvex. Thus, quasi-Fuchsian representations and hyperconvex representations coincide. O. Guichard extended recently this statement for all dimensions ([13]).

2.5.3. *Canonical Anosov flag representations.* Here, we consider another family of (G, Y) -Anosov representations, which are not quasi-Fuchsian. They are obtained from an embedding ρ_0 of $H = \mathrm{SL}(2, \mathbb{R})$ into a subgroup of $G = \mathrm{SL}(V)$ that admits a global fixed point in $P(V) \times P(V^*) \backslash X$.

For the fixed point, we select here the pair (e_2, e_2^*) . The embedding ρ_0 is the representation sending the matrix:

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

to the element

$$\begin{pmatrix} a & 0 & b \\ 0 & 1 & 0 \\ c & 0 & d \end{pmatrix}$$

of $G \approx \mathrm{SL}(3, \mathbb{R})$ (in the identification arising from the canonical basis).

Recall the notation introduced in § 2.1 and 2.3. Observe that $\rho_0(A)$ is contained in Z , and that every $\rho_0(h_\pm^s)$ belongs to P^\pm . The adjoint action of $\rho_0(a^t)$ on the Lie algebra $\mathfrak{sl}(V)$ of G is diagonalizable, and our choice of ρ_0 ensures that the subspace spanned by eigenvectors with positive eigenvalues (resp. negative) for $\mathrm{ad}(a^t)$ is the Lie algebra \mathcal{P}^+ (resp. \mathcal{P}^-) of P^+ (resp. P^-). Their intersection is the Lie algebra of Z .

Theorem 2.11 (Proposition 3.1 of [16]). *Let $\bar{\Gamma}$ be any cocompact subgroup of H . The restriction of ρ_0 to $\bar{\Gamma}$ is (G, Y) -Anosov.*

Proof See Appendix A. ■

Remark 2.12. Of course, we will apply Theorem 2.11 to $\bar{\Gamma} = 1(\Gamma)$. Actually, from the beginning, we could have selected as Γ any discrete cocompact subgroup of H , possibly with torsion, but this level of generality requires a little more caution in the formulation of statements, which we considered unnecessary and slightly uncomfortable. The reader should have no difficulty to extend the results of this paper to this more general context.

Definition 2.13. *The (G, Y) -Anosov representation $\rho_0 \circ 1$ is a canonical Anosov flag representation.*

Remark 2.14. Henceforth, except in Proposition 6.3, we will drop the symbol 1 , considering Γ directly as a discrete subgroup of H .

2.5.4. Invariant curves for canonical Anosov representations. Consider a canonical Anosov flag representation $\rho_0 : \Gamma \rightarrow G$. According to lemma 2.9, there are two Γ -equivariant maps $f_\pm : \mathcal{S}_\pm \rightarrow X$, coming from a Γ -equivariant map $f : H \rightarrow Y$. Here, the map f is defined by: $f(h) = ([\rho_0(h)e_1], [e_2], [\rho_0(h)e_3])$ (see Appendix A). Hence, we have: $f_+(h\mathcal{A}_+) = ([\rho_0(h)e_1], [\rho_0(h)^*e_3^*])$, and $f_-(h\mathcal{A}_-) = ([\rho_0(h)e_3], [\rho_0(h)^*e_1^*])$. These two maps have the same image Λ_0 , which is the set of pairs $([u], [u^*])$ where $[u]$ belongs to the projective line $[(e_2^*)^\perp]$ and $[u^*]$ is a projective line containing $[e_2]$ (in a more symmetric formulation, $[u^*]$ belongs to the projective line $[e_2^\perp]$ in $P(V^*)$).

We denote by L_0 the projective line $[(e_2^*)^\perp]$ of $P(V)$, and by L_0^* the projective line $[e_2^\perp]$ of $P(V^*)$. We can then reformulate the statement above: *a point $([u], [u^*])$ in X belongs to the curve Λ_0 if and only if $[u]$ belongs to L_0 and $[u^*]$ belongs to L_0^* .* Observe that for every $[u]$ in L_0 there is one and only one element $[u^*]$ of L_0^* containing $[u]$: the projective line containing $[u]$ and $[e_2]$.

Observe also that Λ_0 is the closure of the set of attractive fixed points of elements of $\rho_0(\Gamma)$ in X .

Finally, *the image of f is the space of triples $([u], [e_2], [w])$ of elements of $P(V)$ where $[u]$ and $[w]$ are distinct elements of L_0 .*

2.5.5. Canonical Anosov flag manifolds. Consider the orbits of $\rho_0(H)$ in X : there is one 1-dimensional orbit, the curve Λ_0 . There are two 2-dimensional orbits:

- the orbit A_0 containing the points $([u], [u^*])$ where $[u]$ belongs to L_0 and $[u^*]$ does not belong to L_0^* ,
- the orbit A_0^* containing the points $([u], [u^*])$ where $[u]$ does not belong to L_0 and $[u^*]$ belongs to L_0^* ,

There is one open orbit: $\Omega_0 = \{([u], [u^*]) \in X \mid [u] \notin L_0, [u^*] \notin L_0^*\}$.

The closures T_0, T_0^* of A_0, A_0^* are Klein bottles, the intersection $T_0 \cap T_0^*$ is Λ_0 , and Ω_0 is the complement in X of $T_0 \cup T_0^*$.

The action of $\rho_0(H)$ on Ω_0 is simply transitive. It provides an identification $\Omega_0 \approx H$. Therefore, the manifold $M = \Gamma \backslash H$ is homeomorphic to the quotient of Ω_0 by $\rho_0(\Gamma)$. It provides a natural flag structure on M .

Definition 2.15. *The quotient $\rho_0(\Gamma) \backslash \Omega_0$ is a canonical Anosov flag manifold.*

2.6. Existence of deformations. Consider the space $\text{Rep}(\Gamma, G)$ of representations of the surface group into G , modulo inner automorphisms of G on the target.

Theorem 2.16 (Hitchin). *$\text{Rep}(\Gamma, G)$ has three connected components.*

Let's briefly discuss each of these 3 components:

- *The Hitchin component:* this is the component containing the Fuchsian representations. Elements of this component are represented by quasi-Fuchsian representations (see § 2.5.2).

- *The trivial component:* this is the component containing the trivial representation.

- Representations in the third component are characterized by the fact that they do not lift to representations from Γ into the double covering $P^+\text{GL}(V)$ of G .

Canonical Anosov flag representations are not quasi-Fuchsian, and they clearly lift to representations in $\text{GL}(V)$: hence, they belong to the trivial component. It follows immediately that they can be deformed to representations in G which are not canonical! But there are much more elementary ways to prove this statement: any canonical representation ρ_0 can be deformed to a strongly irreducible representation, i.e., with image containing no finite index subgroup stabilizing a point or a projective line in $P(V)$ (see e.g. Proposition 3.11 of [1]).

3. GENERAL PROPERTIES OF (G, Y) -ANOSOV REPRESENTATIONS

Let $\rho : \Gamma \rightarrow G$ be any (G, Y) -Anosov representation. According to Lemma 2.9, there is a G -equivariant map $f : H \rightarrow Y$, inducing G -equivariant maps $f_{\pm} : \mathcal{S}_{\pm} \rightarrow X$.

Lemma 3.1. *For any γ in Γ and any attractive fixed point x of γ in \mathcal{S}_{\pm} , the image of x by f_{\pm} is an attractive (resp, repulsive) fixed point of $\rho(\gamma)$ in X .*

Proof Quite straightforward. See proposition 3.2 of [16]. ■

Proposition 3.2. *The representation ρ is discrete and faithful. For every non-trivial γ in Γ , the image $\rho(\gamma)$ is loxodromic, i.e., admits three eigenvalues with distinct norms.*

Proof Except for the discreteness, the proposition follows immediately from Lemma 3.1, the fact that any non-trivial element of Γ admits an attractive fixed in \mathcal{S}_+ , and the fact that loxodromic elements of $\text{SL}(V)$ are precisely elements admitting an attractive fixed point in X . The discreteness follows by classical arguments. See [16] for more details. ■

Actually, loxodromic elements of G have one and only one attractive fixed point in X . Since attractive fixed points of elements of γ are dense in \mathcal{S}_{\pm} , we obtain:

Corollary 3.3. *The maps f_{\pm} have the same image, which is the closure of the set of attractive fixed points of elements of $\rho(\Gamma)$ in X .* ■

Definition 3.4. *The common image $f_+(\mathcal{S}_+) = f_-(\mathcal{S}_-)$ is denoted by Λ , and called the limit curve.*

Recall that Φ^t is a double covering of the geodesic flow: there is a double covering between the associated leaf spaces $\mathcal{S}_\pm \rightarrow \overline{\mathcal{S}}_\pm$. Let $\bar{\tau}$ be the Galois automorphism of this double covering. For any non-trivial element γ of Γ and any attractive fixed point x of γ in \mathcal{S}_\pm , the image $\bar{\tau}(x)$ is an attractive fixed point of γ . By uniqueness of attractive fixed points in X , and according to Lemma 3.1, we have $f_\pm(x) = f_\pm(\bar{\tau}(x))$. By density of attractive fixed points in \mathcal{S}_\pm , we obtain $f_\pm = f_\pm \circ \bar{\tau}$. Hence:

Corollary 3.5. *The maps f_\pm induce maps $\bar{f}_\pm : \overline{\mathcal{S}}_\pm \rightarrow X$.* ■

According to § 2.4.2, the orbit space $\overline{\mathcal{Q}}$ can be identified with the complement $\overline{\mathcal{Q}}$ in $\overline{\mathcal{S}}_+ \times \overline{\mathcal{S}}_-$ of the graph of a homeomorphism $\alpha : \overline{\mathcal{S}}_+ \rightarrow \overline{\mathcal{S}}_-$. The maps \bar{f}_\pm induce a map $\bar{F} : \overline{\mathcal{Q}} \rightarrow X \times X$.

Lemma 3.6. *The image of \bar{F} is contained in the image \mathcal{Y} of Π .*

Proof Indeed, the maps \bar{f}_\pm arise from a map $f : H \rightarrow Y$. ■

Lemma 3.7. *We have the identity: $\bar{f}_+ = \bar{f}_- \circ \alpha$.*

Proof Let x in $\overline{\mathcal{S}}_+$ and y in $\overline{\mathcal{S}}_- \setminus \{\alpha(x)\}$. Then (x, y) belongs to the open set $\overline{\mathcal{Q}} \subset \overline{\mathcal{S}}_+ \times \overline{\mathcal{S}}_-$. The image of (x, y) by \bar{F} belongs to \mathcal{Y} (lemma 3.6). According to Lemma 2.3, we have $f_-(y) \neq f_+(x)$. Hence, $f_+(x)$ belongs to $\Lambda \setminus f_-(\overline{\mathcal{S}}_- \setminus \{\alpha(x)\})$. But $\Lambda \setminus \bar{f}_-(\overline{\mathcal{S}}_- \setminus \{\alpha(x)\})$ is either empty or reduced to $\{\bar{f}_-(\alpha(x))\}$. The lemma follows. ■

In the proof above, we have shown in particular that $\Lambda \setminus \bar{f}_-(\overline{\mathcal{S}}_- \setminus \{\alpha(x)\})$ is not empty. Hence:

Corollary 3.8. *The maps \bar{f}_+ and \bar{f}_- are injective.* ■

The flag manifold X is a closed subset of $P(V) \times P(V^*)$. Let η_\pm (resp. η_\pm^*) be the composition of \bar{f}_\pm with the projection of X on $P(V)$ (resp. $P(V^*)$).

Lemma 3.9. *The maps $\eta_\pm : \overline{\mathcal{S}}_\pm \rightarrow P(V)$ and the maps $\eta_\pm^* : \overline{\mathcal{S}}_\pm \rightarrow P(V^*)$ are injective.*

Proof We only deal with η_+ ; the other cases are similar. Let x, x' be two elements of $\overline{\mathcal{S}}_+$ with the same image by η_+ : if $[u] = \eta_+(x)$, then $\bar{f}_+(x) = ([u], [u^*])$, and $\bar{f}_+(x') = ([u], [v^*])$ for some $[u^*], [v^*]$ in $P(V^*)$ such that $\langle u | u^* \rangle = \langle u | v^* \rangle = 0$.

Assume $x \neq x'$. The pair (x, y) , with $y = \alpha(x')$, is an element of $\overline{\mathcal{Q}} \approx \overline{\mathcal{Q}}$. Its image by \bar{F} is $(\bar{f}_+(x), \bar{f}_-(\alpha(x')))$, which, according to Lemma 3.7, is equal to $(\bar{f}_+(x), \bar{f}_+(x'))$. On one hand, this pair must belong to the image \mathcal{Y} of Π . On the other hand, it has the form $(([u], [u^*]), ([u], [v^*]))$. From the description of the image of $\mathcal{Y} \subset X \times X$ (Lemma 2.3), we obtain a contradiction.

Hence, $x = x'$. The lemma is proved. ■

Definition 3.10. *Let L be the image of η_+ : this is the image of η_- too. Let L^* be the common image of η_+^* and η_-^* .*

According to Lemma 3.9, L and L^* are closed simple curves.

Lemma 3.11. *The limit curve Λ is the set of pairs $([v], [v^*])$ where $[v]$ belongs to L and $[v^*]$ belongs to L^* .*

Proof One of the inclusion is obvious. Conversely, let $([v], [v^*])$ be an element of X with $[v] \in L$, $[v^*] \in L^*$. Let (x, y) be the element of $\overline{\mathcal{S}}_+ \times \overline{\mathcal{S}}_-$ satisfying:

$$\eta_+(x) = [v], \quad \eta_-^*(y) = [v^*]$$

Since $\langle v \mid v^* \rangle = 0$, the pair $(\bar{f}_+(x), \bar{f}_-(y))$ does not belong to \mathcal{Y} . Therefore, (x, y) cannot belong to $\bar{\mathcal{Q}}$. We have $y = \alpha(x)$.

According to Lemma 3.7, we have $\bar{f}_+(x) = \bar{f}_-(y)$. Hence, the $P(V^*)$ -component of $\bar{f}_+(x)$ is $\eta_-^*(y) = [v^*]$. By construction, the $P(V)$ -component of $\bar{f}_+(x)$ is $[v]$. Therefore, the pair $([v], [v^*]) = \bar{f}_+(x)$ belongs to Λ . The lemma follows. ■

By Lemma 3.11 and Lemma 3.9:

Corollary 3.12. *Every projective line in $P(V)$ belonging to L^* (i.e., of the form $[(u^*)^\perp]$ with $[u^*]$ in L^*) intersects L in one and only one point.* ■

The dual statement, with L and L^* exchanged, is of course true. Hence, the corollary above can be complemented by:

Corollary 3.13. *For every point $[u]$ in L , there is one and only one projective line of $P(V)$ belonging to L^* and containing $[u]$.* ■

Corollary 3.14. *The curve L (respectively L^*) is the closure of the set of attractive fixed points in $P(V)$ (respectively $P(V^*)$) of elements of $\rho(\Gamma)$ (respectively $\rho^*(\Gamma)$).* ■

Remark 3.15. Of course, in the canonical case, i.e., when ρ is the restriction of ρ_0 to $\Gamma \subset H$, the curves L, L^* are the projective lines L_0, L_0^* . In this case, we defined Klein bottles T_0, T_0^* and an open domain Ω_0 (see § 2.5.5). These constructions extend to the general case in the following way: define T (respectively T^*) to be the set of flags $([u], [u^*])$ with $[u] \in L$ (respectively $[u^*] \in L^*$), and let Ω be the complement in X of the union $T \cup T^*$.

According to Lemma 3.11, the limit curve Λ is the intersection $T \cap T^*$, and the complements of Λ in T, T^* are denoted A, A^* .

Remark 3.16. All the results of these sections apply to any (G, Y) -Anosov representation and, in particular, to quasi-Fuchsian representations. In this case, L is the boundary of a $\rho(\Gamma)$ -invariant convex domain C in $P(V)$, and L^* is the boundary of the dual convex C^* : elements in L^* are projective lines in $P(V)$ tangent to L . The sets T, T^* are (topological) tori, and A, A^* are annuli.

The domain Ω in this case has 3 connected components:

- one component is the set of flags $([v], [v^*])$ with $[v] \in C$: this component is canonically identified with the projectivized tangent bundle of C ,
- another component is the set of flags $([v], [v^*])$ with $[v^*] \in C^*$: it is canonically identified with the projectivized tangent bundle of C^* ,
- Finally, there is a third component, consisting of the flags $([v], [v^*])$ with $[v] \notin C, [v^*] \notin C^*$.

The last component, in some way, has a lorentzian flavor. Indeed, when ρ is Fuchsian, i.e., when C and C^* are ellipses, this last component is canonically identified with the projectivized bundles of timelike vectors of de Sitter space.

4. SPECIAL DEFORMATIONS

In this section, we fix the embedding $\Gamma \subset H$, i.e., the canonical morphism $\rho_0 : \Gamma \rightarrow G$. The projection of Γ in \bar{H} is injective; we still denote by Γ the image of this projection. The quotient $\Gamma \backslash \bar{H}$ is the surface Σ .

Definition 4.1. *For any γ in Γ let $r(\gamma)$ be the spectral radius of $\gamma \in H$.*

Up to the sign, the eigenvalues of $\gamma \in H$ are $r(\gamma), r(\gamma)^{-1}$; when γ is non-trivial, we have $r(\gamma) > 1$.

4.1. Λ -preserving deformations.

4.1.1. *A flag version of the geodesic flow.* Consider¹ the following 1-parameter subgroup of G :

$$\varphi^t = \begin{pmatrix} e^{t/3} & 0 & 0 \\ 0 & e^{-2t/3} & 0 \\ 0 & 0 & e^{t/3} \end{pmatrix}$$

It commutes with every $\rho_0(h)$. Hence, it preserves the open $\rho_0(H)$ -orbit Ω_0 in X and induces a flow on the quotient manifold $\rho_0(\Gamma)\backslash\Omega_0$. The projection of Ω_0 on $P(V)$ is contained in the complement of L_0 , an affine plane. Consider the coordinate system (u, v) on this plane such that the coordinates of $[ue_1 + e_2 + ve_3]$ are (u, v) .

Actually, the projection of Ω_0 in $P(V)$ is the complement in this plane of the point $[e_2]$, i.e., the complement of the origin $(u, v) = (0, 0)$. The induced action of φ^t on this projection is the homothety of factor e^t fixing $[e_2] = (0, 0)$.

On the other hand, if h is an element of $H \approx \mathrm{SL}(2, \mathbb{R})$ of the form:

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

then $f_+(h)$ projects to $[\rho_0(h)e_1]$ in $P(V)$, with (u, v) -coordinates (a, c) . Hence, for any t , the (u, v) -coordinates of $f_+(ha^t)$ are $(e^t a, e^t c)$. Since the action of a transformation of X is characterized by its projective action on $P(V)$, we see that, via the identification $\Omega_0 \approx H$ we have selected, *the action of φ^t on Ω_0 coincides with the right action of a^t on H .* Hence, *the flow on $\rho_0(\Gamma)\backslash\Omega_0$ induced by φ^t is conjugate to the flow (M, Φ^t) .*

4.1.2. *Linear deformations.* Let $u : \Gamma \rightarrow \mathbb{R}$ be any morphism. The canonical morphism can be deformed to a new morphism, called the *u-deformation*:

$$\rho_u(\gamma) = \varphi^{u(\gamma)} \circ \rho_0(\gamma)$$

The morphism u is an element of $H^1(\Gamma, \mathbb{R})$. On this cohomology space, with $\rho_0 : \Gamma \rightarrow H$ fixed, we can define the *stable norm* (cf. [4]) as follows: for any hyperbolic element γ of Γ , let $t(\gamma)$ be the double of the logarithm of $r(\gamma)$ (this is the length of the closed geodesic associated to Γ in the quotient of the Poincaré disc by Γ). For any element $\hat{\gamma}$ of $H_1(\Gamma, \mathbb{Z})$, and for any positive integer n , let $t_n(\hat{\gamma})$ be the infimum of the values $\frac{t(\gamma)}{n}$ where γ describes all the elements of Γ representing $n\hat{\gamma}$. The limit of $t_n(\hat{\gamma})$ exists; it is the *stable norm of $\hat{\gamma}$ in $H_1(\Gamma, \mathbb{Z})$* . This norm is extended in a unique way on all $H_1(\Gamma, \mathbb{R})$; the dual of it is the *stable norm of $H^1(\Gamma, \mathbb{R})$* . The stable norm of u in $H^1(\Gamma, \mathbb{R})$ is denoted $|u|_s$.

Theorem 4.2. *The representation ρ_u is (G, Y) -Anosov if and only if $|u|_s < 1/2$.*

Proof Assume that ρ_u is (G, Y) -Anosov. The invariant curve Λ must be the closure of the union of attractive fixed points of $\rho_u(\Gamma)$. In particular, it contains the closure of the attractive fixed points of the commutator subgroup $\rho_u([\Gamma, \Gamma])$. But ρ_0 and ρ_u coincide on $[\Gamma, \Gamma]$, and attractive fixed points of elements of $[\Gamma, \Gamma]$ are dense in $\overline{\mathcal{S}}_{\pm} \approx \mathbb{R}P^1$; hence, $\Lambda = \Lambda_0$.

But it is easy to see that if $|u|_s > 1/2$, there is an element γ of Γ such that the attractive fixed point of $\rho_u(\gamma)$ in $P(V)$ is $[e_2]$. This attractive fixed point does not belong to L , which contradicts Corollary 3.14. Therefore, we must have $|u|_s \leq 1/2$. Since the (G, Y) -Anosov property is open in $\mathrm{Rep}(\Gamma, G)$, the inequality is strict: $|u|_s < 1/2$.

¹The factor $1/3$ arises from the fact that this flow truly lies in $\mathrm{PGL}(V)$: multiplying every coefficient by the inverse of the middle diagonal coefficient provides a more elegant expression. . .

The inverse statement, i.e., the fact that ρ_u is (G, Y) -Anosov if $|u|_s < 1/2$, is proved in Appendix A. \blacksquare

Remark 4.3. As the proof above shows, the limit curve Λ of ρ_u , when $|u|_s < 1/2$, is the limit curve Λ_0 of ρ_0 . It does not depend on the inclusion $\Gamma \subset H$.

4.2. Deformations with L_0 remaining constant. Consider morphisms $\rho : \Gamma \rightarrow G$ of the form:

$$\rho(\gamma) = \begin{pmatrix} e^{u(\gamma)/3}a(\gamma) & 0 & e^{u(\gamma)/3}b(\gamma) \\ \mu(\gamma) & e^{-2u(\gamma)/3} & \nu(\gamma) \\ e^{u(\gamma)/3}c(\gamma) & 0 & e^{u(\gamma)/3}d(\gamma) \end{pmatrix}$$

where:

– $\rho_\lambda : \gamma \mapsto \begin{pmatrix} a(\gamma) & b(\gamma) \\ c(\gamma) & d(\gamma) \end{pmatrix}$ is a Fuchsian representation taking value in $\mathrm{SL}(2, \mathbb{R})$,

i.e., is injective, with discrete image.

– $u : \Gamma \rightarrow \mathbb{R}$ is a morphism.

Such a representation is called a *radial representation*.

When u has stable norm (relatively to ρ_λ) strictly less than $1/2$, ρ is called a *hyperbolic representation*. In this case, $[e_2]$ is a fixed point of saddle type of every non-trivial $\rho(\gamma)$. In [1], the action of Γ on $P(V)$ induced by such a representation is called a *hyperbolic action*.

One of the main results of [1] is:

Theorem 4.4 (Theorem A of [1]). *Let ρ be a hyperbolic representation. The action of $\rho(\Gamma)$ on $P(V)$ preserves a continuous closed simple curve L . This curve, if Lipschitz, is a projective line.*

Actually, the first part of this Theorem is a corollary of the following:

Lemma 4.5. *Hyperbolic representations are (G, Y) -Anosov.*

Proof If the maps μ and ν are trivial, it follows from Theorem 4.2. Then, observe that the composition of a hyperbolic representation by the conjugacy in G by φ^t remains hyperbolic radial, with the same ρ_λ and the same u , but with coefficients μ , ν multiplied by e^{-2t} . Since ρ_u is (G, Y) -Anosov, and by stability of (G, Y) -Anosov representations, the conjugated representation for big t is (G, Y) -Anosov. The lemma follows since conjugacy in G does not affect the (G, Y) -Anosov property. \blacksquare

Remark 4.6. The invariant curve L^* in $P(V^*)$ is obviously $L_0^* = [e_2^\perp]$.

Remark 4.7. The analog of Theorem 4.2 remains true: a radial representation is (G, Y) -Anosov if and only if $|u|_s < 1/2$. The proof is similar; we leave it to the reader.

Proposition 4.8. *Let $\rho : \Gamma \rightarrow G$ be a (G, Y) -Anosov representation. Assume that a finite index subgroup of $\rho(\Gamma)$ preserves a proper subspace of V . Then, up to conjugacy in G , ρ or ρ^* is a hyperbolic representation.*

Proof A proper subspace of V is a line or a 2-plane: replacing ρ by ρ^* if necessary, we can assume that the subspace preserved by a finite index subgroup of $\rho(\Gamma)$ is a line. After conjugacy in G , we can assume moreover that this invariant line is spanned by e_2 . Let $\Gamma' \subset \Gamma$ be the finite index subgroup such that $\rho(\Gamma')$ fixes $[e_2]$. The restriction ρ' of ρ to Γ' is still (G, Y) -Anosov, and its limit curve Λ' is the limit curve Λ of ρ .

According to Proposition 3.2, ρ' is faithful, with discrete image. It follows that ρ' is a radial representation, described as above by a Fuchsian representation $\rho'_\lambda :$

$\Gamma' \rightarrow H$ and a morphism $u' : \Gamma \rightarrow \mathbb{R}$. This morphism is trivial on $[\Gamma', \Gamma']$. It follows that $[e_2]$ is a saddle fixed point of $\rho'(\gamma)$ for every element γ of $[\Gamma', \Gamma']$. Hence, the attractive fixed point of $(\rho')^*(\gamma)$ in $P(V^*)$ belongs to $[e_2^\perp]$. The argument used in the proof of Theorem 4.2 and the identity $\Lambda = \Lambda'$ imply here that the invariant curve L_0^* is $[e_2^\perp]$. In particular, it follows that $[e_2]$ is a saddle fixed point of $\rho(\gamma)$ for every γ in Γ , not only in $[\Gamma', \Gamma']$. Therefore, ρ is a linear u -deformation of a canonical representation. As in the proof of 4.2, we infer $|u|_s \leq 1/2$, and, finally, $|u|_s < 1/2$ (by stability of Anosov representations). ■

5. PROPERNESS OF THE ACTION ON Ω

Let $\rho : \Gamma \rightarrow G$ be a (G, Y) -Anosov representation. Recall that $\rho(\Gamma)$ preserves the open domain Ω , which is the set of flags $([u], [u^*])$ with $[u] \notin L$, $[u^*] \notin L^*$.

Theorem 5.1. *For any (G, Y) -Anosov representation $\rho : \Gamma \rightarrow G$, the action of $\rho(\Gamma)$ on the associated domain Ω is free and properly discontinuous.*

Remark 5.2. This Theorem, when ρ is a linear deformation ρ_u , is a reformulation of Théorème 3.4 of [17]. More generally, Theorem 5.1 is proved in [1] (Proposition 4.19 of [1]) in the case of hyperbolic representations.

Therefore, in this section, we don't consider this particular case. By Proposition 4.8, it means that we assume that no proper subspace of V is $\rho(\Gamma)$ -invariant, i.e., that ρ is *strongly irreducible*.

In Appendix B, we show how the proof developed here can be adapted to the reducible case, achieving a complete proof of Theorem 5.1.

All this section is devoted to the proof of Theorem 5.1. Let ρ be a (G, Y) -Anosov representation. According to Lemma 3.2, every $\rho(\gamma)$ is loxodromic: its fixed points in X all have the form $([u], [u^*])$ where either $[u]$ belongs to L , or $[u^*]$ belongs to L^* . None of them belongs to Ω : the action of $\rho(\Gamma)$ is free.

Our task is to prove that the action is proper. Since $\rho(\Gamma)$ is discrete, it amounts to proving that there is no sequence $(\gamma_n)_{(n \in \mathbb{N})}$ in Γ for which there is a sequence of flags $([u_n], [u_n^*])_{(n \in \mathbb{N})}$ in Ω satisfying:

- (1) the sequence $\rho(\gamma_n)_{(n \in \mathbb{N})}$ escapes from any compact of G ,
- (2) the sequence $([u_n], [u_n^*])_{(n \in \mathbb{N})}$ converge to a point $([\bar{u}], [\bar{u}^*])$ in Ω ,
- (3) the sequence $\rho(\gamma_n)([u_n], [u_n^*])_{(n \in \mathbb{N})} = ([v_n], [v_n^*])_{(n \in \mathbb{N})}$ converges to a flag $([\bar{v}], [\bar{v}^*])$ in Ω .

We argue by contradiction, assuming the existence of such a sequence. We denote $g_n = \rho(\gamma_n)$.

Remember that we equip V, V^* with the euclidean metrics N, N^* for which $([e_1], [e_2], [e_3])$ and $([e_1^*], [e_2^*], [e_3^*])$ are an orthonormal basis. It induces a norm on $\text{GL}(V)$: the operator norm $\| \cdot \|$. Let $\mathcal{A} \subset \text{GL}(V)$ be the unit sphere of this norm.

Consider the Cartan decompositions of g_n, g_n^* according to the canonical basis:

$$g_n = k_n \begin{pmatrix} \lambda_n & 0 & 0 \\ 0 & \mu_n & 0 \\ 0 & 0 & \nu_n \end{pmatrix} l_n^{-1}$$

$$g_n^* = k_n \begin{pmatrix} \lambda_n^{-1} & 0 & 0 \\ 0 & \mu_n^{-1} & 0 \\ 0 & 0 & \nu_n^{-1} \end{pmatrix} l_n^{-1}$$

where k_n, l_n are isometries of N , and $\lambda_n \geq \mu_n \geq \nu_n$, with $\lambda_n \mu_n \nu_n = 1$.

The quotients $\bar{g}_n = \frac{g_n}{\|g_n\|}$ and $\bar{g}_n^* = \frac{g_n^*}{\|g_n^*\|}$ belong to \mathcal{A} . By compactness of this unit sphere, passing to a subsequence if necessary, we can assume that \bar{g}_n and \bar{g}_n^* converge to elements \bar{g}, \bar{g}^* , respectively, of \mathcal{A} . Let I be the image of \bar{g} , I_* the image

of \bar{g}^* , K the kernel of \bar{g} and K_* the kernel of \bar{g}^* . By convention, $[I]$, $[K]$, $[I_*]$ and $[K_*]$ are their projections in $P(V)$, $P(V^*)$. Let $I^\perp, K^\perp \subset V^*$ and $K_*^\perp, I_*^\perp \subset V$ be the dual subspaces, and $[I^\perp]$, $[K^\perp]$, \dots their respective projectivizations.

The g_n all have determinant 1. Hence, the item (1) above implies that, after taking a subsequence, the norms $\|g_n\|$ and $\|g_n^*\|$ tend to $+\infty$. If \bar{g}^t denotes the transpose matrix of \bar{g} , we have:

$$\bar{g}^t \circ \bar{g}^* = 0 = \bar{g}^* \circ \bar{g}^t$$

Therefore:

$$I_* \subset I^\perp, \quad K^\perp \subset K_*$$

Since the actions of g_n and \bar{g}_n on $P(V)$ coincide, as for the actions of g_n^* and \bar{g}_n^* on $P(V^*)$, we obtain:

Lemma 5.3. *The sequence $(\bar{g}_n)_{(n \in \mathbb{N})}$ converge uniformly on compact subsets of $P(V) \setminus [K]$ to the restriction of \bar{g} to $P(V) \setminus [K]$. Similarly, the sequence $(\bar{g}_n^*)_{(n \in \mathbb{N})}$ converge uniformly on compact subsets of $P(V^*) \setminus [K_*]$ to the restriction of \bar{g}^* . ■*

Remark 5.4. The image of the restriction of \bar{g} to $P(V) \setminus [K]$ is $[I]$, and the image of the restriction of \bar{g}^* to $P(V^*) \setminus [K_*]$ is $[I_*]$.

Actually, since the norm of g_n tends to $+\infty$, we see that, for \bar{k} and \bar{l} denoting the limits of \bar{k}_n and \bar{l}_n , the matrices \bar{g} and \bar{g}^* have the expressions:

$$\bar{g} = \bar{k} \begin{pmatrix} \bar{\lambda} & 0 & 0 \\ 0 & \bar{\mu} & 0 \\ 0 & 0 & 0 \end{pmatrix} \bar{l}^{-1}$$

$$\bar{g}^* = \bar{k} \begin{pmatrix} 0 & 0 & 0 \\ 0 & \bar{\mu}' & 0 \\ 0 & 0 & \bar{\nu} \end{pmatrix} \bar{l}^{-1}$$

where $\bar{\lambda}, \bar{\nu}$ are positive, and $\bar{\mu}, \bar{\mu}'$ are non-negative.

We will also need to exchange the roles of $([\bar{u}], [\bar{u}^*])$ and $([\bar{v}], [\bar{v}^*])$; this amounts to replacing the γ_n by their inverses, i.e., to replacing g_n and g_n^* . Let h_n, h_n^* be these inverses:

$$h_n = l_n \begin{pmatrix} \lambda_n^{-1} & 0 & 0 \\ 0 & \mu_n^{-1} & 0 \\ 0 & 0 & \nu_n^{-1} \end{pmatrix} k_n^{-1}$$

$$h_n^* = l_n \begin{pmatrix} \lambda_n & 0 & 0 \\ 0 & \mu_n & 0 \\ 0 & 0 & \nu_n \end{pmatrix} k_n^{-1}$$

Their limits for $n \rightarrow +\infty$ are:

$$\bar{h} = \bar{k} \begin{pmatrix} 0 & 0 & 0 \\ 0 & \bar{\mu}' & 0 \\ 0 & 0 & \bar{\nu}' \end{pmatrix} \bar{l}^{-1}$$

$$\bar{h}^* = \bar{k} \begin{pmatrix} \bar{\lambda} & 0 & 0 \\ 0 & \bar{\mu} & 0 \\ 0 & 0 & 0 \end{pmatrix} \bar{l}^{-1}$$

Lemma 5.5. *The coefficients $\bar{\mu}$ and $\bar{\mu}'$ are both zero.*

Proof Assume that one of them, let's say $\bar{\mu}$, is non-zero. Then, K is the line spanned by $\bar{l}(e_3)$. The inclusion $K^\perp \subset K_*$ shows that K_* has dimension 2. Hence, I_* is 1-dimensional. Therefore, the inclusions are all identities:

$$I_* = I^\perp, \quad K^\perp = K_*$$

Fact 1: $[K]$ belongs to the invariant curve L . If not, the g_n -invariant curve L is contained in $P(V) \setminus [K]$. By Lemma 5.3, it implies that L is equal to the projective line $[I]$, which is therefore a $\rho(\Gamma)$ -invariant projective line. This contradicts the strong irreducibility of ρ .

Fact 1': $[I_*]$ belongs to L^* . Apply the argument used for the proof of Fact 1 to the inverse sequence $(h^* = (g_n^*)^{-1})_{(n \in \mathbb{N})}$. We obtain that if $[I^*]$ does not belong to L^* , then L^* is the projective line $[K_*]$, contradicting once more the irreducibility of ρ .

Fact 2: The invariant curve L is a projective line containing $[K]$. Indeed, according to Fact 1', the projective line $[I]$ belongs to L^* . According to Corollary 3.12, the intersection $[I] \cap L$ is a single point $[p]$. On the other hand, the fibers of the restriction of \bar{g} to $P(V) \setminus [K]$ are the projective lines containing $[K]$, with the point $[K]$ excluded. By Lemma 5.3, since $[I] \cap L$ is a single point, $L \setminus [K]$ is contained in one and only one of these fibers. Fact 2 follows.

The lemma follows by irreducibility of ρ . ■

According to Lemma 5.5, $[K]$ is a projective line, and $[I]$ is a single point. Similarly, $[K_*]$ is a projective line, whereas $[I_*]$ is a single point.

Lemma 5.6. *The points $[I]$, $[I_*]$ belong to L , L^* , respectively.*

Proof Assume that $[I]$ does not belong to L . Then, by Lemma 5.3, L is contained in the projective line $[K]$, which is therefore $\rho(\Gamma)$ -invariant, a contradiction. The proof of the dual statement $[I_*] \in L^*$ is similar. ■

Lemma 5.7. *The points $[K^\perp]$ and $[K_*^\perp]$ belong respectively to L^* , L .*

Proof According to Lemma 5.5, the limit in \mathcal{A} of the \bar{h}_n (i.e., the h_n divided by their norms) is:

$$\bar{h} = \bar{l} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \bar{v} \end{pmatrix} \bar{k}^{-1}$$

Hence, the image of \bar{h} is spanned by $\bar{l}e_3$. On the other hand, the kernel K_* of \bar{g}^* is the 2-plane generated by $\bar{l}(e_1^*)$ and $\bar{l}(e_2^*)$. Therefore, K_*^\perp is the line spanned by $\bar{l}e_3$, i.e., the image of \bar{h} . Applying Lemma 5.6 to \bar{h} , we get that $[K_*^\perp]$ belongs to L .

The same argument, applied to \bar{h}^* , implies $[K^\perp] \in L^*$ ■

End of the Proof of Theorem 5.1 in the strongly irreducible case:

Since $([\bar{v}], [\bar{v}^*])$ belongs to Ω , $[\bar{v}]$ does not belong to L . According to Lemma 5.6, $\bar{v} \neq [I]$. It follows, with Lemma 5.3, that $[\bar{u}]$ belongs to $[K]$. Similarly, we have $[\bar{u}^*] \in [K_*]$. Hence:

- the projective line $[K_*]$ in $P(V^*)$ contains $[\bar{u}^*]$ and $[K^\perp]$ (since $K^\perp \subset K_*$),
- the projective line $[\bar{u}^\perp]$ in $P(V^*)$ contains $[\bar{u}^*]$ (since $\langle \bar{u} \mid \bar{u}^* \rangle = 0$) and $[K^\perp]$ (since $[\bar{u}] \in [K]$) too.

But, according to Lemma 5.7, since $[\bar{u}^*]$ does not belong to L^* , we have $\bar{u}^* \neq [K^\perp]$. Hence the projective lines $[K_*]$ and $[\bar{u}^\perp]$ share two distinct points: they are equal. In other words, $[\bar{u}]$ is equal to $[K_*^\perp]$: according to Lemma 5.7, it belongs to L , a contradiction. ■

Remark 5.8. Theorem 5.1 is true for any (G, Y) -Anosov representations, in particular, it applies to representations in the Hitchin component, i.e., dividing a strictly convex domain C of $P(V)$. In this case, we recover the well-known properness of the action on the projectivized tangent bundle of C . But we also obtain the properness of the action on "timelike directions" over the "de Sitter-like" component $P(V) \setminus \overline{C}$.

Remark 5.9. We can ask how Theorem 5.1 can be extended to any other (G, Y) -Anosov representations, for other pairs (G, Y) . For example, for the pair $(G, Y) = (\mathrm{PSL}(2, \mathbb{C}), \mathbb{C}P^1 \times \mathbb{C}P^1 \setminus \mathrm{diag})$, i.e., the case of conformal quasi-Fuchsian representations, Theorem 5.1 corresponds the well-known properness of the action on the discontinuity domain, i.e., the complement in $\mathbb{C}P^1$ of the limit set.

Remark 5.10. In the wonderful paper [18], R. Schwarz considers some particular actions of $\mathrm{PSL}(2, \mathbb{Z})$ on the flag manifold X . He exhibited invariant curves L, L^* with properties completely similar to the invariant curves L, L^* we have considered here²: compare our corollaries 3.12, 3.13 with Theorem 3.3 of [18]. Our Theorem 5.1 echoes Theorem 4.2 of [18]. Hence, it seems reasonable to qualify these representations of $\mathrm{PSL}(2, \mathbb{Z})$ as (G, Y) -Anosov representations. In fact, they are also deformations of the representation obtained by composing $\rho_0 : H \rightarrow G$ with the inclusion $\mathrm{SL}(2, \mathbb{Z}) \subset \mathrm{SL}(2, \mathbb{R})$. But there is a crucial difference: $\mathrm{PSL}(2, \mathbb{Z})$ is indeed a lattice of $\mathrm{PSL}(2, \mathbb{R})$, but not cocompact! It is therefore presumably possible and interesting to extend the notion of (G, Y) -Anosov representations to nonuniform lattices of H .

6. ANOSOV FLAG MANIFOLDS

Thanks to Theorem 5.1, the definition 2.15 can be extended:

Definition 6.1. *Let $\rho : \Gamma \rightarrow G$ be a (G, Y) -Anosov representation. The quotient by $\rho(\Gamma)$ of the domain Ω is called an Anosov flag manifold.*

6.1. Tautological foliations. The fibers of the projections $X \rightarrow P(V), P(V^*)$ are leaves of foliations by circles on X . They are preserved by G ; therefore, they induce two 1-dimensional foliations on any flag manifold. The foliation corresponding to the projection $X \rightarrow P(V)$ is called the *first tautological foliation*. The other one, corresponding to the projection $X \rightarrow P(V^*)$, is called the *second tautological foliation*. Observe that these foliations are both transversely real projective.

Remark 6.2. When ρ is hyperconvex, Ω has three connected components (see remark 3.16):

- one is the set of flags $([v], [v^*])$ with $[v] \in [C]$. We denote it by Ω_1 . The quotient $M_1 = \rho(\Gamma) \backslash \Omega_1$ is naturally identified with the projectivized tangent bundle of the convex real projective surface $S = \rho(\Gamma) \backslash [C]$. The leaves of the first tautological foliation are the fibers of the bundle map. In particular, they are compact. The second tautological foliation is the foliation supported by the geodesic flow of the Hilbert metric on S , quotiented by the involution sending any tangent vector to its opposite: hence, it is topologically conjugate to the geodesic flow of any hyperbolic metric on the surface Σ quotiented by the antipodal map in the fibers (see [1] for more details).

- another component, Ω_2 , which is the set of flags $([v], [v^*])$ with $[v^*] \in [C^*]$. The quotient $M_2 = \rho(\Gamma) \backslash \Omega_2$ is the projectivized tangent bundle of the dual convex real projective surface $S^* = \rho^*(\Gamma) \backslash [C^*]$. The leaves of the second tautological foliation are the fibers of the bundle map, whereas the first tautological foliation is doubly covered by the geodesic flow on Σ .

²There is a minor difference: the action on X considered in [18] contains *polarities*, i.e., projective transformations followed by the flip $([v], [v_*]) \mapsto ([v^\perp], [v^\perp_*])$.

– the third component, Ω_3 , is the “de Sitter”-like component. It is the set of pairs $([v], [v^*])$, where $[v]$ belongs to the Möbius band \mathcal{A} , which is the complement in $P(V)$ of the closure of $[C]$, and $[v^*]$ belongs to the Möbius band \mathcal{A}^* , which is the complement in $P(V^*)$ of the closure of $[C^*]$. Observe that for any $[v]$ in \mathcal{A} , the intersection $[v^\perp] \cap L^*$ is the union of two points $[v_-^*], [v_+^*]$ (note that $L^* = \partial[C^*]$). Then, $[v_\pm^*]$ is tangent to $L = \partial[C]$ at a point $[v_\pm]$. Let $p([v])$ be the intersection of $[(v^*)^\perp]$ and the projective line $[w^*]$ containing $[v_-]$ and $[v_+]$; it belongs to $[C]$. Hence, the flag $(p([v]), [w^*])$ belongs to Ω_1 . We have thus defined a $\rho(\Gamma)$ -equivariant map from Ω_3 into Ω_1 . It is easy to show that it is a homeomorphism, and that it sends the first tautological foliation of Ω_3 onto the second tautological foliation of Ω_1 .

A similar treatment can be applied to the second tautological foliation on Ω_3 , leading to the following statement: *the tautological foliations of $M_3 = \rho(\Gamma) \backslash \Omega_3$ are both topologically conjugate to the geodesic flow of any hyperbolic metric on the surface Σ quotiented by the antipodal map in the fibers.*

6.2. Canonical Goldman flag manifolds. In [1], Anosov flag manifolds associated to linear u -deformations of canonical representations were defined and called *canonical Goldman flag manifolds*.

We recall that in the case of canonical Goldman flag manifolds, the flow φ^t induces an Anosov flow on the associated canonical flag manifold. More precisely, this flow is a double covering of a *Desarguian Anosov flow* as defined in [10], [2]. The tautological foliations are then the strong stable and unstable foliations. See [1] for more details.

For our purpose here, it is more suitable to drop the identification $G = \text{PGL}(V) \approx \text{SL}(V)$, and to consider u -deformations as morphisms $\lambda : \Gamma \rightarrow \text{GL}_2$, where GL_2 is the group of invertible 2×2 matrices, identified with the stabilizer in G of $[(L_0^*)^\perp] = [e_2]$ and $L_0 = [(e_2^*)^\perp]$:

$$\rho(\gamma) = \begin{pmatrix} & & 0 \\ \lambda(\gamma) & & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

The group GL_2 preserves the annulus $A_{can} = P(V) \setminus (L_0 \cup [e_2])$ (there is also a dual action on the annulus $P(V^*) \setminus (L_0^* \cup [e_2^*])$). The minimality and unique ergodicity of horocyclic flows ([6]) imply that the $\lambda(\Gamma)$ -action on A_{can} is minimal and uniquely ergodic: up to a positive factor, there is one and only Borel measure on A_{can} preserved by $\lambda(\Gamma)$.

Proposition 6.3. *Two inclusions λ_1, λ_2 from a surface group Γ into GL_2 induce conjugate actions on the annulus A_{can} if and only if λ_1, λ_2 are conjugate in GL_2 .*

Proof This is a folkloric fact, but we don’t know any appropriate reference.

One of the implication is clear. Let’s prove the inverse statement: let $f : A_{can} \rightarrow A_{can}$ be a homeomorphism conjugating the actions of Γ on A_{can} via λ_1, λ_2 , respectively. Clearly, f extends to a homeomorphism $f : P(V) \setminus L_0 \rightarrow P(V) \setminus L_0$, still Γ -equivariant. Consider φ^t as a group of projective transformations of A_{can} .

For any nontrivial γ in Γ , $[e_2]$ is saddle fixed point of $\lambda_1(\gamma)$ and $\lambda_2(\gamma)$, and the $\lambda_1(\gamma)$ - or $\lambda_2(\gamma)$ -stable leaf of $[e_2]$ is the union of two orbits of φ^t , with $[e_2]$ itself. Moreover, when γ is varying, these stable leaves form a dense subset among the orbits of φ^t . It follows that f sends φ^t -orbits on φ^t -orbits. Hence, there is a continuous function $\alpha : \mathbb{R} \times A_{can} \rightarrow \mathbb{R}$ such that:

$$f(\varphi^t([u])) = \varphi^{\alpha(t,[u])} f([u])$$

For a fixed t , since φ^t commutes with the Γ -actions, $[u] \mapsto \alpha(t, [u])$ is $\lambda_1(\Gamma)$ -invariant. Since the action of $\lambda_1(\Gamma)$ on the annulus is minimal, it follows that α depends only on t . Since $f^{-1} \circ \varphi^t \circ f$ is a one-parameter subgroup of G , $\alpha : \mathbb{R} \rightarrow \mathbb{R}$ is a morphism: there is a positive constant C such that $\alpha(t) = Ct$.

For some non-trivial γ in Γ , and for $j = 1, 2$, denote by $\pm e^{a_j(\gamma)}$, $\pm e^{b_j(\gamma)}$ the eigenvalues $\neq 1$ of $\lambda_j(\gamma)$, with $b < 0 < a$. Then, the $\lambda_j(\gamma)$ -stable leaf of $[e_2]$ is the fixed point set of $\lambda_1(\gamma)\varphi^{-b_1(\gamma)}$. Its image by f is the fixed point set of $\lambda_2(\gamma)\varphi^{-Cb_1(\gamma)}$. It follows that $b_2(\gamma) = Cb_1(\gamma)$. Similarly, $a_2(\gamma) = Ca_1(\gamma)$.

Consider the projections of $\lambda_j(\gamma)(\Gamma)$ in $\mathrm{PSL}(2, \mathbb{R}) = \overline{H}$. They are Fuchsian subgroups, corresponding to metrics g_1, g_2 on the surface Σ with constant curvature -1 . Then, $a_2(\gamma) = Ca_1(\gamma)$ and $b_2(\gamma) = Cb_1(\gamma)$ imply that *the g_2 -length of a closed geodesic c is C times the g_1 -length of the closed geodesic freely homotopic to c* . This is possible only if $C = 1$ (see for example [9]; on page 137, end of ‘‘Exposé 7’’, it is proved that the hypothesis (H) on page 136, i.e., the claim $C \neq 1$, is impossible).

Therefore, if Tr denotes the trace function on the algebra \mathfrak{gl}_2 of 2×2 matrices:

$$(2) \quad \mathrm{Tr} \circ \lambda_1 = \mathrm{Tr} \circ \lambda_2$$

It is well-known that this implies that λ_1 and λ_2 are conjugate in GL_2 . Let’s recall the argument: since λ_j are irreducible representations, every element g of \mathfrak{gl}_2 can be (non-uniquely) written as a sum $\sum_i \eta_i \lambda_j(\gamma_i)$.

For every g in \mathfrak{gl}_2 , select such a decomposition $g = \sum_i \eta_i \lambda_1(\gamma_i)$. Then define $\phi(g) = \sum_i \eta_i \lambda_2(\gamma_i)$. The key point is that $\phi(g)$ does not depend on the selected decomposition of g . Indeed, if $0 = \sum_i \eta_i \lambda_1(\gamma_i)$, then, for every $g' = \sum_k \nu_k \lambda_2(\gamma_k)$ in \mathfrak{gl}_2 , we have:

$$\begin{aligned} \mathrm{Tr}((\sum_i \eta_i \lambda_2(\gamma_i))g') &= \mathrm{Tr}(\sum_{i,k} \eta_i \nu_k \lambda_2(\gamma_i) \lambda_2(\gamma_k)) \\ &= \sum_{i,k} \eta_i \nu_k \mathrm{Tr}(\lambda_2(\gamma_i \gamma_k)) \\ &= \sum_{i,k} \eta_i \nu_k \mathrm{Tr}(\lambda_1(\gamma_i \gamma_k)) \\ &= \mathrm{Tr}((\sum_i \eta_i \lambda_1(\gamma_i))(\sum_k \nu_k \lambda_1(\gamma_k))) = 0 \end{aligned}$$

Since this holds for every g' , and since $g' \mapsto \mathrm{Tr}(g_0 g')$ can be zero only if $g_0 = 0$, we obtain that $0 = \sum_i \eta_i \lambda_2(\gamma_i)$. As a corollary, ϕ is well-defined.

This map ϕ is obviously an algebra automorphism of \mathfrak{gl}_2 ; but such an automorphism is known to be an inner automorphism. Moreover, $\phi \circ \lambda_1 = \lambda_2$. The proposition follows. \blacksquare

6.3. Goldman manifolds. Consider a u -linear deformation ρ of a canonical Anosov flag representation ρ_0 . Here, we consider that the representations $\Gamma \rightarrow \mathrm{GL}_2$ associated to ρ and ρ_0 are the same, i.e., in the notation of the beginning of § 4.2, $\rho_0(\gamma) = e^{u(\gamma)} \rho_\lambda$.

Proposition 6.4. *The actions of $\rho(\Gamma)$ and $\rho_0(\Gamma)$ on $P(V)$ are topologically conjugate.*

Proof We reproduce the short proof in [1]: there is a (Hölder) continuous and homogeneous degree one map $\delta : \mathbb{R}^2 \rightarrow \mathbb{R}$ such that $[xe_1 + ye_3 + ze_2]$ belongs to the invariant curve L if and only if $z = \delta(x, y)$. The $\rho(\gamma)$ -invariance of L implies:

$$\delta(\rho(\gamma)) = \delta(\gamma) + \tau_1(\gamma)x + \tau_2(\gamma)y$$

where $\tau_1(\gamma) = e^{u(\gamma)}\mu(\gamma)$, $\tau_2(\gamma) = e^{v(\gamma)}\nu(\gamma)$ (for $\mu(\gamma)$, $\nu(\gamma)$ as in § 4.2). It follows immediately that $(x, y, z) \mapsto (x, y, z + \delta(x, y))$ induces on $P(V)$ the required topological conjugacy between $\rho_0(\Gamma)$ and $\rho(\Gamma)$. ■

This proposition implies Theorem 5.1 of [1]: *the first tautological foliations associated to ρ , ρ_0 are topologically conjugate*. It follows also that *the action of $\rho(\Gamma)$ on $P(V) \setminus (L \cup [e_2])$ is minimal and uniquely ergodic*.

6.4. Non hyperconvex Anosov flag manifolds. Anosov flag manifolds with hyperconvex (i.e., quasi-Fuchsian) representations are fairly well understood (see remarks 3.16, 5.8). From now, we exclude this case.

Lemma 6.5. *The complement of L in $P(V)$ is a topological disc.*

Proof If L is a projective line, the lemma is obvious. Assume that L^* is a projective line, i.e., that ρ is a hyperbolic representation: consider an affine domain $U \subset P(V)$ not containing the $\rho(\Gamma)$ -fixed point $[(L^*)^\perp]$. Select a coordinate system on $U \approx \mathbb{R}^2$ such that the intersection with U of projective lines in L^* are vertical lines $\{*\} \times \mathbb{R}$. Then, according to corollary 3.12, $L \cap U$ is the graph of a continuous map from \mathbb{R} into \mathbb{R} . Then, the lemma becomes obvious.

We are left with the case where L, L^* are not projective lines, i.e., the strongly irreducible case (cf. proposition 4.8). Let $S(V)$ be the N -sphere of V : the radial projection $\Pi_S : S(V) \rightarrow P(V)$ is the double covering, with covering automorphisms $\pm id$. The group $SL(V)$ acts naturally on $S(V)$: consider the sphere $S(V)$ as the space of half-lines in V .

The limit curve L is a simple closed curve. There are two cases:

- either $\Pi_S^{-1}(L)$ is a connected simple closed curve \hat{L} ,
- or $\Pi_S^{-1}(L)$ is the union of two connected simple closed curves \hat{L}_+, \hat{L}_- .

In the first case, the lemma follows from Jordan's Theorem, and the $(-id)$ -invariance of \hat{L} . Hence, the proof of the lemma amounts to excluding the second case, more precisely, to proving that in the second case, ρ is hyperconvex.

The antipody $-id$ exchanges \hat{L}_+ and \hat{L}_- . For any element γ of Γ , there is one and only one way to lift $\rho(\gamma)$ to an element $\hat{\rho}(\gamma)$ of $SL(V)$ preserving \hat{L}_+ . It provides a representation $\hat{\rho} : \Gamma \rightarrow SL(V)$.

For any $[u^*]$ in L^* , according to corollary 3.12, the kernel of $u^* \in V^*$ intersects $\hat{L}_+ \cup \hat{L}_-$ in two points, one opposite the other. Since $\hat{L}_- = -\hat{L}_+$, the kernel of u^* intersects \hat{L}_+ in one and only one point $u([u^*])$ of \hat{L}_+ : hence, the sign of u^* on $\hat{L}_+ \setminus \{u([u^*])\}$ is constant. Select $u^* \in S(V^*)$ so that this sign is positive.

This process defines a way to simultaneously lift L and L^* in V, V^* to closed subsets \hat{L}_+, \hat{L}_+^* such that for any (u, v^*) in $\hat{L}_+ \times \hat{L}_+^*$ we have $\langle u | v^* \rangle \geq 0$. In the terminology of [5], this means that Λ is a *positive* subset of X . According to proposition 1.2 of [5], since ρ is strongly irreducible, $\rho(\Gamma)$ preserves a strictly convex domain of $P(V)$, meaning precisely that ρ is hyperconvex, a contradiction. ■

Corollary 6.6. *Every non-hyperconvex Anosov flag representation can be continuously deformed to the trivial representation.*

Proof According to [13], a non-hyperconvex Anosov representation does not belong to the Hitchin component. On the other hand, it lifts to a representation $\hat{\rho} : \Gamma \rightarrow P^+GL(V)$. Indeed, keeping the notation used in the proof of Lemma 6.5, we can select a connected component U^+ of $S(V) \setminus \hat{L}$. We then define $\hat{\rho}(\gamma)$ as the unique lift in $P^+GL(V)$ of $\rho(\gamma)$ preserving U^+ . It provides the required representation $\hat{\rho}$. The corollary then follows from § 2.6. ■

Corollary 6.7. *For every $[u]$ in $P(V)$ there is an element $[v^*]$ in L^* “containing” $[u]$, i.e., such that $\langle u | v^* \rangle = 0$.*

Proof If not, there is an element u_0 of $V \setminus \{0\}$ such that L^* is disjoint from $[u_0^\perp]$. The projective line $[u_0^\perp]$ then lifts to a great circle in $S(V^*)$ avoiding \hat{L}^* . Hence, \hat{L}^* is contained in a hemisphere U . We obtain a contradiction since $\hat{L}^* = -\hat{L}^*$ and $U \cap (-U) = \emptyset$. ■

Compare the following proposition with remark 3.16:

Proposition 6.8. *T and T^* are topological Klein bottles, and A, A^* are Möbius bands. The domain Ω is connected, homeomorphic to the solid torus $\Omega_0 \approx \mathbb{S}^1 \times \mathbb{R}^2$.*

Proof By the Jordan-Schönflies Theorem, there is a homeomorphism f of $P(V)$ mapping the simple closed curve L on the projective line L_0 . The circle bundle $\Pi : X \rightarrow P(V)$ and pull-back bundle $f^*\Pi$ have the same Euler class; hence, f lifts to some homeomorphism F of X into itself, preserving the fibers of Π , and inducing f on $P(V)$. Then, $F(T)$ is the Klein bottle T_0 . The first part of the proposition follows. The complement W_0 of T_0 in $P(V)$ is a solid torus, and $F(T^*) \cap W_0$ is an annulus. For a better visualisation, lift to the 4-sheeted covering $\widehat{X} \subset S(V) \times S(V^*)$: we are led to the well-known fact that the complement, in a compact solid torus with boundary \widehat{W}_0 , of a compact annulus with boundary topologically embedded admitting as boundary two disjoint essential curves of $\partial\widehat{W}_0$, is the union of two solid tori. ■

Corollary 6.9. *The flag manifold $M = \rho(\Gamma) \backslash \Omega$ is homeomorphic to a circle bundle over Σ .*

Proof According to proposition 6.8, if $\widehat{\Gamma}$ denote the fundamental group of M , we have an exact sequence:

$$0 \rightarrow \mathbb{Z} \rightarrow \widehat{\Gamma} \rightarrow \Gamma \rightarrow 0$$

where \mathbb{Z} is the fundamental group of the solid torus Ω . Hence, M is sufficiently large (its first homology group is infinite), is $\mathbb{R}P^2$ -irreducible (its universal covering is homeomorphic to \mathbb{R}^3), and is homotopically equivalent to a circle bundle over Σ . The corollary follows from [20]. ■

6.5. Minimality of tautological foliations. Recall that a 1-dimensional foliation is *minimal* if all the leaves are dense.

Theorem 6.10. *Let ρ be a (G, Y) -Anosov representation, which is not hyperconvex.*

- *If ρ is a canonical flag Anosov representation, then the tautological foliations are both minimal,*
- *If ρ is a linear u -deformation, with $u \neq 0$, then the first tautological foliation is minimal, but the second tautological foliation is not minimal,*
- *If ρ is strongly irreducible, then the tautological foliations are both non-minimal.*

Sketch of proof The first case follows from Proposition 6.4. The first statement in the second case follows from Proposition 6.4 too. The second statement of this case is Theorem C of [1]. The third case can be proved using the arguments of §5.2 of [1]: consider the set of flags $([u], [u^*]) \in \Omega$ such that $[u^*] \cap L$ is reduced to a point. Its closure \mathcal{M} is invariant by the second tautological flow. Arguments of Lemma 5.7 of [1] proves that \mathcal{M} is not all of Ω , and those of Lemma 5.8 of [1] show that if \mathcal{M} were empty, then the second tautological foliation would be expansive: by [7], it would be topologically conjugate to a finite covering of some geodesic flow. This leads to a contradiction, as in [1]. ■

7. THE INVARIANT MÖBIUS BANDS

In this section, we consider a *non hyperconvex* Anosov flag representation ρ . The action of $\rho(\Gamma)$ on Λ is conjugate to the usual projective action on $\mathbb{R}P^1$. The action on Ω is proper, topologically conjugate to the action of Γ on H by left translations, at least when ρ is a deformation of a canonical flag representation inside the space of Anosov flag representations (see question 2 in § 8).

But the dynamics on the invariant Möbius bands are much more subtle. When L is a projective line $[(u_0^*)^\perp]$, it follows from corollary 3.12 that $[u_0^*]$ does not belong to L^* . Hence, the flags $([u], [u_0^*])$ with $[u] \in L$ form a continuous $\rho(\Gamma)$ -invariant curve in A . Conversely:

Proposition 7.1. *Consider the Fuchsian projective action of $\Gamma \subset H$ on the projective line $\mathbb{R}P^1$ and the Γ -action on A induced by ρ . If there is a Γ -equivariant measurable map $\sigma : \mathbb{R}P^1 \rightarrow A$, then the limit curve L is a projective line.*

Proof This is essentially the content of Lemma 4.17 of [1], that we reproduce here: let $\overline{\mathcal{Q}}$ be the complement of the diagonal in $\mathbb{R}P^1 \times \mathbb{R}P^1$ (see § 2.4.2). The diagonal action of Γ on $\overline{\mathcal{Q}}$ is ergodic for some Γ -invariant measure equivalent to the Lebesgue measure and preserved by the flip map $(x, y) \mapsto (y, x)$ (for example, the projection on the orbit space of the geodesic flow of the Liouville measure). We say that a subset of $\overline{\mathcal{Q}}$ is conull if the Lebesgue measure of its complement is zero. The crucial and classical observation is that this ergodicity property implies that there is no measurable equivariant map from $\overline{\mathcal{Q}}$ into a topological space on which Γ acts freely and properly discontinuously.

Decompose $\sigma : \mathbb{R}P^1 \rightarrow A$ into two functions $\eta : \mathbb{R}P^1 \rightarrow L$ and $\eta^* : \mathbb{R}P^1 \rightarrow P(V^*) \setminus L^*$.

Assume that the set of pairs (x, y) for which $\eta(y)$ does not belong to $\eta^*(x)$ is conull. Then, its intersection with its image by the flip map is conull, and its intersection with all its Γ -iterates is too. Thus, there is a conull Γ -invariant subset \mathcal{E} of $\overline{\mathcal{Q}}$ of pairs (x, y) for which the projective lines $\eta^*(x)$ and $\eta^*(y)$ intersect at some point $[u(x, y)]$ different from $\eta(x)$ and $\eta(y)$. We have then two cases: either almost every $[u(x, y)]$ belongs to L , or almost all of them belong to $P(V) \setminus L$. In the first case, $(x, y) \mapsto (\eta_+^{-1}(\eta(x)), \eta_+^{-1}(\eta(y)), \eta_+^{-1}([u(x, y)]))$ is a Γ -equivariant map from \mathcal{E} into the set of distinct triples of points of $\mathbb{R}P^1$. Since the action of Γ on this set of triples is free and properly discontinuous, we obtain a contradiction with the ergodic argument discussed above. In the second case, the map associating to a pair (x, y) the flag $([u(x, y)], \eta^*(x))$ is a Γ -equivariant map from \mathcal{E} into Ω . We obtain once more a contradiction with the ergodic argument by Theorem 5.1.

Therefore, the measure of the set of pairs (x, y) for which the line $\eta^*(x)$ contains $\eta(y)$ is conull. Then, by Fubini's Theorem, for almost every x in $\mathbb{R}P^1$ and for almost all y in $\mathbb{R}P^1$, $\eta(y)$ belongs to $\eta^*(x)$. For such a x , if there exists an element γ of Γ such that $\eta^*(x) \neq \rho^*(\gamma)\eta^*(x)$, then for almost every y , $\eta(y)$ is the unique point of intersection $[u_0]$ between $\eta^*(x)$ and $\rho^*(\gamma)\eta^*(x)$. Hence, $[u_0]$ must be a common fixed point of all $\rho(\Gamma)$, but this is impossible since it belongs to L .

It follows that $\eta^*(x) = \rho^*(\gamma)\eta^*(x)$ for every γ : there is a $\rho(\Gamma)$ -invariant projective line in $P(V)$. Hence, ρ is hyperbolic, and L is the projective line $\eta^*(x)$. ■

Of course, the similar lemma with A replaced by A^* is true.

Corollary 7.2. *If ρ is strongly irreducible, the maps η_\pm are not Lipschitz.*

Proof Assume that η_+ is Lipschitz. Then, it is differentiable almost everywhere. Its differential defines a measurable map $\sigma = (\eta, \eta^*) : \mathbb{R}P^1 \rightarrow X$, where the first component η is η_+ , and the second component η^* is the projective line tangent to the image of the differential of η_+ . According to Proposition 7.1, the second component

$\eta^*(x)$ is an element of L^* for almost every x . According to Corollary 3.12, we have $\eta^*(x) = \eta_-(x)$. More precisely, it follows from this corollary 3.12 that L is locally strictly convex. We obtain a contradiction since ρ was assumed to be non-hyperconvex.

The same proof applies to η_- . ■

8. OPEN QUESTIONS

Let $\rho : \Gamma \rightarrow G$ be a non-hyperconvex Anosov flag representation. Here, we have essentially answered Question 8 of [1] by the Theorem 5.1. We also answered Question 6: the curve Λ is Hölder continuous. But some interesting questions are still open:

Question 1 *Is the circle bundle $\rho(\Gamma)\backslash\Omega$ homeomorphic to $\Gamma\backslash H$, i.e., to the double covering of the unit tangent bundle of $\Sigma = \Gamma\backslash\mathbb{H}^2$?*

According to [13], the space of hyperconvex flag representations is connected: it is the entire Hitchin component. Thus, we can wonder:

Question 2 *Is the space of non-hyperconvex Anosov flag representations connected?*

If the answer to question 2 is yes, then the answer to question 1 is yes.

Question 3 *Let $\rho' : \Gamma \rightarrow G$ be another non-hyperconvex Anosov flag representation. Assume that the $\rho'(\Gamma)$ -action on the invariant Möbius band A' (respectively $(A')^*$) is topologically conjugate to the $\rho(\Gamma)$ -action on A (resp. A^*). Does that imply that ρ' and ρ are conjugate in G ?*

Observe that according to Proposition 6.3, Proposition 6.4 and Proposition 7.1, the answer to this question is yes when ρ is a hyperbolic radial representation!

Question 4 *Are the tautological foliations associated to non-hyperconvex Anosov flag representations uniquely ergodic?*

Concerning dynamical properties of tautological foliations, recall Questions 1 and 5 of [1]:

Question 5 *Can a tautological foliation associated to a non-hyperconvex Anosov flag representation admit a periodic orbit? Can it have non-zero entropy?*

APPENDIX A: THE (G, Y) -ANOSOV PROPERTY FOR LINEAR DEFORMATIONS.

In this Appendix, we prove that canonical representations and their u -deformations are (G, Y) -Anosov. We first consider canonical representations. The proof of Theorem 2.11 we produce here is quite sophisticated, but it is a necessary preparation for the most delicate case of u -deformations.

8.1. Canonical representations. Consider the following map $f : H \rightarrow Y$: $f(h) = ([\rho_0(h)e_1], [e_2], [\rho_0(h)e_3])$. It is a $\rho_0(\Gamma)$ -equivariant map, defining a section s of π_ρ . Our task is to prove that s defines a (G, Y) -Anosov structure.

Consider $f_+ = \pi_+ \circ f$: it provides a section s^+ of $\pi_{\rho_0}^X$. It can be written: $f_+(h) = ([\rho_0(h)e_1], [\rho_0^*(h)e_3^*])$.

We first consider the first component $[\rho_0(h)e_1]$, which defines a section s_P^+ of the flat $P(V)$ -bundle associated to ρ_0 : this bundle $\pi_P : E_{\rho_0}(P) \rightarrow M$ is defined in the same way as the bundles E_{ρ_0} and $E_{\rho_0}(X)$; it also admits a horizontal flow Φ_P^t above Φ^t .

Define $\Psi(h, (\alpha, \beta)) = (h, [\rho_0(h)e_1 + \alpha\rho_0(h)e_2 + \beta\rho_0(h)e_3])$. It is a Γ -equivariant map, when $H \times \mathbb{R}^2$ is equipped with the Γ -action $\gamma(h, (\alpha, \beta)) = (\gamma h, (\alpha, \beta))$, and $H \times P(V)$ is equipped with the action $\gamma(h, [u]) = (\gamma h, [\rho_0(\gamma)u])$. Hence, it induces a fibered map $\tilde{\Psi} : M \times \mathbb{R}^2 \rightarrow E_{\rho_0}(P)$. More precisely, it provides a trivialisation of an open neighborhood W of the image of s_P^+ .

Select any left invariant metric m on H . Equip \mathbb{R}^2 with the euclidean norm $d\alpha^2 + d\beta^2$. The image by Ψ of the product metric is a Γ -invariant metric: it provides a metric on the neighborhood W . Now, we simply observe that the flow Φ_P^t is expressed in the chart $\bar{\Psi}$ by the simple expression:

$$\Phi_P^t(p, (\alpha, \beta)) = (\Phi^t(p), (e^t\alpha, e^{2t}\beta))$$

It immediately follows that the image of s_P^+ is a (exponentially) repeller of Φ_P^t .

Similar reasoning on the second component $[\rho_0^*(h)e_3^*]$ of $f_+(h)$ shows that it provides a section of the flat $P(V^*)$ -bundle associated to ρ_0^* which is also a (exponentially) repeller of the corresponding horizontal flow.

Combining these two facts, we obtain that *the image of s_P^+ is an (exponentially) attractor for Φ_X^t* .

A completely similar argument establishes that the image of the section s^- of $E_{\rho_0}(X)$ furnished by $\pi_- \circ f$ is an (exponentially) attractor for Φ_X^t . Of course, the pairs $(s^+(p), s^-(p))$ all belong to \mathcal{Y}_{ρ_0} . It follows that $s \approx (s^+, s^-)$ satisfies all the criteria required in definition 2.10: it defines a (G, Y) -Anosov structure. Theorem 2.11 is proved.

8.2. Linear u -deformations. For a fixed inclusion $\iota : \Gamma \subset H$ and embedding $\rho_0 : H \rightarrow G$, any $u \in H^1(\Gamma, \mathbb{R})$ defines a representation ρ_u . We assume here that the stable norm $|u|_s$ is less than $1/2$, and we want to prove that ρ_u is (G, Y) -Anosov.

Since φ^t preserves the point $[e_2]$ and acts trivially on $L = [(e_2^*)^\perp]$, the map $f(h) = ([\rho_0(h)e_1], [e_2], [\rho_0(h)e_3])$ is still $\rho_u(\Gamma)$ -equivariant: it provides the suitable section s .

Exactly as we did for canonical representations, we study separately the two components $E_{\rho_u}(X)$. Each of them is then decomposed in bundles $E_{\rho_u}(P)$, $E_{\rho_u}(P^*)$. At the end, we have to consider 4 sections of bundles over M by projective spaces, and we have to prove that the images of these sections are repellers or attractors of the associated horizontal flows.

We only discuss here the section of $E_{\rho_u}(P)$ defined by $h \mapsto [\rho_0(e_1)]$. The other sections can be treated in a similar way left to the reader.

We consider once more the map $\Psi(h, (\alpha, \beta)) = (h, [\rho_0(h)e_1 + \alpha\rho_0(h)e_2 + \beta\rho_0(h)e_3])$. The main difference with the canonical case is that the Γ -action on $H \times \mathbb{R}^2$ to be considered is:

$$(3) \quad \gamma(h, (\alpha, \beta)) = (\gamma h, (e^{-u(\gamma)}\alpha, \beta))$$

The quotient of this action is then an \mathbb{R}^2 -bundle E over M , and we have a fibered map $\bar{\Psi} : E \rightarrow E_{\rho_u}(P)$. The image of $\bar{\Psi}$ is a neighborhood of the Φ_P^t -invariant section to be studied.

The key point is to define a metric on $H \times \mathbb{R}^2$ such that:

- the Γ -action defined by (3) is isometric,
- the null section $h \mapsto (h, 0)$ is a repeller for the horizontal flow $(h, (\alpha, \beta)) \mapsto (h\alpha^t, (e^t\alpha, e^{2t}\beta))$.

Let Σ be the Riemannian surface $\Gamma \backslash \mathbb{H}^2$. The cohomology class $u \in H^1(\Gamma, \mathbb{R})$ can be represented by a 1-form ω on Σ such that the integration of ω along any loop representing an element γ of $\Gamma \approx \pi_1(\Sigma)$ is $u(\gamma)$.

The quotient $\Gamma \backslash \bar{H}$ is naturally identified with the unit tangent bundle of Σ : the orbit space of the left action of $SO(2)$ on \bar{H} is canonically identified with Σ . Denote by $\eta : \Gamma \backslash \bar{H} \rightarrow \Gamma \backslash \bar{H}/SO(2)$ the quotient map, and consider the 1-form

$\eta^*(\omega)$. The assumption $|u|_s < 1/2$ implies the following³: *the absolute value of the integration of $\eta^*(\omega)$ along a periodic orbit of $\overline{\Phi}^t$ with period T is less than CT , for $C = 2|u|_s < 1$. Any orbit of Φ^t can be approximated by periodic orbits. Therefore, if $\bar{\theta} : [0, T] \rightarrow \overline{M}$ is an portion of an orbit of $\overline{\Phi}^t$ (i.e., $\bar{\theta}(t) = \overline{\Phi}^t(p)$ for some p), we have:*

$$\left| \int_{\bar{\theta}} \eta^*(\omega) \right| \leq CT$$

Let $\hat{\omega}$ be the lifting of $\eta^*(\omega)$ to $M = \Gamma \backslash H$: if $\theta(t) = \Phi^t(p)$, we have:

$$(4) \quad \left| \int_{\theta} \hat{\omega} \right| \leq CT$$

Finally, the lifting $\tilde{\omega}$ in H of $\hat{\omega}$ is exact: there is a function $\nu : H \rightarrow \mathbb{R}$ such that $d\nu = \tilde{\omega}$. According to (4), we have:

$$(5) \quad |\nu(ha^t) - \nu(h)| \leq Ct$$

Equip the fiber \mathbb{R}^2 over $h \in H$ with the metric $e^{2\nu(h)} d\alpha^2 + d\beta^2$: it gives a metric on $H \times \mathbb{R}^2$. Since $\nu(\gamma h) - \nu(h) = \int_{[h, \gamma h]} d\nu = \int_{[h, \gamma h]} \tilde{\omega} = u(\gamma)$, the transformations defined in (3) are isometries. Now, it follows from (5) that the horizontal flow $(h, (\alpha, \beta)) \mapsto (ha^t, (e^t\alpha, e^{2t}\beta))$ expands the norm on the fibers by a factor at least $e^{(1-C)t}$. Theorem 4.2 follows.

APPENDIX B: PROPERNESS OF THE ACTION IN THE RADIAL CASE.

In this appendix, we give a proof of Theorem 5.1 in the non strongly irreducible case, i.e., according to remark 5.2, in the case of hyperbolic representations. We use all the notation and conventions introduced in § 5. In this section, we obtained a contradiction many times by exhibiting an invariant projective line. This event cannot anymore be considered as a contradiction in the radial case: it slightly complicates the proof.

On the other hand, in the hyperbolic radial case, we have of course additional properties: every g_n (resp. g_n^*) stabilizes $[e_2]$ (resp. $[e_2^*]$). More precisely, e_2, e_2^* are eigenvectors for the intermediate eigenvalues μ_n, μ_n^{-1} . Hence, the N -isometries k_n, l_n can be selected so that they both preserve $[e_2]$: $k_n e_2 = \pm e_2, l_n e_2 = \pm e_2$.

Let's reproduce the proof in § 5, supplying the needed adaptations. Of course, Lemma 5.3 still applies here. Let's consider the analog of Lemma 5.5:

Lemma 8.1. *The coefficients $\bar{\mu}$ and $\bar{\mu}'$ are both zero.*

Proof. Assume $\bar{\mu} \neq 0$. We prove as in 5.5 the identities:

$$I_* = I^\perp, \quad K^\perp = K_*$$

Moreover, $[K]$ and $[I_*]$ are single points.

Fact 1: $[K]$ belongs to the invariant curve L . If not, the g_n -invariant curve L is contained in $P(V) \setminus [K]$. By Lemma 5.3, it implies that L is equal to the projective line $[I]$, which is therefore a $\rho(\Gamma)$ -invariant projective line. In particular, $[K]$ does not belong to $[I]$. Let D be a small open disk containing $[K]$, but with closure disjoint from $[I] = L$. By Lemma 5.3, the union of the iterates $g_n D$ is the complement of $[I]$ in $P(V)$. Hence, one of them contains the closure of D : it follows that D contains the attractive fixed point of some g_n^{-1} . Taking arbitrarily small D ,

³Recall that $\overline{\Phi}^t$ is the geodesic flow with parametrisation multiplied by 2 (see remark 2.5): this factor 2 compensates for the stable norm $1/2$.

we obtain that $[K]$ is the limit of a sequence of attractive fixed points of elements of $\rho(\Gamma)$. By corollary 3.14, $[K]$ belongs to L , a contradiction.

Fact 1': $[I_*]$ belongs to L^* . Same proof as for Fact 1, applied to the inverse sequence $(h^* = (g_n^*)^{-1})_{(n \in \mathbb{N})}$.

Fact 2: The invariant curve L is a projective line containing $[K]$. Same proof as fact 2 in 5.5.

Fact 2': The invariant curve L^* is a projective line containing $[I_*]$. The proof is the same as for Fact 2, the arguments applied to g_n above to the inverse sequence $h_n^* = (g_n^*)^{-1}$.

Conclusion. By facts 2 and 2', the invariant curves are projective lines. It follows that ρ is actually the u -deformation of a canonical Anosov representation:

$$\rho(\gamma) = \begin{pmatrix} e^{u(\gamma)/3}a(\gamma) & 0 & e^{u(\gamma)/3}b(\gamma) \\ 0 & e^{-2u(\gamma)/3} & 0 \\ e^{u(\gamma)/3}c(\gamma) & 0 & e^{u(\gamma)/3}d(\gamma) \end{pmatrix}$$

where $\rho_\lambda : \gamma \mapsto \begin{pmatrix} a(\gamma) & b(\gamma) \\ c(\gamma) & d(\gamma) \end{pmatrix}$ is a fuchsian morphism into $\mathrm{SL}(2, \mathbb{R})$, and $u : \Gamma \rightarrow \mathbb{R}$ a morphism. According to Proposition 4.8, the stable norm of u satisfies $|u|_s < 1/2$.

Lemma 8.1 is then one of the intermediate results of [17]: indeed, let $l(\gamma)$ be as in [17] the logarithm of the spectral radius of the diagonal matrix appearing in the Cartan decomposition of γ in $\mathrm{SL}(2, \mathbb{R})$. Then, the logarithms of the coefficients λ_n , μ_n and ν_n are, modulo some common additive constant, the quantities $u(\gamma_n) + l(\gamma_n)$, $u(\gamma_n)$ and $u(\gamma_n) - l(\gamma_n)$. In the proof of Théorème 3.4 in [17], it is shown that $l(\gamma_n) - |u(\gamma_n)|$ tends to $+\infty$: it precisely means $\bar{\mu} = 0$. ■

Hence, according to Lemma 8.1, $[K]$ and $[K_*]$ are projective lines, and $[I]$, $[I_*]$ are single points.

Lemma 8.2. *The points $[I]$, $[I_*]$ belong to L , L^* , respectively.*

Proof Assume that $[I]$ does not belong to L . Then, by Lemma 5.3, L is contained in the projective line $[K]$. Hence, $L = [K]$. By considering small discs around $[I]$, we obtain, as in the proof of Fact 1 in Lemma 5.5, that $[I]$ is the limit of a sequence of attractive fixed points of elements of $\rho(\Gamma)$. Together with Corollary 3.14, this leads to a contradiction.

A similar proof gives $[I_*] \in L^*$. ■

Lemma 8.3. *The points $[K^\perp]$ and $[K_*^\perp]$ belong to L^* , L respectively.*

Proof This is the dual version of Lemma 8.2: apply the reasoning above to the inverse sequence h_n . ■

The proof of Theorem 5.1 in the reducible case follows then from Lemma 5.3 and Lemmas 8.2, 8.3, exactly as the irreducible case followed in § 5 from Lemmas 5.6, 5.7 (and Lemma 5.3).

REFERENCES

- [1] T. Barbot, *Flag structures on Seifert manifolds*, *Geom. Topol.*, **5** (2001), 227–266, arXiv:math.DS/0104108.
- [2] T. Barbot, *Plane affine geometry and Anosov flows*, *Ann. Scient. Éc. Norm. Sup.*, **34** (2001), 871–889.
- [3] T. Barbot, *Feuilletages transversalement projectifs sur les variétés de Seifert*, *Ann. Inst. Fourier*, **53** (2003), 1551–1613.
- [4] V. Bangert, *Minimal geodesics*, *Ergod. Th. & Dynam. Sys.* **10** (1989), 263–286.
- [5] Y. Benoist, *Automorphismes des cônes convexes*, *Invent. Math.* **141** (2000), 149–193.

- [6] R. Bowen, B. Marcus, *Unique ergodicity for horocyclic foliations*, Israel J. Math., **26** (1977), 43–67.
- [7] M. Brunella, *Expansive foliations on Seifert manifolds and on torus bundles*, Bol. Soc. Brasil. Math., **24** (1993), 89–104.
- [8] S. Choi, W.M. Goldman, *Convex real projective structures on closed surfaces are closed*, Proc. Am. Math. Soc., **118** (1993), 657–661.
- [9] A. Fathi, F. Laudenbach, V. Poenaru, *Travaux de Thurston sur les surfaces*, Astérisque, **66-67** (1979).
- [10] E. Ghys, *Flots d’Anosov dont les feuilletages stables sont différentiables*, Ann. Sci. École Norm. Sup. (4), **20** (1987), 251–270.
- [11] V.L. Ginzburg, B.Z. Gürel, *A C^2 -smooth counterexample to the Hamiltonian Seifert conjecture in \mathbb{R}^4* , Ann. of Math. (2), **158** (2003), no. 3, 953–976, arXiv:math.DG/0110047.
- [12] W.M. Goldman, *Geometric structures on manifolds and varieties of representations*, in “Geometric of group representations”, Contemp. Math., **74** (1988), 169–198.
- [13] O. Guichard, *Sur les représentations des groupes de surface*, preprint (2004).
- [14] K. Kuperberg, *A smooth counterexample to the Seifert conjecture*, Ann. of Math. (2), **140** (1994), no. 3, 723–732.
- [15] G. Kuperberg, *A volume-preserving counterexample to the Seifert conjecture*, Comment. Math. Helv., **71** (1996), no. 1, 70–97, arXiv:math.DS/9504230.
- [16] F. Labourie, *Anosov flows, Surface Groups and Curves in projective Space*, arXiv:math.DG/0401230.
- [17] F. Salein, *Variétés anti-de Sitter de dimension 3 possédant un champ de Killing non trivial*, C. R. Acad. Sci. Paris Sér. I Math., **324** (1997), 525–530.
- [18] R. Schwartz, *Pappus’s theorem and the modular group*, Publ. Math. I.H.E.S., **78** (1993), 187–206.
- [19] W. Thurston, *Three dimensional geometry and topology*, Princeton University Press (1997).
- [20] F. Waldhausen, *On irreducible 3-manifolds which are sufficiently large*, Ann. of Math. (2), **87** (1968), 56–88.

CNRS, UMPA, UMR 5669
École Normale Supérieure de Lyon,
46, allée d’Italie
69364 Lyon cedex 07, FRANCE