Non subanalyticity of sub-Riemannian Martinet spheres

Emmanuel Trélat

► To cite this version:

Emmanuel Trélat. Non subanalyticity of sub-Riemannian Martinet spheres. Comptes rendus de l'Académie des sciences. Série I, Mathématique, 2001, 332, pp.527–532. hal-00086298

HAL Id: hal-00086298

<https://hal.science/hal-00086298>

Submitted on 18 Jul 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Non subanalyticity of sub-Riemannian Martinet spheres

Emmanuel TRÉLAT ^a

^a Laboratoire de Topologie, Université de Bourgogne, 9 av. A. Savary, BP 47870, 21078 Dijon
Courriel : trelat@topolog.u-bourgogne.fr

(Reçu le ???, accepté le ???)

Abstract. Consider the sub-Riemannian Martinet structure (M, Δ, g) where $M = \mathbb{R}^3$, $\Delta = \text{Ker}(dz - \frac{y^2}{2}dx)$ and g is the general gradated metric of order 0 : $g = (1 + \alpha y)^2 dx^2 + (1 + \beta x + \gamma y)^2 dy^2$. We prove that if $\alpha \neq 0$ then the sub-Riemannian spheres $S(0, r)$ with small radii are not subanalytic. © Académie des Sciences/Elsevier, Paris

Non sous-analyticité des sphères sous-Riemanniennes de Martinet

Résumé. Considérons la structure sous-Riemannienne de Martinet (M, Δ, g) où $M = \mathbb{R}^3$, $\Delta = \text{Ker}(dz - \frac{y^2}{2}dx)$ et g est la métrique générale graduée d'ordre 0 : $g = (1 + \alpha y)^2 dx^2 + (1 + \beta x + \gamma y)^2 dy^2$. On montre que si $\alpha \neq 0$, les sphères sous-Riemanniennes $S(0, r)$ de petit rayon ne sont pas sous-analytiques. © Académie des Sciences/Elsevier, Paris

Version française abrégée

Une *structure sous-Riemannienne* (M, Δ, g) est constituée d'une variété M de dimension finie, d'une distribution Δ sur M , et d'une métrique g sur Δ . Elle est dite analytique si tous les objets sont analytiques. Une courbe γ sur M est dite *horizontale* si elle est tangente en tout point à la distribution Δ . Soient x_0, x_1 des points de M , et $T > 0$. Le *problème sous-Riemannien* est de déterminer une courbe horizontale joignant x_0 à x_1 en temps T et minimisant la longueur (au sens de la métrique g). La *distance sous-Riemannienne* $d_{SR}(x_0, x_1)$ entre x_0 et x_1 est la borne inférieure des longueurs des chemins horizontaux joignant x_0 à x_1 en temps T . La *sphère sous-Riemannienne* $S(x_0, r)$ centrée en x_0 , de rayon r , est l'ensemble des points x de M tels que $d_{SR}(x_0, x) = r$.

Reformulons le problème sous-Riemannien dans le cadre de la théorie du contrôle optimal. Notre point de vue étant local, on peut supposer que $M = \mathbb{R}^n$ et $x_0 = 0$. Supposons que la distribution Δ est engendrée par m champs de vecteurs $X_1, \dots, X_m$. Sans perte de généralité, on peut supposer que $T = 1$. Le problème sous-Riemannien est alors équivalent au problème de *contrôle optimal* pour le système (voir par exemple [5]) :

$$\dot{x}(t) = \sum_{i=1}^m u_i(t) X_i(x(t)), \quad x(0) = 0$$

Note présentée par ???

où la fonction $u = (u_1, \dots, u_m)$, appelée le *contrôle*, doit minimiser la *longueur*

$$l(u) = \int_0^1 \sqrt{g(\dot{x}(t), \dot{x}(t))} dt$$

L'application $E : u \in L^2([0, 1], \mathbb{R}^m) \mapsto x_u(1) \in \mathbb{R}^n$, qui à un contrôle u associe l'extrémité de la solution correspondante x_u , est appelée *application entrée/sortie*. C'est une application lisse.

Si un contrôle u minimise la longueur l , alors il existe un *multiplicateur de Lagrange* $(\psi, \psi^0) \in \mathbb{R}^n \times \mathbb{R}$ défini à scalaire près tel que

$$\psi.dE(u) = -\psi^0 dl(u)$$

Si $\psi^0 \neq 0$, le contrôle (ou la trajectoire associée) est dit *normal* ; sinon il est dit *anormal*. Un contrôle anormal est une singularité de l'application entrée/sortie.

Dans [1] il est prouvé que s'il n'existe pas de trajectoire anormale minimisante non triviale partant de 0 alors la distance sous-Riemannienne à 0, $d_{SR}(0, \cdot)$, est sous-analytique en dehors de 0, et donc les sphères sous-Riemanniennes $S(0, r)$ de petit rayon sont sous-analytiques. Cette situation est générique pour des distributions de rang ≥ 3 , voir [3]. Cependant les anormales minimisantes apparaissent génériquement pour des distributions de rang 2, et on conjecture que dans ce cas les sphères sous-Riemanniennes ne sont jamais sous-analytiques. Remarquons que dans [3] les auteurs prouvent que l'existence d'une anormale minimisante est responsable de la non sous-analyticité du germe en 0 de $d_{SR}(0, \cdot)$. Ici la conjecture concerne les sphères sous-Riemanniennes.

Cette conjecture a été prouvée dans [2] dans le cas dit de *Martinet plat*, mais il n'est pas générique. Ici on montre que cette conjecture est vraie pour une métrique générale graduée d'ordre 0. Le cas Martinet est le cas sous-Riemannien le plus simple pour lequel une anormale minimisante apparaît génériquement. La situation est la suivante. Considérons dans $\mathbb{R}^3$ la distribution $\Delta = \text{Ker}(dz - \frac{y^2}{2}dx)$; elle est engendrée par les deux champs de vecteurs

$$X = \frac{\partial}{\partial x} + \frac{y^2}{2} \frac{\partial}{\partial z}, \quad Y = \frac{\partial}{\partial y}$$

Les variables x, y sont de poids 1, et z est de poids 3 (voir [5]). Soit g une métrique analytique sur Δ ; d'après [2] une forme normale générale graduée d'ordre 0 est :

$$g = (1 + \alpha y)^2 dx^2 + (1 + \beta x + \gamma y)^2 dy^2$$

où $\alpha, \beta, \gamma \in \mathbb{R}$. C'est une perturbation du cas plat où $\alpha = \beta = \gamma = 0$. L'unique direction anormale correspond à l'axe x ; elle est minimisante [4].

THÉORÈME 1. – *Si $\alpha \neq 0$ alors les sphères sous-Riemanniennes $S(0, r)$ de petit rayon ne sont pas sous-analytiques.*

1. Introduction

1.1. Sub-Riemannian geometry

A *sub-Riemannian structure* (M, Δ, g) is a triple, where M is a finite dimensional manifold, Δ is a distribution on M , and g is a metric on Δ . It is said analytic if all objects are analytic. A curve γ on M is called *horizontal* if it is tangent at any point to the distribution Δ . Let x_0, x_1 in M ,

and $T > 0$. The *sub-Riemannian problem* is to determine an horizontal curve joining x_0 and x_1 in time T and minimizing the length (in the sense of the metric g). The *sub-Riemannian distance* $d_{SR}(x_0, x_1)$ between x_0 and x_1 is the infimum of lengths of horizontal paths joining x_0 and x_1 in time T . The *sub-Riemannian sphere* $S(x_0, r)$ centered at x_0 , with radius r , is the set of points x in M such that $d_{SR}(x_0, x) = r$.

1.2. Necessary optimality conditions

In order to determine optimality conditions for an horizontal curve, we shall reformulate the sub-Riemannian problem in optimal control theory. Our point of view is local, hence we can suppose that $M = \mathbb{R}^n$ et $x_0 = 0$. Assume that the distribution Δ is spanned by m vector fields $X_1, \dots, X_m$. Without loss of generality, we may suppose that $T = 1$. Then the sub-Riemannian problem is equivalent to the *optimal control problem* for the system (see for instance [5]) :

$$\dot{x}(t) = \sum_{i=1}^m u_i(t) X_i(x(t)), \quad x(0) = 0$$

where the function $u = (u_1, \dots, u_m)$, called the *control*, has to minimize the *length*

$$l(u) = \int_0^1 \sqrt{g(\dot{x}(t), \dot{x}(t))} dt$$

The mapping $E : u \in L^2([0, 1], \mathbb{R}^m) \longrightarrow x_u(1) \in \mathbb{R}^n$, which to a control u associates the extremity of the corresponding solution x_u , is called *end-point mapping*. It is smooth.

If a control u minimizes the length l , then there exists a *Lagrange multiplier* $(\psi, \psi^0) \in \mathbb{R}^n \times \mathbb{R}$ defined up to a scalar such that

$$\psi \cdot dE(u) = -\psi^0 dl(u)$$

Moreover from the *Pontryagin's Maximum Principle* (see [9]) the trajectory x corresponding to this minimizing control u is the projection of an *extremal*, i.e. a solution of the Hamiltonian system :

$$\dot{x} = \frac{\partial H}{\partial p}(x, p, u), \quad \dot{p} = -\frac{\partial H}{\partial x}(x, p, u), \quad \frac{\partial H}{\partial u}(x, p, u) = 0$$

where $H(x, p, u) = \langle p, \sum_{i=1}^m u_i X_i(x) \rangle + p^0 g(\dot{x}(t), \dot{x}(t))$ is the Hamiltonian function, p is an absolutely continuous function on $[0, 1]$ called *adjoint vector*, and p^0 is a constant. Moreover we have, up to a scalar : $(p(1), p^0) = (\psi, \psi^0)$. If $p^0 \neq 0$, the extremal is said to be *normal*, and we can normalize to $p^0 = -\frac{1}{2}$; otherwise it is said *abnormal*. An abnormal control is a singularity of the end-point mapping. An abnormal trajectory is said to be *strict* if it is not the projection of a normal extremal.

1.3. Abnormal minimizers and subanalyticity

In [1] it is proved that if there exists no (non trivial) abnormal minimizer starting from 0 then the sub-Riemannian distance to 0, $d_{SR}(0, \cdot)$, is subanalytic outside 0, and hence sub-Riemannian spheres $S(0, r)$ with small radii are subanalytic. This happens generically for distributions of rank ≥ 3 , see [3]. However abnormal minimizers may appear generically for rank 2 distributions, and we conjecture that in this case sub-Riemannian spheres are never subanalytic. Note that in [3]

authors prove that the existence of an abnormal minimizer is responsible for non subanalyticity of the germ at 0 of $d_{SR}(0, \cdot)$. Here the conjecture concerns sub-Riemannian spheres.

This conjecture was proved in [2] in the so-called *Martinet flat case*, but this case is not generic. Here we prove that this conjecture is true for a general gradated metric of order 0. The Martinet case is the easier sub-Riemannian case where an abnormal minimizer appears generically. The situation is the following. Consider in $\mathbb{R}^3$ the distribution $\Delta = \text{Ker}(dz - \frac{y^2}{2}dx)$; it is spanned by the two vector fields

$$X = \frac{\partial}{\partial x} + \frac{y^2}{2} \frac{\partial}{\partial z}, \quad Y = \frac{\partial}{\partial y}$$

Variables x, y have weight 1, and z has weight 3 (see [5]). Let g be an analytic metric on Δ ; from [2] a general gradated normal form of order 0 is :

$$g = adx^2 + cdy^2$$

where $a = (1 + \alpha y)^2$, $c = (1 + \beta x + \gamma y)^2$, and α, β, γ are real parameters. It is a perturbation of the flat case where $\alpha = \beta = \gamma = 0$. There exists an unique abnormal direction, corresponding to the axis x ; it is minimizing, see [4]. Moreover it is *strict* if and only if $\alpha \neq 0$ (see [2]).

THEOREM 1. – *If $\alpha \neq 0$ (i.e. in the strict case), the sub-Riemannian spheres $S(0, r)$ with small radii are not subanalytic.*

The main idea of the proof is the following. The interest of the previous metric of order 0 is all in the fact that the problem, formulated in the cotangent space of $\mathbb{R}^3$, projects onto the phase plane of a *one-parameter pendulum*. This fact was already used in [2], [6]. In [2], in order to prove that the sphere in the Martinet flat case is not subanalytic, the authors make a direct computation of the sphere. But this is impossible in the general case because equations of extremals given by the Maximum Principle *are no more integrable*. Actually such computations lead to asymptotic expansions of the sphere near the abnormal direction (see [6]), but these expansions are not precise enough to check non subanalyticity. The new idea here is rather *indirect* and consists in using Lagrange multipliers : then proving the theorem amounts to estimating precisely a *first return time near a saddle point* of this one-parameter pendulum and proving that it is not subanalytic.

2. Proof of Theorem 1

Consider for the system $\dot{q} = uX(q) + vY(q)$, $q(0) = 0$, the abnormal control $u = -r, v = 0$. It is always minimizing, see [4]. In order to prove the theorem we shall prove that *the intersection of the sphere $S(0, r)$ with the plane $y = 0$ is not a subanalytic curve near the abnormal direction (Ox)* . Let $A = (-r, 0, 0)$ be the extremity of the abnormal extremal associated to $u = -r, v = 0$; it belongs to the sub-Riemannian sphere $S(0, r)$ with radius r . Consider near A the curve $\mathcal{C} = S(0, r) \cap (y = 0) \cap (z \leq 0)$. If $\alpha \neq 0$, i.e. if the abnormal is *strict*, this curve is tangent at A to $z = 0$ with a contact of order 2 (see [6] and [11] for a general result). Consider the set of Lagrange multipliers $(\psi(q), \psi^0(q))$ associated to points $q \in \mathcal{C} \setminus \{A\}$. Since these points are reached by normal minimizers, and since Lagrange multipliers are defined up to a scalar, we may suppose that their component ψ^0 is equal to $-1/2$. On the other part we easily get from the definition of Lagrange multipliers that vectors $(\psi_x(q), \psi_z(q))$, $q \in \mathcal{C}$ (where ψ_x, ψ_z denote respectively the x and z -components of ψ), are normal to the curve $\mathcal{C}$ in the plane $y = 0$. Hence to prove that the germ at A of the curve $\mathcal{C}$ is not subanalytic, it is enough to prove that the set $\mathcal{L} = \{(\psi_x(q), \psi_z(q)) / q \in \mathcal{C}\}$ is not subanalytic.

The method is the following. First note that $\psi(q)$ is equal to $p(1)$ where $p(\cdot)$ is the adjoint vector associated to the normal minimizer steering 0 to q . Hence to compute Lagrange multipliers we have

to estimate adjoint vectors at final time $t = 1$. The Maximum Principle gives us a parametrization of normal extremals and leads to the following differential system in the Martinet case :

$$\begin{aligned} \dot{x} &= \frac{1}{a}(p_x + p_z \frac{y^2}{2}), \quad \dot{y} = \frac{p_y}{c}, \quad \dot{z} = \frac{y^2}{2a}(p_x + p_z \frac{y^2}{2}) \\ \dot{p}_x &= \frac{p_y^2 c_x}{2c^2} + \frac{(p_x + p_z \frac{y^2}{2})^2}{2a^2} a_x, \quad \dot{p}_y = \frac{p_y^2 c_y}{2c^2} + \frac{(p_x + p_z \frac{y^2}{2})^2}{2a^2} a_y - \frac{(p_x + p_z \frac{y^2}{2})}{a} p_z y, \quad \dot{p}_z = 0 \end{aligned} \quad (1)$$

We investigate the Lagrange multipliers $(p_x(1), p_z(1))$ as $y(1) = 0$ and $z(1) \leq 0$. Note that p_z is constant along an extremal. Near the point A , it happens that $p_z \rightarrow +\infty$; it is a phenomenon of *non-properness* due to the existence of an abnormal minimizer, see [11]. From [5], extremal controls satisfy $au^2 + cv^2 = r^2$. Then we can set : $r \cos \theta = \sqrt{a}u = \frac{p_x + \frac{y^2}{2}p_z}{\sqrt{a}}$, $r \sin \theta = \sqrt{c}v = \frac{p_y}{\sqrt{c}}$. In particular we introduce θ which is an angular coordinate of the adjoint vector. This (*analytic*) cylindric change of coordinates on the adjoint vector is classical in sub-Riemannian geometry.

Now proving that the set $\mathcal{L}$ is not subanalytic amounts to exhibiting a non subanalytic relation between $\theta(1)$ and p_z as $p_z \rightarrow +\infty$. We proceed as follows. Reparametrizing with $ds = \frac{r\sqrt{p_z}}{\sqrt{a}\sqrt{c}}dt$, first note that, due to the particular form of the metric, the previous equations (1) project onto a *pendulum equation* :

$$\frac{d^2\theta}{ds^2} + \sin \theta + \frac{\alpha^2}{p_z} \sin \theta \cos \theta - \frac{\alpha\beta}{p_z} \sin^2 \theta + \frac{\beta}{\sqrt{p_z}} \cos \theta \frac{d\theta}{ds} = 0 \quad (2)$$

It is a *pendulum perturbed* by the small parameter $\varepsilon = \frac{1}{\sqrt{p_z}}$. In this pendulum representation the plane $y = 0$ projects onto the curve Σ :

$$\frac{d\theta}{ds} = \frac{\alpha}{\sqrt{p_z}} \cos \theta - \frac{\beta}{\sqrt{p_z}} \sin \theta \quad (3)$$

and moreover the abnormal direction projects onto the *saddle point* $\theta = -\pi, \frac{d\theta}{ds} = 0$. Near this saddle point we shall now compute a *first return time* t_f for the section Σ , in function of $\theta(t_f)$ and of the parameter p_z . Then, imposing $t_f = 1$ shall give a relation between p_z and $\theta(1)$, and we shall prove that this relation is not subanalytic.

Remark 2.1. – In the pendulum representation the branch $\mathcal{C}$ corresponds to a *local* computation near a saddle point, whereas the branch $S(0, r) \cap (y = 0) \cap (z \geq 0)$ corresponds to a *global* computation of return mapping near the separatrices of the pendulum. The interest of this branch $\mathcal{C}$ is all in the fact that computations are *localized near a saddle*, hence we shall use *normal forms* according to the following Lemma.

LEMMA 2.1. – Let $(X_\varepsilon)_{0 \leq \varepsilon < 1}$ be a one-parameter analytic family of vector fields in $\mathbb{R}^2$ near 0 such that for any ε : $X_\varepsilon(0) = 0$, and let (u, v) denote standard coordinates in $\mathbb{R}^2$. Let $\mu(\varepsilon)$ and $\nu(\varepsilon)$ be the eigenvalues of $dX_\varepsilon(0)$. We suppose that $\mu(0) = -1$ and $\nu(0) = 1$. Then there exists an analytic germ at 0 from $\mathbb{R}^3$ to $\mathbb{R}^2$:

$$\varphi : (u, v, \varepsilon) \mapsto (u_1, v_1) = \varphi(u, v, \varepsilon) \quad \text{with } \varphi(0, 0, \varepsilon) = 0$$

such that in the new coordinates (u_1, v_1) the system $(\dot{u}, \dot{v}) = X_\varepsilon(u, v)$ can be written near 0 as :

$$\dot{u}_1 = \mu(\varepsilon)u_1(1 + o(\varepsilon)), \quad \dot{v}_1 = \nu(\varepsilon)v_1(1 + o(\varepsilon)) \quad \text{as } \varepsilon \rightarrow 0$$

This Lemma may be proved by adding parameters in the proof of [7], see for instance [8]. To fit in this situation, set $u = \theta + \pi$, $v = \frac{d\theta}{ds}$, and $\varepsilon = \frac{1}{\sqrt{p_z}}$. Then the pendulum equation (2) can be written as the system in $\mathbb{R}^2$:

$$\frac{du}{ds} = v, \quad \frac{dv}{ds} = \sin u - \alpha^2 \varepsilon^2 \sin u \cos u + \alpha \beta \varepsilon^2 \sin^2 u + \beta \varepsilon v \cos u$$

Applying Lemma 2.1, we get in new coordinates (u_1, v_1) :

$$\frac{du_1}{ds} = \mu(\varepsilon)u_1(1 + o(\varepsilon)), \quad \frac{dv_1}{ds} = \nu(\varepsilon)v_1(1 + o(\varepsilon))$$

where $\mu(\varepsilon) = 1 + \frac{\beta\varepsilon}{2} + O(\varepsilon^2)$, $\nu(\varepsilon) = -1 + \frac{\beta\varepsilon}{2} + O(\varepsilon^2)$, and actually : $u = u_1 + v_1 + o(\varepsilon)$, $v = u_1 - v_1 + o(\varepsilon)$. Therefore in the new coordinates the section (3) is $\Sigma : v_1 = u_1 + \alpha\varepsilon + o(\varepsilon)$.

Now the return time s_f can be computed in a standard way :

$$s_f = \int_0^{s_f} ds = \int_{v_1(0)}^{v_1(s_f)} \frac{dv_1}{\nu(\varepsilon)v_1(1 + o(\varepsilon))} = \frac{1}{\nu(\varepsilon)}(1 + o(\varepsilon)) \ln \frac{v_1(s_f)}{v_1(0)} \quad (4)$$

On the one part : $v_1(0) = \alpha\varepsilon + o(\varepsilon)$; on the other part we get by integrating the system up to $O(\varepsilon^{\frac{3}{2}})$: $\frac{dt}{ds} = \frac{\varepsilon}{r}(1 + \alpha y)(1 + \beta x + \gamma y) = \frac{\varepsilon}{r}e^{-\beta\varepsilon s} + O(\varepsilon^{\frac{3}{2}})$. Now claiming that the final t_f is equal to 1, we get :

$$1 = \int_0^{s_f} \frac{dt}{ds} ds = \frac{1 - e^{-\beta\varepsilon s_f}}{\beta r} + O(\varepsilon^2)$$

And thus : $s_f = -\frac{\ln(1-\beta r)}{\beta\varepsilon} + O(\varepsilon)$. Putting into (4) we obtain finally :

$$v_1(s_f) = \frac{\alpha\varepsilon}{\sqrt{1-\beta r}} e^{\frac{\ln(1-\beta r)}{\beta\varepsilon}} (1 + o(1)) \quad \text{as } \varepsilon \rightarrow 0$$

In particular $v_1(s_f)$ is not an analytic function in ε as $\varepsilon \rightarrow 0$.

From Lemma 2.1 we know that $v_1(s_f) = An(u(s_f), v(s_f), \varepsilon) \sim \frac{u(s_f) - v(s_f)}{2}$, where $An(\cdot)$ denotes an analytic germ at 0. Moreover, on the section Σ , we have : $v(s_f) = -\alpha\varepsilon \cos u(s_f) + \beta\varepsilon \sin u(s_f)$. Hence : $v_1(s_f) = An(u(s_f), \varepsilon) = \frac{u(s_f)}{2} + \dots$. From the Implicit Function Theorem in the analytic class we get : $u(s_f) = An(v_1(s_f), \varepsilon)$. Therefore $u(s_f)$ is not an analytic function in ε , which ends the proof.

References

- [1] Agrachev A., Compactness for sub-Riemannian length minimizers and subanalyticity, Rend. Semin. Mat. Torino, Vol. 56, 1998.
- [2] Agrachev A., Bonnard B., Chyba M., Kupka I., Sub-Riemannian sphere in the Martinet flat case, ESAIM/COCV, Vol. 2, 377-448, 1997.
- [3] Agrachev A., Gauthier J.P., On subanalyticity of Carnot-Carathéodory distances, à paraître aux Annales de l'IHP.
- [4] Agrachev A., Sarychev A., Strong minimality of abnormal geodesics for 2-distributions, J. of Dynamical and Control Systems, Vol. 1, No. 2, 1995, pp 139-176.
- [5] Bellaïche A., Tangent space in sub-Riemannian geometry, Sub-Riemannian geometry, Birkhäuser, 1996.
- [6] Bonnard B., Trélat E., On the role of abnormal minimizers in SR-geometry, Preprint Dijon, 1998.
- [7] Hadamard J., Sur l'itération et les solutions asymptotiques des équations différentielles, Bulletin de la SMF, tome 29, 1901 - Oeuvres complètes, tome 3, 1023-1026.
- [8] Moussu R., Sur la dynamique des gradients, Existence de variétés invariantes, Math. Ann. 307, 1997, 445-460.
- [9] Pontryagin L. et al., Théorie mathématique des processus optimaux, Eds Mir, Moscou, 1974.
- [10] Roussarie R., On the number of limit cycles which appear by perturbation of separatrix loop of planar vector fields, Bol. Soc. Br. Mat., Vol. 17, No. 2, 1986.
- [11] Trélat E., Some properties of the value function and its level sets for affine control systems with quadratic cost, J. of Dynamical and Control Systems, Vol. 6, No. 4 (Oct. 2000), pp 511-541.