Mathematical and Numerical Analysis of a Class of Non-linear Elliptic Equations in the Two Dimensional Case

Noureddine Alaa, Jean Rodolphe Roche, A. Cheggour

► To cite this version:

Noureddine Alaa, Jean Rodolphe Roche, A. Cheggour. Mathematical and Numerical Analysis of a Class of Non-linear Elliptic Equations in the Two Dimensional Case. Jul 2006, pp.926-936. hal-00095264

HAL Id: hal-00095264

<https://hal.science/hal-00095264>

Submitted on 15 Sep 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Mathematical and Numerical Analysis of a Class of Non-linear Elliptic Equations in the Two Dimensional Case.

Nour Eddine Alaa¹, Abderrahim Cheggour¹ and Jean R. Roche²

¹ Département de Mathématiques et Informatique, Université des Sciences et Techniques Cadi Ayyad, B.P. 618, Guéliz, Marrakech, Maroc
`alaa@fstg-marrakech.ac.ma`

² I.E.C.N., Université Henri Poincaré, B.P. 239, 54506 Vandoeuvre lès Nancy, France `roche@iecn.u-nancy.fr`

Summary. The aim of this paper is to show the existence and present a numerical analysis of weak solutions for a quasi-linear elliptic problem with Dirichlet boundary conditions in a domain Ω and data belonging to $L^1(\Omega)$. A numerical algorithm to compute a numerical approximation of the weak solution is described and analyzed. Numerical examples are presented and commented.

1 Introduction

The principal objective of this work is to give a result of existence and present a numerical analysis of weak solutions for the following quasi-linear elliptic problem:

$$\begin{cases} -\Delta u(x) + G(x, \nabla u(x)) = F(x, u(x)) + f(x) & \text{in } \Omega, \\ u(x) = 0 & \text{on } \partial\Omega \end{cases} \quad (1)$$

where G, F are Caratheodory non negative functions. The function $f \in L^1(\Omega)$ is given finite non negative. The domain $\Omega \subset \mathbb{R}^N$ is open and bounded. Such problems arise from biological, chemical and physical systems.

The two essential ingredients to the analysis of this problem are the convexity of $s \rightarrow G(x, s)$ and that $G(x, s)$ is sub-quadratic w.r.t. s namely:

$$G(x, s) \leq C(k(x) + \|s\|^2), \quad \text{where } k(x) \in L^1(\Omega) \text{ and } C > 0. \quad (2)$$

Then the problem (1) has a solution in $W_0^{1,q}(\Omega)$ where $1 \leq q < N/(N-1)$, $N \geq 2$, provided that (1) has a super-solution in $W_0^{1,1}(\Omega)$.

In previous work [4] the authors show the existence of a weak solution in the one-dimensional case and with arbitrary growth of the non linearity and data measure.

We study a numerical method to compute the solution of the problem (1). In the first step we compute a super solution using a domain decomposition method. In the second step we compute a sequence of solutions of an intermediate problem obtained by using the Yosida approximation of G . This sequence converges to the weak solution of the problem (1).

2 Statement of the Main Result

Throughout this paper we suppose

$$f \in L^1(\Omega), \quad f \geq 0. \quad (3)$$

The functions $G : \Omega \times \mathbb{R}^N \rightarrow [0, +\infty[$ and $F : \Omega \times \mathbb{R} \rightarrow [0, +\infty[$ are such that:

$$G, F \text{ are measurable, } r \rightarrow G(x, r) \text{ and } u \rightarrow F(x, u) \text{ are continuous.} \quad (4)$$

$$G \text{ is convex in } r \text{ and } F \text{ is nondecreasing in } u, \quad (5)$$

$$G(x, 0) = \min \{G(x, r), r \in \mathbb{R}^N\} = 0, \text{ and } F(x, 0) = 0, \quad (6)$$

$$G(x, r) \leq C(|r|^2 + K(x)), \quad (7)$$

$$F(x, u) \in L^1(\Omega) \text{ for every } u \in \mathbb{R} \quad (8)$$

with a constant $C > 0$ and $K \in L^1(\Omega)$.

We introduce now the notion of weak solutions of problem (1).

Definition 1. A function u is said to be a weak solution of the problem (1), if

$$\begin{cases} u \in W_0^{1,1}(\Omega), \quad G(x, \nabla u) \text{ and } F(x, u) \in L^1(\Omega), \\ -\Delta u + G(x, \nabla u) = F(x, u) + f \text{ in } \mathcal{D}'(\Omega). \end{cases} \quad (9)$$

We will be interested in proving the existence of weak positive solutions of problem (1).

Theorem 1. Under hypotheses (3)–(7), and assuming that there exists w such that

$$\begin{cases} w \in W_0^{1,1}(\Omega), \quad F(x, w) \in L^1(\Omega), \\ -\Delta w = F(x, w) + f \text{ in } \mathcal{D}'(\Omega), \end{cases} \quad (10)$$

the problem (1) has a positive weak solution.

3 Proof of Theorem 1

3.1 Approximation Scheme

We consider the sequence defined by $u_0 = w$ and for $n \geq 0$, u_{n+1} is the solution of the problem

$$\begin{cases} -\Delta u_{n+1} + G_{n+1}(x, \nabla u_{n+1}) = F(x, u_n) + f & \text{in } \mathcal{D}'(\Omega), \\ u_{n+1} \in W_0^{1,1}(\Omega), G_{n+1}(x, \nabla u_{n+1}) \in L^1(\Omega), \end{cases} \quad (11)$$

where $G_n(x, r)$ denotes the Yosida approximation of $G(x, r)$. The function $G_n(x, r)$ is convex in r , increases pointwise to $G(x, r)$ as n tends to ∞ and satisfies

$$G_n \leq G_{n+1} \leq G, \quad \|G_{n,r}(x, r)\|_\infty \leq n, \quad (12)$$

where $G_{n,r}$ denotes a section of subdifferential of G_n with respect to r .

The classical works ([3], [6], [9]) combined with an induction argument can be applied to prove that (11) has a solution such that

$$0 \leq u_{n+1} \leq u_n \leq w. \quad (13)$$

3.2 Estimates and Convergence

Let $\{u_n\}_n$ be a sequence defined as above. By integrating (11) in Ω and using (13) we obtain

$$\int_\Omega G_{n+1}(x, \nabla u_{n+1}) dx \leq \int_\Omega F(x, w) dx + \int_\Omega f(x) dx. \quad (14)$$

Therefore $\|\Delta u_{n+1}\|_{L^1(\Omega)}$ is bounded. Then there exists a subsequence still denoted by u_n for simplicity, such that u_n converges strongly to some u in $W_0^{1,q}(\Omega)$, $1 \leq q < N/(N-1)$, and $(u_n, \nabla u_n)$ converges to $(u, \nabla u)$ almost everywhere in Ω (see [8]).

Let us prove that u is in fact a solution of problem (1). According to the definition 2.1, we only have to show that

$$-\Delta u + G(x, \nabla u) = F(x, u) + f \quad \text{in } \mathcal{D}'(\Omega). \quad (15)$$

We know that $F(x, u_n) \rightarrow F(x, u)$ strongly in $L^1(\Omega)$ and, for almost every x in Ω , there holds $G_{n+1}(x, \nabla u_{n+1}(x)) \rightarrow G(x, \nabla u(x))$.

Then there exists a non-negative measure μ (see [7]) such that

$$-\Delta u_n + G_{n+1}(\nabla u_{n+1}) - F(u_n) - f \rightarrow -\Delta u + G(\nabla u) - F(u) - f + \mu \quad \text{in } \mathcal{D}'(\Omega),$$

as n goes to ∞ .

On the other hand

$$-\Delta u_{n+1} + G_{n+1}(x, \nabla u_{n+1}) = F(x, u_n) + f \rightarrow F(x, u) + f \quad \text{in } L^1(\Omega). \quad (16)$$

Consequently

$$-\Delta u + G(x, \nabla u) \leq F(x, u) + f \text{ in } \mathcal{D}'(\Omega). \quad (17)$$

Therefore to conclude the proof of theorem 1, we must establish the opposite inequality. To this end we introduce the following test function

$$\psi \exp(-Cu_{n+1}) H\left(\frac{u_{n+1}}{k}\right), \quad (18)$$

where $H \in C^1(\mathbb{R})$, $0 \leq H(s) \leq 1$, $H(s) = 0$ if $|s| \geq 1$ and $H(s) = 1$ if $|s| \leq \frac{1}{2}$, C is given by relation (7) and $\psi \leq 0$, $\psi \in H_0^1(\Omega) \cap L^\infty(\Omega)$. We multiply the equation satisfied by u_{n+1} in (11) by this test function and we integrate in Ω , to obtain

$$\int_{\Omega} (f_n + F(x, u_n)) \psi \exp(-Cu_{n+1}) H\left(\frac{u_{n+1}}{k}\right) dx = I_1 + I_2 + I_3,$$

where

$$\begin{aligned} I_1 &= \int_{\Omega} \nabla u_{n+1} \nabla \psi \exp(-Cu_{n+1}) H\left(\frac{u_{n+1}}{k}\right) dx, \\ I_2 &= \frac{1}{k} \int_{\Omega} |\nabla u_{n+1}|^2 \psi \exp(-Cu_{n+1}) H'\left(\frac{u_{n+1}}{k}\right) dx, \\ I_3 &= \int_{\Omega} (G_{n+1}(x, \nabla u_{n+1}) - C |\nabla u_{n+1}|^2) \psi \exp(-Cu_{n+1}) H\left(\frac{u_{n+1}}{k}\right) dx. \end{aligned} \quad (19)$$

By investigating separately each term, we get

$$\begin{aligned} \lim_{n \rightarrow \infty} I_1 &= \int_{\Omega} \nabla u \nabla \psi \exp(-Cu) H\left(\frac{u}{k}\right) dx \\ \text{and } \lim_{k \rightarrow \infty} I_2 &= 0 \text{ uniformly on } n. \end{aligned}$$

Now we investigate the remaining term I_3 . Since G_{n+1} satisfies the inequality (7), $\psi \leq 0$, and by applying Fatou's lemma, we obtain

$$\lim_{n \rightarrow \infty} I_3 \geq \int_{\Omega} (G(x, \nabla u) - C |\nabla u|^2) \psi \exp(-Cu) H\left(\frac{u}{k}\right) dx. \quad (20)$$

Finally we have shown

$$\begin{aligned} &\int_{\Omega} \nabla u \nabla \psi \exp(-Cu) H\left(\frac{u}{k}\right) dx + \int_{\Omega} \psi (G(x, \nabla u) - C |\nabla u|^2) \exp(-Cu) H\left(\frac{u}{k}\right) dx \\ &+ \omega\left(\frac{1}{k}\right) \leq \int_{\Omega} (F(x, u) + f) \psi \exp(-Cu) H\left(\frac{u}{k}\right) dx, \end{aligned}$$

where $\omega(\varepsilon)$ denotes a quantity that tends to 0 when ε tends to 0. Now we choose $\psi = -\varphi \exp(Cu) H\left(\frac{u}{k}\right)$, where $\varphi \geq 0$, $\varphi \in \mathcal{D}(\Omega)$ and we replace ψ by

this value in the previous inequality to get after appropriate calculations and using that the third term is equivalent to $\omega(\frac{1}{k})$

$$\begin{aligned} & - \int_{\Omega} \nabla u \nabla \varphi H\left(\frac{u}{k}\right)^2 dx - \int_{\Omega} \varphi G(x, \nabla u) H\left(\frac{u}{k}\right)^2 dx + \omega\left(\frac{1}{k}\right) \\ & \leq - \int_{\Omega} (F(x, u) + f) \varphi H\left(\frac{u}{k}\right)^2 dx. \end{aligned}$$

We finally pass to the limit as k tends to infinity and we use the fact that $\lim_{k \rightarrow \infty} H\left(\frac{u}{k}\right) = 1$, to conclude for every $\varphi \geq 0, \varphi \in \mathcal{D}(\Omega)$ that

$$\int_{\Omega} \nabla u \nabla \varphi dx + \int_{\Omega} \varphi G(x, \nabla u) dx \geq \int_{\Omega} (F(x, u) + f) \varphi dx.$$

This finishes the proof of the theorem 1.

4 Numerical Method

4.1 Introduction

In this section we present the numerical method to solve the equation (1) in $\mathbb{R}^2$. Formally the algorithm can be formulated in the following way:

1) Find $\bar{w} \in H_0^1(\Omega)$ such that:

$$-\Delta \bar{w}(x) \geq F(x, \bar{w}) + f \text{ in } \Omega. \quad (21)$$

2) Given $u_0 = \bar{w}$ we compute a sequence, $\{u_n\}_n$, solution in $H_0^1(\Omega)$ of the non linear equation:

$$-\Delta u_{n+1}(x) + G_{n+1}(x, \nabla u_{n+1}) = F(x, u_n) + f \text{ in } \Omega. \quad (22)$$

Both problems (21) and (22) are non-linear, and if (21) has a solution, in theorem 1 we have shown that (22) then also has a solution.

4.2 Numerical Algorithm

This subsection summarizes the algorithm introduced in the previous subsection.

1) First step: given $\bar{w}^0 = 0$, iteratively for $k = 1$ until convergence we compute $\bar{w}^{k+1} = \bar{w}^k + \delta$ where at each iteration δ is the solution of the linear problem:

$$\begin{cases} -\Delta \delta(x) - \frac{\partial F(x, \bar{w}^k)}{\partial r} \delta(x) = \Delta \bar{w}^k(x) + F(x, \bar{w}^k) + f \text{ in } \Omega, \\ \delta(x) = 0 \text{ on } \partial\Omega. \end{cases} \quad (23)$$

To solve at each iteration the linear problem (23) we consider the domain decomposition method which will be introduced as follows:

- a) We compute $c_\infty = \left\| \frac{\partial F(\overline{w}^k)}{\partial r} \right\|_\infty$. Determine an overlapping subdomain decomposition Ω_i , $i = 1, \dots, m$ such that $\Omega = \cup_{i=1}^m \Omega_i$ and satisfies:

$$\max\{\text{mes}(\Omega_i), i = 1, \dots, m\} < \min\left(\frac{c_0 \pi^2}{c_\infty}, \frac{\pi}{2\sqrt{c_\infty}}\right). \quad (24)$$

We denote by m the number of subdomains Ω_i and $\partial\Omega_i$ is the boundary of Ω_i .

- b) Iteratively:
for $l = 1, \dots$ until convergence and for $i = 1, \dots, m$ we solve the following subdomain problems:

$$\begin{cases} -\Delta \delta_i^l(x) - \frac{\partial F(x, \overline{w}^k)}{\partial r} \delta_i^l(x) = \Delta \overline{w}^k(x) + F(x, \overline{w}^k) + f & \text{in } \Omega_i, \\ \delta_i^l(x) = \delta_j^{l-1}(x), & \text{on } \partial\Omega_i \cap \Omega_j, \\ \delta_i^l(x) = 0, & \text{on } \partial\Omega \cap \partial\Omega_i. \end{cases} \quad (25)$$

On each subdomain Ω_i we consider a finite element approximation method with N_i elements. At the end of the l -th loop we have computed an approximate discrete solution of the linear indefinite problem (23).

- 2) At this step for $u_0 = \overline{w}$, iteratively for $n = 1$, until convergence we solve the following non-linear problem

$$\begin{cases} -\Delta u_n(x) + G_n(x, \nabla u_n) = F(x, u_{n-1}) + f & \text{in } \Omega, \\ u_n(x) = 0 & \text{on } \partial\Omega. \end{cases} \quad (26)$$

At each n -th step the problem (26) is solved by using a Newton method. The discrete approximation of the solution of (1) is obtained at the end of the n -th loop.

4.3 Convergence of the Domain Decomposition Method

To simplify, without loss of generality, we assume that we can consider a two-domain decomposition $\Omega = \Omega_1 \cup \Omega_2$ such that:

$$\max\{\text{mes}(\Omega_i), i = 1, 2\} < \min\left(\frac{c_0 \pi^2}{c_\infty}, \frac{\pi}{2\sqrt{c_\infty}}\right). \quad (27)$$

Now to prove the convergence of the Schwarz overlapping domain decomposition algorithm applied to problem (23), we consider two problems:

$$\begin{cases} -\Delta v_1(x) + c(x) v_1(x) = h(x) & \text{in } \Omega_1, \\ v_1(x) = 0 & \text{on } \partial\Omega \cap \partial\Omega_1; v_1(x) = v_2(x) & \text{on } \partial\Omega_1 \cap \Omega_2 \end{cases} \quad (28)$$

and

$$\begin{cases} -\Delta v_2(x) + c(x) v_2(x) = h(x) & \text{in } \Omega_2, \\ v_2(x) = v_1(x) & \text{on } \partial\Omega_2 \cap \Omega_1; v_2(x) = 0 & \text{on } \partial\Omega \cap \Omega_2. \end{cases} \quad (29)$$

Let v be

$$v = \begin{cases} v_1 & \text{in } \Omega_1, \\ v_2 & \text{in } \Omega_2, \end{cases} \quad (30)$$

$v_1 = v_2$ in $\Omega_1 \cap \Omega_2$.

With the restriction (27) we can suppose the existence of a solution of (28) in $W_0^{1,q}(\Omega_1)$ and a solution of (29) in $W_0^{1,q}(\Omega_2)$.

Then, if v^0 is an initialization function defined in Ω and vanishing in $\partial\Omega$, we define for $k \geq 0$ two sequences v_i^k , $i = 1, 2$ solving the following problems:

$$\begin{cases} -\Delta v_1^{k+1}(x) + c(x)v_1^{k+1}(x) = h(x) & \text{in } \Omega_1, \\ v_1^{k+1}(x) = 0 & \text{on } \partial\Omega \cap \partial\Omega_1; v_1^{k+1}(x) = v_2^k(x) & \text{on } \partial\Omega_1 \cap \Omega_2 \end{cases} \quad (31)$$

and

$$\begin{cases} -\Delta v_2^{k+1}(x) + c(x)v_2^{k+1}(x) = h(x) & \text{in } \Omega_2, \\ v_2^{k+1}(x) = v_1^k(x) & \text{on } \partial\Omega_2 \cap \Omega_1; v_2^{k+1}(x) = 0 & \text{on } \partial\Omega \cap \Omega_2. \end{cases} \quad (32)$$

Theorem 2. Assume Ω_1 and Ω_2 with the restriction (27). Then the sequence v^k converges to v in $W_0^{1,q}(\Omega_1)$ and $W_0^{1,q}(\Omega_2)$.

Proof. We give here an idea of the proof.

Let $d^k = v_1^k - v$ in Ω_1 and $e^k = v_2^k - v$ in Ω_2 then $d^k \in L^\infty(\Omega_1)$ and $e^k \in L^\infty(\Omega_2)$.

Thanks to the maximum principle we prove the following inequalities:

$$\|d^{k+2}\|_\infty \leq \gamma \|d^k\|_\infty \quad \text{and} \quad \|e^{k+2}\|_\infty \leq \gamma \|e^k\|_\infty \quad (33)$$

where $\gamma < 1$.

But to be able to apply the maximum principle it will be necessary that the subdomains Ω_1 and Ω_2 verify the restriction (27).

4.4 Numerical Results

The algorithm introduced in the previous section has been implemented numerically for the model problem (1) where:

$G(x, r) = |r|^p = (r_1^2 + r_2^2)^{\frac{p}{2}}$ and $r = (r_1, r_2) \in \mathbb{R}^2$ for $1 < p < \infty$.

$F(x, s) = \eta s^q$ where $s \in \mathbb{R}^+$ and $1 < q < \infty$.

$f(x) = x_1^\alpha + x_2^\beta$ where $x = (x_1, x_2) \in \Omega$ and $-1 < \alpha, \beta < \infty$.

The number of subdomains is not fixed, it changes at each iteration according to the criterion (27). In figure 1 we can see the solution shape when the algorithm converges with $m = 36$ subdomains.

To study the convergence history of the numerical simulation plotted in figure 1 we consider two steps. In the first step, where we compute a super-solution, we observe the evolution of the number of subdomains: it goes from $m = 4$ subdomains to $m = 36$ subdomains in seven iterations according to criterion (27). Simulation stops after 34 iterations when the residual is less

Fig. 1. $\eta = 45$, $p = q = 3$, $\alpha = \beta = 2$, $m = 36$

than 10^{-6} . In the second step, starting with the super-solution computed in the previous step we perform nine iterations of the Yosida approximation described in section 3 and the simulation stops when the correction computed is in uniform norm less than 10^{-6} .

References

1. Alaa, N., Maach, F., Mounir, I.: Existence for some quasilinear elliptic systems with critical growth nonlinearity and L^1 data. *Journal of Applied Analysis*, **11**, 81–94 (2005)
2. Alaa, N., Mounir, I.: Global existence for reaction-diffusion systems with mass control and critical growth with respect to the gradient. *Journal of Mathematical Analysis and Applications*, **253**, 532–557 (2001)
3. Alaa, N., Pierre, M.: Weak solution of some quasilinear elliptic equations with measures. *SIAM J. Math. Anal.*, **24**, 23–35 (1993)
4. Alaa, N., Roche, J.R.: Theoretical and numerical analysis of a class of nonlinear elliptic equations. *Mediterr. J. Math.*, **2**, 327–344 (2005)
5. Amann, H., Crandall, M.G.: On some existence theorems for semi linear equations. *Indiana Univ. Math. J.*, **7**, 779–790 (1978)
6. Boccardo, L., Murat, F., Puel, J.P.: Existence de solutions non bornées pour certaines équations quasi-linéaires. *Portugaliae Math.*, **41**, 507–534 (1982)
7. Brezis, H.: *Analyse fonctionnelle appliquée*, Masson, Paris (1980)
8. Brezis, H., W. Strauss, W.: Semilinear elliptic equation in L^1 . *J. Math. Soc. Japan*, **25**, 565–590 (1973)
9. Lions, P.L.: Résolution de problèmes elliptiques quasilinéaires. *Arch. Ration. Mech. Analysis*, **74**, 335–353 (1980)
10. Porretta, A.: Existence for elliptic equations in L^1 having lower order terms with natural growth. *Portugaliae Math.*, **57**, 179–190 (2000)